

REFERENCES

Adbikary A.K., Numaga J., and Kaburaki T. (2001):

Rapid detection and typing of oculopathogenic strain of subgenus D adenoviruses by fiber-based PCR and restriction enzyme analysis. *Invest. Ophthalmol. Vis. Sci.*, 42(9): 2010-2015.

Allard A., Albinson B., and Wadell G. (2001):

Rapid typing of human Adenoviruses by a general PCR combined with restriction end nuclease analysis. *J. Clin. Microbiol.*, 39: 489-505.

Altemeier W.A. (2001):

The importance of adenoviral infections in pediatrics. *Pediatr. Ann.*, 30: 439-452.

Aoki K. and Kunai H. (1999):

The molecular epidemiology of adenovirus. *Nippon-Ganka-Zasshi*, 103(5): 335-336.

Arstila, P.P. and Halonen, E.P. (1988):

Direct antigen detection. In: *Laboratory diagnosis of infectious diseases. Principles and practice. Vol. II Viral, Rickettsial, and Chlamydial diseases.* Lennette, H.E., Halonen, P. and Murphy, A.F. (eds) Springer, Verlag New York Chap. 4, P: 60.

Barraza E.M., Ludwig S.L., Gaydos J.C., and Brundage J.F. (1999):

Reemergence of adenovirus type 4 acute respiratory disease in military trainees: Report of an outbreak during a lapse in vaccination. *J. Infect. Dis.*, 179: 153-1533.

Bateman E.D., Hayashi S. and Kuwano K. (1995):

Latent adenoviral infection in follicular bronchiectasis. Am. J. Respir. Crit. Care Med., 151: 170.

Baum S.G. (2000):

Adenoviridae. In: Mandell G.L., Bennett J.E., and Dolin R. eds. Principles and practice of infectious diseases, 5th ed. New York: John Wiley and Sons, p. 1624-1630.

Bergelson J.M., Cunningham J.A., and Kurt-Jones E. (1997):

Isolation of a common receptors for coxsackie B viruses and adenoviruses 2 and 5. Science, 275: 1320-1323.

Bobo L.D., Novak N., and Munoz B. (1997):

Severe disease in children with trachoma is associated with Chlamydia and persistent Chlamydia trachomatis infection. J. Infect. Dis., 167: 1524-1530.

Brown M., Grydusk J.D. and Fortasas E. (1997):

Structural features unique to entric adenoviruses. Arch. Virol. Suppl., 12: 301-307.

Buus D.R., Pflugfelder S.C., and Schachter J. (1998):

Lymphogranuloma venereum conjunctivitis with a marginal corneal perforation. Ophthalmology, 95: 799-802.

Chang C.H., Sheu M.M., and Chern C.L. (2001):

Epidemic keratoconjunctivitis caused by a new genotype of adenovirus type 8 (Ad 8) – a chronological review of Ad 8 in Southern Taiwan. Jpn. J. Ophthalmol. 45: 160-166.

Chomel J.J., Szymczyszyn P. and Honneger D. (1989):

An epidemic of adenovirus type I conjunctivitis. *Pediatric Inf. Dis. J.*, 8: 884.

Collier L. (2000):

Upper respiratory tract and eye infections due to adenoviruses, coronaviruses and rhinoviruses. In *Human Virology* 2 nd ed. Oxford university press, ch. 8, p: 69-71.

Dolter J., Wong J., and Janda J.M. (1998):

Association of *Neisseria cinerea* with ocular infections in pediatric patients. *J. Infect.*, 36: 49-52.

Eiichi Uchio, Koki Aoki, Waka Saioh, and Norihiko Itoh D.V.M. (1997):

Rapid diagnosis of adenoviral conjunctivitis on conjunctival swabs by 10 minute immunochromatography. P: 1299.

Eiichi Uchio, saloshi Takeuchi and Norihiko Iloh (2000):

Clinical and epidemiological features of acute follicular conjunctivitis with special reference to that caused by herpes simplex virus type 1. *Br. J. Ophthalmol.*, 84: 968-972.

Elfath M Elnifro, Rober J Cooper, Paul, E. and Andrew, S. (1999):

Diagnosis of viral and chlamydial keratoconjunctivitis which laboratory test, 83: 622- 327.

Elnifro E.M., Cooper R.J., Klapper P.E., and Bailey A.S. (1999):

Diagnosis of viral and chlamydial keratococonjunctivitis: which laboratory test? *Br. J. Ophthalmol.*, 83: 622-627.

Enders I.F. and Beu J.A. (1956):

"Adenoviruses": Group name proposed for new respiratory – Tract viruses. Science, 124: 119 -120.

Finn A., Andy E. and Talbot G.H. (1988):

An epidemic of adenovirus to infection in aneonatal nursery: course, morbidity and management. Infect. Control Hosp. Epidemiol., 9: 398.

Flomenberg P., Babbitt J. and Drobyski W.R. (1994):

Increasing incidence of adenovirus disease in bone marrow transplant recipients. J. Infections disease, 169: 775.

Folds J.D., Farrer J.A., and Norman D.E. (2000):

Pocket guide to clinical immunology, P: 132-135.

Ford E., Nelson K.E. and Warren D. (1987):

Epidemiology of epidemic E Kerato conjunctivitis. Epidemiol. Rev., 9: 244.

Friedberg T.N., Stenson S.M. and Orenstven J.M. (1999):

Microsporidian keratoconjunctivitis in acquired immunodeficiency syndrome. arch. Ophthalmol., 108: 504-508.

Gahery S.H., Farace F., and Godfrind D. (1998):

Immune response to recombinant capsid proteins of adenovirus in humans. J. Virol., 72: 2388-2397.

Ginsberg H.S. and Pereird H.G. (1991):

A proposed terminology for the adenovirus antigens and virion morphological subunits virology. 28: 782-783.

Gordon Y.J., Nacsens L., De Clerq E., and Veckeneer M. (1996):

Treatment of adenoviral conjunctivitis with topical cidofovir.
Cornea., 7: 15-17.

Gottsch J.D. (1996):

Surveillance and control of epidemic keratoconjunctivitis. Trans.
Am. Ophthalmol. Soc., 94: 539-584.

Grandien M., Pettersson C.A., Svenssonl. and Uhnöo I. (1987):

Latex agglutination test for adenovirus diagnosis in diarrheal
disease. J. Med. Virol., 23: 311-316.

Harsi C.M., Rolim D.P., and Candeias A.N. (1995):

Adenovirus genome types isolated from stool of children with
gastroenteritis in sao Paulo, Brazil. J. Med. Virol., 45: 127-134.

Hendrix R.M., Lindner J.L., and Benton F.R. (1999):

Large, persistent epidemic of adenovirus type 4-associated acute
respiratory disease in U.S Army trainees. Emerg. Infect. Dis., 5:
798- 801.

Hierholzer J.C. (1995):

Adenoviruses. P 947 In Murray PR, Baron E J, Pfaller MA et al
(eds): manual of clinical Microbiology 6th Ed. ASM press,
Washington DC.

Hierholzer J.C. Halonen P.E. and Dahlen P.O. (1993):

Detection of adenovirus in clinical specimens by polymerase chain
reaction and liquid-phase hybridization quantitated by time
resolved flurometry. J. Clin. Microbiol., 31: 1886-1891.

Hierholzer J.C., Halonen P.E. and Dahlen P.O. (1993):

Detection of adenovirus in clinical specimens by polymerase chain reaction and liquid-phase hybridization unantiated by time-resolved fluorometry. J. Clin. Microbiology., 31: 1886-18891.

Hillenkamp J., Reinhard T., and Ross R.S. (2002):

The effects of cidofovir 1% with or without cyclosporin 1% as a topical treatment of adenoviral keratoconjunctivitis: a controlled clinical pilot study. Ophthalmology, 109: 845-850.

Horwitz M.S. (2001):

Adenoviruses. In: Fields B. N., knipe P.M., and Howely P.M., eds. Fundamental Virology, 4th ed. Philadelphia, Lippincott Williams and Willkins, p. 2301-2326.

Huebner R.J. and Row W.P. (1954):

Adenoidal – pharyngeal conjunctiv agents. N. Eng. J. Med., 251: 1077-1086.

Jawetz E., Hanna L., Sonne M., and Thygeson R. (1994):

A laboratory infection of adenovirus type 8. Am. J. Hyg., 69: 13-20.

Jernigan J.A., Lowry B.S., and Hayden S.A. (1993):

Adenovirus type 8 epidemic keratoconjunctivitis in an eye clinic: risk factors and control. J. Infect. Dis., 167: 1307 – 1313.

John A, Jernigan, Barbara S. Lowry and Frederick G. Hayden (1993):

Adenovirus type 8 Epidemic keratoconjunctivitis in an eye clinic: risk factors and control. Journal of Infecious disease, 167: 1007-1013.

Kaye S.B., Morton D.E., Tong C.Y., and O'Donnell N.P. (1998):

Echovirus keratoconjunctivitis. *Am. J. Ophthalmol.*, 125: 187-190.

Kidd A.H., Chroboczek J., Cusack M., and Ruigrok R.W. (1993):

Adenovirus type 40 virions contain two distinct fibers. *Virology*, 192: 73-84.

Krisher K.K. and Menegus M.A. (1987):

Evaluation of three types of cell culture for recovery of adenovirus from clinical specimens. *J. Clin. Microbiol.*, 25: 1323.

Lennett E.H. (1989):

General principles underlying laboratory diagnosis of viral and rickettsial infections, PP 1-177. In: E.H Lennette and NJ Shmidt (eds): *Diagnostic procedures for viral and rickettsial infections*. American Public Health, Association, New York.

Lew J.F., Moe C.L. and Monroe S.S. (1991):

Astrovirus and adenovirus associated with diarrhea in children in day care setting. *J. Infect. Dis.*, 164: 673.

Lietman T., Brooks D., and Moncada J. (1998):

Chronic follicular conjunctivitis associated with *Chlamydia psittaci* or *Chlamydia pneumoniae*. *Clin. Infect. Dis.*, 26: 1335-1340.

Lukashok S. and Horwitz M.S. (1997):

Adenovirus persistence In: Ahmed R. Chen I, eds. *Persistent viral infection*. Chichester., UK: John Wiley Sons, 147-164.

Madeley C.R. (1986):

The emerging role of adenoviruses as inducers of gastroenteritis. *Pediat. Infect. Dis.*, 5: 563.

Madeley C.R. and Peiris S.M. (1997):

Adenoviruses, respiratory diseases, conjunctivitis. Gut Infection, 43: 394-398.

Mann D., Moreb J., Smith S., and Gian V. (1998):

Failure of intravenous ribavirin in the treatment of invasive adenovirus infection following allogenic bone marrow transplantation: A case report. J. Infect., 36: 227-228.

Maria C.M., Fabiano M.S., Caroline S.S., and Eduardo M.V. (2003):

Adenoviruses isolated from civilian and military personnel in the city of Rio de Janeiro, Brazil. Rev. Inst. Med. Trop.; P: 1-8.

Mc Donnel J.M., Mc Connell P.K. and Green W.R. (2000):

Demonstration of papilloma virus capsid antigen in human conjunctival neoplasia. Arch. Ophthalmol., 104: 1801-1805.

Mei Y.F. and Wadell G. (1993):

Hemagglutination properties and nucleotide sequence analysis of the fiber gene of adenovirus genome types 11a and 11p. Virology, 194: 453-462.

Meur man O., Lausanne O. and Sarkkinen H. (1983):

Immunoassay diagnosis of adenovirus infections in children. J. Clin. Microbial., 18: 1190.

Mistchenko A.S., Diez K.A. and Mariani A.L. (1994):

Cytokines in adenoviral disease in children Association of interleukin. interleukin-8 and tumor neerosis factor alpha levels with clinical outcome. J. Pediatr., 124: 714.

- Mombacerts I.M., Maudgal P., and Knockaerts D.C. (1991):**
Bilateral follicular conjunctivitis as a manifestation of Lyme Disease. Am. J. Ophthalmol., 112: 96-97.
- Morris D.J., Klapper P.E., and Cleator G.M. (1995):**
Viral cross infection in ophthalmology. Postgrad. Doctor Caribbean, 11: 152-160.
- Nandy A., Addy M., and Chowdhury A.B. (1995):**
Leishmainial Blepharoconjunctivitis. Trop. Geogr. Med., 43: 303-306.
- Norrby E., Bartha A. and Boulanger P. (1976):**
Adenoviridea. Intervirology; 7: 117-125.
- O'Brien T.P. (2000):**
Conjunctivitis. In: Mandell G., Benett J.E., and Dolin R.principles and Practice of infectious Diseases. 5th ed., p. 1251-1256.
- Pavan- Langston D. (1997):**
Major ocular viral infections: Herpes simplex, Adenovirus, Epstein-Barr virus, Pox. In: Galasso G., Merigan T., whitely R. (eds.): Antiviral Agents and viral Dieases of Man. 4th ed. Philadelphia, Lippincott-Raven, P. 187-228.
- Peter K., Wishart and Charles James (1982):**
Prevalence of acute conjunctivitis caused by Chlamydia, adenovirus, and herpes simplex virus in an ophthalmic casualty department.

Slack J.W., Hyndiuk R.A., and Harris G.T. (1995):

Blastomycosi of the eyelid and conjunctiva. Ophthal Plast.
Reconstr. Surg., 8: 143-149.

Stewart P.L., Burnett R.M., Cyrklaff M. and Fuller S.D. (1991):

Image reconstruction reveals the complex molecular organization
of adenovirus cell, 67: 145.

Toogood C.I., Crompton J. and Hay R.T. (1992):

Antipeptide antisera define neutralizing epitopes on the adenovirus
hexon. J. Gen. Virol., 73: 1429-1435.

Ursula Weitgasser, Eva Maria Haller and Yosuf El-Sharbrawi (2002):

Evaluation of polymerase chain reaction for the detection of
adenoviruses in conjunctival swabs specimens using degenerate
primers in comparison with direct immunofluorescence.
Ophthalmologica, 217: 329-332.

***Van Ogtrop M.L., Van Zoeren – Groggen D., and van Boven C.P.
(1997):***

Serratia marcescens infections in neonatal departments:
Description of an outbreak and review of literature. J. Hosp.
Infect., 36: 95-103.

Van Oostrum J. and Burnett R.M. (1985):

Molecular composition of the adenovirus type 2 virion. J. Viral,
56: 439-448.

Ward J.B., Siojo L.G. and Waller S.G. (1993):

Clinical trial of trifluridine, dexamethasone and artificial tears in
the treatment of epidemic keratoconjunctivitis. Cornea, 12: 216.