

REFERENCES

REFERENCES

- Abd El - Aziz, A.A., Shaarawy, M., El Mallah, S.Y. and Radwan, H.M. (1993):**
 Clinical value of plasma fibronectin, protein C and pregnancy specific B1 glycoprotein Sp1 in PIH.
 Obstet. Gynecol. XXV International conference of pathophysiology of pregnancy Cairo- Egypt November 23- 26
 Abstracts
- Abilgaard, V. (1981):**
 Antithrombin III and related inhibitors of coagulation.
 Recent advances in blood coagulation. (3:151).
 Edipollerl. (churchill livingston, london)
- Aghai, E., Brunstein, J. Quitt, M., Abramovici, H. and Froom, P. (1990):**
 Antithrombin III levels in pre-eclampsia.
 ISr. J. Med. Sci. Mar., 26 (3): 142-3.
- Alexander, S.J., Colonna, G. and Edethoch, H. (1979):**
 The structure and stability of human plasma cold-insoluble globulin.
 J. Biol. Chem., 254:1501.
- Ali, I.U. and Hynes, R.O. (1978):**
 Effects of LETS glycoprotein on cell motility.
 Cell 14:439.
- Anne, A.K., Carolyn, M.C., Gabriel, M.R., Wilfred, M., Cari, S.A. and Peter, B. (1992):**
 Rapid expression of fibronectin in the rabbit heart after myocardial infarction with and without reprofusion.
 J. Clin. Invest., 89:1060-1068.
- Anunciado, A.N., Stubbs, T.M., Pepkowitz, S.H., Lazarchick, J. and Miller, C.M. (1987):**
 Altered villus vessel fibronectin in preeclampsia.
 Am. J. obstet. Gynecol. Apr., 156 (4): 898-900.
- Aoki, N. (1979):**
 Natural inhibitors of fibrinolysis.
 Prog. Cardiovasc. Dis., 21:267-286.

Aoki, N. (1984):

Genetic abnormalities of the fibrinolytic system.
Semin. Thromb. Hemost., 10:42.

Aoki, N., Moroi, M. and Tachiya, K. (1978):

Effects of α_2 -plasmin inhibitor on fibrin clot lysis.
Its comparison with α_2 - macroglobulin.
Thromb. Haemostas., 39:22.

Aoki, N., Sakato, Y. Ichinose, A. (1983):

Fibrin-associated plasminogen activation in α_2 plasmin inhibitor.
Blood, 62:1118.

Aoki, N., Moroi, M., Matsuda, M. and Tachiya K. (1977):

The behavior of α_2 - plasmin inhibitor in fibrinolytic states.
J. Clin. Invest., 60:361.

Aoki, N., Sakata, Y., Matsuda, M. and Tateno, K. (1980):

Fibrinolytic states in a patient with congenital Deficiency of α_2 - plasmin inhibitor.
Blood, 55 (3): 483.

Atha, D.H., Stephens, A.W. and Rosenberg, R.D. (1984):

Evaluation of critical groups required for binding heparin to antithrombin.
Proc. Natl. Acad. U.S.A., 81:1030.

Azar, R. and Turpin, D. (1989):

Plasma fibronectin a marker of arterial hypertension in pregnancy.
J. Gynecol. obstet. Biol. Repord., 18 (7): 863.

Balian, G., Click, E.M., Crouch, E., Davidson, J.M. and Bornstein, P. (1979):

Isolation of a collagen-binding fragment from fibronectin and cold-insoluble globulin.
J. Biol. Chem., 254:1429.

Ballegeer, V., Spitz, B., Kieckens, L., Moreau, H; Van-Assche, A. and Collen, D. (1989):

Predictive value of increased plasma levels of predictive value of increased plasma levels of fibronectin in gestational hypertension.

Am. J. Obstet. Gynecol, Aug, 16 (2): 432-6.

Ballegeer, V.C., Spitz, B, De. Baene, L.A., Van-Assche, A.F., Hidajat, M. and Criel, A.M. (1992):

Platelet activation and vascular damage in gestational hypertension.

Am. J. Obstet. Gynecol. Feb., 166 (2): 629-33.

Barham, D. and Trinder, P. (1972):

Enzymatic determination of uric acid.

Analyst., 97:142-145.

Bell, W.R. (1977):

Haematologic abnormalities in pregnancy.

Med. Clin. North. Am., 61:179.

Beller, F.K. and Uszynki, M. (1974):

Disseminated intravascular coagulation in pregnancy.

Clin. Obstet. Gynecol., 17:250.

Berardi, J.C., Rouchon, M.H., Laloux, V., Akli, J. and Berardi, G.L. (1984):

A study of the mean concentration of plasma antithrombin III in arterial hypertension in pregnancy.

Rev. Fr. Gynecol. obstet. Nov., 79 (11): 709-12.

Bhatia, R.K., Saleh, A.A., Bottoms, S.F. and Mammen, E.F. (1989):

Elevated fibronectin levels and preeclampsia.

Am. J. Obstet. Gynecol., 160 (4):1022.

Bjork, I., olson, T. and shore, J.D. (1989):

Molecular mechanisms of the accelerating effect of heparin on the reactions between antithrombin and clotting proteinases. In lane D.A. and lindahi U (eds) Heparin: Chemical and Biological properties, Clinical Applications (229). London Edward Arnold.

- Bjork, I., Danielsson, A., Fenton, J.W. and Jonvall, H. (1981):**
The site in human antithrombin for functional proteolytic cleavage by human thrombin.
Fed. Eur. Biochem. soci. Let., 126:257.
- Bjork, I., Jackson, C.M., Jonvall, H., Lavine, K.K., Nording, K. and Salsgiver, W.J. (1982):**
The active site of antithrombin III. Release of the same proteolytically cleaved form of the inhibitor from complexes with Factor Xa and thrombin.
J. Biol. Chem., 257:2406.
- Blumberg, P.M. and Robbins, P.W. (1975):**
Effect of proteases on activation of resting chick embryo fibroblasts and on cell surface proteins.
Cell, 6: 137.
- Blumenstock, F.A., Saba, T.M., Weber, P. and Laffin, R. (1978):**
Biochemical and immunological characterization of human opsonic α_2 -S.B. glycoprotein. Its identity with cold-insoluble globulin.
J. Biol. Chem., 253:4287.
- Blystone, S.D. and Kaplan, J.E. (1990):**
Fibronectin amino-terminal domains enhance macrophage fibrin binding. Evidence for an amino-terminal fibronectin receptor.
Circ. Shock, 31: A 26.
- Blystone, S.D., Weston, L.K. and Kaplan, J.E. (1991):**
Fibronectin dependent macrophage fibrin binding.
Blood, 78: No. (11):2900
- Bock, S.C., Marrinan, J.A. and Radziejewska, E. (1988):**
Antithrombin III Utah: proline 407 to leucine mutation in a highly conserved region near the inhibitor reactive site.
Biochem., 27:6171.
- Bock, S.C., Harris, J.F., Balazs, I. and Trent, J.M. (1985):**
Assignment of the human antithrombin III structural gene to chromosome 1q 23-25.
Cytogenetics cell Genetics. 39:67.

- Boer, K., Cate, J.W., Sturk, A., Borm, J.J. and Treffers, P.E. (1989):**
Enhanced thrombin generation in normal and hypertensive pregnancy.
Am. J. Obstet. Gynecol. Jan., 160 (1): 95-100.
- Bonnar, J. (1987):**
Hypertension in pregnancy.
Recent advances in obstet. and Gynecol, 15:3-13.
- Bonnar, J., McNicol, G.P. and Douglas, A.S. (1971):**
Coagulation and fibrinolytic systems in pre-eclampsia and eclampsia.
Br. Med. J., 11:12.
- Borg, J.Y., Owen, M.C., Soria, C., Caen, J., and Carrel, R.W. (1988):**
Proposed heparin binding site in antithrombin based on Arginine 47.
J. Clin. Invest., 81:1292-1296.
- Boss, J.H. (1963):**
On the pathogenic power of placental extract. A toxemia of pregnancy.
Obstet. Gynecol., 59:478.
- Brady, K., Duff, P. and Yancey, M.K. (1990):**
Plasma fibronectin concentrations during normal term labour.
Obstet. Gynecol., 75:619-21.
- Brennan, S.O., George, P.M. and Jordan, R.E. (1987):**
Physiological variant of antithrombin III lacks carbohydrate side chain at Asn 135.
Fed. Eur. Biochem. Soci.Let., 219:431-436.
- Brod. J. and Sirota, J.H. (1948):**
Detemination of creatinine in blood.
J. Clin. Invest., 27:645.
- Brossens, I.A. (1977):**
Morphological changes in the uteroplacental bed in pregnancy hypertension.
Clin. obstet. Gynecol., 4:573.

Brubaker, D.B., Ross, M.G. and Marinoff, D. (1992):

The function of elevated plasma fibronectin in preeclampsia.
Am. J. Obstet. Gynecol., 166:526-31.

Buller, H.R., Weenink, A. H., Treffers., P.E., Kahle, L.H., Otten, H.A. and Tencate, J.W. (1980):

Severe antithrombin III deficiency in a patient with pre-eclampsia. Observations on the effect of human AT III concentrate transfusion.
Scand. J. Haematol., 25:81.

Busch, C. and Owen, W.G. (1982):

Identification in vitro of an endothelial cell surface cofactor for antithrombin III.
J. Clin. Invest. 69:726.

Calvin, S., Weinstein, L., Witte, M.H. and Finely, P.R. (1990):

Plasma levels of fibronectin and prostacyclin metabolite in peripartum pre-eclamptic women.
Am. J. perinatol., 7 (2): 125.

Cararelli, P.M., Blumenstock, F.A., Mckeown-Longo, P.J., Saba, T.M., Mazurkiewicz, J.E. and Dias, J.A. (1990):

High-affinity binding of fibronectin to cultured kupffer cells.
J. Leuk. Biol., 48:426.

Carlson, T.H., Atencio, A.C. and Simon, T.I. (1984):

In vivo behavior of radioiodinated rabbit antithrombin III: demonstration of a non circulating vascular compartement.
J. Clin. Invest., 74:191.

Carlson, T.H., Atencio, A.C. and Siman, T.I. (1985):

Comparison of the behavior in vivo of two forms of antithrombin III.
Biochem. J., 225:557.

Carlson, T.H., Simon, T.I. and Atencio, A. C. (1985):

In vivo behavior of human radioiodinated Antithrombin III : distribution among three physiologic pools.
Blood, 66 (1) : 13-19.

Carlson, T.H., Babcock, T., Atencio, A.C., Levinson, C. and Mora, H.R. (1988):

Behaviour of antithrombin III isoforms on immobilized heparin. Evidence that the isoforms bind to different numbers of low affinity heparin sites.
J. Biol. Chem., 263:2187.

Caron, C., Gouemand, J., Marey, A., Beague, D., Ducroux, G. and Drouvin, F. (1991):

Are hemostatic and fibrinolytic parameters predictors of preeclampsia in pregnancy associated hypertension.
Throm. haemost. Oct., 66 (4): 410-4.

Carrell, R.W. and Travis, J. (1985):

α_1 - Antitrypsin and the serpins: variation and countervariation.
Trends Biochem. Sci., 10:20.

Carrell, R.W., Owen, M., Brennan, S. and Vaughan, L. (1979):

Carboxy terminal fragment of human α_1 - antitrypsin from hydroxylamine cleavage: homology with antithrombin III.
Biochem. Biophys. Res. Commun., 91:1032.

Carter, W. G. and hakomori, S. (1979):

Isolation of galactoprotein from hamster embryo fibroblasts and characterization of the carbohydrate unit.
Biochem., 18:730.

Castaneda, H., Garcia-Romero, H. and Clanto, M. (1970):

Severe oedema- proteinuria-hypertension-gestosis.
Am. J. obstet. Gynecol., 107:58.

Castro, L., Arora, C., Roll, K. and Parvez, S. (1988):

The presence of atrial natriuretic peptide (ANP) in extra cardiac tissues in the rat fetus and neonate. International society for the study of hypertension in pregnancy. Scientific program and abstract 6th.
International congress, Monthreal., Canada, May, 22:26-40.

Cauchi, M.N. (1984):

Coagulation disorders of pregnancy.
In the clinical pathology of pregnancy. and the newborn
infant first (ed) PP: 118 Edward Arnold.

Chandra, T., Stackhouse, R., Kidd, V. J., Robson, J.H. and woo, S.L. (1983):

Sequence homology between human α_1 - antichymotrypsin,
 α_1 - antitrypsin and antithrombin III.
Biochem., 22:5055.

Chang, J.Y. and Tran T.H. (1986):

Antithrombin III Basel, Identification of a pro to leu
substitution in a hereditary abnormal antithrombin with
impaired heparin cofactor activity.
J. Biol. Chem., 261:1174.

Chen, A.B., Amroni, D.L. and Mosesson, M.W. (1977):

Heterogenicity of fibronectin of human plasma, a circulating
cell-surface protein.
Biochem. Biophys. Acta, 493:310.

Chen, A.B., Mosesson, M.W. and Solish, G.I. (1976):

Identification of the cold-insoluble globulin of plasma in
amniotic fluid.
Am. J. Obstet. Gynecol., 125:958.

Chesley, J. (1985):

Diagnostic criteria for preeclampsia.
obstet. Gynecol., 65:423.

Chihata, Y.M. and Elyan, A. (1993):

Confirmation of chihata's theory: The disease of surface
tension of blood is the cause of preeclampsia.
Obstet. Gynecol. XXV International conference of
pathophysiology of pregnancy.
Cairo-Egypt November 23-26. Abstracts.

Choay, J., Petitou, M., Lormeau, J.C., Sinay, P., Casu, B. and Gatti, G. (1983):

Structure-activity relationship in heparin: a synthetic pentasaccharide with high affinity for antithrombin III and eliciting high affinity for antithrombin III and eliciting high anti-factor X a activity.

Biochem. Biophys. Res. Commun., 116:492.

Clark, R.A., Quinn, J.H., Winn, H.J., Lanigan, J.M., Dellepella, P. and Calvin, R.B. (1982):

Fibronectin is produced by blood vessels in response to injury.

J. Exp. Med., 156:646.

Clemmensen, I. (1979):

Different molecular forms of alpha-2- antiplasmin, in collen D, wiman B. verstratete, M, (eds):

The physiological inhibitors of coagulation and fibrinolysis.

Amsterdam, Elsevier/ North-Holland : 131

Clemmensen, I., Thorsen, S., Mullertz, S. and Peterse, L.C. (1981):

Properties of three different molecular forms of the α_2 - plasmin inhibitor.

Eur. J. Biochem., 120:105.

Collen, D. and Lijnen, H.R. (1986):

The fibrinolytic system in man.

Crit. Rev. Oncol. Hematol., 4 (3): 249-301.

Collen, D. and Wiman, B. (1978):

Fast-acting plasmin inhibitor in human plasma.

Blood, 51:563.

Collen, D. and Wiman, B. (1979):

Turnover of antiplasmin, the fast-acting plasmin inhibitor of plasma.

Blood, 53 (2) : 313.

- Colvin, R.B., Gardner, P.I., Roblin, R.O., Verderberg, E. L. and Mosesson, M.W. (1979):**
Cell surface fibrinogen/fibrin receptors on cultured human fibroblasts association with cold insoluble globulin (fibronectin) and loss in SV-40 transformed cells.
Lab. Invest., 41:464.
- Conard, J., Horellou, M.H. and Samoma, M. (1988):**
Incidence of thromboembolism associated with congenital disorders in coagulation and fibrinolysis.
Acta chirurgica scandinavica supplement, 543:15.
- Conard, J., Brosstad, F., Larsen, M.L., Samama, M. and Abildgaard, U. (1983):**
Molar antithrombin concentration in normal human plasma.
Haemostasis, 13:363-368.
- Condie, R. G. and Ogston, D. (1976):**
Sequential studies on components of the haemostatic mechanism in pregnancy with particular reference to the development of preeclampsia.
Br. J. obstet. Gynecol., 83:938.
- Cosio F.G. and Bakalitz, A. (1986):**
Binding of fibronectin to Ag. Ab complex.
Lab. Clin. Med., 103:613.
- Cosio, F.G., Eddy, A., Monster, M.I. and Bergstein, J.M. (1985):**
Decreased plasma fibronectin levels in children with hemolytic uraemic syndrome.
Am. J. Med., 78:549-554.
- Courtoy, P.J. and Boyles, J. (1983):**
Fibronectin in the microvasculature: Localization in the pericyte-endothelial interstitium.
J. Ultrastruct. Res., 83:258.
- Crouch, E., Balian, G., Holbrook, K., Duskin, D. and Bornstein, P. (1978):**
Amniotic fluid fibronectin. Characterization and synthesis by cells in culture.
J. Cell, Biol., 78:701.

Cunningham, F.G., MacDonald, P.C. and Gant, N.F. (1989):

Hypertensive disorders in pregnancy.

In Williams obstetrics 18 th (ed.) vol. II: 653-695 Appleton and lange.

Dacie, J.V. and Lewis, S.M. (1975):

Basic haematological techniques. In practical Haematology 6th ed: 41-43. (ed) churchil living stone.

Davey, D.A. (1988):

Hypertensive disorders of pregnancy.

In: studd J;ed. progress in obstertics and Gynecology volume 5. Edinburgh Living stone, London, 89-104.

De-Boer, K., Buller, H.R., Ten-Cate, J.W. and Treffers, P.E. (1991):

Coagulation studies in the syndrome of hemolysis, elevated liver enzymes and low platelets.

Br.J.obstet.Gynecol., 98 (1) : 42-7.

Dejgrad, A., Anderson, T. and Gluide, F. (1986):

The influence of insulin on the raised plasma fibronectin concentration in human obesity.

Acta. Med. Scand., 220 (3): 269.

Dessau, W., Adelman, B.C., Timpl, R. and Martin, G.R. (1978):

Identification of the sites in collagen α -chains that bind serum anti-gelatin factor (cold-insoluble globulin).

Biochem. J., 169:55.

Devraj-Kizuk, R., Chui, D.H., prochonik, E., V., Carter, C., J., Ofosu, F., A. and Blajch man M.A. (1988):

Antithrombin III- Hamilton: a gene with opoint mutation (quanine to adenine) in codon 382 causing impaired serine protease reactivity.

Blood, 72:1518.

Dian, P., Torbergesen, T., Mathesen, E. and Borud, O. (1988):

Preeclampsia. A mitochondrial disease.

International society for the study of hypertension in pregnancy. Scientific program and abstracts. 6 th.

International congress Montreal Canada May, 22:26-55.

Diluzio, N.R., Pisano, J.C. and Salky, N.K. (1977):

Macrophage recognition mechanisms, clinical and experimental consideration, in Diluzio N.R. (ed) *The Reticuloendothelial system and Immune phenomena*. New York, N.Y., plenum., 373.

Dixit, V.M., Haverstick, D.M., O'Rourke, K., Hennessy, S.W., Brockelmann, T.J., McDonald, J.A., Grant, G.A., Santoro, S.A. and Frazier, W.A. (1985):

Inhibition of platelet aggregation by a monoclonal antibody against human fibronectin.

Proc. Natl. Acad. U.S.A., 82:3844.

Dixon, H.G. (1978):

preeclampsia and eclampsia.

Brown antenatal care. Browne, J.C.M.C. and Dixon G. (eds) 11 th (edn). Elbs and churchill livingstone: 198.

Domula, M., Bykowska, K., Weszryniewicz, Z., Lopaciuk, C.S. and Kopic, M. (1985):

Plasma fibronectin concentration in healthy and septic infants. J. pediatr., 144 (1) : 49.

Duchange, N., Chasse, J.F., Cohen, G.N. and Zakin, M.M. (1986):

Identification of a point mutation leading to an arginine to cysteine replacement in a silent deficiency.

Nucleic acids Research, 14:2408.

Duckert, F. (1972):

Documentation of the plasma factor XIII deficiency in man. Ann NY Acad. Sci., 202:190.

During, R. and Schwarzlos, G. (1990):

[Antithrombin III, level in normal pregnancy and patients with pregnancy induced hypertension].

Zentralbl Gynakol., 112 (6): 351-4 Quoted from computer abstracts.

Edsall, J.T., Gilbert, G.A. and Scheraga, H.A. (1955):

The non clotting component of the human plasma fraction 11 (cold insoluble globulin).

J. Am. Chem. Soc., 77:157.

Ellis, V., Scully, M.F. and Kakkar, V.V. (1983):

The effect of divalent metal cations on the inhibition of human coagulation factor Xa by plasma proteinase inhibitors. *Biochimica Biophysica, Acta*, 747:123.

Engelmann, L., Ruckhaberle, K.E., Engelmann, B., Scheel, H., Vogtmann, C. and Deckert, F. (1993):

Gestosis. Thrombophilia and pulmonary embolism in a primipara with twin pregnancy.

Gesamte, Z., Inn. Med., 48 (4): 190-6.

Quoted from computer abstract.

Engesser, L., (1988):

Thrombophilia: disorders of blood coagulation and fibrinolysis.

Thesis university of leiden.

Engvall, E. and Ruoslahti, E. (1977):

Binding of soluble form of fibronectin surface protein, fibronectin, to collagen.

Int. J. Cancer, 20:1.

Engvall, E., Ruoslahti, E. and Miller, E.J. (1978):

Affinity of fibronectin to collagens of different genetic types and to fibrinogen.

J. Exp. Med., 147:1584.

Erdjument, H., Lane, D.A., Panico, M., Di-Marzo, V. and Morris, H.R. (1988):

Single amino acid substitutions in the reactive site of antithrombin Northwick park and to histidine in antithrombin Glasgow.

J. Bio. Chem., 265:5589-5593.

Eriksen, H.D., Hansen, P.K., Brocks, V. and Jensen, B.A. (1987):

Plasma fibronectin concentration in normal pregnancy and pre-eclampsia.

Acta Obstet. Gynecol. Scand, 66 (1): 25-8.

Fallon, H.J. (1982):

Liver disease.

Medical complication during pregnancy, 2nd Edition W.B. Saunders company. U.S.A., 278.

Ferris, T.F. and Francisco, L.L. (1982):

Toxaemia and hypertension during pregnancy.

Bull. Ny Acad. Med., 58:178.

Ffrench-Constant, C.I., Van-De-Water, H.F., Dverak, N. and Hynes, R.O. (1989):

Reappearance of an embryonic pattern of fibronectin splicing during wound healing in the adult rat.

J. Cell. Biol., 109:903-911.

Fish, W.W., Orre, K. and Bjork, I. (1979):

The production of an inactive form antithrombin through proteolysis by thrombin.

F.E.B. Soci, Letters, 98:103.

Foda, M.S., Youssef, A., Elsadek, M., Shafeek, A. and El-Minawi, M. (1979):

Hypertensive states of pregnancy.

Manual of obstet. P., 65, 3 rd. ed.

Scientific book centre. Cairo.

Foidart, J.M., Nochy, D., Mahieu, P.R., Lambotte, R. and Briety, J. (1982):

Accumulation of laminin, fibronectin, fibrin, basement membrane proteoglycan and several collagens in glomeruli of patients with preeclampsia.

8th European congress of perinatal Medicine Brussels Sept. 7-10.

Foidart, J.M., Nochy, D., Nusgens, B., Foidart, J.B., Mahieu, P.R., Lapiere, C.M., Lambotte, R. and Bariety, J. (1983):

Accumulation of several basement membrane proteins in glomeruli of patients with pre-eclampsia and other hypertensive syndromes of pregnancy possible role of renal prostaglandins and fibronectin.

Lab. Invest. Sep., 49 (3): 250-9.

Franzen, L.E., Svensson, S. and Larm, O. (1980):

Structural studies on the carbohydrate portion of human antithrombin III.

J. Biol. Chem., 255:5090-5093

Fuchen, H., Nakabayashi, M., Stock, D. and Sakamotoes, E. (1980):

Urinary fibrinolysis in toxemia of pregnancy.

Acta obstet. Gynecol. Scandina, 59:499-505.

Fukuda, M. and Hakomori, S. (1979):

Proteolytic and chemical fragmentation of galactoprotein a, a major transformation-sensitive glycoprotein released from hamster embryo fibroblasts.

J. Biol. Chem., 254:5442.

Fukuda, M. and Hakomori, S. (1979):

Carbohydrate structure of galactoprotein a, a major transformation sensitive glycoprotein released from hamster embryo fibroblasts.

J. Biol. Chem., 254:5451.

Furcht, L.T., Mosher, D.F. and Wendelschafer-crabb, G. (1978):

Effects of cell density and transformation on the formation of a fibronectin extracellular filamentous matrix on human fibroblasts.

Cancer Res., 38:4618.

Furcht, L.T., Mosher, D.F. and Wendelschafer-crabb, G. (1978):

Immunocytochemical localization of fibronectin (LETS protein) on the surface of 16 myoblasts. light and electron microscopic studies.

Cell, 13:263.

Furcht, L.T., Wendelschafer-crabb, G., Mosher, D.F., Arneson, M., Hammerschmidt, D. and Voodbridge P. (1979):

The role of fibronectin in normal platelet aggregation and demonstration of an inherited disease with defective platelet aggregation correctable with normal fibronectin.

J. Cell, Biol., 83 (part 2) 61.

Furie, M. and Rifkin, D.B. (1978):

Ordering of proteolytic fragments of human cold insoluble globulin.(CIG)
J. Cell. Biol., 79 (part. 2): 54.

Fyrand, O. (1980):

Studies on fibronectin in the IV indirect immunofluorescence.
Studies in lupus erythematosus.
Br. J. Dermatol., 102:167.

Gahmberg, C.G., Keihn, D. and Hakomori, S. (1974):

Changes in a surface labelled galactoprotein and in glycolipid concentrations in cells transformed by a temperature-sensitive polyoma virus mutant.
Nature, 248:413.

Ganrot, P.O. (1967):

Inhibition of plasmin activity by α_2 -macroglobulin.
Clin. Chem. Acta, 16:328.

Ganrot, P.O. (1972):

Variation of the concentrations of some plasma proteins in normal adults, in pregnant women and in newborns.
Scand. J. Clin. Lab. Invest., 29 suppl. 124:83.

Gans, H. and Tan B.H. (1967):

α_1 -Antitrypsin, an inhibitor for thrombin and plasmin.
Clin. Chem. Acta, 17:111.

Gant, N.F. and Worley, R.J. (1980):

Hypertension in pregnancy concepts and management
Appleton century crofts, New York, 100-115.

Gerbas, F.R., Bottoms, S., Farag, A. and Mammen, E.F. (1990):

Changes in hemostasis activity during and the immediate postpartum period.
Am. J. Obstet. Gynecol. May, 162 (5): 1158-63.

Gettins, P. (1987):

Antithrombin III and its interaction with heparin.
 A comparison of the human, bovine and porcine, Proteins by
 HNMR spectroscopy.
 Biochem., 26:1391-1398.

Gettins, P. and Wooten, E.W. (1987):

On the domain structure of antithrombin III.
 Tentative localization of the heparin binding region using
 HNMR spectroscopy.
 Biochem., 26:4403-4408.

Gilabert, J., Fernandez, J.A., Espana, F., Aznar, J. and Estelles, A. (1988):

Physiological coagulation inhibitors (proetin S, protein C and
 antithrombin III) in severe preeclamptic states and users of
 oral contraceptives.
 Thromb. Res. Feb., 49 (3): 319-29.

Ginsberg, M.H. and Henson, P.M. (1978):

Enhancement of platelet response to immune complexes and
 IgG aggregates by lipid A rich bacterial lipopolysaccharides.
 J. Exp. Med., 147:207.

Ginsberg, M.H., Pointer, R.G., Forsyth, J., Birdwell, C. and Plow, E.F. (1980):

Thrombin increases expression of fibronectin antigen on the
 platelet surface.
 Proc. Natl. Acad. Sci. U.S.A., 77:1409.

Ginsberg, M.H., Forsyth, J., Lightsey, A., chediak, J. and plow, E., F., (1983):

Reduced surface expression and binding of fibronectin by
 thrombin-stimulated thromboasthenic platelets.
 J. Clin. Invest., 71:619.

Girolami, A., Lazzaro, A.R. and Simioni, P. (1988):

The relationship between defective heparin cofactor activities
 and thrombotic phenomena in AT III abnormalities.
 Thromb. and Haemostasis, 59 : 121.

Gow, L., Campbell, D.M. and Ogston, D. (1983):

Blood levels of proteinase inhibitors in preeclampsia.
Br. J. Obstet. Gynecol. Oct., 90 (10): 950-2.

Grinnell, F. (1976):

The serum dependence of baby hamster kidney cell attachment to substratum.
Exp. cell Res., 97:265.

Grinnell, F., Hays, D. G. and Minter, D. (1977):

Cell adhesion and spreading factor partial purification and properties.
Exp. Cell Res. 110:175.

Hahn, L.H. and Yamada, K. (1979):

Identification and isolation of a collagen-binding fragment of the adhesive glycoprotein fibronectin.
proc. Natl. Acad. Sci. U.S.A., 76:1160.

Hahn, E., Wick, G., Pancev, D. and Timpl, R. (1980):

Distribution of basement membrane proteins in normal and fibrotic human liver, collagen type IV, laminine and fibronectin.
Gut, 21:63.

Harker, L.A. and Finch, C.A. (1969):

Thromokinetics in man.
J. Clin. Invest., 48:963.

Haversitck, D.M., Cowan, J.F., Yamada, K.M. and Santoro, S.A. (1985):

Inhibition of platelet adhesion to fibronectin, fibrinogen, and von willebrand factor substrates by a synthetic tetrapeptide derived from the cell-Binding Domain of fibronectin.
Blood, 66 N 4 (Oct.) : 946.

Hayman, E.G. and Ruoslahti, E. (1979):

Distribution of fetal bovine serum fibronectin and endogenous rat cell fibronectin in extracellular matrix.
J. Cell Biol., 83:255.

Heilmann, L., Hojnacki, B. and Spanuth, E. (1991):

Haemostasis and pre-eclampsia.

Geburtshilfe-frauenheilkd. Mar, 51 (3): 223-7.

Quoted from computer abstracts.

Hess, L.W., O'Brien, W.F., Holmberg, J.A., Winkel, C.A., Monaghan, W.P. and Hemming, V.G. (1986):

Plasma and amniotic fluid concentrations of fibronectin during normal pregnancy.

Obstet. Gynecol., 68:25

Highsmith, R.F. and Rosenberg, R.D. (1974):

The inhibitor of human plasmin by human antithrombin heparin cofactor.

J. Biol. Chem., 249:4335-4338.

Ho, C.H., Yang, Z.L., (1992):

The predictive value of the hemostasis parameters in the development of pre-eclampsia.

Throm-Haemost. Feb 3., 67 (2): 214-8.

Hofler, H. and Kingemann, H. (1984):

Fibronectin and factor VIII related antigen in liver cirrhosis and acute liver failure.

J. Clin. Chem. Clin Biochem., 22 (1): 14.

Hogstrop, H. and Saldeen, T. (1982):

Synthesis of alpha 2- antipiasmin by rat liver cells.

Thromb. Res., 28:19.

Holmes, W.E., Lijnen, H.R. and Nelles, L. (1987-a):

α_2 - antipiasmin Enschede: alanine insertion and abolition of plasmin inhibitory activity.

Science, 238:209.

Holmes, W.E., Nelles, L., Lijnen, H.R. and Collen, D. (1987):

Primary structure of human α_2 -Antipiasmin, a serine protease Inhibitor (serpin).

J. biol. Chem., 262 (4): 1659.

Hormann, H. and Seidl, M. (1980):

Affinity chromatography on immobilized fibrin monomer III, the fibrin affinity center of fibronectin.
Hoppe seyle's Z physiol. Chem., 361:1449.

Hoylaerts M., Owen, W.G. and Collen, D. (1984):

Involvement of heparin chain length in the heparin catalyzed inhibition of thrombin by antithrombin III.
J. Biol. Chem., 259:5670.

Hunt L.T. and Dayhoff, M.O. (1980):

A surprising new protein superfamily containing ovalbumin, antithrombin III and α_1 -proteinase inhibitor.
Biochem. Biophysical Research Communication, 95:864.

Hussain R. and Goecke, C. (1993):

New approaches to the aetiology of hypertension in pregnancy.
Obstet Gynecol. XXV International conference of pathophysiology of pregnancy.
Cairo-Egypt Nov, 23-26 Abstracts.

Hynes, R.O. (1986):

Fibronectins.
Scientific American, 254:32-42.

Hynes, R.O. and Bye, J.M. (1974):

Density and Cell Cycle dependence of cell surface proteins in hamster fibroblasts.
Cell, 3:113.

Hynes, R.O., Ali, I.U., Destree, A.T., Matuner, V., Perkins, M.E., Senger, D.R., Wagener, D.D. and Smith K.K. (1978):

A large glycoprotein lost from the surfaces of transformed cells.
Ann. N.Y. Acad. Sci., 312:317.

Ichinose, A., Tamaki, T. and Aoki, N. (1983):

Factor XIII-mediated cross-linking of NH₂-terminal peptide of α_2 -plasmin inhibitor to fibrin.
FEBS let., 153:369.

Ichinose, A., Mimuro, J., Koide, T. and Aoki, N. (1984):

Histidine-rich glycoprotein and α_2 - plasmin inhibitor in
inhibitor of plasminogen binding to fibrin.
Thromb. Res., 33:401.

Iioka, H., Akada, S., Yamada, Y., Moriyama, I.S., and Ichijom, M. (1993):

Effect of lipid peroxidation on platelet aggregation inhibiting
activity of human placental chorioepithelial brush border
membrane vesicles.

Obstet. Gynecol. XXV International conference of
pathophysiology of pregnancy.

Cairo-Egypt, Nov., 23-26 Abstracts.

Ismail, Z.A., Soltan, M.H., Sammour, M.B., Kafafi, S.M., Abdalla, K.A. and Aballa H.H. (1993):

The role of some clinical and laboratory tests in the prediction
of preeclampsia.

Obstet. Gynecol. XXV International conference of
pathophysiology of pregnancy.

Cairo-Egypt Nov., 23-26 Abstracts.

Jack, A.P. and Paul, C.M., (1980):

Hypertensive disorders in pregnancy. William Obstet. P: 665-
700. 16 th Ed. Appleton century-crofts, New York, U.S.A.

Jaffe, E.A. and Mosher, D.F. (1978):

Synthesis of fibronectin by cultured human endothelial cells.
J. Exp. Med., 147:1779.

Jallat, S., Carvallo, D. and Tessier, L.H. (1986):

Altered specificities of genetically engineered $\alpha 1$ - antitrypsin
variants.

Protein Engineering, 1:29-35.

Jelick, F. and Hormann, H. (1979):

Fibronectin (cold insoluble globulin) Influence of heparin and
hyaluronic acid on the binding of native collagen.

Hoppe scyler's Z physiol. Chem., 360:597.

Quoted from computer abstract.

Jileck, F. and Hormann, H. (1977):

Cold-insoluble globulin III. Cyanogen bromide and plasminolysis fragments containing a label introduced by transamidation.

Hoppe seyley's Z physiol. Chem., 358:1165.

Qouted from computer abstract.

Johansson, S., Rubin, K., Hook, M., Ahlgren, T., and Seljelid, R., (1979):

In vitro biosynthesis of cold-insoluble globulin (fibronectin) by mouse peritoneal macrophages FEBS lett, 105:313.

Quoted from the Journal of the American society of Hematology, 56 (2): 148-158.

John, F.T. and Eric, J. (1986):

Interaction of culture derived macrophage with the fibroblast-binding domain of fibronectin is necessary but insufficient signal for fibronectin enhancement of CRI-mediated - phagocytosis.

J. of Immunology, 13 (10) : 3793.

Jordan, R.E., Oosta, G.M., Gardner, W.T. and Rosenberg R.D. (1980):

The kinetics of hemostatic enzyme-antithrombin interactions in the presence of low molecular weight heparin.

J. Biol. Chem., 255:10081.

Jorgensen, A.M., Borders, C.L. and Fish, W.W. (1985):

Arginine residues are critical for the heparin cofactor activity of antithrobin III.

Biochem. J., 231:59-65.

Kaplan, J.E. (1980):

Maintainance of fibrin solubility by plasma fibronectin.

J. lab. Clin. Invest., 96:1054.

kidd, V.J. and Woo, S.L. (1986):

Molecular analysis of the serine proteinase inhibitor gene family. In Barrett A.J. and salvesen G. (eds) proteinase inhibitors, 421 Amsterdam. Elsevier science publishers B.V.

- Killam, A.P., Dillard, S.H., Patton, R.C. and Peerson, P.R. (1975):**
Pregnancy induced hypertension complicated by acute liver disease and disseminated intravascular coagulation.
Am. J. Obstet. Gynecol., 123:823.
- Kleinman, H.K., Wilkes, C.M. and Martin, G.R. (1981):**
Interaction of fibronectin with collagen fibrils.
Biochem., 20:2325.
- Kleinman, H.K., McGoodwin E.B., Martin, G.R., Klebe R.J. Fietzek, P. and Woolley, D.E. (1978):**
Localization of the binding site for cell attachment in the α_1 (1) chain of collagen.
J. Biol. Chem., 253:5642.
- Kluft, C., Los, P. and Jie, A.F. (1984):**
The molecular form of alpha 2- antiplasmin with affinity for plasminogen is selectively bound to fibrin by factor XIII.
Thromb. Res., 33:419.
- Kluft, C., Vellenga, E., Brommer, E.J. and Wijngaards, G. (1982):**
A familial hemorrhage diathesis in a Dutch family: An inherited deficiency of alpha-2 antiplasmin.
Blood, 59:1169.
- Kluft, C., Jie, A.F., Los, P., Dooijewaard, G. and Traas, D.W. (1983):**
Expression of the two forms of α_2 - antiplasmin in functional and immunochemical assays. In progress in fibrinolysis. vol.(6).
J.F. Davidson, F. Bachmann, C.A. Bouvier, and E.K.O. Kruithof, editors, churchill livingstone, Edinburgh. 385-387.
- Kluft, C., Los, P., Jie, A.F., VanHins, bergh, V.W., Vellenga, E., Jespersen, J., and Henny C.P.(1986):**
The mutual relationship between the two molecular forms of the major fibrinolysis inhibitor alpha-2- antiplasmin in blood.
Blood, 67 (3) : 616.

- Kluft, C., Nieuwenhuis, H.K., Rijken, D.C., Groeneveld, E., Wijngaards, G., Van Berkel, W., Dooijewaard G. and Sixma, J.J. (1987):**
 α_2 - Antiplasmin Enschede: dysfunctional α_2 - Antiplasmin molecule associated with an autosomal recessive hemorrhagic disorder.
 J. Clin. Invest., 80:1391.
- Knot, E.A., Dejong, E. and ten cate, J.W. (1986):**
 Purified radiolabeled antithrombin III metabolites in three families with hereditary AT III deficiency:
 Application of a three compartment model.
 Blood, 67:93-98.
- Kobayashi, T. and Terao, T. (1987):**
 Preeclampsia as chronic disseminated intravascular coagulation. Study of two parameters: thrombin- antithrombin III complex and D-dimers.
 Gynecol. Obstet. Invest., 24 (3): 170-8.
- Koh, S.C., Anandakumar, C., Montan, S. and Ratnam, S.S. (1993):**
 Plasminogen activators, plasminogen activator inhibitors and markers of intravascular coagulation in preeclampsia.
 Gynecol. Obstet. Invest., 35 (4): 214-21.
- Koide, T., Odani, S., Takahashi, K., Ono, T. and Sukuragawa, N. (1984):**
 Antithrombin III Toyama: Replacement of arginine 47 by cysteine in hereditary abnormal antithrombin that lacks heparin binding activity.
 Proc. Natl. Acad. Sci., 81:289.
- Kokot, F. and Cekanski, H. (1972):**
 Plasma renin activity in peripheral and uterine venous blood in pregnant and non pregnant women.
 J. Obstet. Gynecol. of Br. Common wealth, 79:72-76.
- Kornblihtt, A.R., Umezawa, K., Vibe-pedersen, K. and Baralle, F.E. (1985):**
 Primary structure of human fibronectin: Differential splicing may generate at least 10 polypeptides from a single gene.
 EMBO. J., 4 : 1755-9.

- Koullapis, E.N., Nicolaides, K., Collins, W.P., Rodack, C.H. and Campbell S. (1982):**
Plasma prostanoids in pregnancy induced hypertension.
Br.J. Obstet. Gynecol., 89:617.
- Kraayenbrink, A.A., Dekker, G.A., Van-Kamp, G.J. and Van-Geijn, H.P. (1993):**
Endothelial vasoactive mediators in preeclampsia.
Am. J. Obstet. Gynecol., 169:160-5.
- Krulthof, E.k.O., Than, C.T., Ransijn, A. and Bachmann, F. (1984):**
Demonstration of a fast-acting inhibitor of plasminogen activators in human plasma.
Blood, 64 (4): 907-913.
- Kuroiwa, A., Igisu, K., Yano, T., Okada, N. and Okada, H. (1988):**
Fibronectin enhances respiratory burst of phagocytes stimulated by zymosan and immune complexes.
Immunology, 65:177.
- Lane, D.A. and Caso, R. (1989):**
Antithrombin: structure, genomic organization, function and inherited deficiency.
Baillieres's clinical Haematology., 2 (4): 961.
- Lane D.A., Erdjument, H. and Thompson, E. (1989):**
A novel amino acid substitution in a congenital variant antithrombin: antithrombin pescara, Arg 393 to pro, caused by a CGT to CCT mutation.
J. Biol. chem., 264:10200.
- Lane, D.A., Flynn, A.M. and Ireland, H. (1987):**
Antithrombin Northwick park: demonstration of an inactive high M.W. complex with increased affinity for heparin.
B.J. Haema., 65:451.
- Lane D.A., Denton, J., Flynn, A.M., Thunberg, L. and Lindahl, U. (1984):**
Anticoagulant activities of heparin oligosaccharides and their neutralization by platelet factor 4.
Biochem. J., 218:725.

- Lane, D.A., Pejler, G., Flynn, A.M., Thompson, E. and Lindahl, U. (1986):**
Neutralization of heparin related saccharides by histidine rich glycoprotein and platelet factor (4).
J. Biol. Chem., 261:3980.
- Lanser, M. and Saba, T. (1981):**
Fibronectin as co-factor necessary for optimal granulocyte phagocytosis of staphaureus.
J. Reticuloendothelial society, 30:415.
- Lao, T.T., Yin, J.A., Ng, W.K. and Yuen, P.M. (1990):**
Relationship between maternal antithrombin III and protein C protein S levels before, during and after delivery.
Gynaecol. Obstet. Invest., 30:87.
- Laurence A., Sherman, L. and Jaan, L. (1982):**
Fibronectin: Blood turnover in normal animals and during intravascular coagulation.
Blood, 60 (3): 558-563.
- Lazarchick, J., Stubbs, T.M., Romein L., Van-Dorsten, J.P. and Loadholt, C.B. (1986):**
Predictive value of fibronectin levels in normotensive gravid women destined to become preeclmaptic.
Am. J. Obstet., Gynecol., 154:1050.
- Leiberman, J.R., Hagay, Z.J., Mazor, M., Wiznitzer, A., Aharon, M., Nathan, I. and Dvilansky, A. (1988):**
Plasma antithrombin III levels in pre-eclampsia and chronic hypertension.
Int. J. Gynecol. Obstet. Aug., 27 (1): 21-4.
- Lijnen, H. R., Holmes, W.E., Van-Hoef, B., Wiman, B., Rodriguez, H. and Collen, D. (1987):**
Amino-acid sequence of human α_2 -Antiplasmin.
Eur. J. Biochem., 166:565.
- Lindheimer, M.D. and Katz, A.I. (1977):**
Kidney function and disease in pregnancy. 82-85 Quoted from Recent Advances In Obstet. and Gynecol., 15:3-32.

Lockwood C.J. and Peters, J.H. (1990):

Increased plasma levels of ED1+ cellular fibronectin precede the clinical signs of preeclampsia.
Am.J. Obstet. Gynecol., 162:358.

Loebermann, H., Tokuoka, R., Deisenhofer, J. and Huber, R. (1984):

Human α_1 -proteinase inhibitor. Crystal structure analysis of two crystal modifications, molecular model and preliminary analysis of the implications for function.
J. Mol. Biol., 177:531.

Loller, P. and Owen, W.G. (1980):

Clearance of thrombin from circulation in rabbits by high-affinity binding sites on endothelium.
J. Clin. Invest., 66:1222.

Loller, P., MacIntosh, S.C. and Owen, W.G. (1984):

Reaction of antithrombin III with thrombin bound to the vascular endothelium.
J. Biol. Chem., 259:4335.

Longas M.O. and Finaly, T.H. (1980):

The covalent nature of human antithrombin III-thrombin bond.
Biochem. J., 189:481.

Luding, H. (1971):

Surface structure of the human term placenta and the uterine wall postpartum in the screen scan.
Am. J. Obstet. Gynecol., 111:328.

MacGillivray, I. (1981):

Raised blood pressure in pregnancy.
Aetiology of preeclampsia.
Br. J. Hosp. Med. Aug, : 110.

Mancini, G., Carbonare, A.O. and Hermens, J.F. (1965):

Immunochemical quantitation by single radial immunodiffusion.
Immunochem., 2:235.

- Marcum, J.A. and Rosenberg, R.D. (1984):**
Anticoagulant active heparin-like molecules from vascular tissue.
Biochem., 23:1730.
- Marcum, J.A., Mckenney, J.B. and Rosenberg, R.D. (1984):**
Acceleration of thrombin-antithrombin complex formation in rat hind-quarters via heparin-like molecules bound to the endothelium.
J. Clin. Invest., 74:341.
- Marcum, J.A., Fritze, L.M., Gall, S.J., Karp, G. and Rosenberg, R.D. (1983):**
Microvascular heparin like species with anticoagulant activity.
Am. J. Physiol., 245:11725.
- Marcum, J.A., Atha, D.H., Fritze, L.M., Naworoth, P., Stern, D. and Rosenberg, R.D. (1986):**
Cloned bovine aortic endothelial cells synthesize anticoagulant active heparin sulphate proteoglycans.
J. Biol. Chem., 261:7507.
- Matthews, C.M. (1957):**
Theory of tracer experiments with ^{131}I - Labelled plasma proteins.
Phys. Med. Biol., 2:36.
- McCafferty, M., Lepow, M., Saba, T., Cho, E., Menwissen, H. and White, J. (1983):**
Normal fibronectin level as a function of age in the pediatric population.
Ped. Res., J. 17 (6):482.
- McGann, B.C. and Triplett, M.J. (1982):**
Plasma AT III concentration and its function in normal persons.
Blood, 64:877.
- Menzel, E. and Dunky, A. (1986):**
Plasma fibronectin in the aged.
J. Gerontol., 19 (3): 152.

- Micheli, M., Defranco, A., Nitti, M., Siro-Brigiani, G., Lucisano, F., Gargane, G., Greco, P. and Pietropado, F. (1989):*
Fibronectin levels in pregnancy and puerperium.
Clin. Exp. Obstet. Gynecol., 16 (2-3): 59.
- Miles, L.A., Plow, E.P., Donnelly, K.J., Hougie, C. and Griflin, J.H. (1982):*
A bleeding disorder due to deficiency in alpha-2- antiplasmin.
Blood, 59:1246.
- Millertz, S. and Clemmensen, I. (1976):*
The primary inhibitor of plasmin in human plasma.
Biochem. J., 159:545.
- Mimuro, J., Koike, Y., Sumi, Y. and Aoki, N. (1987):*
Monoclonal antibodies to discrete regions in α_2 - plasmin in inhibitor.
Blood, 69 (2): 446-453.
- Miura, O., Sugahara, Y., Aoki, N., (1989):*
Hereditary α_2 - plasmin inhibitor deficiency caused by a transport-deficient mutation (α_2 -PI-okinowa). Deletion of Glu¹³⁷ by a trinucleotide deletion blocks intracellular transport.
J. Biol. Chem., 264:18213.
- Mizuochi, T., Fujji, J., Kurachi, K. and Kobta, A. (1980):*
Structural studies of the carbohydrate moiety of antithrombin III.
Archives, Biochem. and Biophysic., 203:458.
- Moodley, J., Norman, R.J. and Reddik, J.B. (1984):*
Central venous concentrations. of immunoreactive prostaglandin E., and 6 keto-prostaglandin F1 in preeclampsia.
Br. Med. J., 288:1487-89.
- Moon, D.G., Kaplan, J.E., and Mazurkewicz, J.E. (1986):*
The inhibitory effect of plasma fibronectin on collagen induced platelet aggregation.
Blood, 67:450.

Moroi, M., and Aoki, N. (1976):

Isolation and characterization of α_2 -plasmin inhibitor from human plasma.

J. Biol. Chem., 251:5956.

Moroi, M. and Aoki, N. (1977):

Inhibition of proteases in coagulation, kinin-forming and complement systems by α_2 -plasmin inhibitor.

J. Biochem., 82:969.

Morrison, P.R., Edsall, J.T. and Miller, S.G. (1948):

Preparation and properties of serum and plasma proteins X VIII. The separation of purified fibrinogen from fraction 1 of Human plasma.

J. Am. Chem. Soc., 70:3013.

Quoted from the Journal of the American society of Hematology (Ed): Mosesson, M.W and Amrani, 56 (2): 145-158 (1980).

Mosesson, M.W. and Amrani, D.L. (1980):

The structure and biological activities of plasma fibronectin.

The Journal of the American society of Hematology, 56 (2): 145-158.

Mosesson, M.W. and Umfleet, R.A. (1970):

The cold-insoluble globulin of human plasma. I; purification, primary characterization, and relationship to fibrinogen and other cold-insoluble fraction components.

J. Biol. Chem., 245:5728.

Mosesson, M.W., Coman, R.W. and Shen, S. (1968):

Chronic intravascular coagulation syndrome, report of a case with special studies of associated plasma cryoprecipitate ("cryofibrinogen").

N. Engl. J. Med., 278:815.

Quoted from the Journal of American society of Hematology (Ed): Mosesson, M.W. and Amrani, D. 1(1980) 56 (2): 145-158.

Mosesson, M.W., Chen, A.B. and Huseby, R.M. (1975):

The cold-insoluble globulin of human plasma studies of its essential structural features.
Biochem. Biophys. Acta, 386:509.

Mosesson, M.W., Wolfenstein-Todel, C., Levin, J. and Characterization, L. (1979):

Characterization of amebocyte coagulogen from the horseshoe crab.
Thromb. Res., 14:765.

Mosher, D.F. (1975):

Cross-linking of cold-insoluble globulin by fibrin stabilizing factor.
J. Biol. Chem., 250:6614.

Mosher, D.F. (1976):

Action of fibrin stabilizing factor on cold insoluble globulin and α_2 -macroglobulin in clotting plasma.
J. Biol. Chem., 251:1639.

Mosher, D.F. (1977):

Labeling of a major fibroblast surface protein (fibronectin) catalyzed by blood coagulation factor XIII.
Biochem. Biophys. Acta, 491:205.

Mosher, D.F. and proctor, R.A. (1980):

Binding and factor XIIIa - mediated cross-linking of a 27 kilodalton fragment of fibronectin to staphylococcus aureus.
Science, 209:927.

Mosher, D.F. and Williams, E.M. (1978):

Fibronectin concentration is decreased in plasma severely ill patients with disseminated coagulation.
J. Lab. Clin. Med., 91:729.

Mosher, D.F., Schad, P.E. and Kleinman, H.K. (1979):

Cross-linking of fibronectin to collagen by blood coagulation factor XIIIa.
J. Clin. Invest., 64:781.

Mosher, D.F., Schad, P.E. and Vann, J.M. (1980):

Cross-linking of collagen and fibronectin by factor XIIIa. Localization of participating glutamyl residues to a tryptic fragment of fibronectin.

J. Biol. Chem., 255:1181.

Mostafa, H.H. and Mostafa, S. (1990):

Urinary calcium creatinine ratio, microalbuminuria, serum procollagen III peptide and plasma fibronectin in prediction of preeclampsia.

Egypt. Obstet. Gynecol. May, XVI (2): 15:22.

Mostafa, H.H., Roshdi, M., Mostafa, S., Eldawakhli, A.E., El ghonemi, S.T. and Sharawi, S. N. (1991):

Serum amino-terminal propeptide of type III procollagen and plasma fibronectin in normal and complicated pregnancy.

The J. of the Egypt. Society of obstet Gynecol. sept. XVII (3): 41.

Mutoh, S., Kobayashi, M., Hirata, J., Itoh- N., Maki-M., Komatsu, Y., Yoshida, A., Sasa, H., Kuroda, K. and Kikuchi, Y. (1992):

Studies of blood coagulation-fibrinolysis regarding kallikrein-kinin system in severe pre-eclampsia.

Agents-Actions- Suppl., 38 (pt 2) : 342.

Nordenman, B. and Bjork, I. (1981):

Influence of ionic strength and PH on the interaction between high-affinity heparin and antithrombin.

Biochimica Biophysica Acta, 672:227.

Oian, R., Kjeldsen, S.E., Eide, I. and Norman, N. (1984):

Increased plasma epinephrine correlates with blood pressure in preeclampsia.

Clin. Exp. Hpert., 83:61-73.

Oishi, M., Yamamoto, M., Yoshida, T., Sade, M. and Matsumoto, K. (1986):

Opsonic activity of plasma fibronectin for staph aureus by human alveolar macrophages.

J. Exp. Med., 149 (1) : 112.

Okajima, K., Abe H., Maeda, S., Motomura, M., Tsyjihata, M., Nagataki, S., Okabe, H. and Takatsuki, K. (1993):

Antithrombin III Nagasaki (ser 116-pro) : A heterozygous variant with Defective Heparin Binding Associated with thrombosis.

Blood, 81, No 5 (March 1): 1300.

Okamiya, H. (1990):

Blood coagulation and fibrinolytic studies in patients with toxemia of pregnancy.

Nippon Sanka fujinka Gakkai. Zasshi Nov. 42 (11): 1557-64.

Quoted from computer abstracts

Olden, K., Pratt, R.M. and Yamada, K.M. (1978):

Role of carbohydrates in protein secretion and turnover: effects of tunicamycin on the major cell surface glycoprotein of chick embryo fibroblasts.

Cell, 1:161.

Olden, K., Pratt, R.M. and Yamada, K.M. (1979):

Role of carbohydrate in biological function of the adhesive glycoprotein fibronectin.

Proc. Natl. Acad. Sci. U.S.A., 76:3343.

Olson, S.T. and Shore, J.D. (1982):

Demonstration of a two-step reaction mechanism for inhibition of a thrombin by antithrombin III and identification of the step affected by heparin.

J. Biol. Chem., 257:14891-14895.

Olson, S.T., Srinivasan, K.R., Bjork, I. and Shore, J.D. (1981):

Binding of high-affinity heparin to antithrombin III. Stopped flow kinetic studies of the binding interaction.

J. Biol. Chem., 256:11073.

Pearlstein, E. (1976):

Plasma membrane glycoprotein which mediates adhesion of fibroblasts to collagen.

Nature, 62:497.

- Pearlstein, F., Gold, I.I. and Garcia-pardo, A. (1980):**
Fibronectin, A review of its structure and biological activity.
Mol.cell Biochem., 29:103.
- Pena, S.D. and Hughes, R.C. (1978):**
Fibroblast to substratum contacts mediated by the different forms of fibronectin.
Cell, Biol. Internatl. Rep., 2:339.
- Perkins, M.E., Ji, T. H. and Hynes, R.O. (1979):**
Cross-linking of fibronectin to sulfated proteoglycans at the cell surface.
Cell, 16:941.
- Perri, R., Key, N. and MCCartly, J. (1982):**
Fibronectin enhances in vitro monocyte macrophage mediated tumouricidal.
Blood, Aug., 60:430.
- Perry-K.G. and Martin, J.N. (1992):**
Abnormal hemostasis and coagulopathy in pre-eclampsia.
Clin.obstet-Gynecol-Jun., 35 (2): 338-50.
- Peters, M., Breederveld, C., Kahle, L.H. and Tencate, J.W. (1982):**
Rapid microanalysis of coagulation parameters by automated chromogenic substrate methods, application in neonatal patients.
Thromb. Res., 28:773-781.
- Petersen, L.C. and Clemmensen, I. (1981):**
Kinetics of plasmin inhibiton in the presence of a synthetic tripeptide substrate.
Biochem. J., 199:121.
- Petersen, T.E., Dudek-Wojciechowska, G., Sottrup-Jensen, L. and Magnusson, S. (1979):**
Primary structure of antithrombin III (heparin cofactor). partial homology between α_1 - antitrypsin and antithrombin III. In collen, D, wiman, B., verstraetete M (eds) the physiological inhibitors of blood coagulation and fibrinolysis, pp 43-54 Amsterdam: Elsevier science publishers.

- Peterson, C.B. and Blackburn, M.N. (1985):**
Isolation and characterization of an antithrombin III variant with reduced carbohydrate content and enhanced heparin binding.
J. Biol. Chem., 260:610.
- Peterson, C.B. and Blackburn, M.N. (1987):**
Antithrombin conformation and the catalytic role of heparin Does cleavage by thrombin induce structural changes in the heparin-binding region of antithrombin,
J. Biol. Chem., 262:7552.
- Peterson, C.B., Noyes, C.M., Pecon, J.M., Chrch, F.C. and Blackburn, M.W. (1987):**
Identification of a lysyl residue in antithrombin which is essential for heparin binding.
J. Biol. Chem., 26:8061-8065.
- Pierschbacher, M.D. and Ruoslahti, E. (1984):**
Cell attachment activity of fibronectin can be duplicated by small synthetic fragments of the molecule.
Nature 309:30.
- Pignot, S. and Busine, A. (1989):**
Fibronectin an early marker of pre-eclampsia.
J. Gynecol. obstet. Biol. Reprod., 18 (7): 867.
- Pinto, S., Abbate, R. Rostagnon, C., Bruni, V., Rosati, D. and Nerisneri, G., G. (1988):**
Increased thrombin generation in normal pregnancy.
Acta Eur. Fertil. sep., 19 (5): 263-7.
- Pipkin, F.P. (1984):**
Medical statistics made easy. Churchill livingstone.
Edinburgh-london.
Melbourne and New York.
- Plow, E.F. and Ginsberg, M.H. (1981):**
Specific and saturable binding of plasma fibronectin to thrombin-stimulated human platelets.
J. Biol. Chem., 256:9477.

Plow, E.F., Birdwell, C. and Ginsberg, M.H. (1979):

Identification and quantitation of platelet-associated fibronectin antigen.
J. Clin. Invest., 63:540.

Plow, E.F., Marguerie, G.A. and Ginsberg, M.H. (1985):

Fibronectin Binding to thrombin-stimulated platelets.
Dependent pathways.
Blood, 66 No. (1) :26:32.

Plow, E.F., Srouji, A.H., Meyer, D., Margucrie, G.A. and Ginsberg, M.H. (1984):

Evidence that three adhesive proteins interact with a common recognition site on activated platelets.
J. Biol. Chem., 259:5388.

Pomerantz, M.W. and Owen, W.G. (1978):

A catalytic role heparin. Evidence for a ternary complex of heparin cofactor, thrombin and heparin.
Biochemica. Biophysical Acta, 535:66.

Posttethwaite, A.E., Koski-oja, J., Balian, G. and Kang. A.J. (1981):

Induction of fibroblast chemotaxis by fibronectin.
Localisation of the chemotactic region to a 140,000 M.W. non-gelatin binding fragment.
J. Exp. Med., 153:494

Potempa, J., Shieh, B.H. and Travis, J. (1988):

Alpha-2-antiplasmin, a serpin with two separate but overlapping reactive sites.
Science, 241:699.

Prochownik, E.V., Markham, A.F., and Orkin, S.H. (1983).

Isolation of a DNA clone for human antithrombin III.
J. Biol. Chem., 258:8389.

Proctor, R.A., Prendergast, E. and Mosher, D.F. (1982):

Fibronectin mediates attachment of staphylococcus aureus to human neutrophils.
Blood, 59 (4): 681.

- Pulkkinen, M.D., Lehto, M., Jalkanen, M. and Nanto, S.K. (1984):**
Collagen types and fibronectin in the uterine muscle of normal and hypertensive pregnant patients.
Am. J. Obstet. Gynecol. Aug., (1), 149 (7): 711-7.
- Rajaraman, R., MacSween, J.M. and Murdock, C.A. (1983):**
Exogenous fibronectin requirement for adhesion by neoplastic human cells.
Exp. Cell. Biol., 51:9-18.
- Ramon, R. (1983):**
Preeclampsia, eclampsia and other gestational edema, proteinuria and hypertension disorders.
Current obstetric and Gynecology diagnosis and treatment, 4th. Ed. Lange Medical Duliban. P., 738.
- Rathaus, M.Kovatz, S., Aderet, N.B. and Bernheim, J. (1982):**
Renal prostaglandins in preeclampsia.
Clin. Nephrol., 18:216.
- Reinthalier, A., Mursch-Edlmayer, G. and Tatra, G. (1990):**
Thrombin-antithrombin III Complex levels in normal pregnancy with hypertensive disorders and after delivery.
Br. J. Obstet. Gynecol. Jun., 97 (6): 506. 10.
- Reitman, S. and Frankel, S. (1957):**
Colorimetric, determination of GOT and GPT activity
Am. J. Clin. Path., 28:56.
- Rijken, D.C., Groeneveld, E., Kluft, C. and Nieuwenhuis, H. K. (1988):**
Alpha 2- antiplasmin Enschede is not an inhibitor, but a substrate, of plasmin.
Biochem. J., 255:609.
- Robertson, E.G. (1971):**
The natural history of oedema during pregnancy.
J. Obstet. Gynecol. Br. Common wealth, 78:520-529.
- Romero, R. and Duffy, T.P. (1980):**
Platelet disorders in pregnancy.
Clinics in perinatology, 7:327.

Romero, R., Lockwood, C.J. and Oyarsun, E. (1988):
 Toxemia: New concepts in an old disease.
 Semin. Perinatol, 12:302-23.

Rosenberg, R.D. and Damus, P.A. (1973):
 The purification and mechanism of action of human
 antithrombin-heparin cofactor.
 J. Biol. Chem., 248:6490.

Rosenberg, R.D. and Jordan, R.E. (1977):
 The naturalization of thrombin and other serine proteases of
 the hemostatic mechanism.
 Chemistry and Biology of thrombin.
 Edited by Lundblad R.L., Fenton, J.W. 375-393.

**Ruelland, A., Chevrantbreton, O., LeGras, B., Clerc, C. and Cloarec, L.
 (1985):**
 Amniotic fibronectin in normal pregnancy, and pregnancy
 with fetal anencephaly.
 J. Gynecol. Obstet. Biol. Reprod., 14 (3): 305-8.

Ruggeri, Z.M., Bader, R. and de Marco, L. (1982):
 Glanzmann thrombasthenia. Deficient binding of von Wille-
 brand factor to thrombin stimulated platelets.
 Proc. Natl. Acad. Sci. U.S.A., 79:6038.

Ruoslahti, E., Engvall, E., (1978):
 Immunochemical and collagen-binding properties of
 fibronectin.
 Ann. Ny Acad. Sci, 312:178.

Ruoslahti, E. and Vaheri, A. (1974):
 Novel human serum protein from fibroblast plasma
 membrane.
 Nature, 248:790.

Ruoslahti, E. and Vaheri, A. (1975):
 Interaction of soluble fibroblast surface antigen with
 fibrinogen and fibrin. Identity with cold insoluble globulin of
 human plasma.
 J. Exp. Med., 141:497.

Ruoslahti, E. Vuento, M. and Engvall, E. (1978):

Interaction of fibronectin with antibodies and collagen in radiomimunoassay.

Biochem. Biophys. Acta, 534:210.

Ruoslahti, E., Engvall, E., and Hayman, E. G. (1981):

Fibronectin current concepts of its structure and functions.

Collogen Res., 1:95-128.

Ruoslahti, E., Vaheri, A., Kuusela, P. and Linder, E. (1973):

Fibroblast surface antigen: A new serum protein.

Biochim, Biophys. Acta, 322:352.

Ruoslahti, E., Hayman, E. G., Kuusla, P., Shively, J.E. and Engvall, E. (1979):

Isolation of tryptic fragment containing the collagen binding site of plasma fibronectin.

J. Biol. chem., 254:6054.

Saba, T.M. (1970):

Physiology and physiopathology of the reticuloendothelial system.

Arch. Intern. Med., 126:1031.

Saba, T.M. and Jaffe, E. (1980):

Plasma fibronectin (opsonic glycoprotein) Its synthesis by vascular endothelial cells and role in cardiopulmonary integrity after trauma as related to reticuloendothelial function.

Am. J. Med., 68 : 577.

Saba, T., Blumenstock, F., Shah, D.M., Candaburn, R.H., Hrinda, M.E. and Deno, D.C. (1986):

Reversal of opsonic deficiency in surgical trauma and burn patients by infusion of human plasma fibronectin, correlation with experimental observation.

Am. J. Med., 80 (2): 129.

Saito, H., Goodnough, L.T., Knowles, B.B. and Aden, D.P. (1982):

Synthesis and secretion of α_2 - plasmin inhibitor by established human liver cell lines.

Proc. Natl. Acad. Sci. U.S.A., 79:5684.

Sakata, Y., Mimuro, J. and Aoki, N. (1984):

Differential binding of plasminogen to crosslinked and non cross-linked fibrin: Its significance in hemostatic defect in factor XIII deficiency.

Blood, 63:1393.

Sakata, Y., Tatena, K., Tamaki, T. and Aoki, N. (1979):

Calcium dependent binding of α_2 -plasmin inhibitor to fibrin.

Thromb. Res., 16:279.

Saleh, A.A., Bottoms, S.F., Norman, G., Farag, A. and Mammen, E.F. (1988):

Hemostasis in hypertensive disorders of pregnancy.

Obstet. Gynecol. May, 71 (5): 719-22.

Saleh, A.A., Bottoms, S.F., Farag, A.M., Dombrowski, M.P., Welch, R.A., Norman, G. and Mammen, E.F. (1992):

Markers for endothelial injury, clotting and platelet activation in preeclampsia.

Arch. Gynecol. Obstet., 251 (3): 105-10.

Salem, H.T., Mostafa, S.A., Abdel-Moniem, N., Kamel, H.S. and Mohamed, S.A. (1993):

Roll-over test, plasma fibronectin, plasma antithrombin III and serum iron leveles in prediction of preeclampsia

Obstet. Gynecol. XXV International conference of pathophysiology of pregnancy.

Cairo-Egypt November 23-26 Abstracts.

Samson, D., Stirling, Y., Woolf, L., Howarth, D., Segatchian, M.J. and dechazal, R. (1984):

Management of a planned pregnancy in a patient congenital antithrombin deficiency.

Br. J. Haemato., 56:243.

Santoro, S.A. (1983):

Inhibition of platelet aggregation by fibronectin.

Biochem. Biophys. Res. Commun., 116:135.

Sas, G., Koves, A., Peto, I., and Domjan, G. (1978):

On the inheritance of the abnormal antithrombin III (AT III, Budapest') .
Thromb. Haemio., 39:530-532.

Sasaki, T., Morita, T. and Twanagea, S. (1986):

Identification of the plasminogen binding site of human α_2 -plasmin inhibitor.
J. Biochem., 99:1699.

Seki, H. (1986):

Studies on plasma proteins in normal pregnancy and gynecological malignancy.
Nippon. Sanka, Fujinka. Gakkai, Zasshi. Mar., 38 (3): 317-26.
Quoted from compouter abstract.

Sekiguchi, K., Fukuda, M. and Hakomori, S. (1981):

Domain structure of hamster plasma fibronectin isolation and characterization of four functionally distinct domains and their unequal distribution between two subunit polypeptides.
J. Biol. Chem., 256:6452.

Shaarawy, M., El-Mallah, S.Y. and El-Temamy, M.M. (1989):

Urinary IgG, plasma Antithrombin- III and serum spl glycoprotein levels: new multiparametric biochemical profile for the assessment of the severity of pre-eclampsia and the prediction of fetal outcome.
J. Egypt Soci. Obstet. Gynecol., X V, (1): 37.

Sherlock, S. (1985):

Hepatic cirrhosis, In: Diseases of liver and Biliary system: Sherlock. S., (7th) ed; 335.
Blackwell scientific publication oxford.

Smith, J.W. and Knauer, D.J. (1987):

A heparin binding site in antithrombin III.
J. Biol. Chem., 262:11964-11972.

Smith, R.T. and Vonkroff, R. (1957):

A heparin-precipitable fraction of human plasma .

Isolation and characterization of fraction.

J. Clin. Invest., 36:596.

Quoted from the Journal of the american society of Hematology (Ed). Mosesson M. Wan Amrani, 56 (2) 145-158, (1980).

Sprengers, E.D. and Kluft, C. (1987):

Plasminogen activator inhibitors.

Blood, 69 (2) : 381-387.

Stathakis, N.E. and Mosesson, M.W. (1977):

Interactions among heparin, cold insoluble globulin, and fibrinogen in formation of the heparin precipitable fraction of plasma.

J. Clin. Invest., 60:55.

Stathakis, N.E., Founta, S.A. and Tsianos, E. (1981):

Plasma fibronectin in normal subjects and in various disease states.

J. Clin. Pathol., 34:504-508.

Stenman, S. and Vaheri, A. (1978):

Distribution of a major connective tissue protein, fibronectin, in normal human tissue.

J. Exp. Med., 147:1054.

Stephens, A.W., Siddiqui, A. and Hirs, C.H. (1988):

Site-directed mutagenesis of the reactive center (serine 394) of antithrombin III.

J. Biol. Chem., 263:15849.

Stubbs, T.M., Lazarchick, J. and Horger, E.D. (1984):

Plasma fibronectin levels in preeclampsia. a possible biochemical marker for vascular endothelial damage.

Am. J. Obstet. Gynecol. Dec., 150 (7): 885-7.

Suarez, C.R., Walenga, J., Mangagna, L.C. and Fareed, J. (1985):

Neonatal and maternal fibrinolysis: activation at time of birth.

Am. J. Hematol. Aug., 19 (4): 365-72.

- Sumi, Y., Ichikawa, Y., Nakamura, Y., Miura, O. and Aoki, N. (1989):**
Expression and characterization of pro α_2 - antiplasmin inhibitor.
J. Biochem., 106:703.
- Tamaki, T. and Aoki, N. (1981):**
Cross-linking of α_2 -plasmin inhibitor and fibronectin to fibrin by fibrin-stabilizing factor.
Biochem. Biophys. Acta, 661:280.
- Tamaki, T. and Aoki, N. (1982):**
Cross-linking of α_2 - plasmin inhibitor to fibrin catalyzed by activated fibrin-stabilizing factor.
J. Biol. Chem., 257:14767.
- Tamas, P., Feledi, E., Ertl, T., Kett, A. and Werling, J. (1992):**
Maternal plasma fibronectin and neonatal birth weight.
Gynecol. Obstet. Invest., 33 (2) : 124-5.
- Taylor, R.N., Casal, D.C., Jones, L.A., Varma, M., Martin, J.N. and Roberts, J.M. (1991):**
Selective effects of pre-eclamptic sera on human endothelial cell procoagulant protein expression.
Am. J. Obstet. Gynecol. Dec., 165 (6 pt.1): 1705-10.
- Taylor, R.N., Crombleholme, W.R., Friedman, S.A., Jones, L.A., Casal D.C. and Roberts, J.M. (1991):**
High plasma cellular fibronectin levels correlate with biochemical and clinical features of preeclampsia but cannot be attributed to hypertension alone.
Am. J. Obstet. Gynecol. Oct., 165 (4pt 1) : 895-901.
- Teger-Nilsson, A.C. (1979):**
Use of chromogenic substrates for screening of inhibitors of coagulation and fibrinolysis in patients. In chromogenic peptide Substrates: chemistry and clinical usage (Scully, M.F., and Kakkar, V.V. Eds), churchill Livingstone, Edinburgh PP: 269-276).
- Teisner, B., Davery, M.W. and Grudzinskas, J.G. (1982):**
Circulating antithrombins in pregnancy.
Br. J. Obstet. Gynecol., 89:62-64.

Terao, T., Maloi, M., Ikenae, T., Gotoh, K., Murato, M., Iwasaki, H., Shibata, J., Nakabayashi, M., Muraoka, M. and Takeda, Y. (1991):

The relationship between clinical signs and hypercoagulable state in toxemia of pregnancy.

Gynecol.Obstet.Invest., 31 (2): 74:85.

Terao, T., Kobayashi, T., Imai, N., Oda, H. and Karasawa, T. (1989):

Pathological state of the coagulatory and fibrinolytic system in preeclampsia and the possibility of its relation with AT III concentrate.

Asia Oceania, J. Obstet. Gynecol. Mar., 15 (1): 25-32.

Thaler, E. and Lechner, K. (1981):

Antithrombin III deficiency and thromboembolism.

Clinics in haemato., 10:369.

Thiagarajha, S., Bourgeois, F.J., Barbert, G.M. and Caulde, M.R., (1984):

Thrombocytopenia in preeclampsia: associated abnormalities and management principles.

Am. J. Obstet. Gynecol., 150:1-6.

Thurnau, G.R., Morgan, M.A., Brubaker, D., Romine, M., Blackshaw C., Fishburne, J.I. and Fishburn, I.J. (1987):

Plasma fibronectin levels in normal pregnancy and pre-eclampsia a preliminary report.

Int. J. Gynecol. Obstet., Dec., 25 (6): 441-4.

Tong, Y. (1991):

Changes of plasma fibronectin and antithrombin III in pregnancy induced hypertension.

Chung-Hua-Fu-Chan-Ko-Tsa-Chin. Jan.; 26 (1) : 66-68.

Quoted from computer abstracts.

Travis, J. and Salvesen, G.S. (1983):

Human plasma proteinase inhibitors.

Annu. Rev. Biochem., 52:655.

Trinder, P. (1969):

Determination of glucose in blood using glucose oxidase with an alternative oxygen acceptor.
Ann. Clin. Biochem., 6:24-27.

Vaheri, A., Kurkinen, M., Lchto, V.P., Linder, E.J. and Timpl, R. (1978):

Co-distribution of pericellular matrix proteins in cultured fibroblasts and loss in transformation: fibronectin and procollagen.
Proc. Natl. Acad. Sci., U.S.A., 75:4944.

Vanwersch, J.W. and Ubachs, J.M. (1991):

Blood coagulation and fibrinolysis during normal pregnancy.
Eur. J. Clin. Chem. Clin. Biochem. Jan., 29 (1): 45-50.

Visser, W., Derkx, F., Markovitz, J., Debruin, R. and Wallenburg, H. (1988):

Cardiac production of natriuretic peptide in preeclampsia.
International society for study of hypertension in pregnancy.
Scientific program and abstracts 6 th. International congress.
Montreal, Canada. May, 22:26-63.

Wagner, D.D. and Hynes, R.O. (1979):

- Domain Structure of fibronectin and its relation to function.
Disulfide and sulfhydryl groups.
J. Biol. Chem. 254:6746.

Wagner, D.D. and Hynes, R.O. (1980):

Topological arrangement of the major structural features of fibronectin.
J. Biol. Chem., 255:1672.

Walker, J.E., Campbell, D.M. and Ogston, D. (1982):

Blood levels of proteinase inhibitors in pregnancy.
Br.J. Obstet. Gynecol., 89:208.

Wallmo, L., Gyzander, E., Karlsson, K., Lindstedt, G., Radherg, T. and Teger-Nilsson, A.C. (1982):

Alpha 2- antiplasmin and alpha 2- macroglobulin the main inhibitors of fibrinolysis during the menstrual cycle, pregnancy, delivery, and treatment with oral contraceptives.
Acta Obstet. Gynecol., Scand, 61 (2): 417.

Weenink, G.H., Borm, J., Ten-Cate, J.W. and Treffers, P.E. (1983):

Antithrombin III (AT III) levels in normotensive and hypertensive pregnancy.
Gynecol. Obstet. Invest., 16 (4): 230-42.

Weenink, G.H., Treffers P.E., Vijn, P., Smorenberg-Schoorl, M.E. and Ten-cate J.W. (1983-a):

Plasma antithrombin III levels in preeclampsia.
Clin. Exp. Hypert., (B) 2 (1) : 145-62.

Weenink, G.H., Treffers, P.E., Vijn, P., Smorenberg-schoorl, M.E. and Ten-cate, J.W. (1984):

Antithrombin III levels in preeclampsia correlate with maternal and fetal morbidity.
Am.J. Obstet. Gynecol. Apr., 15:148 (8): 1092 -7.

Weiner, C.P. (1988):

The mechanism of reduced antithrombin III activity in women with preeclampsia.
Obstet. Gynecol., Dec., 72 (6): 847-9

Weiner, C.P. and Brandt, J. (1980):

Plasma antithrombin III activity in normal prgnancy.
Obstet. Gynecol., 56:601.

Weiner, C.P. and Brandt, J. (1982):

Plasma antithrombin III activity an aid in the diagnosis of pre-eclampsia-eclampsia.
Am.J. obstet. Gynecol., Feb., 142 (3): 275-81:

Weiner, C.P. and Kwaan, H.C. (1985):

Antithrombin III activity in women with hypertension during pregnancy.
Obstet., Gynecol., 65:301.

Weiner, C.P. and Bonstib, S.M. (1990):

Relationship between renal biopsy and plasma antithrombin III activity in women with early onset preeclampsia.
Am. J. Perinatol. Apr., 7 (2) : 139-43.

Weiner, C.P., Kwaan, H.C., Xu, C., Paul, M., Burmeister, L. and Hauck, W. (1985):

Antithrombin III activity in women with hypertension during pregnancy.
Obstet. Gynecol., 56:601.

Weiner, C.P., Herrig, J.E., Pelzer, G.D. and Heilskv, J. (1990):

Elimination of antithrombin III concentrate in healthy pregnant and preeclamptic women with an acquired, antithrombin. III deficiency.
Thromb. Res. May, 15, 58 (4) : 365-401.

Weir, R.J., Brown, J.J., Fraser, R., Lever, A.F. and McIlwain, E. (1973):

Plasma renin, renin substrate, angiotensin II and aldosterone in hypertensive diseases of pregnancy.
Lancet, 1: 291-294.

Weiss, H.J. (1978):

Drug therapy: Antiplatelet therapy.
N. Engl. J. Med., 298:1344.

Weiss, M.A., Oai, B.S. and Oai, Y.M. (1979):

Immunofluorescent localization of fibronectin in the human kidney.
Lab. Invest., 41:340.

Weissmann, G., Pearlstein, E., Perez, H.D., Falkow, S., Goldstein, I.M. and Hoffstein, S.T. (1980):

Neutrophils synthesize and deposit fibronectin on surfaces to which they attach.
Trans. Assoc. Am. physicians, 93:72.

Welch, M.P., Odland, G.F. and Clark, R.A., (1990):

Temporal relationships of F actin bundle formation, collagen and fibronectin matrix assembly and fibronectin receptor expression to wound contraction.

J. Cell Biol., 110:133-145.

Williams, E., Janmey, P. and Peny, J. (1982):

Confirmational studies of fibronectin effects PH, ionic strength and collagen binding.

J. Biol. chem., 527:1473.

Wiman, B. and Collen, D. (1978):

On the kinetics of the reaction between human antiplasmin and plasmin.

Eur. J. Biochem., 84:573.

Wiman, B. and Collen, D. (1979):

On the mechanism of the reaction between human alpha 2-antiplasmin and plasmin.

J. Biol. Chem., 254:9291.

Wiman, B., Boman, L. and Collen, D. (1978):

On the kinetics of the reaction between human antiplasmin and a low molecular weight form of plasmin.

Eur. J. Biochem., 87:143.

Wiman, B., Nilsson, T. and Collen, D. (1980):

On the mechanism of the reaction between plasmin and α_2 -antiplasmin. In protides of the Biological fluids. H. Peterse. . editor pergamon press. Ltd., Oxford, 379-382.

Wiman, B., Nilsson, T. and Cedegren, B. (1982):

Studies on a form of α_2 - antiplasmin in plasma which does not interact with the lysine binding sites in plasminogen.

Thromb. Res., 28:193.

Winter, J.H., Bennett, B. and Watt, L.J. (1982):

Confirmation of linkage between antithrombin III and Duffy blood group and assignment of AT III to 1q22-q25.

Ann. Hum. Genetics, 46 : 29.

- Winter, J.H., Fenech, A., Mackie, M., Bennett, B. and Douglas, A., S. (1982).**
Treatment of venous thrombosis in antithrombin III deficient patients with concentrates of antithrombin III.
Clin. Lab. Haemat., 4:101.
- Wodzicki, A.M. and Coopland, A.T. (1977):**
Coagulation and the hypertensive disease of pregnancy.
Am. J. Obstet. Gynecol., 129 : 393.
- Wolff, I., Timpl, R., Pecker, I. and Steffen, C. (1967):**
A two- component system of human serum agglutinating gelatin Coated erythrocytes.
Vox. Sang., 12:443.
- Wong, R.E., Chang, T. and Feinman, R.D. (1982):**
Reaction of antithrombin with proteases. Nature of the reaction with trypsin:
Biochem., 21 : 6.
- Xie, Y.S. (1989):**
Predictive and diagnostic value fibronectin levels in pregnancy-Induced hypertension.
Chung-Hua. Fu. Chanko. Tsachin. Mar. 24 (2): 70.
Quated from computer. abstracts.
- Yamada, K.M. and Kennedy, D.W. (1979):**
Fibroblast cellular and plasma fibronectins are simillar but not identical.
J.Cell Biol., 80:492.
- Yamada, K.M. and Weston, J.A. (1974):**
Isolation of major cell surface glycoprotein from fibroblasts.
Proc. Natl. Acad. Sci. U.S.A., 71:3492.
- Yamada, K.M. and Kennedy, D.W. (1984):**
Dualistic nature of adhesive protein function. Fibronectin and its biologically active peptide fragments can autoinhibit fibronectin function.
J.Cell Biol., 99:29.

Yamada, K.M., Yamada, S.S. and Pastan, I. (1976):

Cell surface protein partially restores morphology, adhesiveness, and contact inhibition of movement to transformed fibroblasts.

Proc. Natl. Acad. Sci. U.S.A., 73:1217.

Yamada, K.M., Schlesinger, D.H., Kennedy, D.W. and Pastan, I. (1977):

Characterization of a major fibroblast cell-surface glycoprotein.

Biochemistry, 16 : 5552.

Yamaguchi, Y., Isemura, M., Kosakai, M., Stato, A., Suzuki, M., Kan, M. and Yosizawa, Z. (1984):

Characterization of fibronectin from fetal human plasma in comparison with adult plasma fibronectin.

Biochem. Biophys. Acta, 790:53.

Yin, E.T., Wessler, S. and Stoll, P.J. (1971):

Identity of plasma activated factor X inhibitor with antithrombin III and heparin cofactor.

J. Biol. Chem., 246:3712-3719.

Yoshimura, T., Ito, M., Nakamura, T. and Okamura, H. (1992):

The influence of labor on thrombotic and fibrinolytic systems.

Eur. J. Obstet. Gynecol. Reprod. Biol. May, 13:44 (3): 195-9.

Yuasa, S., Ishizawa, M., Yuki, Y., Minoura, T. Fjuita, N., Takahashi, H., Yoshimura, M., Yamamoto, T. and Okada, H. (1992):

Coagulation and fibrinolysis in pregnancy.

Rinsho. Byori Dec., 40 (12): 1287-91. Quoted from compouter abstracts.

Zlatopolskil, A.D., Chubukina, A.N. and Zaidenbeg, M.A. (1989):

Influence of fibronectin fragments on the proliferative activity of fibroblasts.

Biochemistry, 54:55-60.

Zucker, M.B., Mosesson, M.W., Brockman, M.J. and Kaplan, K.L. (1979):

Release of platelet fibronectin (cold-insoluble globulin) from alpha granules induced by thrombin or collagen.

Lack of requirement for plasma fibronectin in ADP-induced platelet aggregation.

Blood, 54:8.