

REFERENCES

References

- Abdallah,N.M., Nassar,M.I., AbdEl-Hameed,G. and Marcell,N.G.
(1990):
Carcinoembryonic antigen in bilharzial polypi.
2nd Int. Conference of Tumor Markers Oncology, Cairo,
Egypt.
- Akiyama,T.,Sudo,C.,Ogawara,H.,Totoshima,K. and Yamamoto,T.
(1986):
The ptdoct of human *c-erbB-2* gene : A 185 Kilodalton
glycoprotein with tyrosine kinase activity.
Science , 232 : 1646 .
- Ali,I.U., Campbell,G, Lidereau,R. and Callahan,R. (1988) :
Lack of evidence for the prognostic significance of *c-erbB-2*
amplification in human breast carcinoma.
Oncogene Res. , 3 : 139.
- Arends, J.W., Groniowski, M.M., De Koning Gans,H.J.,and Bosman,
F.T.(1983) :
Immunohistochemical study of the distribution of secretory
component and IgA in normal and diseased uterine mucosa .
Int. J. Gynecol. Pathol. , 2 : 171 .
- Arias-Stella ,J. (1959) :
Atypical endometrial changes associated with the presence of
chorionic tissue .
AMA Arch. Pathol. , 58 : 112.
- Arnon,R., Bustin,M. and Calef,E. (1976) :
Cross-reactivity of antibodies to a synthetic undecapeptide
analogous to the amino terminal segment of the human sera.
Proc. Natl. Acad. Sci. USA, 73 : 2123 .

- Atkin , N.B. , Richards , B.M. , and Ross , A.J. (1959) :**
The deoxyribonucleic acid content of carcinoma of the uterus.
Br. J. Cancer , 13 : 773 .
- Atkin , N.B. (1976) :**
Prognostic significance of ploidy level in human tumors. I :
Carcinoma of the uterus.
J. Natl. Cancer Inst. , 56 : 909 .
- Atkin,N.B. (1985):**
Antioncogenes .
Lancet , ii : 1189 .
- Ausems ,E.W.,Vander Kamp, J. K.and Book ,J. P. (1985):**
Nuclear morphometry in the determinations of the prognosis of
marked atypical endometrial hyperplasia.
Int .J. Gynecol. pathol . , 4 : 180 .
- Baisch,H., Beck,H.P. and Christensen,I.J. (1982):**
A comparison of mathematical methods for the analysis of
DNA histograms obtained by flow cytometry.
Cell Tissue Kinet. , 15 : 235 .
- Baltimore, D. (1986):**
Oncogenes : The beginning of the molecular description of a
cancer .
Cell Biophys. , 9 (1,2) : 17 .
- Banjo,C., Gold,P. , Freedman,S.O. and Krupey,J. (1972) :**
Immunologically active heterosaccharides of carcinoembryonic
antigen of human digestive system .
Nature , 238 : 183 .
- Bargmann.C.I., Hung,M.C. and weinberg,R.A. (1986 A) :**
Multiple independent activations of the neu oncogene by a
point mutation altering the transmembrane domain of p185.
Cell , 45 : 649 .
- Bargmann,C.I. , Hung ,M.C. and Weinberg,R.A. (1986 B):**
The neu oncogene encodes an epidermal growth factor receptor
-related protein.
Nature , 319 : 226 .

- Bargmann,C.I. and Weinberg ,R.A. (1988):**
Oncogenic activation of the neu-encoded receptor protein by point mutation and deletion.
EMBO J. , 7 : 2043.
- Barlogie , B. , Gohde,W., Dosik,G. and Schumann , j. (1980) :**
Cellular DNA content as a marker of neoplasia in man.
Am. J. Med. , 69 : 195 .
- Barlogie, B., Ruber,M .N., Schumann, J . , Johnston, T.D. , Derwinko,B.,Swartzendruber,D.E. and Gonde,W. (1984):**
Flow cytometry in clinical cancer research .
Cancer Res. , 43 : 3982 .
- Barnes,D.M., Lammie,G.A., Millis,R.R., Gullick,W.L., Allen,D.S. and Altman ,D.G.(1988):**
An immunohistochemical evaluation of *c-erbB-2* expression in human breast carcinoma.
Br. J. Cancer , 58 : 448.
- Beatty,J.D., Romero,G.,Brown,P.W.,Lawrence,W. and Terz,J.J. (1979):**
Clinical value of carcinoembryonic antigen. Diagnosis, prognosis and follow up of patients with cancer.
Arch. Surg.,114: 563.
- Begent, R.H. (1987) :**
The value of CEA in clinical practice .
Br. J. Hosp. Med. , 37 (4) : 335 .
- Benjamin, I. and Block, R.E. (1977) :**
Endometrial response to estrogen and progesterone therapy in patients with gonadal dysgenesis.
Obstet. Gynecol. , 50 : 136 .
- Berchuck ,A . , Kamel,A. whitaker, R. et al. , (1990) :**
Overexpression of HER-2/neu is associated with poor survival in advanced epithelial ovarian cancer .
Cancer Res. , 50 : 4087 .

- Berchuck,A., Rodriguez,G. Kinney ,R.B., Soper,J.T., Dodge,R.K.,
Clarke Pearson,D.L. and Bast,R.C. (1991):**
Overexpression of HER-2/neu in endometrial cancer is
associated with advanced stage disease.
Am.J. Obstet. Gynecol. , 164 : 15 .
- Berger, M. S., Locher,G. W., Saurer,S., Gullick,W. J., Waterfield, M. D.,
Groner,M. and Hynes,N.E.(1988):**
Correlation of *c-erbB-2* gene amplification and protein
expression in human breast carcinomas with nodal status and
nuclear grading.
Cancer Res. , 48 : 1238 .
- Beutler, H. K., Dokerty, M. B. and Randall, L. M. (1963):**
Precancerous lesions of the endometrium .
Am. J. Obstet. Gyencol. , 86 : 433 .
- Bigsby, R.M., Alixin, L., Bomalaski, J., Stehman,F.B., Look,K.Y. and
Sutton, G.P. (1992):**
Immunohistochemical study of *HER-2/neu*, epidermal growth
factor receptor and steroid receptor expression in normal and
malignant endometrium .
Obstet. Gynecol. , 79 : 95 .
- Biunno, I. , Pozzi, M.R. , and Terrier, M.J. (1988) :**
Structure and expression of oncogenes in surgical specimens of
human breast carcinomas.
Br. J. Cancer , 57 : 464 .
- Bocking , A. , Chatelain ,R. , Bisterfeld ,S. , Noll, E. Biesterfeld , D.
Wohltmann ,D. and Goecke, C. (1989) :**
DNA grading of malignancy in breast cancer : prognostic
validity, reproducibility and comaprison with other
classifications.
Analyt. Quant. Cytol. Histol. , 11 : 73 .
- Bonte, J . (1986):**
Diagnosis and treatment of precancerous endometrial lesions.
**In:Schultz(ed) Endometrial cancer . Int Symp Marburg , p
26 (Quoted from Trope and Lindahl , 1992) .**

- Borst, M.P., Baker, V.V., Dixon, D., Hatch, K.D., Shingleton, H.M., and Miller, D.M. (1990) :**
Oncogene alterations in endometrial carcinoma.
Gynecol. Oncol. , 38 : 364 .
- Bradford , M. M. (1976) :**
A rapid and sensitive method for quantitation of microgram quantities of protein utilizing the principle of protein -dye binding.
Anal. Biochem., 72 : 248 .
- Brescia, R.J, Barakat, R. A. , Beller, U. , Frederickson, G. and Demopoulos, R.I. (1990) :**
The prognostic significance of nuclear DNA content in malignant epithelial tumors of the ovary .
Cancer , 65 : 141 .
- Brumm, C., Riviere, A., Wilckens, C. and Loning, T. (1990) :**
Immunohistochemical investigation and northern blot of *c-erbB-2* expression in normal , premalignant and malignant tissues of the corpus and cervix uteri.
Virchows Arch. A. Pathol. Anant. Histopathol. , 417(6) : 477.
- Buchberg , A.M. and Viskochil , D. (1990) :**
Deletions and translocations interrupt a cloned gene at the neurofibromatosis type I locus .
Cell , 62 : 187 .
- Bulletti, C., Jasonni, V. M., Tabanelli, S., Balducci, M., Fuschini, G. and Flamigni, C. (1987):**
Danazol reverses endometrial hyperplasia to normal endometrium.
Acta. Eur. Fertil., 18 : 185 .
- Burt, R.W., Ratcliffe, J.G. and Stack, B.H.R. (1978) :**
Serum biological markers in lung cancers.
Br.J. cancer, 37: 714.

- Capon , D.L. , Chen , E.Y. and Ievinson,D. H. (1983) :**
Complete nucleotide sequences of the T24 human bladder carcinoma oncogene and its normal homologue .
Nature , 302 : 33 .
- Caspersson , T. O. (1987) :**
History of the development of cytophotometry from 1935 to the present .
Analyt. Quant. Cytol. Histol. , 9 (1) : 2 .
- Chamlian , D.L. and Taylor , H.B. (1970) :**
Endometrial hyperplasia in young women .
Obstet Gynecol. , 36 : 659.
- Chan,S. and Sikora,K. (1987):**
The potential of oncogene products as tumor markers.
Cancer Surv. , 6 (2) : 185 .
- Chin,J.L., Huben,R.P. and Nava,E. (1985):**
Flow cytometric analysis of DNA content in human bladder tumors and irrigation fluids.
Cancer , 56 : 1677 .
- Christopherson,W.M. (1986) :**
The significance of pathologic findings. In : **Endometrial cancer. Clinincs in Obestetrics and Gynecology.**
Creasman, W.T. (ed.). W.P. Saunders . London, ch.13 ;P. 673 .
- Christopherson, W. M. and Gray, L. A .(1992):**
Premalignant lesions of the endometrium In : **Gynecologic Oncology.** Edited by Malcom Coppleson ,2nd edn. Churchill Livingstone ,London. volume 2 ch. 46, p. 731.
- Christopherson, W.M. (1993):**
Cytologic detection and diagnosis of cancer : Its contributions and Limitations.
Cancer, 51 :1201.

- Cohen , S. , Carpenter , G. and King , L.J. (1980) :**
Epidermal growth factor receptor protein kinase interaction .
J. Biol. Chem. , 225 : 4834 .
- Coleman,R.L., Schink, J.C. , Miller,D.S., Bauer, K. D., August, C. Z., Rademaker,A.W. and Lurain,J.R. (1993) :**
DNA flow cytometric analysis of clinical stage I endometrial carcinoma with lymph node metastases .
Gynecol. Oncol. , 50 : 20 .
- Coligan,J.E., Henkart,P.A. and Todd,C.W. (1973):**
Heterogeneity of carcinoembryonic antigen.
Immunochemistry , 10 : 591.
- Coligan,J.E. and Todd,C.U. (1975):**
Structural studies on carcinoembryonic antigen periodate oxidation.
Biochemistry,14 : 805.
- Coligan,J.E., Egan,M.L. and Guyer,R.L.(1975):**
Structural studies on carcinoembryonic antigen.
Ann. Ny. Acad. Sci. , 259 : 355 .
- Collard,J.G., DeBoer,P.A.J., Janssen, J.W.G., Schijven,J.F. and DeJong,B.(1985):**
Gene mapping by chromosome spot hybridization.
Cytometry , 6 : 179 .
- Constanza,H.E.,Saro,J.D.,Nathanson,L.,Rule,A. and Schwartz,R.S. (1973):**
Carcinoembryonic antigen : Report of screening study.
Cancer, 33 : 583 .
- Coulter,W.H. (1956):**
High speed automatic blood cell counter and cell size analyzer.
Proc. Natl. Electronics Conf. , 12 : 1034 .
- Coussens,L.,Yang-Feng,T.L.,Liao.Y.C.,Chen,E.and Ullrich,A.(1985):**
Tyrosine kinase receptor with extensive homology to EGF receptor shares chromosomal location with neu oncogene.
Science , 230 : 1132 .

- Covone,A.E., Johnson,P.M., Mutton,D., and Adinolf,M.(1984):**
Trophoblast cells in peripheral blood from pregnant women.
Lancet , ii : 841 .
- Cowell,J.K. and Wigley,C.B.(1980):**
Changes in DNA content during in vitro transformation of
mouse salivary gland epithelium.
J. Natl. Cancer Inst. , 64 : 1443 .
- Cowell, J.K. (1982) :**
Double minutes and homogeneously staining regions : gene
amplification in mammalian cells .
Annu. Rev. Genet. , 16 : 21 .
- Creasman, W.T. and Weed , T.C. (1976) :**
Screening techniques in endometrial cancer .
Cancer , 38 : 436 .
- Creasman, W.T.,Soper,J.T.,McCarthy jr, K.S.,Hanshow, W.and
Clarke-Pearson,D.L. (1985) :**
Influence of cytoplasmic steroid receptor content on prognosis of
early stage endometrial carcinoma.
Am. J .Obstet. Gynecol . , 151 : 922 .
- Creasman , W. T. (1991) :**
Estrogen replacement therapy in previously treated cancer
patient is a contra- indication .
Obstet. Gynecol. , 77 : 308 .
- Cutler, B.S. . Forbes ,A.P. Ingersoll, F.M. ans Scully ,R.E. (1972) :**
Endometrial cancer after Stilbosterol therapy in gonadal
dysgenesis .
N. Eng. J. Med. , 287 : 628 .
- Czerniak,B.,Papenhausen,P.R.,Herz,F. and Koss ,L.G. (1985):**
Flow cytometric identification of cancer cells in sffusions with
Cal monoclonal antibody.
Cancer , 55 :2783 .
- Dabbs, D.J., Geisinger, K.R. and Norris, H.T. (1986) :**
Intermediate filament in endometrial and endocervical
carcinoma .
Am.J. Surg.Pathol. , 10 : 568 .

- Dallenbach-Helweg, F.G. (ed.) (1981) :**
In : Histopathology of the endometrium. 3rd edn. Springer,
Verlag ,Berlin, Heidelberg , New York . P . 137 .
- Daunter,B. (1990) :**
Tumor markers in gynecologic oncology.
Gynecol. oncol. , 39 : 1 .
- Dean,P.N., Dolbeare,F., Gratzner,H., Rice,G.C. and Gray,J.W. (1984) :**
Cell cycle analysis using a monoclonal antibody to Brd Urd.
Cell Tissue Kinet. , 17 : 427 .
- Dent, P. B. , McCulloch, P. B., Wesley- James, G., Maclaren, P., Muirhead, W. and Dunnett,C.W. (1978):**
Measurement of CEA in patients with bronchogenic carcinoma.
Cancer , 42 : 1484.
- DeVita , V.T. , Hellmons , S. and Rosenberg , S. (1993) :**
In : cancer , principles and practice of oncology ,
DeVita,V.T. (ed.) 4th edn., Lippincott Co. , Philadelphia , P.
163 .
- Dewhurst ,C.J. , Dekoos, E.B. and Haines,R.M. (1975):**
Replacement hormone therapy in gonadal dysgenesis.
Br. J. Obstet. Gynecol., 82 : 412 .
- DiFiore, P.P., Pierce, J.H., Kraus, M.H., Segatto ,O., King, C.R. and Aaronson,S.A.(1987) :**
erbB-2 is a potent oncogene when overexpressed in NIH/3T3 cells.
Science , 237 : 178.
- Disaia , P.J. and Creasman , W.T. (eds.)(1993) :**
Endometrial hyperplasia and estrogen therapy . In : Clinical
Gynecologic Oncology . Mosby ,St. Louis , Toronto, 4th
edn. P. 122 .

- Dolbeare ,F.A.and Smith,R.E. (1979):**
Flow cytoenzymology: Rapid enzyme analysis of single cells.
In: Flow cytometry and sorting.Melamed,M.R.,Mullaney,P.F.
and Mendelsohn,M.L.(eds.) John Wiley & Sons.New York , P.
317 .
- Doolittle , R.F. , Hunkapiller, M.W. and Hood ,L.E. (1983) :**
Simian sarcoma virus oncogene , *V-sis* ,is derived from the
genes encoding a platelet derived growth factor .
Science , 221 : 275 .
- Dyas,C.H., Simmons,T.K., Ellis,C.N., Austin,J.M. , Partridge,E.E.,
Kilgore,L.C. reading,C., Nelson,K. and Blakemore,W.S.
(1992):**
Effect of deoxyribonucleic acid ploidy status on survival of
patients with carcinoma of the endometrium .
Surg. Gynecol. Obstet. , Feb 174(2) : 133 .
- Dyson,J.E.D.,Joslin,C.A.F.,Quirke,P.,Rothwell,R.I.and
Bird,C.C.(1985):**
Quantitation by flow cytofluorometry of response of tumors of
the uterine cervix to radiotherapy.
Br. J. Radiol. , 58 : 41 .
- Eichner,E. and Abellera,M. (1987) :**
Endometrial hyperplasia treated by progestins.
Obstet. Gynecol., 38 : 739 .
- Eissa , S. And El-Sharkawy , T. (1993) :**
Correlation of *c-erbB-2* oncoprotein and DNA cycle analysis in
colorectal carcinoma .
Egypt. J. Biochem. , 11 (2) : 273 .
- El-Ahmady, O. ,Sakr ,E. , Kandil. O. ,Khalifa,A. , Eissa, S. and El-
Ansary,T. (1989) :**
Predictive value of tumor markers in endometrial hyperplasia
and carcinoma .
J. Tum. Mark. Oncol. , 4 : 1 .

- Elliot, A. (1978) :**
The Immunochemical complexity of CEA: A golden dream or molecular nightmare.
Cancer ,42: 1585.
- El-Toni,N.F., Hassan,S., Yousef,F.,, Labib,N., Ali,N . and Hathout , H. (1985):**
The effect of tamoxifen on endometrial hyperplasia .
J. Kuveit. Med .Assoc. , 19 : 175 .
- Emdin ,S.O., Stenling,R. and Roos,G. (1987):**
Prognostic value of DNA content in colorectal carcinoma .
Cancer ,60 : 1282 .
- Evenson,D.P., Darzynkiewicz,Z. and Melamed,M.R. (1982):**
Simultaneous measurement by flow cytometry of sperm cell viability and mitochondrial membrane potential related to cell motility.
J. Histochem. Cytochem. , 30 : 279 .
- Fahmy,E.,Abaza,H., El-Sawy,M., Moussa,L., Moussa,S.M. and Salim,M. (1990):**
Fibronectin,alpha-fetoprotein and malignant liver disease.
2nd Int. conference of tumor marker oncology, Cairo, Egypt.
- Feichter, G.E. , Hoffken, H. and Heep, J. (1982) :**
DNA flow cytometric measurements on the normal, atrophic, hyperplastic and neoplastic human endometrium. **Virchows Arch. , 398 : 53 .**
- Fenoglio, C.M., Crum,C.P., and Ferenczy,A. (1982):**
Endometrial hyperplasia and carcinoma .Are ultrastructural, biochemical and immunohistochemical studies useful in distinguishing between them ?
Pathol. Res. Pract., 174 : 257.
- Ferguson, A. M. and Fox , H. (1984) :**
The expression of CEA antigen in normal , hyperplastic and neoplastic endom etrium .
Brit. J. Obstet. Gynecol. , 91 : 1042 .

Flanagan, J.G. and Leder, P. (1988) :

Neu proto-oncogene fused to an immunoglobulin heavy chain gene requires immunoglobulin light chain gene for cell surface expression and oncogenic transformation.

Proc. Natl. Acad. Sci. USA , 85 : 8057 .

Fox,H. and Buckley,C.H. (1982):

The endometrial hyperplasia and their relationship to endometrial neoplasia.

Histopath. , 6: 493 .

Freed,G.J.I. and Taylor,G. (1972) :

Carcinoembryonic antigen in feces.

Br. Med. J. , 1 : 85 .

Friedlander,M.L., Hedley, D.W. and Taylor, I.W. (!984) :

Clinical and biological significance of aneuploidy in human tumors .

J. Clin. Pathol. , 37 : 961 .

Fujimoto,S , Kitsukawa ,Y. and Itoh, K. (1979) :

CEA in gasteric juice and feces as an aid in the diagnosis of gastrointestinal cancer.

Ann. Surg. ,189: 34.

Fukushige, S.I. , Matsubara, K.I. , and Yoshida, M. (1986) :

Localization of a novel *v-erbB*-related gene, *c-erbB-2* , on human chromosome 17 and its amplification in a gastric cancer cell line .

Mol. cell Biol. , 6 : 958 .

Gal , D. (1986) :

Hormonal therapy for lesions of the endometrium.

Seminars Oncology , 13 : 33 .

Geisinger, K.R. , Kute, T.E., Marshall, R.B. , Homesley, H.D. and Morgan, T.M. (1986) :

Endometrial adenocarcinoma. A multiparameter clinicopathological analysis including the DNA profile and the sex steroid hormone receptors.

Cancer , 58 : 1518 .

- Gilbert, S. (1991) :**
Oncogenes and cell growth . In :Developmental biology . 3rd
edn. W.P. Saunders. London, P. 736.
- Go,V.L., Ammon,H.V., Holfermuller, K.H.,Krag,E. and Phillips, S.F.
(1975) :**
Quantification of carcinoembryonic antigen-like activities in
normal human gastrointestinal secretion.
Cancer , 36 : 2346 .
- Gold ,P. and Freedman,S.O.(1965):**
Determination of tumor specific antigen in human colonic
carcinomata by immunologic tolerance and absorption
techniques .
J. Exp. Med. ,121 : 329 .
- Gold,P. (1972);**
Tumor specific antigen in GI cancer.
Hosp. Practice , 7: 79 .
- Gompel, C. and Silverberg, S. G. (1985):**
In :Pathology in Gynecology and Obstetrics. Ed.3 , chapter 4 :
The corpus Uteri , pp. 149- 227 . Philadelphia : Lippincott, ..
- Gordon,J. ,Regan,J.W.,Finkle,W.D.and Ziel,H.k.(1977):**
Estrogen and endometrial carcinoma ,an independent pathology
review supporting original risk estimate.
N. Eng. J. Med. , 297 : 570 .
- Gore,H. and Hertig,A.T. (1962):**
Premalignant lesions of the endometrium.
Clin. Obstet. Gynecol., 5 : 1148 .
- Gore and Hertig,A.T. (1966):**
Carcinoma in situ of endometrium.
Am. J. Obstet. Gynecol. , 94 : 134 .
- Graham,F.L. , and Van der Erb ,A.J. (1973) :**
A new technique for the assay for infectivity of human
adenovirus 5 DNA
Virology , 52 : 456.

- Gray, L.A. and Barnes , M.L. (1964) :**
Histogenesis of endometrial carcinoma.
Ann. Surg. , 159 : 976 .
- Greenblatt, R.B. , Gambrell, R.D. and Stoddard, L.D. (1982) :**
The protective role of progesterone in the prevention of
endometrial cancer.
Pathol. Res. Pract. , 174 : 127.
- Gucker,F.T. O'Konski,C.T. and Pickard,H.B. (1947):**
A photoelectric counter for colloidal particles .
J. Am. Chem. Soc. , 69 : 2422 .
- Gullick,W.J.,Berger,M.S.,Bennett,P.L.P.,andWaterfield,M.D.
(1987):**
Expression of *c-erbB-2* protein in normal and transformed cells.
Int. J. Cancer , 40 : 246.
- Gusberg,S.B. (1947):**
Estrogens and adenomatous hyperplasia.
Am.J.Obstet.Gynecol. , 54 : 905 .
- Gusberg, S.B. , Moore ,D.B. and Martin .F. (1954) :**
Precursors of corpus cancer,estrogen and adenomatous hyper-
plasia .
Am. J. Obstet. Gynecol. , 68 : 1472 .
- Gusberg,S.B. and Kaplan,M. (1963) :**
Precursors of corpus cancer.
Am. J. Obstet .Gynecol. , 87: 662 .
- Gustafson,H.,Tribukait,B. and Esposti,P.L. (1982):**
The prognostic value of DNA analysis in primary carcinoma in
situ of the urinary bladder.
Scand. J. Nephrol. ,16 : 141 .
- Gusterson.B.A.,Machin,L.G.,Gullick,W.J.,Gibbs,N.M.,Powles,T.J.,
Elliot,C., Ashlev,S.,Monaghan,P. and Harrison,S. (1988):**
c-erbB-2 expression in benign and malignant breast disease.
Br. J. Cancer , 58 : 453 .

- Guyton , A.G. (1991) :**
Cell production. In : Textbook of medical physiology . 8th edn. ,W.B. Saunders company.London, ch. 7 , P. 31 .
- Haluska , F.G. ,Tsujimoto,Y. and Corce,C.M. (1987):**
Oncogene activation by chromosome translation in human malignancy.
Annu. Rev. Genet. , 21 : 321 .
- Hammarberg,C., Slezak,P. and Tribukait,B. (1984):**
Early detection of malignancy in Ulcerative Colitis : A flow-cytometric DNA study
Cancer , 53 : 291 .
- Hammarstrom, S., Engvall, E., Johansson, B, G., Sevensson, S., Sundblad, G. and Goldstein,I.J. (1975):**
Nature of the tumor-associated determinant(s) of carcino-embryonic antigen.
Proc. Natl. Acad.Sci. USA, 72: 1528 .
- Hanson ,D .J. (1959):**
Studies of the endometrial stroma in cystic glandular hyperplasia.
Am. J. Clin. Pathol. ,32: 1952.
- Hattori,T.,Hosokawa,Y., Fukuda,M. ,Sugihara,H., Hamada, S., Takamatsu, T.and Fujita,S. (1984) :**
Analysis of DNA ploidy in early gastric carcinomas of Japanese.
Cancer , 54 : 1591 .
- Hayes,D.F., Carney,W., Tondini,C., Petit,D., Henderson,C. and Kufe,D.W. (1989):**
Elevated circulating c-neu oncogene product in patients with breast cancer.
Abstract presented at the 12th Annual San Antonio Breast Cancer Symposium (December-8-9, 1989).

- Hedley,D.W., Friedlander ,M . L . , Taylor,I. W. , Rugg, C. A. and Musgrove,E.A.(1983):**
Method for analysis of cellular DNA content of paraffin-embedded pathological material using flow cytometry.
J. Histochem. Cytochem. , 31 : 1333 .
- Hedley,D.W., Friedlander,M.L. and Taylor,I.W. (1985):**
Application of DNA flow cytometry to paraffin-embedded archival material for the study of aneuploidy and its clinical significance.
Cytometry , 6 : 327 .
- Helle,M.,Hellin,H.,Ashorn,P.,Putkinen,E.L.,Krohn,K.,and Wahlsrrm,T. (1992):**
The expression of CEA , CA 19-9 and HMFG antigens in ovarian clear cell and endometriod carcinomas.
Pathol. Res. Pract. , Feb 188 (1-2) : 74 .
- Hendrickson , M.R. and Kempson , R.L. (1980) :**
The differential diagnosis of endometrial adenocarcinoma .
Pathology , 12 : 35 .
- Hertig,A.T., Sommers,S.C. and Bengloff,H. (1949) :**
Genesis of endometrial carcinoma. Carcinoma in situ.
Cancer, 2 : 946 .
- Hiddemann,W., Schumann,J. and Andreeff,M. (1984):**
Convention on nomenclature for DNA cytometry.
Cytometry , 5 : 445 .
- Hill, A.B. (ed.) (1979) :**
In : A short textbook of medical statistics . English Language Book Society , London. p. 86 .
- Holland,F.J. and Emil,F. (1982):**
Carcinoembryonic antigen.
Cancer Medicine , 4 : 15 .

- Holmes, W.E., Sliwkowski, M.X., Akita, R.W., Henzel, W.J., Lee, J.W., Yansura, Abadi, N., Raab, H., Lewis, G.D., Shepard, H.M., Kuang, W.J., Wood, W.I., and Vandlen, R.L. (1992): Identification of Heregulin, a specific activator of p 185 *erbB2*. *Science*, 256 : 1205.
- Holyoke, E. D. , Reynoso ,G. and Chu, T. (1972) : Carcinoembryonic antigen in patients with carcinoma of digestive tract, Proceeding of second conference on embryonic fetal antigens in cancer. National Technical Information Service, USA Department of commerce , Springfield P. 215 .
- Hunter, T. (1984) : The proteins of oncogenes . *Sci. Amer.* , 252 (2) : 60 .
- Hunter, T. and Cooper J. A. (1985) : Protein tyrosine kinase. *Annu. Rev. Biochem.* , 54 : 897 .
- Hustin, J. (1978) : Immunohistochemical demonstration of several tumor markers in neoplastic and preneoplastic states of the uterine mucosa . *Gynecol. Obstet. Invest.* , 9 : 3 .
- Ikeda, M. , Watanabe, Y. , Nanjoh, T. and Noda, K. (1993) : Evaluation of DNA ploidy in endometrial carcinoma . *Gynecol. Oncol.* , 50 : 25 .
- Isaacs, J.T., Wake, N., Coffey, D.S. and Sandberg, A. (1982): Genetic instability coupled to clonal selection as a mechanism for tumor progression in Dunning R-3327 rat prostatic adenocarcinoma system. *Cancer Res.* , 42 : 2353 .
- Iversen ,O.E. and Laerum ,O.D. (1985) : Ploidy disturbances in endometrial and ovarian carcinomas : A review . *Analyt. Quant. Cytol. Histol.* , 7 : 327 .

- Iversen,O.E. (1986) :**
Flow cytometric deoxyribonucleic acid index : a prognostic factor in endometrial carcinoma .
Am. J. Obstet. Gynecol. , 155 : 770 .
- Iversen ,O.E. , Utaaker, E. and Skaarland, E. (1988):**
DNA ploidy and steroid receptors as predictors of disease course in patients with endometrial carcinoma .
Acta. Obstet. Gynecol. Scand. , 67 : 531 .
- Jakobsen,A.,Baek Kristensen,P. and Kvist Poulsen, H. (1983):**
Flow cytometric classification of biopsy specimens from cervical intraepithelial neoplasia.
Cytometry , 4 : 166 .
- Johnson,T.S., Katz,R. and Perhouse,M. (1988) :**
Flow cytometry applications in cytopathology.
Analyt. Quant. Cytol. and Histol.,10 : 423.
- Jones, H. (1988) :**
Endometrial hyperplasia and Endometrial carcinoma (chapter, 28 and 29).In: **Novak's textbook of gynecology . Jones, H.,Wentz, A.C. and Burnett, L.S. (eds.)edn. 11. Williams and Wilkins Baltimore USA P. 716 .**
- Kallioniemi,O.P., Blanco,G.,Alavaikko,M.,and Koivula,T. (1988):**
Improving the prognostic value of DNA flow cytometry in breast cancer by combining DNA index and S-phase fraction .
Cancer , 62 : 2183 .
- Karlan ,B.Y. (1993) :**
The emerging genetics of ovarian cancer : Biologic implications and therapeutic applications of **HER-2/neu** oncogene over-expression .
The second international Cancer Molecular Biology Symposium , Cairo, Egypt P.35 .
- Kashimura ,M. , Saito , T. and Hachisuga ,T. (1988) :**
Cytological finding in endometrial hyperplasia .
Acta. Cytol. , 32 : 335 .

- Katayama,K.T. and Jones,H.W. (1967):**
Chromosomes of hyperplasia and carcinoma of the endometrium.
Am. J. Obstet. Gynecol. , 97 : 978 .
- Katz, R.L. , Tohnson , T.S. and Williamson , K.D. (1985) :**
Comparison of cytologic and acridine-orange flow cytometric detection of malignant cells in human body cavity fluids .
Analyt. Quant. Cytol. Histol. , 7 : 227 .
- Kaufman,R.H., Abbott,J.P. and Wall,J.A.(1959):**
The endometrium before and after wedge resection of the ovaries in the Stein-Leventhal Syndrome.
Am. J. Obstet. Gynecol , 77: 1271 .
- Kaupila,A., Kujansuu,E. and Vikko,R. (1982) :**
Cytosol estrogen and progestin receptors in endometrial carcinoma of patients treated with surgery,radiotherapy and progestin.
Cancer, 50: 2157.
- Kelly ,J.D. , Raines, E.W. and Ross, R. (1985) :**
The B chain of PDGF is sufficient for mitogenesis.
EMBO J. , 3 : 3399 .
- Khalifa , A. (1991) :**
Tumor Markers . In : **Applied Biochemistry and Clinical Pathology . A. Khalifa (ed.) Published by Scientific Book Center , Cairo Egypt.**
- Khoo,S.K. and Mackay, E.V.(1973) :**
Carcinoembryonic antigen in cancer of the female reproductive system: Its detection in whole serum by microradioimmunoassay.
J. Clin. Path. , 26 : 470 .
- King,A.,Seraj,I.M.and Wagner,R.J.(1984):**
Stromal invasion in endometrial carcinoma.
Am.J.Obstet.Gynecol. ,149: 10 .

- King,C.R., Kraus,M.H. and Aaronson,S.A. (1985):**
Amplification of a novel v-erbB-related gene in a human mammary carcinoma.
Science , 229 : 974 .
- Kistner, R.W. (ed.) (1964) :**
Endometrial hyperplasia .In: **Gynecology. Principle and practice**,1st edn.Year Book Publishers, London,p. 278.
- Kistner,R.W. (1970):**
The effects of progestational agent on hyperplasia and carcinoma in situ of the endometrium.
Int. J. Gynecol. obstet. , 84 : 561 .
- Kistner , R.W. (ed.) (1990) :**
Endometrial hyperplasia .In: **Gynecology . Principle and practice . R.W. Kistner (ed.) 5th edn. Year book Publishers, London , P. 228.**
- Kjorstad, K.E.,Welander, C.,Halvorsen, T. ,Grude,T.and Onsrud,M. (1978):**
Progestogens as primary treatment in pre-malignant changes of the endometrium . In : **Endometrial cancer** ,Bush,King and Taylor (eds) . Bailliere Tindall. London , p 188.
- Kleinman,M.S., Turner,M D. and Thayer,W.(1972):**
Serum CEA in patients with inflammatory bowel disease.
Gastroenterology , 62 : 772 .
- Knudson, A.G. (1985) :**
Hereditary cancer oncogenes and anti-oncogenes .
Cancer Res. ,45 : 1437 .
- Korentz,K., Schlag,P. and Moller,P. (1990):**
Expression of epidermal growth factor receptor in normal colorectal mucosa, adenoma and carcinoma.
Virchows Arch. A. Pathol. Anat. , 416 : 343 .

Kouyoumdjian, J., Rymer ,J. , Philippon, C. and Braconnier , F. (1993) :

Determination o the *c-erbB-2 (HER -2-neu)* oncoprotein in normal , benign and malignant tissues of breast origin .

The second International Cancer Molecular Biology Symposium , Cairo, Egypt , P.8 .

Kraus, F.T. (1977) :

Female genitalia . In **Pathology**. Edited by W.A.D. Anderson and John M. Kissane. 7th edn. vol. II .The C.V. Mosby co. St. Louis ,Toronto, P. 1640.

Kraus,M.H.,Popescu,N.C.,Amsbaugh,S.C.and King,C.R. (1987):

Overexpression of the EGF receptor-related proto-oncogene *erbB-2* in human mammary tumor cell lines by different molecular mechanisms.

EMBO . J. , 6 : 605.

Krishan,A. and Ganapathi, R. (1979) :

Laser flow cytometry and cancer chemotherapy : Detection of intracellular anthracycline by FCM .

J. Histochem. Cytochem. , 27 : 1655.

Kurman R.J. and Norris H.J. (1982) :

Evaluatiuon of criteria for distinguishing atypical endometrial hyperplasia from well differentiated carcinoma.

Cancer , 49 : 2547 .

Kurman R.J., Karrinski,P.F. and Norris, H.J. (1985) :

The behavior of endometrial hyperplasia. A long term study of untreated hyperplasia in 170 patients.

Cancer , 56 : 403 .

Kurman R.J. and Norris H.J. (1987) :

Endometrial Carcinoma . In : Blaustein's Pathology of the Female Genital Tract . R.J. Kurman (ed.) ,3rd edn. Springer-Verlag . New York p 324.

Kurman R.J. (1988):

Endometrial hyperplasia and its relation to certain types of carcinoma. In : **Surgey in the treatment of gynecologic cancer** .Proceedings of the international symposium university Press , Antwerp, sep 30- oct 1 p 11 (Quoted from Trope and Lindahl , 1992) .

Kute,T.E., Linville,C. and Barrows,G.(1983):

Cytofluorometric analysis for estrogen probes.
Cytometry , 4 : 132 .

Kute ,T.E. , Gregory,B. , Galleshaw, J., Hopkins, M. and Buss, D. (1988):

How reproducible are flow cytometry data from paraffin embedded blocks ?

Cytometry , 9 : 494 .

Kynast,B.,Binder,L.,Marx,D.;Zoll,B.,Schmoll,H.J.,Oellerich,M. and Schauer,A. (1993):

Determination of a fragment of the *c-erbB-2* translational product p185 in serum of breast cancer patients .

J. Cancer Research Clinical Oncology , 119 : 249.

Kysela,B., Siracky,J., Redecha,M. and Bardos,A. (1990):

Flow cytometry (FCM) analysis of endometrial hyperplasia and carcinoma.

Neoplasma. , 37 (5) : 489 .

Lacal , J.C. , Anderson , P.S. and Aronson ,S.A. (1986) :

ras-p21 protein with high or low GTPase activity can effeciently transform NIH/3T3 cells .

Cell , 44 : 609 .

Landy, A., Ohlsson-Wilhelm,B. and Giorgi,J.V. (1990):

Application of flow cytometry to the study of HIV infection.

Aids , 4 : 479 .

Leiserowitz, G. S., Harris,S.A., Subramaniam, M., Keeney, G.L., Podratz, K. C., and Spelsberg, T.C. (1993):

The protooncogene *c-fms* is overexpressed in endometrial cancer.

Gynecol. Oncol. , 49 : 190 .

Libermann, T. A. , Nusbaum, H. R. and Razon, H. (1985):

Amplification, enhanced expression and possible rearrangement of EGF receptor gene in primary human brain tumors of glial origin .

Nature , 313 : 144 .

Lindahl,B., Alm,P., Ferno,M., Killander,D., Langstrom,E., Norgen, A. and Trope,C. (1987):

Prognostic value of flow cytometrical DNA measurement in stage I -II endometrial carcinoma . Correlations with steroid receptor concentration, tumor myometrial invasion and degree of differentiation .

Anticancer Res. , 7 : 781.

Lindahl,B. , Alm,P., Ferno,M. and Norgren ,A. (1988):

Endometrial hyperplasia. II : Endocrine factors.

Cancer Res. , 48 : 1236 .

Lindahl, B., Alm, P., Ferno, M. and Norgen, A. (1990) :

Endometrial hyperplasia : A prospective randomized study for histopathology tissue steroid receptors and plasma steroids after curettage .

Anticancer Res. ,10 : 725.

Lindahl, B. and Alm , P.(1991) :

Flow cytometrical DNA measurements on endometrial hyperplasias. A prospective follow up study after curettage only or addition of high dose gestagen treatment

Anticancer Res. , 11 : 391 .

Linial , M. , Gunderson ,N. and Groudine ,M. (1985) :

Enhanced transcription of *c-myc* in bursal lymphoma cells requires continuous protein synthesis .

Science , 32 : 1126 .

LoGerfo,P., Krupey, J. and Hansen, H. (1971) :

Demonstration of an antigen common to several varieties of neoplasia.

N. Engl. J. Med. , 285 : 138.

Loken,M.R. and Stall,A.M. (1982) :

Flow cytometry as an analytical and preparative tool in immunology .

J. Immunol. Methods , 50 : 85 .

Look,A.T.,Hayes,F.A., Nitschke,R.,McWilliams,N.B. and Green,A.A. (1984):

Cellular DNA content as a predictor of response to chemotherapy in infants with unresectable neuroblastoma .

N. Engl. J. Med. , 311 : 231 .

Lovett , E.J. , Schnitzer , B. and Keren , D. (1984):

Application of flow cytometry to diagnostic pathology .

Lab. Invest. , 50 : 115 .

Lowenstein,M.S. and Zamcheck,N. (1978):

Carcinoembryonic antigen (CEA) levels in benign gastrointestinal disease states.

Cancer , 42 : 1412.

Lupu,R., Colomer,R., Zugmaier,G., Sarup,J., Shepard,M., Slamon,D and Lippman,M.E.(1990):

Direct interaction of a ligand for the *erbB2* oncogene product with the EGF receptor and p185*erbB2*.

Science , 249 : 1552.

Lurie ,B.B., Lowenstein,M.S. and Zamcheck,N.(1975) :

Elevated circulating CEA levels in benign extrahepatic biliary tract obstruction and inflammation .

JAMA , 233 : 326 .

MacDonland , P.C. and Sittrei, P. K. (1974):

Relationship between extraglandular production of estrone and the occurrence of endometrial neoplasia .

Gynecol. Oncol. , 2 : 2159.

Maguire, H.C. and Greene , M.I. (1989) :

The *neu (c-erbB-2)* Oncogene.

Seminars in Oncology , 16 : 148 .

- Maguire,H.C., Jaworsky,C., Cohen,J.A.,Hellman,A.B.,Weiner,D.B. and Greene,M.I. (1990):**
Distribution of neu (*c-erbB-2*) protein in human skin .
J. Invest. Dermatol. , 89 : 786 .
- Mansell,H. and Hertig,A.T. (1955):**
Granulosa-Theca cell tumors and endometrial carcinoma. A study of their relationship and a survey of 80 cases.
Obstet. Gynecol. , 6 : 385.
- Martin ,G.S. (1970):**
Rous sarcoma virus, a function required for the maintenance of the transformed state .
Nature , 227 : 1021 .
- Martin, F. and Martin,M.S. (1972):**
Radioimmunoassay of carcinoembryonic antigen in extracts of human colon and stomach.
Int. J. Can. , 9 : 641.
- Martin,E.M.,James,K.K.,Lurtubise,P.E.,Cataland,P. and Minton,J.P. (1977):**
The use of CEA as an early indicator for gastrointestinal tumor recurrence and second-look procedure.
Cancer , 39 : 440.
- Martin-Zanca, D. , Hughes ,S.H. and Brabacid ,M. (1986) :**
A human oncogene formed by the fusion of truncated tropomyosine and protein tyrosine kinase sequence .
Nature , 319 : 743 .
- Mauro, F. , Teodori , L. Schumann, J. and Gohde,W. (1986) :**
Flow cytometry as a tool for the prognostic assessment of human neoplasia .
Int. J. Radiat. Oncol. Biol. Phys. , 12 : 625 .
- Mayers,R.E., Sutherland,D.J.,Kellen,J.A. and Malkin,A.(1978):**
Carcinoembryonic antigen in breast cancer .
Cancer , 42 : 1520.

- McBride, J.M. (1959):**
Premenopausal cystic hyperplasia and endometrial carcinoma.
J. Obst. Gynecol. Br. Commonw. , 66 : 288.
- McCartney, J.C. and Camplejohn, R.S. (1986):**
DNA flow cytometry of non-Hodgkin's lymphomas.
J. Pathol. , 148 : 122 A .
- McGuire, W.L. and Dressler, L.G. (1985):**
Emerging impact of FCM in predicting recurrence and survival
in breast cancer patients.
J. Natl. Cancer Inst. , 75 : 405 .
- McKenzie, S. J., Marks , P.J., Lam, T., Morgan, J., Panicali,
D.L., Trimpe, K.L. and Carney, W.P. (1989):**
Generation and characterization of monoclonal antibodies
specific for the human neu oncogene product, p185.
Oncogene , 4 : 543 .
- Meeker, W.R. and Kashniri, R.L. (1973):**
Clinical evaluation of carcinoembryonic antigen test.
Arch. Surg. , 107 : 266.
- Melamed, M.R. and Klein, F.A. (1984):**
Flow cytometry of urinary bladder irrigation specimens.
Hum. Pathol. , 15 : 302 .
- Mencaglia , L. , Valle , R.F. , Perino , A. and Gilardi , G. (1990) :**
Endometrial carcinoma and its precursors : early detection and
treatment
Int. J. Gynecol. Obstet. , 31 : 107 .
- Moberger, B., Auer, G., Forsslund, G. and Moberger, G. (1984):**
The prognostic significance of DNA measurements in
endometrial carcinoma .
Cytometry , 5 : 430 .
- Moberger, B. , Sennerstam , R. , and Auer , G. (1989) :**
DNA and nuclear protein characteristics in non-neoplastic and
neoplastic endometrial tissue .
Anal. Cell Pathol. , 2 (1) : 15 .

- Moldavn,A. (1934) :**
Photoelectric technique for the counting of microscopial cells.
Science , 80 : 188 .
- Molnar,I.G.,Vandevoerde,J.P. and Gitanick,G.L. (1976) :**
CEA levels in fluids bathing gastrointestinal tumors.
Gastroenterology , 70 : 513.
- Moore,T.,Dhar,P.,Zamcheck,N.L.,Keely,A.,Gottlieb,L. and Kupchik,H.Z. (1971):**
Carcinoembryonic antigen(s) in liver disease.Clinical and morphological studies.
Gastroenterology , 63 : 88 .
- Moore,T.,Dhar,P.,Zamcheck,N.L. and Kupchik,H.Z. (1972) :**
CEA in benign conditions.
Gastroenterology , 73 : 120.
- Morse , A. R. and Curran , G. J. (1985) :**
Distribution of epithelial membrane antigen in normal and abnormal endometrial tissue .
Brit. J. Obstet. Gynecol. , 92 : 1286 .
- Muhonen,T.,Pyrhonen,S.,Laasonen,A., and Asko-Seljavaara, S. (1991) :**
DNA aneuploidy and low S-phase fraction as favourable prognostic signs in metastatic melanoma.
Br. J. Cancer , 64 : 749 .
- Murray , R. K. (1993) :**
Cancer , oncogenes and growth factors. In :
Harper'Biochemistry.Murray, R.K. , Granner, D.K. , Mayes, P.A. and Rodwell,V.W. (eds.), 23rd edn.
Appleton & lange .California USA. Ch. 62 P. 710.
- Nakopoulou, L., Minaretzis, D.,Tsionou,C. H.and Mastrominas, M. (1989):**
Value of immunohistochemical demonstration of several epithelial markers in hyperplastic and neoplastic endometrium.
Gynecol. Oncol. , 37 : 346 .

- Naus ,G.J. , DeVere, R. ,Richart, R.M. and Deitch ,A.D. (1985) :**
Predictive value of flow cytometric analysis of paraffin embdded tissue in endometrial carcinoma.
Lab. Invest. , 52 : 48 .
- Neiman , P. (1987) :**
Oncogenes and neoplastic diseases . In : **Harrison's principles of internal medicine** . 11th edn. , vol. 1 chapter 59 Int. edn. , PP. 310 - 317 .
- Newbury,R. ,Schuerch,C. ,Fanning,J. ,Glidewell,O. and Evans,M. (1990):**
DNA content as a prognostic factor in endometrial carcinoma.
Obstet. Gynecol. , 76 : 251.
- Norris,H.J.,Tavasoli,F.A. and Kurman,R.J. (1983) :**
Endometrial hyperplasia and carcinoma. Diagnostic considerations .
Am. J. Surg. Pathol. , 7 : 839 .
- Novak,E.R. and Yui,E. (1936) :**
Relation of hyperplasia to adenocarcinoma of uterus.
Am. J. Obstet. Gynecol. , 32 : 674.
- Novak, E.R. and Rutledge , F. (1948) :**
Atypical endometrial hyperplasia simulating adenocarcinoma.
Am. J. Obstet. Gynecol. , 55 : 46 .
- Novak,E.R. and Woodruff, E. (eds.) (1979) :**
Histology of the endometrium, . In : **Novak's Gynecologic and Obstetric Pathology with endocrinal relations** . 8th ed. W.B. Saunders. Philadelphia. ch. 7, P. 347.
- Nowell ,P.C. and Hungerford , D.A. (1960) :**
A minute chromosome in human chronic granulocytic leukemia.
Science , 132 : 1497 .
- Nowell , P.C. (1976) :**
The clonal evolution of tumor cell populations.
Science , 194 : 23 .

Nugent, F.W. and Hansen, E.R. (1971) :

Radioimmunoassay of CEA as a diagnostic test for cancer of colon.

Lahey Clin. Found. Bull. , 20 : 85.

Nugent, A., McDermott, E., Duffy, K., O'Higgins, N., Fennelly, J.J. and Duffy, M.J. (1992) :

Enzyme-linked immunosorbent assay of *c-erbB-2* oncoprotein in breast cancer.

Clinical Chemistry , 38 : 1471.

Oram , D. (1990) :

The management of cancer of the uterine corpus . In : **Clinical Gynecological Oncology** J.H. Shepherd and J.M. Monaghan (eds.) , 2nd edn. Blackwell. London PP. 115 - 137 .

Oud, P.S., Henelerik, J.B.J. and Beck, L.M. (1985) :

Flow cytometric analysis and sorting of human endometrial cells after immunocytochemical labelling for cytokeratin using a monoclonal antibody.

Cytometry , 6 : 159 .

Padhy, L.C., Shih, C., Cowing, D., Finkelstein , R. and Weinberg, R.A. (1982):

Identification of a phosphoprotein specifically induced by the transforming DNA of rat neuroblastomas.

Cell , 28 : 865 .

Page , D.L. (1991) :

Atypical hyperplasia , narrowly and broadly defined .

Human Path. , 22 : 631 .

Payne , F.L. (1936) :

The clinical significance of endometrial hyperplasia.

Am. J. Obstet. Gynecol. , 34 : 762 .

Penman , S. and Williams , J.G. (1975) :

Messenger RNA sequences in growing and resting mouse fibroblasts.

Cell , 6 : 197 .

Perkins, A .S. and VandeWoude, G.F. (1993) :

Principles of molecular cell biology ; Oncogenes . In : **Cancer , principles and practice of oncology**. DeVita V. T. (ed.) 4th edn. Lippincott Co.Philadelphia , chapter 3 ,P. 35 .

Plunkett, E.R. and Wolfe , B.M. (1992) :

Prolonged effect of a novel ,low dosage continuous progestin cyclic estrogen replacement program in postmenopausal women .

Am. J. Obstet. Gynecol. , 166 :117 .

Plow, E. F. and Edgington , T.S. (1975) :

Isolation and characterization of a homogenous isometric species of CEA.

Int. J. Cancer , 15 : 748 .

Press , M. (1993) :

HER/2neu gene amplification and expression in human carcinomas : Implications for clinical management.

The second International Cancer Molecular Biology Symposium , Cairo , Egypt ,P.96 .

Primus , F.J., Goldenberg, D.M. and Hansen ,H.J. (1973) :

Metabolism of carcinoembryonic antigen in human tumor hamster host model .

Fed. Proc. , 32 : 834 .

Pritchord, D. and Todd, C. (1979) :

Purification of CEA by removal of contaminating mucopolysaccharides.

Cancer Res. , 36 : 4699 .

Quillamor, R.M., Furlong ,J.W. , Hoschner, J.A. and Wynn, R.M. (1988) :

Relative prognostic significance of DNA flow cytometry and histologic grading in endometrial carcinoma .

Gynecol. Obstet. Invest. , 26 : 332 .

Quirke,P. and Dyson, J.E.D. (1986) :

Flow cytometry : Methodology and Applications in pathology .

J. Pathol. ,149 : 79 .

Qurike, P., Fozard, J.R.J., Dixon, M.F., Dyson, J.E.D., Gites, J.R. and Bird, C. C. (1986) :
DNA aneuploidy in colorectal adenomas .
Br. J. Cancer , 53 :: 477 .

Ramzy, I. (ed.) (1990) :
Endometrial hyperplasia . In : **Essential of gynecology and obstetrics pathology**. 2nd edn. Appleton Century Crofts ,
Sydney . Canada .London. P 153.

Ronstrom, L. , Tribukait,B. and Esposti,P.L.(1981) :
DNA pattern and cytological findings in fine needle aspirates of
untreated prostatic tumors.
Prostate , 2 : 79 .

Rosenwaks,Z . , Wentz, A.C. , Urban , M.D. and Jones, G.S. (1979) :
Endometrial pathology and estrogens .
Obstet. Gynecol. , 53 : 403 .

Rous, P. (1911) :
A sarcoma of fowl transmissible by an agent separable from
tumor cells.
J. Exp. Med. , 12 : 397 .

Rowley, J.D. (1973) :
A new consistent chromosomal abnormality in chronic
myelogenous leukemia identified by quinacrine fluorescence
and Giemsa staining .
Nature , 243 : 290 .

**Rubin,S.C., Finstad,C.L., Wong,G.Y., Plante,M., and Lloyed,K.O.
(1993):**
Prognostic significance of **HER-2/neu** expression in advanced
epithelial ovarian cancer : A multivariate analysis .
Am. J. Obstet. Gynecol. , 168 : 162 .

Sakamoto, C. (1984) :
Sonographic criteria of phasic changes in human endometrial
tissues .
Int. J. Gynecol. Obstet . , 23 : 7.

- Sargent,E.R., Gomella,L.G.,Belledgrun,A. ,Linehan,W.M. and Kassid,A. (1989):**
Epidermal growth factor gene expression in normal human kidney and renal cell carcinoma .
J. Urol. , 142 : 1364 .
- Sasano ,H. and Garrett , C.T. (1992) :**
Oncogenes in gynecological tumors .
Curr. Top. Pathol. , 85 : 357 .
- Sato,S., Ito,K., Ozawa,N., Yajima,A. and Sasano,H. (1991) :**
Expression of c-myc, epidermal growth factor receptor and *c-erbB-2* in human endometrial carcinoma and cervical adenocarcinoma.
J. Exp. Med. ,165(2) : 137.
- Scarselli, G., Tantini, C. ,Colafrancesca, M., Taddei, G.L. ,Bargelli, G., Venturini, N. and Branconi ,F. (1988) :**
Levo-Norgesterel Nova T and precancerous lesions of the endometrium .
Eur. J. Gynecol. Oncol. , 9 : 284 .
- Schechter,A.L.,Hung,M.C., Vaidyanathan,L., Weinberg,R.A., Yang-Feng,T.L.Francke,U.,Ullrich,A. and Coussens,L. (1985) :**
An erbB-homologus gene distinct from and unlinked to the gene encoding the EGF receptor.
Science , 229 : 976 .
- Schiff, I. , Sela, H.K. , Cramer, D., Tulchinsky, D. and Ryan, K. (1982) :**
Endometrial hyperplasia in women on cyclic or continuous estrogen regimen .
Fertil. Steril. , 37 : 79 .
- Schubert,D. , Heinemann, S.and Carlisle ,W. (1974):**
Clonal cell lines from the rat central nervous system.
Nature , 249 : 224 .

- Schwab,M. ,Alitalo,K. , Klempnauer, K.B. ,Varmus,H.E. , Bishop, J.M. and Gelbent ,F. (1983) :**
Amplified DNA with limited homology to myc cellular oncogene is shared by human neuroblastoma cell lines and a neuroblastoma tumor .
Nature , 305 : 245 .
- Seeger ,R.C. , Brodeur ,G.M. and Sather , H. (1985) :**
Association of multiple copies of the N-myc oncogene with rapid progression of neuroblastoma tumor .
N. Engl. J. Med . , 313 : 1111 .
- Semba, K. , Kamata ,N. and Toyoshima, K. (1985) :**
A *v-erbB*-related proto-oncogene, *c-erbB-2* , is distinct from the *c-erbB-1* EGFR gene and is amplified in a human salivary gland adenocarcinoma .
Proc. Natl. Acad. Sci. USA , 82 : 6497 .
- Shapiro,J.R. and Shapiro,W.R. (1985) :**
The subpopulations and isolated cell types of freshly resected high grade human gliomas
Cancer and Metastasis Reviews , 4 : 107 .
- Shapiro, H.M. (1991) :**
Flow cytometry : Past , present and future . In : **Diagnostic flow cytometry .** Coon J.S., and Weinstein R.S. (eds.) Williams & Wilkins.Balltimore USA , P.1 .
- Sherman, A.I. (1978) :**
Precursors of endometrial cancer .
Israel J. Med. Science , 14 : 370 .
- Shih, C. , Shilo ,B. Z., and Goldfarb, M.P. (1979) :**
Passage of phenotypes of chemically transformed cells via transfection of DNA and chromatin.
Proc. Nat .Acad. Sci. USA , 76 : 5714 .
- Shih, C., Padhy, L.C. and Murray, M. (1981) :**
Transforming genes of carcinomas and neuroblastomas introduced into mouse fibroblasts .
Nature , 290 : 261 .

Shih , C. and Weinberg ,R.A. (1982) :

Isolation of a transforming sequence from a human bladder carcinoma cell line.

Cell , 29 : 161 .

Shuster, J. , Silverman, M. and Gold , P. (1973) :

Metabolism of human CEA in Xenogenic animals .

Cancer Res . , 33 : 65 .

Slamon,D.J.,Clark,G.M.,Wong,S.G.,Levin,W.J., and McGuire,W.L. (1987):

Human breast cancer: Correlation of relapse and survival with amplification of the HER-2/neu oncogene.

Science , 235 : 177 .

Slamon, D.J., Gogolphin, W., Jones, L.A.,Levin, W.J., Stuart, S.G., Udove, J., Ullrich, and Press,M.F. (1989) :

Studies of the HER-2/neu proto-oncogene in human breast and ovarian cancer.

Science , 244 : 707 .

Smith , J.J. and Schulman ,H. (1985) :

Current dilatation and curettage practice : A need for revision .

Obstet . Gynecol. , 65 : 516 .

Soszka, T. and Olszewski, K. (1986) :

Plasminogen activators and their inhibitors in normal, hyperplastic and carcinomatous human endometrium .

Thromb. Res. , 42 : 835 .

Stanley , M.A. (1969) :

Chromosome constitution of human endometrium .

Am. J. Obstet. Gynecol. , 104 : 99 .

Stehelin,D. ,Varmus,H.E. , Bishop,J.M. and Vogt,P.K. (1976) :

DNA related to the transforming genes of avian sarcoma viruses is present in normal avian DNA .

Nature , 260 : 170 .

- Stendahl,U., Wagenius,G., Strang,P. and Tribukait,B. (1988):**
Flow cytometry in invasive endometrial carcinoma .
Correlations between DNA content S-phase rate and clinical parameters.
In Vivo , Mar-Apr 2(2) : 123 .
- Stern,D.F. and Kamps, M.P. (1988):**
EGF-stimulated tyrosine phosphorylation of p185neu: a potential model for receptor interactions.
EMBO J. , 7 : 995.
- Stone,K.R.,Craig,R.B.,Palmer,J.O.,Rivkin,S.E. and McDivitt, R. W. (1985) :**
Short-term cryopreservation of human breast carcinoma cells for flow cytometry.
Cytometry , 6 : 357 .
- Stonehill, E.H. and Bendich , A. (1970) :**
Retrogenetic expression . The appearance of embryoneal antigens in cancer cell.
Nature , 228 : 370 .
- Susini,T., Rapi, S. , Savino,L., Berti,P. and Massi, G. (1994) :**
Prognostic value of flow cytometric deoxyribonucleic acid index in endometrial carcinoma: comparison with other clinical -pathological parameters.
Am. J. Obstet. Gynecol. , 170 (2) : 527 .
- Svenberg, T. (1976) :**
Carcinoembryonic antigen-like substances of human bile .
Int. J. Cancer , 17 : 588 .
- Symonds , D.A. (1990) :**
Prognostic value of pathologic features and DNA analysis in endometrial carcinoma .
Gynecol. Oncol. , 39 : 272 .
- Taketani,Y. and Mizuno, M. (1988) :**
Cyclic changes in epidermal growth factor receptor in human endometrium during menstrual cycle.
Endocrinol. Jpn. , 35 : 19 .

- Tal,M., Wetzler,M., Josefberg,Z., Deutch,A., Assaf,D., Kris,R., Shiloh,Y., Givol,D., and Schlessinger,J. (1988):**
Sporadic amplification of the HER-2/neu protooncogene in adenocarcinomas of various tissues.
Cancer Res. , 48 : 1517 .
- Tandon,A.K.,Clark,G.m.,Chamness,G.C.andMcGuire,W.L.(1989):**
HER-2/neu oncogene protein and prognosis in breast cancer.
J.Clin. Oncol. , 8 : 1120.
- Tavassoli , F. and Kraus , F.T. (1978) :**
Endometrial lesions in uteri resected for atypical endometrial hyperplasia .
Am. J. Clin. Path. , 70 : 770 .
- Taylor , H.C. (1932) :**
Endometrial hyperplasia and carcinoma of body of uterus.
Am. J. Obstet. Gynecol. , 23 : 309 .
- Taylor, I.W. (1980) :**
A rapid single step staining technique for DNAanalysis by flow microfluorimetry.
J. Histochem. Cytochem. , 28 : 1021 .
- Terakawa, N. , Inove ,M. , Shimuzu, I. , Ikegami, H. , Mizutani, T., Sakata, M., Tanizawa,Q. and Matsemoto ,K. (1988) :**
Preliminary report on the use of *Danazol* in the treatment of endometrial adenomatous hyperplasia .
Cancer , 62 : 2618 .
- Thomas,P. and Hems, D.A. (1975) :**
The hepatic clearance of circulating native and sialo CEA by the rat .
Biochem. Biophys. Res. Comm. , 67 : 1205 .
- Thornton, J.G., Quirke, P. and Wells, M. (1989) :**
Flow cytometry of normal , hyperplastic and malignant human endometrium .
Am. J. Obstet. Gynecol. , 161 : 487 .

- Toppozada , M. K. , Ismail , A. A. , Hamed , R. S. , Ahmed , K. S. and El-Faras, A. (1988) :**
Progesterone challenge test and estrogen assays in menopausal women with endometrial adenomatous hyperplasia .
Int. J. Gynecol. Obstet. , 26 : 115 .
- Trent ,J.M. and Davis, J.R. (1979) :**
D-group chromosome abnormalities in endometrial cancer and hyperplasia.
Lancet , ii : 361 .
- Tribukait,B., Gustason,H. and Esposti,P.L.(1982):**
The clinical significance of ploidy and proliferation in the clinical and biological evaluation of bladder tumors.
Br. J. Urol. , 54 : 130.
- Trope, C. and Lindahl, B. (1992) :**
Premalignant lesions of the endometrium . **In : Gynecologic Oncology .Malcolm Coppleson (ed.) 2nd edn .Churchill Livingstone , London, Vol.2 ch. 47, P 747 .**
- Tsionou, C.H., Minaretzis, D., Papageorgiou, I.,Nakopoulou, L., Michalas, S. and Aravantinos,D. (1991):**
Expression of carcinoembryonic antigen and ferritin in normal , hyperplastic and neoplastic endometrium.
Gynecol. Oncol. , 41 (3) : 193 .
- Tsou , K. , Hong, D. and Varello ,M.A. (1985) :**
Flow cytometric DNA and 5'-nucleotide phosphodiesterase in endometrium .
Cancer , 56 : 2340 .
- Tsuda, H., Hirohashi, S. , Yamamoto , H., Yokota, J.,Yoshida, T., Sakamoto,H., Terada,M.and Sugimura,T.(1989):**
Correlation between long-term survival in breast cancer patients and amplification of two putative oncogene coamplification units:hst-1/int-2 and *c-erbB-2*/ear-1.
Cancer Res. ,49 : 3104.

- Valet,G.,Ormerod,M.G.,Warnecke,H.H.,Benker,G.and
Ruhenstroth - Bauer,G. (1981) :
Sensitive three-parameter flow cytometric detection of
abnormal cells in human cervical cancers : a pilot study .
J. Cancer Res. in Clin. Oncol. , 102 : 177 .
- Valet,G., Raffael,A. and Russmann,L. (1985) :
Determination of intracellular calcium in vital cells by FCM.
Naturwissenschaften , 72 : 600 (Quoted from Quirke and
Dyson , 1986).
- Van Dam,P.A., Lowe,D.C., Shepherd,J.H., Vergote,I.B. and
Watson,J.V. (1990):
Multiparameter flow cytometric quantification of membrane
proteins in long-term cryopreserved solid tumors.
Lancet , ii : 689 .
- Van Dam,P.A., Lowe,D.G., Watson,J.V., James,M., Chard,T.,
Hudson,C.N. and Shepherd,J.H. (1991) :
Multiparameter flow cytometric quantitation of epidermal
growth factor receptor and *c-erbB-2* oncoprotein in normal and
neoplastic tissues of the female genital tract .
Gynecol. Oncol. , 42 : 256 .
- Van de Vijver, M. J., Berstlarr, R. and Devilee, P. (1987) :
Amplification of the neu (*c-erbB-2*) oncogene in human
mammary tumors is relatively frequent and is often
accompanied by amplification of the linked c-erbA oncogene.
Mol. Cell Biol. , 7 : 2019.
- Van de Vijver,M.J., Peterse ,J.L., Wisman,P.,Lomans,J.,Dalesio,O.
and Nusse,R. (1988):
Neu-protein overexpression in breast cancer.
N. Engl. J. Med. , 319 : 1239 .
- Van den Ingh,H.F., Griffioen,G. and Cornelisse,C.J. (1985) :
Flow cytometric detection of aneuploidy in colorectal
adenomas.
Cancer Res. , 45 : 3329 .

Van Kleist , S. and Butn , P. (1969) :

Cellular localization of an embryonic antigen in human colonic tumors .

Int. J. Can. , 4 : 874 .

Vayliang, G.O. and Zamcheck, N. (1982) :

The role of tumor markers in the management of colorectal cancer .

Cancer , 50 : 2618 .

Vellios, F. (1972) :

Endometrial hyperplasia ; precursors of endometrial carcinoma.

Path. Ann. , 7 : 201 .

Vellios , F. (1974) :

Endometrial hyperplasia and carcinoma in situ .

Gynecol. Oncol. , 2 : 152 .

Venter,D,J., Tuzi,N.L., Kumar,S. and Gullick,W.J.(1987) :

Overexpression of *c-erbB-2* oncoprotein in human breast carcinomas: Immunohistological assessment correlates with gene amplification .

The Lancet , 2 : 69.

Visser,J.W.M., Jongeling,A.A.M. and Tanke,H.J. (1979):

Intracellular pH determination by fluorescence measurements.

J. Histochem. Cytochem. , 27 : 32 .

Wagner ,D. , Richart , R.M. and Turner , J. Y. (1967) :

Deoxyribonucleic acid content of presumed precursors of endometrial carcinoma.

Cancer , 20 : 2067 .

Walle, A.J. and Niedermayer, W. (1985) :

Aneuploidy as a marker of minimal residual disease in leukemia.

Cancer Detection and Prevention , 8 : 303 .

- Wang,D.P., Fujii,S., Konishi,I., Nanbu,Y., Iwai,T., Nonogaki,H. and Mori,T.(1992):**
Expression of *c-erbB-2* protein and epidermal growth factor receptor in normal tissues of the female genital tract in the placenta.
Virchows. Arch. A. Pathol. Anat. Histopathol. , 420 (5) : 385 .
- Wantzin,G.L., Larsen,J.K., Christensen,I.J., Ralfkiaer,E. and Thomsen,K. (1984):**
Early diagnosis of cutaneous Y-cell lymphoma by DNA flow cytometry on skin biopsies.
Cancer , 54 : 1348 .
- Watson , J.V. (1986) :**
Oncogenes, cancer and analytical cytology .
Cytometry , 7 : 400 .
- Watson , J.D. (ed.) (1992) :**
Oncogenes and anti-oncogenes . In : **Recombinant DNA , 2nd**
W.B. Saunders company, Philadelphia , chapter 3 p. 35 .
- Waymouth,C.(1982) :**
Methods for obtaining cells in suspension from animal tissues.
In : Cell separation.Methods and selected applications,
Pretlow,T.G., Pretlow,T.P. (ed.) vol. 1. New York : Academic
press, P. 1 .
- Weaver , R.S. (ed.)(1989) :**
Genes and cancer . In: **Genetics, W.N. Brown publisher .**
2nd edn. chapter 17 p. 995 .
- Weinberg , R.A. (1989) :**
Oncogenes, antioncogenes and the molecular bases of multistep carcinogenesis .
Cancer Res. , 49 : 3713 .
- Weinstein,I.B. (1989) :**
The origins of human cancer : Molecular mechanisms of carcinogenesis and their implication for cancer prevention and treatment.
Cancer Res . , 48 : 4135 .

- Welch, W. R. and Scully, R. E. (1977) :**
Precancerous lesions of the endometrium .
Hum. Pathol. , 8 : 503 .
- Wentz , W.B. (1964) :**
Effect of a progestational agent on endometrial hyperplasia and
endometrial carcinoma .
Obstet. Gynecol. , 24 : 370 .
- Wentz , W.B. (1974) :**
Progestin therapy in endometrial hyperplasia.
Gynecol. Oncol. , 2 : 362 .
- Wentz , W.B. (1985) :**
Progestin therapy in lesions of the endometrium .
Seminar Oncology , 12 : 23 .
- Wheless, L.L. (1991) :**
Flow instrumentation and data analysis. **In : Diagnostic Flow
Cytometry** ,edited by Coon J.S. and Weinstein R.S.;**Williams
and Wilkins; Batlimore USA .P. 17 .**
- White , L.N. (1993) :**
An overview of screening and early detection of gynecologic
malignancies .
Cancer , 71 : 1400 .
- Whitefield , C. R. (ed.)(1988) :**
**In : DewHurst's Textbook of Obestetrics and Gynecology
For Postgraduate . 4th edn. (ELBS edn.) Blackwell
Scientific Publications . Oxford, London . P. 635.**
- Wied,G.L.,Messina,A.M.,Manglano,J.I.,Meier,P.and Blough,R.R.
(1964):**
Fluorospectrophotometric analysis of endometrial and
endocervical epithelial cells.
Acta. Cytol. , 8 : 408 .
- Wiernik,P. H. and Serpick, A.A. (1973) :**
Plasma CEA in patients with advanced cancer .
Cancer Res. , 14 : 14 .

- Wijkstrom, H., Gustafson, H. and Tribukait, B. (1984) :**
DNA analysis in evaluation of transitional cell carcinoma before cystectomy.
J. Urol. , 132 : 894 .
- Williams , P.L. and Warwick , R. (eds.) (1980) :**
Cytology .In:Gray's anatomy. 36th edn. Churchill Livingstone,London,P.23 .
- Wolley,R.C., Hery,F. and Koss,L.G. (1982) :**
Caution on the use of lymphocytes as standards in the flow cytometric analysis of cultured cells .
Cytometry , 2 : 370 .
- Wu,J.T., Astill, M.E. and Zhang,P. (1993) :**
Detection of the extracellular domain of *c-erbB-2* oncoprotein in sera from patients with various carcinomas : correlation with tumor markers.
J. Clinical Laboratory Analysis , 7 : 31 .
- Yamamoto, T., kawa, S., Akiyama, T., Semba, K., Nomura, N., Miyajima, N., Saito, T. and Toyoshima,K. (1986 A) :**
Similarity of protein encoded by the human *c-erbB-2* gene to epidermal growth factor receptor.
Nature , 319 : 230 .
- Yamamoto, T., Kamata, N. and Kawano, H. (1986 B) :**
High incidence of amplification of the epidermal growth factor receptor gene in human squamous carcinoma cell lines .
Cancer Res. , 46 : 414 .
- Yokoto, J. , Yamamoto, T. and Toyoshima, K. (1986) :**
Amplification of *c-erbB-2* oncogene in human adenocarcinomas in vivo .
Lancet , i : 765 .
- Zaino ,R.J. (1992) :**
Endometrial carcinoma : the need to pursue a new pathology .
Gynecol . Oncol. , 45 : 233 .