

REFERENCES

Albrecht JL and Tomich PG (1996): The maternal and neonatal outcome of triplet gestation. *Am. J Obstet Gynecol.*, 174: 1551-1556.

Al-Mufti R, Hambley H, Albaiges G, Lees C and Nicolaides KH (2000): Increased fetal erythroblasts in women Who Subsequently develop preeclampsia. *Hum. Reprod.*, 15: 1624-1628.

Amicucci P, Gennarelli M, Novelli G, and Dallapiccola B (2000): Prenatal Diagnosis of myotonic dystrophy using fetal DNA obtained from maternal plasma. *Clin Chem.*, 46: 301-302.

Ananth CV, Smulian JC and Vintzileos AM (1999): Incidence of placental abruption in relation to cigarette smoking and hypertensive disorders during pregnancy: A Meta-analysis of observational studies. *Obstetrics and Gynecology*, 93: 622-628.

Anker P, Mulcahy H, Chen XQ, and Stroun M (1999): Detection of circulating tumour DNA in the blood (plasma/serum) of cancer patients. *Cancer Metastasis Rev.*, 18: 65-73.

Ariga, H, Ohto, H, Busch MP, Imamura S, Watson R W, and Lee TH (2001): Kinetics of fetal cellular and cell-free DNA in the maternal circulation during and after pregnancy: implications for noninvasive prenatal diagnosis. *Transfusion*, 41, 1524-1530.

Artlett CM, Smith JB, and Jimenez SA (1998): Identification of fetal DNA and cells in skin lesions from women with systemic sclerosis. *N. Engl. J. Med.*, 338:1186-1191.

Ashkenas J and Werb Z (1996): Proteolysis and the biochemistry of life-or-death decisions. *J. Exp. Med.*, 183: 1947-1951.

Bertero MT, Camaschella C, Serra A, Bergui L and Caligaris-Cappio F (1988): Circulating 'trophoblast' cells in pregnancy have maternal genetic markers. *Prenat. Diagn.*, 8 : 585-590.

Bianchi DW (1997): Progress in the genetic analysis of fetal cells circulating in maternal blood. *Curr. Opin. Obstet. Gynecol.*, 9: 121-125.

Bianchi DW (1999): Fetal cells in maternal circulation: feasibility for prenatal diagnosis. *Br. J. Haematol.* , 105:574-583.

Bianchi DW, and Lo YM (2001): Fetomaternal cellular and plasma DNA trafficking. *Ann. N.Y. Acad. Sci. USA*, 945:119-131.

Bianchi DW, Flint AF, Pizzimenti MF, Knoll JH, and Latt SA (1990): Isolation of fetal DNA from nucleated erythroblasts in maternal blood. *Proc. Nat. Acad. Sci. USA*, 87: 3279-3283.

Bianchi DW, Simpson JL, Jackson LG, Elias S, Holzgreve w, Evans MI, Dukes KA, Sullivan LM, Klinger KW, Bischoff FZ, Hahn S, Johnson KL, Lewis D, Wapner RJ and Cruz F (2002): Fetal gender and aneuploidy detection using fetal cells in maternal blood: analysis of NIFTY I data. *Prenat. Diagn.*, 22:609-615.

Bianchi DW, Williams JM, Sullivan LM, Hanson FW, Klinger KW and Shuber AP (1997): PCR quantitation of fetal cells in maternal blood in normal and aneuploid pregnancies. *Am. J. Hum. Genet.* , 61:822-29

Bianchi DW, Zickwolf GK, Weil GJ, Sylvester S and DeMaria MA (1996): Male fetal pro progenitor cells persist in maternal blood for as long as 27 years postpartum. *Proc. Natl. Acad. Sci. USA*, 93:705-708.

Bischoff F, Neguten D, Marquez Do D, Moise K, Simpson JL, and Elias S (1999): Non invasive determination of fetal RhD status using fetal DNA in maternal serum and PCR. *J. Soc. Gynecol. Invest.*, 6: 64-69.

Bischoff F, Sinacori MK, Do DM, Dang DB, Horni C, Lewis DE, and Simpson JL (2002): Cell free fetal DNA and intact cells in maternal blood circulation: implications for first and second trimester non invasive prenatal diagnosis. *Human Reproduction Update*, 8: 493-500.

Botezatu I, Serdyuk O, Potapova G, Shelepov V, Alechina R and Molyaka Y (2000): Genetic analysis of DNA excreted in urine: a new approach for detecting specific genomic DNA sequences from cells dying in an organism. *Clin. Chem.*, 46: 1078-1084.

Chan FY, Cowley NM, Wolter L, Stone M, Carmody F, Saul A and Hyland CA (2001): prenatal RHD gene determination and dosage analysis by PCR: Clinical evaluation. *Prenat. Diagn.*, 21:321-326.

Chan KC, Zhang J, Hui AB, Wong N, Lau TK, Leung TN, Lo K, Huang DW and Lo YM (2004): Size distributions of maternal and fetal DNA in maternal plasma. *Clin. Chem.*, 50: 88 - 92.

Chen CP, Chern SR and Wang W (2000): Fetal DNA in maternal plasma: The prenatal detection of a paternally inherited fetal aneuploidy. *Prenat. Diagn.* 20: 355-357.

- Chen H, Griffin DK, Jestice K, (1998):** Evaluating of the culture of fetal erythroblasts from maternal blood for non-invasive prenatal diagnosis. *Prenat. Diag.*, 18: 883-892.
- Chen XQ, Stroun M, Magnenat JL, Nicod LP, Kurt AM Lyautey J, Lederrey C and Anker P. (1996):** Microsatellite alterations in plasma DNA of small cell lung cancer patients. *Nat. Med.* 2: 1033-1035.
- Cheung MC, Goldberg JD, and Kan YW (1996):** Prenatal diagnosis of sickle cell anemia and thalassemia by analysis of fetal cells in maternal blood. *Nat. Genet.*, 14: 264-268.
- Chiu R, Poon L, Lau T, Leung T, Wong E, and Lo YM (2001):** Effects of blood-processing protocols on fetal and total DNA quantification in maternal plasma. *Clin. Chem.*, 47: 1607-1613.
- Ciaranfi A, Curchod A, and Odartchenko N (1977):** Post-partum survival of fetal lymphocytes in the maternal blood]. *Schweiz. Med. Wochenschr.*, 107 : 134 - 138.
- Costa JM, Benachi A, Gautier E, Jouannic JM, Ernault, P, and Dumez Y (2001):** First-trimester fetal sex determination in maternal serum using real-time PCR. *Prenat. Diagn.*, 21: 1070-1074.
- Cunningham FG (2001):** Hypertensive disorder in pregnancy. In *Williams Obstetrics*, 21st ed. New York: McGraw-Hill, pp. 567-618.
- Cunningham FG (2001):** Maternal adaptations to pregnancy. In *Williams Obstetrics*, 21st ed., New York: McGraw-Hill, pp. 167-200.

De Grouchy J, and Trebuchet C (1971): Fetomaternal transfusion of blood lymphocytes and identification of the sex of the fetus. *Ann. Genet.*, 14:133-137.

Dekker GA and Sibai BM (1998): Etiology and pathogenesis of preeclampsia: current concepts. *Am. J. Obstet. Gynecol.*, 179: 1359-1375.

Difederico E, Genbacew O, and Fisher SJ (1999): Preeclampsia is associated with widespread apoptosis of placental cytotrophoblasts within the uterine wall. *Am. J. pathol.* 155: 293-301.

Duley L (2002): Preeclampsia and hypertension. *Clinical Evidence*, (7): 1296-1309.

Elias S, Lewis DE, and Simpson JL (1996): Isolation of fetal nucleated red blood cells from maternal blood: persistence of cells from prior pregnancy is unlikely to lead to false positive results. *J. Soc. Gynecol. Invest.*, 3: 359.

Emlen W and Mannik M (1978): Kinetics and mechanism for removal of circulating single- stranded DNA in mice. *J. Exp. Med.*, 147: 684 – 689.

Farina A, Caramelli E, Concu M, Sekizawa A, Ruggeri R, Bovicelli L, Rizzo N and Carinci P (2002): Testing normality of fetal DNA concentration in maternal plasma at 10-12 completed week's gestation: a preliminary approach to a new marker for genetic screening. *Prenat. Diagn.*, 22: 148-152.

- Farina A, LeShane ES, Lambert MG, Canick JA, Lee T , Neveux LM, Palomaki GE and Bianchi DW (2003):** Evaluation of cell-free fetal DNA as a second-trimester maternal serum marker of Down syndrome pregnancy. Clin. Chem., 49: 239-242.
- Farina A, Sugito Y, Matsuoka R, Manko I Saito H and Okia T (2004):** Proteinuria and hypertension are independent factors affecting fetal DNA values: A retrospective analysis of affected and unaffected patients. Clin. Chem., 50 : 221-224.
- Fleischhacker M (1999):** First international symposium on circulating nucleic acids in plasma/serum. Eur. J. Med. Res., 4: 488-490.
- Fournie GJ, Martres F, Pourrat JP, Alary C, and Rumeau M (1993):** Plasma DNA as cell death marker in elderly patient. Gerontology, 39: 215 – 221.
- Frisch SM and Ruoslahti E, (1997):** Integrins and anoikis. Curr. Opin. Cell Biol., 9: 701- 706.
- Ganshirt D, Smeets FW, Dohr A, Walde C, Steen I, Lapucci C, Falcinelli C, Sant R, Velasco M , Garritsen HS, and Holzgreve W. (1998):** Enrichment of fetal nucleated red blood cells from the maternal circulation for parental diagnosis: experiences with triple density gradient and MACS based on more than 600 cases. Fetal Dign. Ther., 13: 276 – 286.
- Hahn S, and Holzgreve W (2002):** Fetal cells and cell free fetal DNA in maternal blood: new insight in preeclampsia . Hum. Reprod. Update, 8: 501-508.

- Hahn S, Sant R, and Holzgreve W (1998):** Fetal cells in maternal blood: current and future perspectives. *Mol. Hum. Reprod.*, 4: 515-521.
- Hahn S, Zhong XY, Burk MR, Troeger C, and Holzgreve W (2000):** Multiplex and real time quantitative PCR on fetal DNA in maternal plasma. A comparison with fetal cells isolated from maternal blood. *Ann. N.Y. Acad. Sci.* , 906 :148-152.
- Haque KA, Pfeiffer RM, Beerman MP, Struewing JP, Chanock JP, and Bergen AW (2003):** Performance of high throughput DNA quantification methods. *BMC Biotechnol.*, 3:20-24.
- Hawes CS, Suskin HA, and Kalionis B (1994):** Detection of paternally inherited mutations for beta-thalassemia in trophoblast isolated from peripheral maternal blood. *Ann. N. Y. Acad. Sci.*, 731:181-185
- Hawes CS, Suskin HA, Petropoulos A, Latham SE and Muller UM (1994):** A morphologic study of trophoblast isolated from peripheral blood of pregnant women. *Am. J. Obstet. Gynecol.*, 170:1297-1300.
- Herzenberg LA, Bianchi DW, and Schroder J (1979):** Fetal cells in the blood of pregnant women: detection and enrichment by fluorescence-activated cells sorting. *Proc. Natl. Acad. Sci. USA* , 76:1453-1455.
- Holzgreve W, and Hahn S. (1999):** Novel molecular biological approaches for the diagnosis of preeclampsia. *Clin. Chem.*, 45: 451-452.

Holzgreve W, Ghezzi F, DiNaro E, Ganshirt D, Maymon E and Hahn S (1998): Disturbed feto-maternal cell traffic in pre-eclampsia. *Obstet. Gynecol.*, 91: 669-672.

Honda H, Miharu N, Ohashi Y, Samura O, Kinutani M, Hara T and Ohama K (2002): Fetal gender determination in early pregnancy through qualitative and quantitative analysis of fetal DNA in maternal serum. *Hum. Genet.*, 110: 75-79.

Houfflin DV, O'Donnell H, Overton T, Bennett PR and Fisk NM (2000): High sensitivity of fetal DNA in plasma compared to serum and nucleated cells using un nested PCR in maternal blood. *Fetal Diagn. Ther.*, 15: 102-107.

Jimenez DF and Tarantal AF (2003): Quantitative analysis of male fetal DNA in maternal serum of Gravid Rhesus Monkeys. *Pediatric Research*, 53: 18-23.

Johansen M, Redman CW, Wilknis T and Sargent IL (1999): Trophoblast deportation in human pregnancy – its relevance for pre-eclampsia. *Placenta*, 20: 531-539.

Johnson K, Dukes K, Vidaver J, Leshane JS, Ramirez I, Weber W, Bischoff F, Sinuh H, Sharma, D, Hire L, Bianchi D, Simpson J, Holzgreve W, Elias S and Klinger W (2004): Inter laboratory comparison of fetal male DNA detection from common maternal plasma samples by real –time PCR. *Clin. Chem.*, 50: 516-521.

Kao SM, Tang GC, Hsieh TT, Young KC, Wang HC and Pao CC (1992): Analysis of peripheral blood of pregnant women for the presence of fetal Y chromosome-specific ZFY gene deoxyribonucleic acid sequences. *Am. J. Obstet. Gynecol.*, 166:1013-1019.

Khosrotehrani K, and Bianchi DW (2005): Multi-lineage potential of fetal cells in maternal tissues: a legacy in reverse .Cell Sci., 118:1559-1563.

Kolialexi A, Tsangaris G, Antsaklis A and Mavroua A (2004): Rapid clearance of fetal cells from maternal circulation after delivery. Ann. N. Y. Acad. Sci.,1022:113-118.

Lambert NC, Lo YM, Erickson TD, Tylee TS, Guthrie KA, Furt DE and Lee NJ (2002): Male microchimerism in healthy women with scleroderma: cells or circulating DNA? Aquantitative answer .Blood, 100:2845-2851.

Lau TW, Leung TN, Chan LY, Lau TK, Chan KC , Tam WH and Lo YM (2002): Fetal DNA clearance from maternal plasma is impaired in preeclampsia. Clin. Chem., 48: 2141-2146.

Leon SA, Shapiro B, Sklaroff DM, and Yaros MJ (1977): Free DNA in serum of cancer patients and the effect of therapy. Cancer Res., 37: 646-650.

Leung TN, Smith SC, TO KE, Sahota DS and Baker PN (1999): Increased apoptosis in pregnancies complicated by pre-eclampsia. J. Soc. Gynecol. Invest ., 6: 127.

Leung TN, Zhang J, Lau TK, Chan YS, and Lo YM (2001): Increased maternal plasma fetal DNA concentrations in women who eventually develop preeclampsia. Clin. Chem., 47: 137 – 139.

Leung TN, Zhang J, Lau TK, Hjelm NM and Lo YM (1998): Maternal plasma fetal DNA as a marker for preterm labour. Lancet, 352:1904-1905.

Levine RJ, Hauth JC, Curet LB, Sibai BM, Catalano PM, Morris CD, DerSimonian R, Esterlitz JR, Raymond EG, Bild DE, Clemens JD and Cutler JA (1997): Trial of calcium to prevent preeclampsia N. Engl. J. Med., 337: 69-76.

Levine RJ, Qian C, LeShane ES, Yu KF, England LJ Schisterman EF, Watagnara R, Romero Rand Bianchi DW (2004): Two-stages elevation of cell-free fetal DNA in maternal sera before onset preeclampsia. Am.J. Obstet. Gynecol., 190: 707-713.

Liou JD, Hsieh TT and Pao CC (1994): Persistence of cells of fetal origin in maternal circulation of pregnant women. Fetal Cell in Maternal Blood: Prospects for Noninvasive Prenatal Diagnosis (Edited by: Edited by Simpson JL Elias S). New York Academy of Sciences, 237

Liou JD, Pao CC ,Hor JJ and Kao SM (1993): Fetal cells in the maternal circulation during first trimester in pregnancies. Hum. Genet. , 92: 309-311

Lo YM (2000a): Fetal DNA in maternal plasma. Ann. N.Y. Acad. Sci., 906: 141-147.

Lo YM, Corbetta N, Chamberlain PF, Rai V, Sargent IL, Redman CW and Wainscoat JS (1997): Presence of fetal DNA in maternal plasma and serum. Lancet, 350: 485-487.

Lo YM, Hjelm NM, Fidler C, Sargent IL, Murphy MF, Chamberlain PF, Poon PM, Redman CW and Wainscoat JS (1998b): Prenatal diagnosis of fetal RhD status by molecular analysis of maternal plasma. *N. Engl. J. Med.*, 339:1734-1738.

Lo YM, Lau TK, Chan LY, Leung TN and Chang AM (2000b): Quantitative analysis of the bi-directional fetomaternal transfer of nucleated cells and plasma DNA. *Clin. Chem.*, 46: 1301-1309.

Lo YM, Lau TK, Zhang J, Leung TN, Chang AM, Hjelm NM, Elmes RS, and Bianchi DW (1999a): Increased fetal DNA concentrations in the plasma of pregnant women carrying fetuses with trisomy 21. *Clin. Chem.*, 45: 1747-1751.

Lo YM, Leung TN, Tein MS, Sargent IL, Zhang J and Lau TK (1999b): Quantitative abnormalities of fetal DNA in maternal serum in preeclampsia. *Clin. Chem.*, 45: 184-188.

Lo YM, Patel P, Baigent CN, Gillmer MD, Chamberlain P, Travi M, Sampietro M, Wainscoat JS and Fleming KA (1993): Prenatal sex determination from maternal peripheral blood using the polymerase chain reaction. *Hum. Genet.*, 90: 483 – 488.

Lo YM, Rainer TH, Chan LY, Hjelm NM, and Cocks RA (2000a): Plasma DNA as a prognostic marker in trauma patients. *Clin. Chem.*, 46: 319-323.

Lo YM, Tein MS, Lau TK, Haines CJ, Leung TN, Poon PM, Wainscoat JS, Johanson PJ, Chang AM and Hjelm NM (1998a): Quantitative analysis of fetal DNA in maternal plasma and serum: implications for noninvasive prenatal diagnosis. *Am. J. Hum. Genet.*, 62: 768-775.

- Lo YM, Tein MS, Pang CC, Yeung CK, Tong KL, and Hjelm NM (1998c):** Presence of donor-specific DNA in plasma of kidney and liver-transplant recipients. *Lancet*, 351: 1329 – 1330.
- Lo YM, Zhang J, Leung TN, Lau TK, Chang AM and Hjelm NM (1999c):** Rapid clearance of fetal DNA from maternal plasma. *Am. J. Hum. Genet.*, 64: 218-224.
- Lui YY, Woo KS, Wang AY, Yeung CK, Li PK, Chan E, Ruygrok P and Lo YM (2003):** Origin of plasma cell free fetal DNA after solid phase organ transplantation. *Chin. Chem.*, 49:495-496.
- Moreau P, Teyssier M Kirszenbaum M, Gluckman E, Gouarnd L, Carosella E and Dausset J (1994) :** HLA-G mRNA forms in human trophoblasts and peripheral blood lymphocytes: potential use in prenatal diagnosis. *Folia. Biol.*, 40:431-438.
- Nelson JL (1996):** Maternal-fetal immunology and autoimmune disease: is some autoimmune disease auto-alloimmune or allo-autoimmune? *Arthritis Rheum.*, 39:191-194.
- Nelson JL, Furst DE Maloney S, Gooly T, Evans PC, Smith A, Bean MA, Ober C and Bianchi DW (1998):** Microchimerism and HLA-compatible relationships of pregnancy in scleroderma. *Lancet* , 351:559-562.
- Ng EK, Leung TN, Tsui NB, Lau TK, Panesar NS Chiu RW and Lo YM (2003):** the concentration of circulating corticotropin releasing hormone mRNA in maternal plasma is increased in preeclampsia. *Clin. Chem.*, 49:727-731.

Ng EK, Tsui NB, Lau TK, Leung TN, Panear NS, Lit LC, Chan KW and Lo YM (2003): mRNA of placental origin is readily detectable in maternal plasma. *Proc. Natl. Acad. Sci. USA*, 100:4748-4753.

Oettle C, Hall D, Roux A and Grove D (2005): Early onset severe pre-eclampsia: expectant management at a secondary hospital in close association with a tertiary institution. *BJOG*, 112: 84-88.

Perloff D (1998): Hypertension and pregnancy –related hypertension. *Cardiology Clinics*, 16 : 79-101.

Pertl B and Bianchi DW (2001): Fetal DNA in maternal plasma: emerging clinical application. *Obstet. Gyn.*, 98: 483- 490.

Pertl B, Sekizawa A, Samura O, Orescovic I, Rahaim PT, and Bianchi DW (2000): Detection of male and female fetal DNA in maternal plasma by multiplex fluorescent PCR amplification of short tandem repeats. *Hum. Genet.*, 106: 45-49.

Poon LM, Leung TN, Lau TK, and Lo YMD (2000): Presence of fetal RNA in maternal plasma. *Clin. Chem.*, 46: 1832-1834.

Poon LM, Leung TN, Lau TK, Chow KC and Lo YM (2002): Differential DNA methylation between fetus and mother as a strategy for detecting fetal DNA in maternal plasma. *Clin. Chem.*, 48, 35-41.

Price JO, Elias S, Wachtel SS, Klinger K, Dockter M, Tharapel A, Shulman LP, Phillips OP, Meyers CM, and Shook D (1991) : Prenatal diagnosis with fetal cells isolated from maternal blood by multi parameter flow cytometry. Am. J. Obstet. Gynecol , 165:1731-1737.

Redman CW and Sargent IL (2005): Latest advances in understanding preeclampsia. Science, 308: 1592-1594.

Roberts JM and Cooper DW (2001): Pathogenesis and genetics of pre-eclampsia. Lancet, 357: 53-56.

Saito H, Sekizawa A, Morimoto T, Suzuki M and Yanaihara T (2000): Prenatal DNA diagnosis of a single gene disorder from maternal plasma .Lancet , 356: 1170.

Sambrook J, Fritsch EF and Maniattis T (2001): Molecular cloning a laboratory manual. Cold spring Harbor Laboratory Press.

Samura O, Miharu N, Hyodo M, Honda H, Ohashi Y, Honda N, Hara Ohama K, (2003): Cell free fetal DNA in maternal circulation after amniocentesis Clin. Chem., 49: 1193-1195.

Samura O, Sekizawa A , Zhen DK, Falco VM and Bianchi DW (2000): Comparison of fetal cell recovery from maternal blood using a high-density gradient for the initial separation step: 1.090 versus 1.119 g/ml. Prenat. Diagn., 20:281-286

Sekizawa A, Farina A, Zhen DK, Wang JY, Falco VM, Elmes S and Bianchi DW (1999): Improvement of fetal cell recovery from maternal blood suitable density gradient for FACS separation. Fetal Diagn. Ther., 14: 229-233.

Sekizawa A, Farina A, Sugito Y, Matsuoka R, Iwasaki N, Saito H and Okai T, (2004): Proteinuria and hypertension are independent factors affecting fetal DNA values: A retrospective analysis of affected and unaffected patients. Clin. Chem., 50: 221-224.

Sekizawa A, Jimbo M, Saito H, Iwasaki M, Matsuoka R, Okai T and Farina A (2003): Cell –free fetal DNA in the plasma of pregnant women with severe fetal growth restriction. Am. J. Obstet. Gynecol, 188:480-484.

Sekizawa A, Jimbo M, Saito H, Iwasaki M, Sugito Y and Yukimoto Y (2002): Cell-free fetal DNA is increased in plasma of women with hyperemesis gravidarum. Clin. Chem., 47: 2164-2165.

Sekizawa A, Kondo T, Iwasaki M, Watanabe A, Jimbo M Saito H and Okai T (2001): Accuracy of fetal gender determination by analysis of DNA in maternal plasma. Clin. Chem., 47: 1856-1858.

Sekizawa A, Samura O, Zhen DK, Flaco V, Farina A and Bianchi DW (2000): Apoptosis in fetal nucleated erythrocytes circulating in maternal blood. Prenat. Diagn., 20: 886-889.

Sekizawa A, Yokokawa K, Sugito Y, Iwasaki M, Yukimoto Y, Ichizuka K, Saito H and Okai T (2003): Evaluation of bi-directional transfer of plasma DNA through placenta. Hum. Genet., 113: 307-310.

Shimazaki C, Ochiai N, Uchida R, Okano A, Fuchida S, Aslihara E, Inaba T Fujita N, Maruya E, and Nakagawa M (2003): Non- T cell depleted HLA haploidentical stem cell transplantation in advanced hematological malignancies based on the fetomaternal microchimerism. *Blood*, 101: 3334-3336.

Shulman LP (2003): Fetal cells in maternal blood. *Current Women's Health Reports*, 3: 47-54.

Shulman LP, Phillips OP, Tolley E, Sammons D, and Wachtel SS (1998): Frequency of nucleated red blood cells in maternal blood during the different gestational ages. *Hum. Genet.*, 103:723-726

Sibai B, Dekker G and Kupferminc M (2005): Preeclampsia. *Lancet*, 365: 789-799.

Sibia BM (1996): Treatment of hypertension in pregnant women. *New Engl. J. Med.*, 335: 257-264.

Siva S, Johanson S, McCracken S and Morris J (2003): Evaluation of the clinical usefulness of isolation of fetal DNA from the maternal circulation. *Australian & New Zealand Journal of obstetric & gynecology*, 43: 10-15.

Skjaerven R, Wilcox AJ, and Li RT (2002): The interval between pregnancies and the risk of preeclampsia. *New England Journal of Medicine*, 346: 33-38.

- Smid M, Lagona F, De Benassuti L, Ferrari A, Ferrari M and Cremonesi L (1999):** Evaluation of different approaches for fetal DNA analysis from maternal plasma and nucleated blood cells. Clin. Chem., 45: 1570 – 1572.
- Smith MA (2003):** Placental mRNA in maternal plasma: Prospects for fetal screening. PNAS, 100:4360-4362.
- Snedecor G, Cochran W and Cox D (1989):** Statistical Methods (8th edition). The Iowa State University Press.
- Stemmer M, Faller M, Pencreac'h E, Guerin E, Schneider A, Jaqmin D, Quoix E, Gaub MP and Oudet P (2003):** Use of magnetic beads for plasma cell free DNA extraction: Toward automation of plasma DNA analysis for molecular diagnostics. Clin. Chem., 49: 1953-1955.
- Stenman J and Orpana A (2001):** Correspondence accuracy in amplification. Nature Biotechnol., 19: 1011-1012.
- Stroun M, Anker P, Maurice P and Gaban PB (1977):** Circulating nucleic acids in higher organisms. Int. Rev. Cytol., 51: 1-48.
- Swinkels DW, Dekok JB, Hendriks JC, Wiegerinck E, Zusterzeel PL and Steegers EA (2002):** Hemolysis, elevated liver enzymes, and low platelet count (HELLP) syndrome as a complication of preeclampsia in pregnant women increased the amount of cell-free fetal and maternal DNA in maternal plasma and serum. Clin. Chem., 48: 650-653.
- Swinkels DW, Wiegerinck E, Steegers E and De Kok JB (2003):** Effects of blood processing protocols on cell free DNA quantification in plasma. Clin. Chem., 49 : 525-526.

Tang NL, Leung TN, Zhang J, Lau TK, and Lo YM (1999): detection of fetal derived paternally inherited X-chromosome polymorphisms in maternal plasma. Clin. Chem.,45:2033-2035.

Thomas MR, Williamson R Craft I, Yazdani N, and Rodeck CH (1994): Y chromosome sequence DNA amplified from peripheral blood in women in early pregnancy. Lancet , 343:413-414.

Tyndall A, and Gratwohl A (1998): Microchimerism: friend or foe? Nat. Med., 4:386-388.

Van Wijk IJ, De Hoon AC, Jurhawan R, Tjoa ML, Griffioen S, Mulders MA, van Vugt JM and Oudejans CB (2000): Detection of apoptotic fetal cells in plasma of pregnant women. Clin. Chem., 46: 729-731.

Van Wijk IJ, Van Vugt JM, Mulders MA, Konst AA, Weima SM, and Oudejans CB (1996): Enrichment of fetal trophoblast cells from the maternal peripheral blood followed by detection of fetal deoxyribonucleic acid with a nested X/Y polymerase chain reaction. Am. J. Obstet. Gynecol , 174:871-878.

Von Dadelszen P, Magee LA and Roberts JM (2003): Sub classification of preeclampsia. Hypertension Pregnancy, 22: 143: 148.

Wachtel SS, Sammons D and Twitty G (1998): Charge flow separation: quantification of nucleated red blood cells in maternal blood during pregnancy. Prenat. Diagn., 18:455-463.

- Wachtel SS, Sammons D, Manley M, Wachtel G, Twitty G, Utermohlen J, Phillips OP, Shulman LP, Taron DJ, Muller UR, Koeppen ©, Ruffalo TM, Addis K, Porreco R, Collins JM, Parker NB, and McGavran L (1996):** Fetal cells in maternal blood :Recovery by charge flow separation . Hum. Genet., 98:162-166.
- Wald NJ and Hackshaw AK (1997):** Combining ultrasound and biochemistry in first trimester screening for Down's syndrome. . Prenat. Diagn., 17: 821-829.
- Walknowska J, Conte FA and Grumbach MM (1969):** Practical and theoretical implications of fetal maternal lymphocyte transfer. Lancet , 1: 1119-1122.
- Wataganara T, LeShane ES, Farina A, Messerlian GM, Lee T, Canick JA, and Bianchi DW (2003):** Maternal serum cell-free fetal DNA levels are increased in cases of trisomy 13 but not trisomy 18. Hum. Genet., 112: 204-208.
- Wataganara T, Peter I, Messerlian GM, Borgatta L and Bianchi DW (2004):** Inverse correlation between maternal Weight and second trimester circulating cell free Fetal DNA levels. Obst. Gynecol, 104:545-550.
- Wong IH, Lo YM, Zhang J, Liew CT, Ng MH, Wong N, Lai PB, Lau WY, Hjelm NM, and Johnson PJ (1999):** Detection of abnormal p16 methylation in the plasma and serum of liver cancer patients. Cancer Res., 59: 71-73.
- Zamerowski S, Lumley M, Arreola R, Dukes K and Sullivan L (2001):** Favorable attitudes toward testing for chromosomal abnormalities via analysis of fetal cells in maternal blood. Genet. Med., 3: 301-309.

Zhong XY, Burk MR, Troeger C, Jackson LR Holzgreve W, and Hahn S (2000): Fetal DNA in maternal plasma is elevated in pregnancies with aneuploid fetuses. *Prenat. Diagn.* 20: 795-798.

Zhong XY, Holzgreve W and Hahn S (2000): Detection of fetal Rhesus D and sex using fetal DNA from maternal plasma by multiplex polymerase chain reaction. *Br. J. Obstet. Gynaecol.* 107: 766-769.

Zhong XY, Holzgreve W and Hahn S (2001): Circulating fetal and maternal DNA in pregnancies at risk and those affected by preeclampsia. *Ann. N. Y. Acad. Sci.*, 945: 138-140.

Zhong XY, Holzgreve W and Hahn S (2002): Cell free fetal DNA in the maternal circulation does not stem from the transplacental passage of fetal erythroblasts. *Mol. Hum. Reprod.*, 8: 864-870.

Zhong XY, Holzgreve W, Hoesli I and Hahn S (2005): Circulating corticotrophin-releasing hormone mRNA concentration are increased in women with preterm delivery but not in those who respond to tocolytic treatment. *Clin. chem.*, 51: 635-636.

Zhong XY, Holzgreve W, Li JC, Aydinli K and Hahn S (2000): High levels of fetal erythroblasts and fetal extra cellular DNA in the peripheral blood of a pregnant woman with idiopathic polyhydramnios: Case report. *Prenat. Diagn.*, 20: 838-841.

Zhong XY, Laivuori H, Livingston JC, Ylikorkala O, Sibai BM and Holzgreve W (2001): Elevation of both maternal and fetal extra cellular circulating deoxyribo-nucleic acid concentrations in the plasma of pregnant women with preeclampsia. *Am. J. Obstet. Gynecol.* 184: 414-419.