

REFERENCES

- Abdel-Aziz, M.T.; Abdel-Kader, M.M.; Shalaby, E.;  
Tawadrous, G.A.; Salam, N.; Shawky, W.;  
El-Meleigy, S.; and Abdel-Aziz, A. (1980):  
Effect of DL-Carnitine on plasma lipids in  
normal and obese human females.  
Egypt. J. Physiol. Sci., 7 (1): 11-20.
- Abdel-Aziz, M.T.; Tawadrous, G.A.; Soliman, K.M. ;  
Mustafa, S.; Abdan, M.S.; Table, A. (1984):  
Effect of carnitine on blood lipid pattern  
in diabetic patients .  
Nutr. Rep. Intern., 29 (5) : 1071-1079.
- Abrahms, J.J.; Ginsberg, H. and Grundy, S.M. (1982) :  
Metabolism of cholesterol and plasma trigly-  
cerides in non-ketatic diabetes mellitus.  
Diabetes 31 : 903-910.
- Albers, J.J.; Adolphson, J.L.; and Hazzard, W.R.(1977):  
Radioimmunoassay of human plasma Lp (a)  
lipoprotein.  
J. Lipid Res., 18 : 331-338.
- Albrink M.J.; Laviates, P.H., and Man, E.B. (1963) :  
Vascular disease and serum lipids in diabetes  
mellitus : observations of 30 years.  
Ann. Intern. Med. 58 : 305-23.

Ames, R.P. (1983) :

Metabolic disturbances increasing the risk  
of CHD during diuretic-based antihyperten-  
sive therapy.

Am. Heart J. 106 : 1207.

Ashley, F.W.; Kannel, W.B. (1974) :

Relation of weight change to changes in  
atherogenic traits : the Framingham study .

J. Chronic Dis. 27 : 103.

Assmann, G.; and Brewer, H.B. (1974) :

Lipid-protein interactions in HDL.

Proc. Natl. Acad. Sci. (U.S.A.), 71 : 1534-1538.

Assmann, G. ; Funke, H. and Schriewer, H. (1982) :

The relation of HDL-apolipoprotein A<sub>1</sub> and  
HDL-cholesterol to risk factors of coronary  
heart disease : Initial results of the prospec-  
tive epidemiological study in company employees  
in westfalia.

J. Clin. Chem. Clin. Biochem., 20 (5)- 287-289.

Avogaro, A ; Bon, G.B.; Cazzolato, G.; et al. (1978):

Variations in apolipoprotein B and A<sub>1</sub> , during  
the course of myocardial infarction.

Eur. J. Clin. Invest., 8 : 121-9.

Avogaro, P.; Bon, G.B.; Cazzolato, Quinci, G.B.;

Belussi, F. (1978) :

Plasma levels of apolipoprotein A<sub>1</sub> and  
apolipoprotein B in human atherosclerosis.  
Artery, 4 : 385-94.

Avogaro, P.; Bon, G.B.; Cazzolato, G.; Quinci,

G.B. (1979) :

Are apolipoproteins better discriminators  
than lipids for atherosclerosis ?  
Lancet I : 901-3.

Bagdad, J.D.; Porte, D. Jr.; and Bierman, E.L. (1967):

Diabetic lipaemia. A form of acquired fat  
induced lipoaemia.

N. Engl. J. Med., 276 : 427-433.

Bagdad, J.D.; Porte, D. Jr.; and Bierman, E.L. (1968):

Acute insulin withdrawal and the regulation  
of plasma triglycerides removal in diabetic  
subjects.

Diabetes, 17 : 127-132.

Ballosse, E.O.; Bier, D.M.; and Havel, R.J. (1972) :

Early effects of anti-insulin serum on  
hepatic metabolism of plasma free fatty acids  
in dogs.

Diabetes, 21 : 280-288.

- Barclay, M.; Barclay, R.K.; Skipski, V.P.; Terebus-  
kekish, O.; Muller, C.H.; Shah, E.; and  
Elkins, W.L. (1965) :  
Fluctuation in human serum lipoproteins  
during the normal menstrual cycle.  
Biochem. J., 96 : 205.
- Baron, H.; Chen, Y.I.; and Reaven, G.M. (1981) :  
Evidence for a new cause of defective plasma  
removal of very low density lipoproteins in  
insulin deficient rats.  
Diabetes, 30 : 496-499.
- Basso, J.V.; and Havel, R.J. (1970) :  
Hepatic metabolism of free fatty acids in  
normal and diabetic dogs.  
J. Clin. Invest., 49 : 537-47.
- Basu, S.K.; Brown, M.S.; Ho, Y.K.; Havel, R.J.; and  
Goldstein, J.L. (1981) :  
Mouse macrophage synthesized and secrete a  
protein resembling opoprotein E.  
Proc. Natl. Acad. Sci., (USA) 78 : 7545-7549.
- Beard, J.T. (1983) :  
Serum uric acid and coronary heart disease.  
Am. Heart J., 106 : 397.

Bengtsson, C. (1973 ) :

Ischaemic heart disease in women : a study based on a randomized population sample of women and women with myocardial infarction in Gotenberg, Sweden.

Acta. Med. Scand . (suppl.) 549.

Bennion, L.J.; and Grundy, S.M. (1977) :

Effects of diabetes mellitus on cholesterol metabolism in man.

N. Engl. J. Med., 296 : 1365.

Berg, K. (1963) :

Lipoproteins and arteriosclerosis.

Acta Pathol. Microbiol. Scand. 59, suppl., 166-167, 369-382.

Berg, K.; Dahlen, G.; and Frick, M.H. (1974) :

Lp (a) lipoprotein on pre-beta lipoprotein in patients with coronary artery disease.

Clin. Genet., 6 : 230-235.

Berkowitz, D.; Glassman, S. (1965) :

Effects of hypertriglyceridaemia on urinary uric acid outputs.

Circulation, 32 ( suppl. 2) : 2.

Bihari-V.M.; Szekely- J. Gruber, E. (1981) :

Plasma high density lipoproteins in coronary, cerebral and peripheral vascular disease.

Atherosclerosis 40 (3-4) : 337-45.

Blackburn, H. (1980) :

Epidemiologic evidence . In : American health foundation, plasma lipids, optimal levels for health.  
New York Academic Press, 7.

Blix, G.; Tiselius, A. and Svensson, H. (1941) :

Separation of plasma lipoproteins.  
J. Biol. Chem., 137 : 485-494.

Blue, M.L.; Williams, D.L.; Zacker, S.; Ali Khan, S.;  
and Blum, C.B. (1983) :

Apolipoprotein E synthesis in human kidney, adrenal gland and liver.  
Proc. Natl. Acad. Sci. (USA) 80 : 283-287.

Boguski, M.S.; Elshou bagy, N.; Taylor, J.M.; and  
Gordon, J.I. (1984) :

Apolipoprotein A IV contains 13 tandem repetitions of 22-amino acid sequent with amphipathic helical potential.  
Proc. Natl. Acad. Sci. (USA) 81 : 5021-5025.

Bolinder, J.; Lithell, H.; Sharfars, E. and Amer, P.  
(1986) :

Effects of obesity, hyperinsulinaemia, and glucose intolerance on insulin action in adipose tissue of sixty-year-old man.  
Diabetes 35 : 282-290.

Bonora, E.; Zavavoni, I.; Coscelli, C. and Butturini, U. (1983) :

Decreased hepatic insulin extraction in subjects with mild glucose intolerance. Metabolism 32 (5) : 438-446.

Bradley, R.F. (1971) :

Cardiovascular disease : In Joslin's Diabetes Mellitus. Marble, A., White, P.; Bradley, R.F.; and Krall, L.P.; Eds. Philadelphia, Lea & Febiger, p. 417-77.

Brennan, P.J.; Simpson, J.M.; Blacket, R.B.; Gilchrist, C.A. (1980) :

The effects of body weight on serum cholesterol, serum triglycerides serum urate and systolic blood pressure. Aust. NZJ. Med., 10 : 15.

Briones, E.R.; Mao, S.J.T.; Palumbo, P.J., Fallon, W.M.O., Chemoweth, W. and Kottke, B.A. (1984) :

Analysis of lipids and Apolipoproteins in Insulin. Dependent and Noninsulin Dependent Diabetics. Metabolism 33 (1) : 42-49.

Brischetto, C.S.; Conner, W.E.; Connor, S.L.; and Matarazzo, J.D. (1983) :

Plasma Lipid and Lipoprotein Profiles of Cigarette Smokers from Rondonly Selected Families; Enhancement of Hyperlipidaemia and Depression of High Density Lipoprotein. Am. J. Cardiol., 52 : 675.

- Brown, M.S.; and Goldstein, J.L. (1983) :  
Lipoprotein Metabolism in the Macrophage  
Implications for cholesterol deposition in  
atherosclerosis.  
Annu. Rev. Biochem., 52, 223.
- Brown, M.S.; and Goldstein, J.L. (1984) :  
How LDL receptors influence cholesterol and  
atherosclerosis.  
Sci. Am., 251 : 58.
- Brown, M.S.; Kovanen, P.T.; and Goldstein I.L. (1981) :  
Regulation of plasma cholesterol by lipo-  
protein receptors.  
Science, 212 : 628-635.
- Brown, M.S.; Goldstein, J.L.; Fredrickson, D.S. (1983):  
Familial type 3 hyperlipoproteinaemia (dysbeta-  
lipoproteinaemia). In : Stanobury, J.B.;  
Wyngaarden, J.B.; Fredrickson, D.S.; Goldstein,  
J.L.; Brown, M.S., eds. The metabolic basis of  
inherited disease. 5th ed. New York : McGraw-  
Hill, 655-71.
- Brunzell, J.D.; Porto, D.; Jr.; and Bierman, E.L. (1973):  
Evidence for a common, saturable triglyceride  
removal mechanism for chylomicrons and very  
low density lipoproteins in man.  
J. Clin. Invest., 52 : 1578-1585.

- Brunzell, J.; Porte, D.; and Bierman, E. (1975) :  
Reversible abnormalities in postheparin  
lipolytic activity during the late phase  
of release in diabetes mellitus.  
Metabolism, 24 : 1123-37.
- Brunzell, J.D.; Chait, A. and Bierman, E.L. (1978):  
Pathophysiology of lipoprotein transport.  
Metabolism 27 (9) : 1109.
- Brunzell, J.D.; Porte, D., Jr.; and Bierman, E.L.  
(1979) :  
Abnormal lipoprotein lipase mediated plasma  
triglyceride removal in untreated diabetes  
mellitus associated with hypertriglyceri-  
daemia.  
Metabolism, 28 : 902 - 907.
- Buccolo, B.; Duncan; and Gerardo, H. (1984) :  
Non enzymatic glycosylation of proteins, a  
new tool for assessment of cumulative hyper-  
glycaemia in epidemiologic studies, post and  
future.  
Am. J. Epidemiol., 20 (2) 177-182.
- Buccolo, G.; and David, H. (1973) :  
Quantitative determination of serum trigly-  
ceride by the use of enzymes.  
Clin. Chem., 19 : 476-482.

- Calvert, G.D.; Graham, J.J.; and Mannik, T. (1978):  
Effect of therapy on plasma high density  
lipoprotein-cholesterol concentration in  
diabetes mellitus.  
Lancet, 2: 66-68.
- Case, R.B.; Black, G.W.; Chadwick, J.N. (1985) :  
Type A behaviour and survival after acute  
myocardial infarction.  
N. Eng. J. Med., 312 : 737.
- Castelli, W.P.; and Anderson, K. (1986) :  
A population at risk : prevalence of high  
cholesterol levels in hypertensive patients  
in the framingham study.  
Am. J. Med., 80 (suppl 2 A) : 23-32.
- Castelli, W.P.; Doyle, J.T.; Gordon, T.; Hames, C.G.;  
Hjortland, M.C.; Hulley, S.B.; Kagan, A.; and  
Zukel, W.J. (1977) :  
HDL-cholesterol and other lipids in coronary  
heart disease.  
Circulation, 55 : 767.
- Cheung, P.; Kao, F.T.; Law, M.L.; Jones, C.; Puck, T.T.;  
and Chan, L. (1984) :  
Localization of the structural gene for human  
apolipoprotein A<sub>1</sub> on the long arm of human  
chromosome.  
Proc. Natl. Acad. Sci. (USA), 81 : 508-511.

- Chiba, K.; Koizumi, A.; Kumai, M.; Watanaba, T. and Keda, M. (1983) :  
Nation wide survey of high density lipoprotein cholesterol among farmers in Japan.  
Prev. Med., 12 (4) : 508-522.
- Coronary Drug Project Research Group (1976) :  
Serum uricacid : Its association with other risk factors and with mortality in coronary heart disease.  
J. Chronic Dis., 29 : 557.
- Crall, F.V.; and Roberts, W.C. (1978) :  
The extramural and intramural coronary arteries in juvenyl diabetes mellitus.  
Analysis of nine necropsy patients aged 19 to 38 with onset of diabetes before age 15 years.  
Am. J. Med., 64 : 221-230.
- Dahlen, G.; Berg, K.; and Frick, M.H. (1976) :  
Lipoprotein, prebetalipoprotein, serum lipids and atherosclerotic disease.  
Clin. Genet., 9 : 556-558.
- Dawber, T.R. (1958) :  
An epidemiologic study of heart disease : The Framingham Study.  
Nutr. Rev. (Jan), 16 : 1.

Dawber, T.R. (1963) :

Coronary heart disease.  
Bibl. Cardiol., 13 : 9.

DeBacker, G.; Rosseneu, M.J. and Deshypere, J.P. (1982):

Discriminative value of lipids and apoproteins  
in coronary heart disease.  
Atherosclerosis (Shannon, Irel.), 42 : 197.

Deeley, R.G.; Tam, S.P.; and Archer, T.K. (1985) :

The effects of estrogen on apolipoprotein  
synthesis.  
Can. J. Biochem. Cell Biol., 63 : 882.

DeFronzo, R.A. and Ferrannini, E. (1982) :

The pathogenesis of non-insulin dependent  
diabetes : an update Medicine (Baltimore).  
61 : 125-140.

Deutscher, S.; Epstein, F.H.; Kjelsberg, M.O. (1966) :

Familial aggregation of factors associated  
with coronary heart disease.  
Circulation, 33 : 911.

Dolphin, P.J. (1985) :

Lipoprotein metabolism and the role of apo-  
lipoproteins as metabolic programmers.  
Can. J. Biochem. Cell Biol. 63 : 850.

- Ducimetiere, D.; Eschwege, E.; Papoz, L.; Richard, J.L.; Claude, J.R.; and Rosselin, G.(1980):  
Relationship of plasma insulin levels to the incidence of myocardial infarction and coronary heart disease mortality in middle aged population.  
Diabetologia, 19 : 205-10.
- Dunn, F.L.; Bilheimer, D.W.; and Grundy, S.M. (1980):  
Pathogenesis of hypertriglyceridemia in diabetes mellitus . The effect of improved glucose regulation.  
Clin. Res., 28 : 391 A.
- Dyer, A.R.; Stamler, J.; Berkson, D.M.; and Lindberg H.A. (1975) :  
Relationship of relative weight and body mass index to 14 year mortality in the Chicago Peoples Gas Company.  
J. Chronic Dis., 28 .1.109.
- Egusa, G.; Beltz, W.F.; Grundy, S.M.; and Howard, B.V. (1985) :  
Influence of obesity on the metabolism of apolipoprotein B in humans.  
J. Clin. Invest., 76 : 596-603.
- Eisenberg, S. (1983) :  
Lipoproteins and lipoprotein metabolism.  
Klin. Wochenschr, 61. 119-132.

- Eisenberg, S.; Bilheimer, D.W.; and Levy, R.I. (1973):  
On the metabolic conversion of human plasma  
VLDL to LDL.  
Biochem. Biophys. Acta, 326 : 361-377.
- Enger, S.C.; Herbjrsen, K.; Erikssen, J.; Frettand,  
A. (1977) :  
High density lipoproteins and physical acti-  
vity : the influence of physical exercise ,  
age and smoking on HDL-cholesterol and the  
HDL/total cholesterol ratio.  
Scand., J. Clin. Lab. Invest. 37 (3) : 251-5.
- Epstein, F.H. (1964) :  
Hereditary aspects of coronary heart disease.  
Am. Heart J. 67 : 445.
- Erikssen, J.; and Enger, S.C. (1978) :  
The effect of smoking on selected coronary  
heart disease risk factors in middle aged men.  
Acta. Med. Scand. 203 (1-2) : 27-30.
- Fielding, C.J.; and Havel, R.J. (1977) :  
Lipoprotein lipase.  
Arch. Pathol. Lab. Med., 101 : 225-229.
- Flegg, H.M. (1973) :  
An investigation of the determination of serum  
cholesterol by an enzymatic method.  
Annu. Clin. Biochem., 10 : 79.

Fredrickson, D.S.; Levy, R.I.; and Lees, R.S. (1967):  
 Fat transport in lipoproteins-an integrated  
 approach to mechanisms and disorders.  
 N. Engl. J. Med., 276 : 148-56.

Fredrickson, D.S.; Goldstein, J.L. and Brown, M.S.(19780):  
 The familial hyperlipoproteinaemia. In  
 standburg, J.B.; Wyngaarden, J.B.; and  
 Fredrickson, D.S. (eds.) : Metabolic Basis  
 of Inherited Diseases. 4th edition, New York  
 McGraw-Hill, PP. 604-655.

Fredrickson, D.S.; Goldstein, J.L.; and Brow, M.S.(1980):  
 The familial hyperlipoproteinaemias. In  
 standburg, J.B.; Wyngaarden, J.B. and Fredrickson,  
 D.S. (eds.) : Metabolic basis of inherited  
 Diseases. 4th edition, New York, McGraw - Hill,  
 pp. 604-655.

Friedberg. C.K. (1966) :  
 Pathogenesis of Atherosclerosis. In diseases  
 of the heart, 3rd edition. WB Sanders Company,  
 Philadelphia, pp. 651-675.

Friedman, M. St.; George, S.; Byers, S.O.; and Roseman,  
 R.H. (1960) :  
 Excretion of catecholamines, 17-ketosteroids,  
 17-hydroxycorticoids and 5-hydroxyindole in  
 men exhibiting a particular behaviour pattern  
 (A) associated with high incidence of clinical  
 coronary artery disease.  
 J. Clin. Invest., 39 : 758.

Friedman, M.; Thoresen, C.E.; and Gill, J. (1980):  
 Type A Behavior : Its Possible Role, Detection,  
 and Alteration in Patients with Ischaemic  
 Heart Disease, in J.W.Hurst (ed.), "Update V :  
 The Heart " McGraw-Hill, New York, p. 87.

Froelicher, V.F.; and Oberman, A. (1972) :  
 Analysis of Epidemiologic Studies of Physical  
 Inactivity as a Risk Factor for Coronary  
 Artery Disease, Prog. Cardiovasc. Dis., 15 : 41.

Fruchart, J.C. (1982) :  
 Rev. France des Laboratoires., 103 : 7.

Fuller, J.H.; Shipley, M.J.; Rose, G.; Jarrett, R.J.;  
 and Keen, H. (1980) :  
 Coronary heart disease risk and impaired  
 glucose tolerance.  
 Lancet, 1 : 1373 - 1376.

Ganda, O.M.P. (1980) :  
 Pathogenesis of macrovascular disease in the  
 human diabetic.  
 Diabetes 29 : 931-942.

Garcia, M.J.; McNamara, P.M.; Gordon, T.; Kannel, W.B.  
 (1979) :  
 Morbidity and mortality in diabetes in the  
 frarningham population. Sixteen year follow-up  
 study.  
 Diabetes, 23 : 105-111.

- Gidez, L.I.; Miller, G.J.; Burstein, M.; Slagle, S.;  
Eder, H.A. (1982) :  
Separation and quantitation of subclasses  
of human plasma high density lipoproteins  
by a simple precipitation procedure.  
J. Lipid Res., 23 : 1206-23.
- Glass, A.R.; Burman, K.D.; Dahms, W.T. and Boehm ,  
T.M. (1981) :  
Endocrine function in human obesity.  
Metabolism 30 : 89-104.
- Glueck, G.J., Taylor , H.I.; Jacobs, D. et al. (1980):  
Plasma high density lipoprotein cholesterol.  
Association with measurement of body mass.  
The Lipid Research Clinics Program Prevalence  
Study.  
Circulation 62 (Suppl. 4 ) : 62-69.
- Goldstein, J.L.; and Brown, M.S. (1977) :  
The LDL pathway and its relation to atheros-  
clerosis.  
Annu-Rev. Biochem., 46 : 897-930.
- Goldstein, J.L.; and Brown, M.S. (1983) :  
Familial hypercholesterolaemia. In : The  
metabolic basis of inherited disease. .  
New York, McGraw-Hill Book Co., Inc., p . 672.

- Goldstein, J.L., Hazzard, W.R.; Schrott, H.G.;  
Bierman, E.L.; Motulsky, A.G. (1973) :  
Hyperlipidaemia in coronary heart disease.  
Lipid levels in 500 survivors of myocardial  
infarction.  
J. Clin. Invest., 52 : 1533-43.
- Gonen, B.; Baenzrger, J.; and Schonfield, G. (1981):  
Non enzymatic glycosylation of low density  
lipoproteins in vitro. Effects on cell  
interactive properties.  
Diabetes, 30 : 875-878.
- Gordon, T.; Kannel , W.B.; Dawber, T.R.; McGee, D.  
(1975) :  
Changes associated with quitting smoking :  
The Framingham Study.  
Am. Heart J., 90 : 322.
- Gordon, T.; Castelli, W.P.; Hjortland, M.C.; Kannel,  
W.B.; and Dawber, T.R. (1977) :  
High density lipoprotein as a protective  
factor against coronary heart disease : The  
Framingham Study.  
Am. J. Med. 62 : 707-714.
- Gotto, A.M. (1986) :  
Interactions of the major risk factors for  
coronary heart disease.  
The Am. J. Med., 80 (Suppl. 2 A ) : 48-55.

- Greenfield, M.; Kolterman, D.; and Olefsky, J. (1980):  
Mechanism of hypertriglyceridaemia in  
diabetic patients with fasting hyperglycaemia.  
Diabetologia, 18 : 441-446.
- Grimm, R.H.; and Hunninghake, D.B. (1986) :  
Lipids and Hypertension : implications of new  
guidelines for cholesterol management in the  
treatment of hypertension.  
Am. J. Med., 80 (Suppl 2 A ) : 56-63.
- Havekes, L.; Van Gent, C.M.; Stegerhock, C.L.; Arntzenius,  
A.C.; and Hesse, L.W. (1981) :  
High density lipoprotein cholesterol and apo-  
lipoprotein A-I levels in 32-33 year-old women  
on steroid contraceptives - differences between  
two frequently used low oestrogen pills.  
Clin. Chem. Acta., 116 : 223.
- Havel, R.J. (1976) :  
Lipid transport and the availability of insulin.  
Horm. Metab. Res., 8 (suppl 4) : 51-53.
- Havel, R.J. (1982) :  
Familial dysbetalipoproteinaemia. New aspects of  
pathogenesis and diagnosis.  
Med. Clin. North Am., 66, 441-454.
- Havel, R.J.; Eder, H.A.; and Bragdon, J.H. (1955) :  
The distribution and chemical composition of  
ultracentrifugally separated lipoproteins in  
human serum.  
J. Clin. Invest., 34 : 1345-1353.

- Haynes, S.G.; Feinleib, M.; and Kannel, W.B. (1980) :  
The Relationship of Psychosocial Factors to  
Coronary Heart Disease in the Framingham  
Study : III. Eight-Year Incidence of Coronary  
Heart Disease.  
Am. J. Epidemiol., 3 : 37.
- Hill, A.B. (1979) :  
In " A short Text book of Medical Statistics  
10th edition" The English Language Book  
Society and Hodder and Stoughton.
- Howard, B.V.; Savage, P.J.; Bennion, L.J.; and Bennett,  
P.H. (1978) :  
Lipoprotein concentration in diabetes mellitus.  
Atherosclerosis, 30 : 153-62.
- Hubert, H.B.; Feinleib, M.; McNamara, P.M.; Castelli,  
W.P. (1983) :  
Obesity as an Independent Risk Factor for  
Cardiovascular Disease : A 26-year follow-up  
of Participants in the Framingham Heart Study.  
Circulating, 67 : 968-877.
- Hulley, S.B.; Rosenman, R.H.; Bawol, R.D.; and Brand,  
R.J.; (1980) :  
The association between triglyceride and  
coronary heart disease.  
N. Engl. J. Med., 302 : 1383-1389.

Idem ( 1984) :

Persistent problems in the pathogenesis of atherosclerosis.

Arteriosclerosis 4, 443-451.

Innerarity, T.L.; Pitas, R.E., and Mahley, R.W.(1982):

Modulating effects of canine HDL on cholesterol ester synthesis induced by B-VLDL in macrophages.

Arteriosclerosis 2 : 114.

Ishikawa, T.; Fidge, N.; Thelle, D.S.; Forde, O.H.;

Miller, N.E. (1978) : The Tromso Heart Study,

Serum apolipoprotein A I concentration in relation to future coronary heart disease.

Eur. J. Clin. Invest., 8 : 179-82.

Janus, E.D.; Nicoll, A.; Wooton, R.; Turner, P.R.;

Magill, P.J.; and Lewis, B. (1980) :

Quantitative studies of VLDL conversion to LDL in normal controls and primary hyperlipidemic states.

Eur. J. Clin. Invest., 10 : 149-159.

Jarrett, J. (1977) :

Diabetes and the heart : coronary heart disease.

Clin. Endocrinol. Metabol., 6 : 389-402.

Jarrett, R.J. (1984) :

Type 2 (non-insulin-dependent) diabetes  
mellitus and coronary heart disease-  
chicken, egg or neither ?  
Diabetologia 26 : 99-102.

Jenkins, C.D. (1976) :

Recent Evidence Supporting Psychologic  
and Social Risk Factors for Coronary  
Disease, N. Engl  
N. Engl. J. Med., 294 : 987, 1033.

Kannel, W.B.; and McGee, D.L. (1979) :

Diabetes and cardiovascular disease : The  
Framingham study.  
JAMA, 241 : 2035-8.

Kannel, W.B.; Hjortland, M.C.; McNamara, P.M.; and  
Gordon, T. (1976) :

Menopause and risk of cardiovascular  
disease.  
Ann. Intern. Med., 85, 447-452.

Kannel, W.B.; Castelli, W.P.; Gordon, T. (1979-a):

Cholesterol in the prediction of atheros-  
clerotic disease. New perspectives based  
on the Framingham Study.  
Ann. intern. Med., 90 : 85-91.

- Kannel, W.B.; Gordon, T.; and Castelli, W.P. (1979-b):  
Obesity, lipids and glucose intolerance. The  
Framingham Study.  
Am. J. Clin. Nutr., 32 : 1238-1245.
- Karakash, C.; Assimaco-Poulos-Jeanne, F.; Jeanrenaud,  
B. (1976) :  
An anomaly of insulin in perfused livers of  
obese hyperglycemic mice.  
J. Clin. Invest., 57 : 1117-1124.
- Karathanasis, S.K.; Zannis, V.I.; Breslow, J.L. (1983):  
Isolation and characterization of the human  
apoprotein A I gene.  
Pro. Natl. Acad. Sci., (USA) 8 : 6147-6151.
- Kennedy, A.L.; Lippin, T.R.; and Lavery, T.D. (1978) :  
Relation of high density lipoprotein cholesterol  
concentration to type of diabetes and  
its control.  
Brit. Med. J., 2 : 1191-1194.
- Kissenbah, A.H.; Alfarsi, S.; Evans, D.J. and Adams  
P.W. (1982) :  
Integrated regulation of very low density  
lipoprotein in triglyceride and apolipoprotein  
B kinetics in non insulin dependent diabetes  
mellitus.  
Diabetes 31 : 217-25.

- Klein, R.; Klein, B.E.; Coroni, J.C.; Moready, J.;  
Cassel, J.C.; Tyroles, H.A. (1973) :  
Serum uric acid. Its relationship to  
coronary heart disease risk factors and  
cardiovascular disease, Evans County,  
Georgia. Arch. Intern. Med., 132 : 401.
- Kunkel, H.G.; and Slater, R.J. (1952) :  
Separation of lipoprotein using electro-  
phoresis.  
J. Clin. Invest., 31, 677-684.
- Kushwaha, R.S.; Hazzard, W.R.; Gagne, G.; Chait, A.;  
and Albers, J.J. (1977) :  
Type III hyperlipoproteinaemia : Pasadoxical  
hypolipidaemic response to estrogen.  
Ann. Intern. Med., 87, 515-525.
- Leren, P.; and Helgeland, A. (1986) :  
Coronary heart disease and treatment of hyper-  
tension : Som Oslo Study Data.  
Am. J. Med., 80 (Supple 2 A ) : 3.
- Leren, P.; Helgeland, A.; Hjermann, I.; Holme, I. (1983) :  
The Oslo study : CHD risk factors, socioecono-  
mic influences, and intervention.  
Am. Heart J., 106 : 1200.
- 7

Levy R.I. (1983) :

Proceedings of the Symposium Hyperlipoproteinemia : Dietary and Pharmacologic Intervention, Washington, DC, May 23, pp.1-4.

Lewis, B.; Mancini, M.; Mattock, M.; Chait, A., and

Froser, T.R. (1972) :

Plasma triglyceride and fatty acid metabolism in diabetes mellitus.

Eur. J. Clin. Invest. 2 : 445-53.

Lindgren, F.T. (1980) :

The plasma lipoproteins : Historical developments and nomenclature.

Ann. N.Y. Acad. Sci., 348, 1-15.

Lippel, K.; Tyroler, H.A.; Gotto, A. Jr.; et al.(1983) :

Workshop on apolipoprotein quantification.

Arteriosclerosis, 3 : 452-64.

Maciejko, J.J.; Holmes, D.R.; Kottke, B.A.; Zinsmeister, A.R.; Dinh, D.M.; Mao, S.J.T. (1983) :

Apolipoprotein A-I as a marker of angiographically assessed coronary artery disease.

N. Engl. J. Med., 309 : 385-9.

MacMahon, S.W.; MacDonald, G.J.; and Blacket, R. B.(1985):

Plasma lipoprotein levels in treated and untreated hypertensive men and women. The National Heart Foundation of Australia Risk Factor

Prevalence Study.

Arteriosclerosis 5 (4) : 391-396.

Mahley, R.H. (1981) :

Cellular and molecular biology of lipoprotein metabolism in atherosclerosis.

Diabetes 30 (Suppl. 2 ) 60-65.

Mahley, R.W. (1982) :

Atherogenic hyperlipoproteinaemia.

Med. Clin. North, Am., 66 : 375.

Mahley, R.W. (1983) :

Development of accelerated atherosclerosis.

Arch. Pathol. Lab. Med., 107 : 393.

Mahley, R.W. (1985) :

Atherogenic lipoproteins and coronary artery disease.

Circulation 72 (5) : 943-948.

Mahley, R.W.; and Innerarity, T.L. (1983) :

Lipoprotein receptors and cholesterol homeostasis.

Biochem. Biophys. Acta, 737 : 197.

Mahley, R.W.; and Angelin, B. (1984) :

Type III hyperlipoproteinemia.

Adv. Intern. Med., 29 : 385.

- Mahley, R.W.; Innerarity, T.L.; Rall, S.C. Jr.; and Weisgraber, K.H. (1984) :  
Plasma lipoproteins : apolipoprotein structure and function.  
J. Lipid Res., 25 : 1277.
- Mahley, R.W.; Innerarity, T.L.; Rall, S.C. Jr.; and Weisgraber, K.H. (1985) :  
Lipoproteins of special significance in atherosclerosis.  
Ann. N.Y. Acad. Sci., 454 : 209.
- Mazzaferri, E.L.; Starich, G.H.; Lardinois, C.K., and Bowen, D. (1985) :  
Gastric inhibitory polypeptide responses to nutrients in Caucasians and American Indians with Obesity and Non-insulin dependent diabetes mellitus.  
J. Clin. Endocrinol. Metabol.; 61 (2) : 313-32).
- McGuire, E.A.; Tobin, J.D.; Berman, M. and Andres, R. (1979) :  
Kinetics of native insulin in diabetic obese and aged men.  
Diabetes 28 : 110-120.
- McNamara, J.J.; Malot, M.A.; Stermple, J.F.; and Cutting, R.T. (1971) :  
Coronary artery disease in combat casualties in Vietnam.  
JAMA 216 : 1185-1187.

- Meade, T.W.; Chakrabarti, R.; Haines, A.P.; North  
W.R.S.; Stirling, Y. (1979) :  
Characteristics affecting fibrinolytic  
activity and fibrinogen concentrations.  
Br. Med. J., 1 : 152.
- Meadelson, J.H.; Mello, N.K. (1973) :  
Alcohol-induced hyperlipidaemia and beta-  
lipoproteins.  
Science 180 : 1372.
- Miller, N.E. (1982) :  
Coronary atherosclerosis and plasma lipo-  
proteins : epidemiology and pathophysiologic  
considerations.  
J. Cardiovasc. Pharmacol., 4 (suppl. 2) :  
5195-5195.
- Miller, N.E.; Hammett, F.; Saltissi, S., et al. (1981):  
Relation of angiographically defined coronary  
artery disease to plasma lipoprotein subfrac-  
tions and apolipoproteins.  
Br. Med. J., 282 : 1741-4.
- Morton, N.E., Gulbrandsen, C.L.; Rao, D.C.; Rhoads,  
G.G.; Kagan, A. (1980) :  
Determinants of blood pressure in Japanese ;  
American families.  
Hum. Genet., 53 : 261-6.

Multiple Risk Factor Intervention Trial Research  
Group (1982) :

Multiple risk factor intervention trial :  
Risk factor changes and mortality results.  
JAMA , 248 : 11465.

Murray, R.G.; Tweddel, A.; Third, J.L.H.C.; Hutton,  
I.; Hillis, W.S.; Lorimer, A.R.; and Lawrie,  
T.D.V. (1975) :  
Relation between extent of coronary artery  
disease and severity of hyperlipoproteinaemia  
Brit. Heart J., 37-1205.

Myers, A.R.; Epstein, F.H.; Dodge, H.S.; Mikkelsen,  
W.M. (1968) :  
The relationship of serum uric acid to risk  
factors in coronary heart disease.  
Am. J. Med., 95 : 520.

Myers, L.H.; Phillips, N.R.; and Havel, R.J. (1976) :  
Mathematical evaluation of methods for  
estimation of the concentration of the major  
lipid components of human serum lipoproteins.  
J. Lab. Clin. Med., 88 : 491-505.

National Diabetes Data Group (1979) :  
Classification and diagnosis of diabetes  
mellitus and other categories of glucose  
intolerance.  
Diabetes 28 : 1039-59.

- Nayak, R.V.; Feldman, E.B.; and Carter, A.C. (1962) :  
Adipokinedic effect of intravenous cortisol  
in human subjects.  
Proc. Soc. Exp. Biol. Med., 111 : 682-686.
- New, M.I.; Roberts, T.W.; Bierman, E.L.; and Reader,  
G.G. (1963) :  
The significance of blood lipid atterations  
in diabetes mellites.  
Diabetes 12 : 208-12.
- Nikkila, E.A. (1973) :  
Triglycride metabolism in diabetes.  
Prog. Biochem. Pharmacol., 8 : 271-299.
- Nikkila, E.A. (1981) :  
High density lipoproteins in diabetes.  
Diabetes 30 : 82 - 87.
- Nikkila, E.A. and Kekki, M. (1973) :  
Plasma triglycerides transport kinetics in  
diabetes mellitus.  
Metabolism 22 : 1-22.
- Nikkila, E.A.; Hormila, P. (1978) :  
Serum lipids and lipoproteins in insulin  
treated diabetes.  
Diabetes ; 31 (Suppl 5) : 1-81.

Nikkila, E.A.; Huttunen, J.K.; and Ehnholm, C.(1977):  
 Posthepatin plasma lipoprotein lipase and  
 hepatic lipase in diabetes mellitus.  
 Relationship to plasma triglyceride metabolism.  
 Diabetes 26 : 11-21.

Noel, S.P.; Wong, P.; Dolphin, P.J.; Dory, L.; and  
 Rubinstein, D. (1979) :  
 Secretion of cholesterol-rich lipoproteins by  
 perfused livers of hypercholesterolaemic rats-  
 J. Clin. Invest., 64 : 674.

Noppa, H. (1980) :  
 Body weight changes in relation to incidence  
 of ischaemic heart disease and change in risk  
 factors for ischaemic heart disease.  
 Am. J. Epidemiol., 111 : 693.

O'Hara, D.D.; Porte, D.; and Williams, R.H. (1966 ) :  
 The effect of diet and thyrosin on plasma  
 lipids in myxedema.  
 Metab. Clin. Exp., 15 : 123.

Olefsky, J.M. (1980) :  
 Obesity in : Isselbacher, K.J., Adams, R.D.,  
 Braunwald, E.; Wilson, J.D. (eds).  
 Harison's principle of internal medicine,  
 International students edition, 9th ed.  
 McGrow-Hill, Koga Kusha, 411.

Olefsky, J.M. (1981) :

Insulin resistance and insulin action in an  
in vitro and in vivo perspective.

Diabetes 30 : 148-162.

Orchard, T.J.; Rodgers, M.; and Hedley, A.J. (1981):

Serum lipids in a teenage population :  
Geographic , seasonal , and environmental  
factors.

Int. J. Epidemiol., 10 : 161-170.

Paisey, R.; Elkeles, R.S.; and Hanobley, J. (1978) :

The effects of chlorpropomide and insulin  
on serum lipids, lipoproteins and fractional  
triglyceride removal.

Diabetologia 15 : 81-89.

Pan, W.H.; Cedres, L.B.; Liu, K.; Dayer, A.; James, A.;

Schoenberger, J.A.; Shekelle, R.B.; Stamler,  
R.; Smith, D.; Collette, P.; and Stamler J.

(1986) :

Relationship of clinical diabetes and asymptomatic  
hyperglycaemia to risk of coronary heart  
disease mortality in men and women.

Am. J. Epidemiol., 123 : 604-516.

Pearson, T.A.; Buikley, B.H.; Achaff, S.C.; Kwiterovsch,  
P.O.; Gordis, L. (1979) :

The association of low levels of HDL cholesterol  
and arteriographically defined coronary artery  
disease.

Am. J. Epidemiol., 100 : 285-95.

Phillips, G.B. (1984) :

Evidence for Hyperestrogenemia . As the link between Diabetes Mellitus and Myocardial Infarction.

Am. J. Med., 76 : 1041-48.

Phillips, N.R.; Havel, R.J.; and Kane, J.P. (1981):

Levels and interrelationships of serum lipoprotein cholesterol and triglycerides.

Association with adiposity and consumption of ethanol, tobacco and beverages, containing coffeine.

Arteriosclerosis 1 : 13-24.

Philips, G.B.; Castelli, W.P.; Abbot, R.D.; and McNamara, P.M. (1983) :

Association of hyperestrogenemia and coronary heart disease in man in Framingham Cohort.

Am. J. Med., 74, 863-869.

Pietri, A.; Dunn, F.L.; and Raskin, P. (1980) :

The effects of diabetic control on plasma lipid and lipoprotein levels : A comparison of conventional therapy and continuous subcutaneous insulin infusion.

Diabetes 29 : 1001-1005.

Pitas, R.E.; Innerarity, T.L.; and Mahley, R.W. (1983):

Foam cells in explants of atherosclerotic rabbit aortas have receptors for B-VLDL.

Arteriosclerosis 3 : 2.

Pyorala, K. (1985) :

Diabetes and coronary heart disease.  
Acta endocrinologica , 110 (Suppl. )  
272 : 11-19.

Rabinwitz, D. (1970) :

Some endocrine and metabolic aspects of  
obesity.  
Ann. Rev. Med., 21 : 241-258.

Reaven, G.M.; Greenfield, M.S.(1981) :

Diabetic hypertriglyceridaemia : Evidence  
for three clinical syndromes.  
Diabetes, 30 : 66-75.

Rechritzer, P.A. et al. (1983) :

Relation of exercise to the recurrence rate  
of myocardial infarction in men : Ontario  
Exercise Heart Collaborative Study.  
Am. J. Cardiol., 51 : 65.

Rissanen, A.M. (1979) :

Familial aggregation of coronary heart disease  
in a high incidence area (North Karelia,  
Finland).  
Br. Heart. J., 42 : 294-303.

Rissanen, A.M., Nikkila, E.A. (1977) :

Coronary artery disease and its risk factors  
in families of young men with angina pectoris  
and in controls.  
Br. Heart J., 39 : 875.

Rissanen, A.M.; Nikkila, E.A. (1979) :

Aggregation of coronary risk factors in families of men with fatal and non fatal coronary heart disease.

Br. Heart J., 42 : 373-80.

Ross, R. (1986) :

Coronary Heart Disease : Factors influencing atherosclerosis, in Hurst, J.W. : the heart McGraw-Hill Co., New York, 801-816.

Ross, R.; and Glomset, J.A. (1976) :

Pathogenesis of atherosclerosis.

N. Engl. J. Med., 295 : 369-420.

Santen, R.J.; Williams, P.W.; and Fajans, S.S. (1972) :

Atherosclerosis in diabetes mellitus : correlation with serum lipid levels, adiposity and serum insulin levels.

Arch. Intern. Med., 130 : 833-43.

Saudek, C.D.; and Yong, N.L. (1981) :

Cholesterol metabolism in diabetes mellitus.

The role of diet.

Diabetes 30 (Suppl : 2) : 26-81.

Schonfeld, G.; Birge, C.; Miller, J.P.; Kessler, G.

and Santiago, J. (1974) :

Apolipoprotein B levels and altered lipoprotein composition in diabetes.

Diabetes 23 : 827-834.

Schrott, H.G.; Clarke, W.R.; Wiebe, D.A.; Connor, W.E.; Lauer, R.M. (1979) :  
Increased coronary mortality in relatives of hypercholesterolaemic school children : The Muscatine study.  
Circulation 59 : 320.

Segrest, J.P.; Jackson, R.L.; Morrisett, J.D.; and Gotto, A.M. (1979) :  
A molecular theory of lipid-protein interactions in the plasma lipoproteins.  
FEBS Lett., 38 : 247-253.

Shaw, L.W. (1981) :  
Effects of a prescribed supervised exercise program on mortality and cardiovascular morbidity in patients after a myocardial infarction : The National Exercise and Heart Disease Project.  
Am. J. Cardiol., 48 : 39.

Shen, B.W.; Scanu, A.M.; and Kezoly, F.J. (1977)  
Structure of human serum lipoproteins inferred from compositional analysis.  
Pro. Natl. Acad. Sci. (USA), 74 : 837-841.

Shennan, N.M.; Seed ?. and Wynn, V. (1985) :  
Variation in serum lipid and lipoprotein levels associated with changes in smoking behaviour in non-obese caucasian males.  
Atherosclerosis 58 (1-3). 17-25.

Sholtz, R.I.; Rosenman, R.H.; Brand, R.J. (1975):

The relationship of reported parental history to the incidence of coronary heart disease in the Western Collaborative Group Study.  
Am. J. Epidemiol., 102 : 350 - 6.

Simpson, R.W.; Mann, J.I.; Hockaday, T.D.R.; Hockaday, J.M.; Turner, R.C.; and Jelts, R. (1979):

Lipid abnormalities in untreated maturity onset diabetics and the effect of treatment.  
Diabetologia 16 : 101-6.

Slack, J.; Evans, K.A. (1966) :

The increased risk of death from ischaemic heart disease in first degree relatives of 121 men and 96 women with ischaemic heart disease.  
J. Med. Genet., 3 : 239.

Slone, D. (1981) :

Risk of myocardial infarction in relation to current and discontinued use of oral contraceptives.  
N. Engl. J. Med., 305 : 402.

Sniderman, A.; Teng, B.; Jerry, M. (1975) :

Determination of B protein of low density lipoprotein directly in plasma.  
J. Lipid Res., 16 : 465-7.

Sniderman, A.; Shapiro, S.; Marpole, D.; Skinner, B.;  
Teng, B.; Kwiterovich, P.O.Jr. (1980) :  
Association of coronary atherosclerosis with  
hyperapobeta-lipoproteinaemia (increased  
protein but normal cholesterol levels in  
human plasma low density (B) lipoproteins).  
Proc. Natl. Acad. Sci.(USA) 77 : 604-8.

Sniderman, A.D.; Wolfson, C.; Teng, B.; Franklin, F.A.;  
Bachorik, P.S.; Kwiterovich, P.O.Jr. (1982) :  
Association of hyperapobetalipoproteinaemia  
with endogenous hypertriglyceridaemia and  
atherosclerosis.  
Ann. Intern. Med., 97 : 833-9.

Sokolow, M. and McIlroy, M.B. (1986) :  
Clinical cardiology 4th edition.  
Lange Medical Publications, p. 132-208.

Sosenko, J.M.; Breslow, J.L.; Miettinen, O.S.; and  
Gabbay, K.H. (1980) :  
Hyperglycaemia and plasma lipid levels.  
N. Engl. J. Med., 302 : 650-654.

Stamler, J. (1979) :  
Research related to risk factors.  
Circulation 160 : 575.

- Stanford, B.A., Malter, S.; Fell, R.D.; Sady, S.;  
Papanek, P.; Cresanta, M. and Louisville,  
K.Y. (1984) :  
Cigarette smoking exercise and high density  
lipoprotein cholesterol.  
Atherosclerosis 52 (1) : 73-83.
- Steinberg, D. (1983) :  
Lipoproteins and atherosclerosis : a look  
back and a look ahead.  
Arteriosclerosis 3 : 383.
- Sterky, G.; Larsson, Y.; and Persson, B. (1963)  
Blood lipids in diabetic and non diabetic  
school children.  
Acta Paediatr. Scand., 52 : 11-21.
- Strong, J.P.; and McGill, H.C.Jr. (1982) :  
The natural history of coronary atherosclerosis.  
Am. J. Pathol., 40 : 37-49.
- Surgeon General (1983) :  
The health consequence of smoking U.S.  
Department of Health and Human Services  
No. DHAS (PHS) 89-50204 : 6-7.
- Takkunen, H.; Reunanen, A. (1977) :  
Hyperuricaemia and other cardiovascular risk  
factors.  
Adv. Exp. Med. Biol., 76 B : 238.

Tan, M.H.; Macintosh, W.; Weldon, K.L.; Kapoor, A.;  
 Chandler, B.M.; and Hindmarsh, T.J. (1980):  
 Serum HDL cholesterol in patients with  
 abnormal coronary arteries.  
 Atherosclerosis 37 : 187-188.

Taskinen, M.R.; and Nikkila, E.A. (1979) :  
 Lipoprotein lipase activity of adipose tissue  
 and skeletal muscle in insulin deficient hum  
 diabetics. Relation to high density and very  
 low density lipoproteins and response to  
 treatment.  
 Diabetologia 17 : 315-316.

Thompson, G. (1984) :  
 Apoproteins : determinants of lipoprotein  
 metabolism and indices of coronary risk.  
 Br. Heart. J., 51 : 585-588.

Trinder, P. (1969) :  
 Determination of glucose in blood using  
 glucose oxidase with an alternative oxygen  
 acceptor.  
 Ann. Clin. Biochem., 6 : 24.

University Group Diabetes Program (1982) :  
 Effects of hypoglycemic agents on vascular  
 complications in patients with adult-onset  
 diabetes.  
 Diabetes 31 (Suppl 5 ) : 1 - 81.

Uterman, G.; Hees, M.; and Steinmetz, A. (1977)

Polymorphism of apoprotein E and occurrence of dysbetalipoproteinaemia in man.

Nature (London) 269 : 604-607.

Wallentin, L.; and Sundin, B. (1985) :

Aberrations in lipoprotein lipids in men with coronary artery disease and the influence of obesity, smoking and beta-adrenceptor blocking drugs.

Atherosclerosis. Mar. 54 (3) : 241-9.

Warnick, G.R.; Mayfield, C.; Albers, J.J.; Hazzard, W.R. (1979) :

Gel isoelectric focusing method for specific diagnosis of familial hyperlipoproteinaemia Type 3.

Clin. Chem., 25 : 279-84.

Weidman, S.W.; Ragland, J.B.; Fisher, J.N.; Kitabchi, A.E.; and Sabesin, S.M. (1982) :

Effects of insulin on plasma lipoproteins in diabetic ketoacidosis : evidence for a change in high density lipoprotein composition during treatment.

J. Lipid Res., 23 : 171.

Wenger, N.K.; and Sclanl, R.C. (1986) :

Prevention of Coronary Atherosclerosis, in Hurst, J.W.; the heart. McGraw-Hill Co., New York., 817-838.

- Whayne, T.F.; Alaupovic, P.; Curry, M.D.; Lee, E.T.;  
Anderson, P.S.; Schechler, E. (1981) :  
Plasma apolipoprotein B and VLDL, LDL., and  
HDL-cholesterol as risk factors in the  
development of coronary artery disease in  
male patients examined by angiography.  
Atherosclerosis 139 : 411-24.
- William, P.N.; David, S.F.; Antonie, W.V.; Paul, D.G.;  
Sathanur, R.S.; James, L.C.; et al. (1986) :  
Relation of serum lipoprotein levels and  
early atherosclerosis.  
N. Engl. J. Med., 16 : 138-144.
- Williams, P.; Robinson, D.; and Bailey, A. (1979) :  
High density lipoprotein and coronary risk  
factors in normal men.  
Lancet 1 (8107) : 72-5.
- Wilson, D.E.; Schreiberman, P.H.; Day, V.C.; and Arky.  
R.A. (1970) :  
Hyperlipidemia in an adult diabetic population  
J. Chronic. Dis., 23 : 501-6.
- Witzman, J.L.; Mahoney, E.M. and Branks, M.J. (1982):  
Non enzymatic glycosylation of low density  
lipoprotein alters its biologic activity.  
Diabetes 31 : 283-291.
- Wynn, V.; Doar, J.W.H.; Mills, G.L.; and Stokes, T.  
(1969) :  
Fasting serum triglyceride, cholesterol and

lipoprotein levels during oral contraceptive therapy.

Lancet ii : 756-760.

Yamamoto, T.; Davis, C.G.; Brown, M.S.; Schneider, W.J.; Casey, M.L.; Goldstein, J.L.; and Russel, D.W. (1984) :

The human LDL receptor. A cysteine-rich protein with multiple alu sequence in its mRNA.

Cell 39 : 27.

Yano, K.; Rhoads, G.; Kagan, A. (1977) :

Epidemiology of serum uric acid among 8000 Japanese-American men in Hawaii.

J. Chronic Dis., 30 : 171.

Yarnel, J.W.; Milbark, J.; Walker, C.L.; Feily, A.M.; Hayes, T.M. (1982) :

Determinant of high density lipoprotein and total cholesterol in women.

J. Epidemiol. Community Health 36(3) :167-71.