

REFERENCES

- 1- *Alicandri, C.; Fariello, R. and Boni, E. (1987) :*
 Left ventricular hypertrophy and ambulatory monitoring of blood pressure.
J. Clin. Hypertens, 3 : 197-202.

- 2- *Badawi, H. : (1991) :*
 Hypertension in Egypt : Control of hypertension in developing countries.
Tropical cardiology, 9 : 27 - 31.

- 3- *Baumgart, P. (1992) :*
 Ambulatory blood pressure monitoring indications, procedure, evaluation, relevance, costs and benefit".
Niern. Hockdruak. Kran. Khcitten, 21 : 5223-8.

- 4- *Bauwens, F.R.; Duprez, D. and De Buyzer, M. (1991) :*
 Influence of the arterial BP and non hemodynamic factors on left ventricular hypertrophy in essential hypertension.
Am. J. Cardiol; 68 925 - 929.

- 5- *Bernardi Ricordi, L.; Lazzari, P.; Solda, P.; Calciati, A.; Ferrari, M.R.; Vandiea, I.; Finardi, G. and Fratino, P. (1992) :*
 Impaired circadian modulation of sympatho-vagal activity in diabetes : a possible explanation for altered temporal anset of cardiovascular disease.
Circulation, 86 : 1443-1452.

- 6- *Bevan, A.J.; Homour, A.J. and Scott, F.H. (1969) :*
 Direct arterial blood pressure recording in unrestricted man.
Clinsci, 36 : 329 - 334.

- 7- *Bikkina, M.; Larson, M.G. and Levy, D. (1993) :*
 A symptomatic ventricular arrhythmias and mortality risk in subjects with left ventricular hypertrophy.
J. Am. Coll. Cardiol., 22 : 1111 - 1116.

- 31- *Devereux, R.B.; Reichek, N. (1977) :*
Echocardiographic determination of left ventricular mass in man anatomic validation of the method.
Circulation; 55 : 613-18.
- 32- *Devereux, R.B.; Savage, D.D.; Sachs, I.; Laragh, J.H. (1983) :*
Relation of hemodynamic load to left ventricular hypertrophy and performance in hypertension.
Am. J. Cardiol, 51 : 171 - 176.
- 33- *Dimond, P.; Drayer, JIM.; Gardin, J.M. and Brewer, D.D. (1997) :*
Ambulatory blood pressure monitoring and left ventricular mass in patients with and without LVH.
Hypertension, 9 : 1161-1167.
- 34- *Dilworth, A.B.; Diloranzo, L. Moccia, D.; Costantino, G. and Bounissimo, S. (1987) :*
Echocardiographic assessment of left ventricular hypertrophy by using body mass as one factor for detection of left ventricular mass index.
Br. Heart, J. Vol. 96, pp. 93-98.
- 35- *Dominiczak, A.F.; Jeffs, B. and Connell, J.M. (1998):*
New genetic concepts in hypertensive cardiovascular disease.
Curr. Opin. Cardiol., 13 (5): 304-11.
- 36- *Douglas, M.L.; Gardin, J.M.; Brewer, D.D. and Weber, M.A. (1988):*
Methods of recognition and assessment of left ventricular hypertrophy.
Am. J. Cardiol, 21 : 161 - 166.
- 37- *Douglas, M.L.; Savage, D.D. Sachs, I. And Laragh, J.H. (1983) :*
Relation of hemodynamic load to left ventricular hypertrophy and performance in hypertension.
Am. J. Cardiol, 51 : 171 - 6.
- 38- *Drayer, JIM, Cardin, J.M.; Brewer, D.D. and Weber, M.A. (1987) :*
Disparate relationships between blood pressure and left ventricular mass in patients with and without left ventricular hypertrophy.
Hypertension, 9 (Supl. II) : 1161 - 1164.

- 39- *Drayer, JIM; Weber, A.M. and Nakamura, D.K. (1985) :*
Automated ambulatory blood pressure monitoring : a study in
age matched normotensive and hypertensive men.
Am. Heart, J. June, Vol. 109, No. 6, pp. 1334 - 1338.
- 40- *Ducaillar, G.; Ribstein, J.; Daures, J.P. and Mimran, A. (1992) :*
Sodium and left ventricular mass in untreated hypertensive and
normotensive subjects.
Am. J. Physiol., 263 : H177-H181.
- 41- *Dunn, F.G.; Oigman, W. and Sundgaard-Rise, K. (1983) :*
Racial difference in cardiac adaptation to essential hypertension
determined by echocardiographic indexes.
J. Am. Coll. Cardiol., 1 : 1348-1351.
- 42- *Egashira, K.; Inou, T. and Hirooka, Y. (1993) :*
Impaired coronary blood flow response to a cetylcholine in
patients with coronary risk factors and proximal atherosclerotic
lesions.
J. Clin. Invest., 91 : 29 - 37.
- 43- *El-Demerdash, F. and Maati, A. (1994) :*
Doppler assessment of right ventricular function in patients
with essential hypertension.
Am. J. Hypertens., 78: 66 - 68.
- 44- *Fagard, R.; Staessen, J.A. and Thijis, L. (1998) :*
The relationship between left ventricular mass and day time
and night time blood pressures.
J. Hypertens, 13 : 823-829.
- 45- *Ford, El : (1976) :*
Heart Size.
Circulation, 39 : 297 - 2 - 303.
- 46- *Fouad, F.M.; Tarazi, R.C. and Gallagher, J.H. (1980) :*
Abnormal left ventricular relaxation in hypertensive patients
Clin. Sci, 59 : 411 - 414.
- 47- *Fouad, F.M.; Tarazi, R.C. and Liebson, P.R. (1987) :*
Echocardiographic studies of regression of left ventricular
hypertrophy in hypertension.
Hypertension (Suppl. II) : 1165-1168.

- 48- **Freis, E.D. (1960) :**
Hemodynamics of hypertension.
Physiol. Rev. 40 : 27 - 52.
- 49- **Frohlich, E.D.; Apstein, C. and Chobanian, A.V. (1992) :**
The heart in hypertension.
N. Engl. J. Med.; 327 : 998 - 1008.
- 50- **Frohlich, E.D. (1987) :**
Overview of haemodynamic and non hemodynamic factors
associated with LVH.
J. Mol. Cell. Cardiol., 21 : 3 - 10.
- 51- **Frohlich, E.D. and Tarazi, R.C. (1979) :**
Is arterial pressure the sole factor responsible for hypertensive
cardiac hypertrophy?
Am. J. Cardiol, 44 : 959 - 63.
- 52- **Fukamizu, A. (1998):**
Functional role of angiotensin II predicted from the phenotypes
of transgenic and knockout mice.
Nippon. Yajirigaku. Zasshi., 112 (3): 213-20.
- 53- **Furlan, R.; Guzzettis, Crivellaro, W.; Dassi, S.; Tinelli, M.; Baselli,
Cerutti, S.; Sleight, P. and Malliani, A. (1990) :**
Continuous (24 hours) assessment of the neural regulation of
systemic arterial pressure and RR variabilities in ambulant
subject.
Circulation, 81 : 537-547.
- 54- **Ganau, A.; Devereux, R.B. and Roman, M.J. (1992):**
Patterns of left ventricular hypertrophy and geometric
remodeling in essential hypertension.
J. Am. Coll. Cardiol., 19 : 1550 - 1558.
- 55- **Ganau, A.; Devereux, R.B.; Roman, M.J. and Schnall, P.;
Pickering, T.G. and Laragh, J.H. (1990) :**
Concentric remodeling of the left ventricle in hypertensive
patients with normal ventricular mass.
JACC, February, Vol. 13, No. 2, p. 245.

- 48- **Freis, E.D. (1960) :**
Hemodynamics of hypertension.
Physiol. Rev. 40 : 27 - 52.
- 49- **Frohlich, E.D.; Apstein, C. and Chobanian, A.V. (1992) :**
The heart in hypertension.
N. Engl. J. Med.; 327 : 998 - 1008.
- 50- **Frohlich, E.D. (1987) :**
Overview of haemodynamic and non hemodynamic factors
associated with LVH.
J. Mol. Cell. Cardiol., 21 : 3 - 10.
- 51- **Frohlich, E.D. and Tarazi, R.C. (1979) :**
Is arterial pressure the sole factor responsible for hypertensive
cardiac hypertrophy?
Am. J. Cardiol, 44 : 959 - 63.
- 52- **Fukamizu, A. (1998):**
Functional role of angiotensin II predicted from the phenotypes
of transgenic and knockout mice.
Nippon. Yajirigaku. Zasshi., 112 (3): 213-20.
- 53- **Furlan, R.; Guzzettis, Crivellaro, W.; Dassi, S.; Tinelli, M.; Baselli,
Cerutti, S.; Sleight, P. and Malliani, A. (1990) :**
Continuous (24 hours) assessment of the neural regulation of
systemic arterial pressure and RR variabilities in ambulant
subject.
Circulation, 81 : 537-547.
- 54- **Ganau, A.; Devereux, R.B. and Roman, M.J. (1992):**
Patterns of left ventricular hypertrophy and geometric
remodeling in essential hypertension.
J. Am. Coll. Cardiol., 19 : 1550 - 1558.
- 55- **Ganau, A.; Devereux, R.B.; Roman, M.J. and Schnall, P.;
Pickering, T.G. and Laragh, J.H. (1990) :**
Concentric remodeling of the left ventricle in hypertensive
patients with normal ventricular mass.
JACC, February, Vol. 13, No. 2, p. 245.

- 56- *Gardin, J.M.; Henery, W.L. Savage, D.D.; Ware, J.H.; Burn, C. and Borer, J.S. (1979) :*
Echocardiographic measurements in normal subjects :
evaluation of an adult population without clinically apparent
heart disease.
J. Clin. Ultrasound, 7: 439 - 447.
- 57- *Gardin, J.M.; Savage, D.D.; Ware, J.H. and Henry, W.L. (1987) :*
Effect of age, sex and body surface area on echocardiographic
left ventricular wall mass in normal subjects.
Hypertension, 9 (Suppl. II) : 1136 - 1139.
- 58- *Ghali, J.K. Kadakia, S.; Cooper, R.S. and Liao, Y.L. (1991) :*
Impact of left ventricular hypertrophy an ventricular
arrhythmias in the absence of coronary artery disease.
J. Am. Coll. Cardiol., 17 : 1277 - 1282.
- 59- *Glock, F.L.; Cox, J.; O'Malley, K.; Atkins, N. and O'Brien, E. (1956) :*
Reproducibility and variability of casual blood pressure data.
J. Hypertension, December, Vol. 9, Suppl. 3, pp. S73-S78.
- 60- *Goose, P.; Ansoborlo, P. Lemetayer, P. and Clementy, G. (1998) :*
Day-time and night - time blood pressure monitoring in
hypertensive patients.
Am. J. Hypertens, 9 : 269 - 272.
- 61- *Gordon, T.; Sorlie, P. and Kannel, W.B. (1976) :*
Problems in the assessment of blood pressure : the framingham
study.
Int. J. Epidemiol, 5 : 327-334.
- 62- *Grossman, G.; Roudaut, R.; Reynaud, P.; Jullien, E. and
Dalocchio, M. (1975) :*
Methods of recognition and assessment of left ventricular
hypertrophy.
Am. J. Hypertens, 6 : 731 - 734.
- 63- *Hamond, I.W.; Devereux, R.B. and Alderman, M.H. (1986) :*
The prevalence and correlates of echocardiographic left
ventricular hypertrophy among employed patients with
uncomplicated hypertension.
J. Am. Coll. Cardiol., 7 : 639 - 650.

- 64- *Hamond, I.W.; Giorgi, D.M.A.; Omboni, S.; Parati, G. and Mancina, G. (1990) :*
Persistent blood pressure increase induced by increases salt intake.
J. Hypertens. Vol. 6, pp. 495 - 499.
- 65- *Hartford, M.; Wikstrand, J.C.; Wallentin, I.; Ljungman, S.M. and Berglund, G.L. (1985) :*
Left ventricular wall stress and systolic function in untreated primary hypertension.
Hypertension, 7 : 97 - 104.
- 66- *Hatle, L. (1993) :*
Deoppler echocardiographic evaluation of diastolic function in hypertensive cardiomyopathies.
Eur. Heart. J., 14 (Suppl. J) : 88 - 94.
- 67- *Henry, S.; Roudaut, R.; Reynoud, P. and Jullien, E. (1978) :*
Relationship between left ventricular mass, body surface area and other parameters.
Am. J. Hyperten, 7 : 431 - 434.
- 68- *Hoegholm, A.; Kristensen, K.S.; Bang, L.E.; Nielsen, J.W.; Nielsen, W.B. and Madsen, N.H. (1993) :*
Left ventricular mass and geometry in patients with established hypertension and white coat hypertension.
Am. J. Hypertens. 6 : 282 - 286.
- 69- *Houghton, J.L.; Frank, M.J.; Carr, A.A.; Von Dohlen, T.W. and Prisant, L.M. (1990) :*
Relation among coronary flow reserve, left ventricular hypertrophy and thaliw perfusion defects in hypertensive patients without obstructive coronary artery disease.
J. Am. Coll. Cardiol., 15 : 43 - 51.
- 70- *Ibrahim, M.M. (1998):*
Preliminary results of NHP.
J. of the Egyptian hypertension society. Volume 4, No. 2.
- 71- *Imai, Y.; Abek and Miuro, Y. (1988) :*
Hypertensive episodes and circadian fluctuations of blood pressure in patients with pheochromocytoma".
Hypertnesion, 6 : 9 - 15.

-
- 72- *Imani, Y.; Abek, Munakata, M.; Sakuma, H.; Hashimoto, J.; Imaik, Sekino, H. and Yoshinaga, K. (1990) :*
Circadian blood pressure variations under different pathophysiological conditions.
J. Hypertension, 8 (Suppl. 9) 5125 - 321.
- 73- *Inada, Y.; Murakami, M.; Kaido, K. and Nakao, K. (1999):*
Effects of the new angiotensin II type 1 receptor antagonist KRH-594 on several types of experimental hypertension.
Arzneimittelforschung, 49 (1): 13-21.
- 74- *Inouye, L.; Massie, B. and Loge, D. (1984) :*
Abnormal left ventricular filling : an early finding in mild to moderate systemic hypertension.
Am. J. Cardiol., 53 : 120-126.
- 75- *Julius, S.; Ellis, C.N.; Pascual, A.V.; Matice, M.; Hanson, L. and Hanyor, S.N. (1974) :*
Home blood pressure determination value in borderline hypertension.
JAMA, 229 : 663 - 666.
- 76- *Julius, S.; Meija, A.; Jones, K.; Krause, L.; Schork, N. and Van de Ven, C. (1990) ;*
White coat versus sustained borderline hypertension in Tecumseh, Michigan.
Hypertension, Vol. 16, pp. 617-623.
- 77- *Kamel, W.B.; Devereux, R.B.; Herold, E.M.; Jason, M.; Fisher, J. and Laragh, J. (1993) :*
Relation of concentric left ventricular hypertrophy and extra cardiac target organ damage to supra normal left ventricular performance in established essential hypertension.
Am. J. Cardiol., 62 : 446 - 453.
- 78- *Kannel, W.B.; Gordon, T. and Offutt, D. (1969) :*
Left ventricular hypertrophy by electrocardiogram. Prevalence, incidence and mortality in the Framingham study.
Ann. Intern. Med., 71 : 89 - 105.
-

- 79- Kannel, W.B.; Horan, M.J.; Sprague, M.K.; Padgett, N.E. and Shriver, K.K. (1970) :
Left ventricular hypertrophy and prevalence of mortality in hypertensive patients.
Circulation, 16 : 73 - 76.
- 80- Kannel, W.B.; Kimball, Thomas, R. and Daniels, Stephen, R. (1993):
Target organ damage in hypertensive patients as a predictor of mortality rate.
J. Hypertension, 81 : 109 - 113.
- 81- Kannel, W.B.; Sorlie, P. and Gordon, T. (1991) :
Left ventricular hypertrophy in hypertensive patients is a predictor of poor prognosis and increased mortality rate.
Circulation, 97 : 1080 - 1084.
- 82- Kario, K.; Matsuo, T.; Kobayashi, H.; Imiya, M.; Matsuo, M. and Shimada, K. (1998) :
Nocturnal fall of blood pressure and silent cerebrovascular damage in elderly hypertensive patients. Advanced silent cerebro-vascular in extreme dippers.
Hypertension, Jan. 67 : 130 - 135.
- 83- Kawano, Y.; Tochikubo, O.; Minamisawa, K.; Miyajima, E. and Ishii, M. (1994) :
Circadian variations of hemodynamics in patients with essential hypertension comparison between early morning and evening.
J. Hypertens, Dec., 12 : 12, 1405 - 12.
- 84- Kennedy, H.L.; Horan, M.J.; Sprague, M.K.; Padgett, N.E. and Shriver, K.K. (1983) :
Ambulatory blood in healthy normotensive males.
Am. Heart J. October, Vol. 106, pp. 717-722.
- 85- Kronig, B. (1989) :
Nochtliche Hypertonie : Epidemiologie, Kardiovascularized Risiko. In : Herausgeber, JHP Penzel, T.; Cassel, W.; Von Wichert, P. Eds. Schalf - Atmung - Kreislauf, Berlin : Springer - Verlag (in Press).

- 86- *Kupari, M.; Koskinen, P. and Virolainen, J. (1994) :*
Correlates of left ventricular mass in a population sample aged
36 - 37 years. Focus on lifestyle and salt intake.
Circulation, 89 : 1041 - 1050.
- 87- *Laevie, C.J.; Schmieder, R.E. and Messerli, F.H. (1988) :*
Ambulatory blood pressure monitoring : practical
considerations.
Am. Heart, J. 116 : 1146 - 1151.
- 88- *Lakatta, E.G. (1979) :*
Alterations in the cardiovascular systems that occur in
advanced age.
Fed. Proc. 38 : 163 - 167.
- 89- *Lang, R.M.; Borow, K.A. and Weinert, L. (1988) :*
Circadian blood pressure patterns in hypertensive patients.
Importance of nocturnal blood pressure.
Circulation, 78 : 517 - 521.
- 90- *Ledingham, J.G.G. (1992) :*
Definitions of Hypertension in relation to risk; The role of
ambulatory monitoring.
Eur. Heart. J. 13 : 35 - 38.
- 91- *Leeneen, H.H. (1989) :*
Left ventricular hypertrophy in hypertensive patients.
Am. J. Med., 86 : 63 - 65.
- 92- *Leo Nova, M.V.; Belousovl, B.; Semenchuk, G.A.; Malysheva,
E.A.; Upniskii, A.A. and Istratoveslu (1997) :*
An analysis of the indices of ambulatory circadian blood
pressure monitoring in patients with arterial hypertension.
Ter. Arkh, 69 : 35 - 38.
- 93- *Levy, D.; Anderson, K.M.; Savage, D.D.; Balkus, S.A.; Kannel,
W.B. and Castalli, W.P. (1987) :*
Risk of ventricular arrhythmias in left ventricular hypertrophy :
The framingham Heart Study.
Am. J. Cardiol., 60 : 560 - 565.

-
- 94- Levy, D.; Anderson, K.M.; Savage, D.D.; Kannel, W.B.; Christiansen, J.C. and Castelli, W.P. (1988) :
Echocardiography detected left ventricular hypertrophy :
prevalence and risk factors. The Framingham Heart Study.
Ann. Intern. Med., 108 : 7 - 13.
- 95- Levy, D.; Garrison, R.J.; Savage, D.D.; Kannel, W.B. and Castelli, W.P. (1990) :
Prognostic implications of echocardiographically determined
left ventricular mass in the Framingham Heart Study.
N. Engl. J. Med., 322 : 1561 - 1566.
- 96- Liebson, P.R.; Grandits, G. and Prineas, R. (1993) :
Echocardiographic correlates of left ventricular structure
among 844 mildly hypertensive Men and Women in the
treatment of mild hypertension study (TOMHS).
Circulation, 87 : 476 - 486.
- 97- Liebson, P.R.; Lewis, J.F. and Maron, B.J. (1986) :
Echocardiographic detection of left ventricular hypertrophy.
Am. J. Cardiol. 78 : 620 - 625.
- 98- Lima, E.G.; Spritzer, N.; Herkenhoff, F.L. and Vasquez, E.C
(1995):
Ambulatory blood pressure values in relation to left ventricular
hypertrophy in hypertensive patients.
Hypertension, 26 : 121-4.
- 99- Linzbach, A.J. (1969) :
Heart failure from the point of view of quantitative anatomy.
Am. J. Cardiol. 5 : 370 - 82.
- 100- Littler, W.A. (1979) :
Sleep and blood pressure : "Further observations".
Am. Heart, J., 97 : 35 - 37.
- 101- Lreell, B.H. and Grossman, W. (1987) :
Cardiac hypertrophy : the consequences for diastole.
J. Am. Coll. Cardiol., 9 : 1189 - 1193.
- 102- Lurbe, A.; Redon, J. and Rascual, J.M. (1993) :
Altered blood pressure during sleep in normotensive subjects
with type I diabetes.
Hypertension, 21(2) : 227 - 235.
-

- 103- *Mallion, J.M.; De Gaudemaris, R.; Siche, P.J.; Maitre, A. and Pitiot, M. (1990) :*
Day and night blood pressure levels in normotensive and essential hypertensive subjects.
J. Hypertension, 8 : 49 - 56.
- 104- *Mancia, G.; Gamba, P.L.; Omboni, S.; Placariff, G.; Rega, R. and Zanchetti, A. (1997):*
Ambulatory blood pressure monitoring.
J. Hypertens, 31:81-86.
- 105- *Mancia, G. (1990):*
Ambulatory blood pressure monitoring research and clinical applications.
J. Hypertens, 8 : 51 - 513.
- 106- *Mancia, G.; Ferrari, A.; Gregorini, L.; Parat, G.; Pomidossi, G.; Bertinieri, G. and Grassi, G. (1983) :*
Blood pressure and heart rate variabilities in normotensive and hypertensive human beings.
Circ. Res. 53 : 96 - 104.
- 107- *Mancia, G.; Ferrari, A.; Gregorini, L.; Parati, G.; Pomidossi, G. and Bertinieri, G. (1980) :*
Blood pressure variability in man : its relation to high blood pressure, age and baroreflex sensitivity.
Clin. Sci., 59 : 401 - 404.
- 108- *Mancia, G.; Parati, G.; Pomidossi, G.; Grassi, G.; Casadei, R. and Zanchetti, A. (1987) :*
Alerting reaction and rise of blood pressure during measurement by physician and nurse.
Hypertension, 9 : 209 - 215.
- 109- *Mann, S.; Altrnan, D.G. and Raftery, E.B. (1983) :*
"Circadian variation of blood pressure in autonomic failure"
Circulation, 68 : 477 - 483.
- 110- *Marcus, M.L.; Harrison, D.G. Chilian, W.M. (1987) :*
Alterations in the coronary circulation in the hypertrophied ventricles.
Circulation, 75 : 1110-1125.

- 111- Martell, N.; Rodrigo, J.L. and Fernandez-Pinilla, C. (1991) :**
Sodium intake and atrial natriuretic factor as determinants of left ventricular dimensions : the Torrejon Study.
J. Hypertens, 9 : 258 - 259.
- 112- Mayet, J.; Elizari, D.J.; Dunn, F.G. and Frolich, E.D. (1995) :**
Ambulatory blood monitoring is a predictor of target organ damage in comparison with office blood pressure readings.
Circulation, 98 : 176 - 184.
- 113- McDonough, J.R.; Grrison, G.E. and Hames, C.G. (1964) :**
Blood pressure and hypertensive disease among negroes and whites.
Ann. Intern. Med., 61 : 208-228.
- 114- McLenachan, J.M.; Henderson, E.; Morris, K.I. and Dargie, J.H. (1987) :**
Ventricular arrhythmias in patients with hypertensive left ventricular hypertrophy.
N. Engl. J. Med., 317 : 787 - 792.
- 115- Meija, S.; Ruff, S.; Scuttle, K.L. and Tofler, G.H. (1990) :**
Ambulatory blood pressure monitoring is superior casual blood pressure in office.
J. Hypertens, 12 : 51 - 54.
- 116- Mejer, J.L.; Ardesch, H.G.; Van Rooyien, J. and Debruij, J.H.B. (1986) :**
Low dose captopril twice daily lowers blood pressure without disturbance of the normal circadian rhythm.
Postgrad. Med. J. 61 : 101 - 109.
- 117- Messerli, F.H. (1996) :**
Multifactorial pathogenesis of left ventricular hypertrophy.
Medicographia, 29 : 123 - 127.
- 118- Messerli, F.H. and Aepfelbacher, F.C. (1995) :**
Left ventricular hypertrophy : Pathophysiologic sequelae and therapeutic considerations.
Medicographia, 17 : 30 - 34.

- 119- **Messerli, F.H. and Devereux, R.B. :**
Left ventricular hypertrophy good or evil?
Am. J. Med. 75 : 1-3.
- 120- **Messerli, F.H.; Garavaglia, G.E.; Schmieder, R.E.; Sundgaard-Riisek, Nunez, B.D. and Amodeo, C. (1987) :**
Disparate cardiovascular findings in men and women with essential hypertension.
Ann. Intern. Med., 107 : 158 - 161.
- 121- **Messerli, F.H. and Ketelhut, R. (1991) :**
Left ventricular hypertrophy : an independent risk factor.
J. Cardiovasc. Pharmacol., 17 (Suppl. 4) : 259 - 267.
- 122- **Messerli, F.H.; Nunez, B.D.; Ventura, H.O. and Snyder, D.W. (1987) :**
Overweight and sudden death : increased ventricular ectopy in cardiomyopathy of obesity.
Arch. Intern. Med., 147 : 1725 - 1728.
- 123- **Messerli, F.H.; Oren, S. and Grossman, E. (1988) :**
Left ventricular hypertrophy and antihypertensive therapy.
Circulation, 68: 27-33.
- 124- **Meyer Sabellek, W.; Schuttle, K.L. and Gotzen, R. (1990) :**
Non invasive ambulatory blood pressure monitoring : techchmical possibilities and problems.
J. Hypertens, 8 : 3-10.
- 125- **Middeke, M.; Ruff, S. and Kluglich, M. (1992) :**
What does general practice hypertension mean?
Z. Gesamnte inn. Med., 47 : 558-561.
- 126- **Morgan (1995) :**
Circadian variation and triggers of onset of a cute cardiovascular disease".
Circulation, 79 : 733 - 743.
- 127- **Muller, J.E.; Tofler, G.H.; Stone, P.H. (1989):**
Circadian variation and triggers of onset of acute cardiovascular disease.
Circulation, 79: 733-743.

- 128- *Munakata, M.; Imai, Y.; Abe, K. and Sasaki, S. (1988) :*
Involvement of the hypothalamo-pituitary - adrenal axis in the control of circadian blood pressure rhythm.
J. Hypertension, 6 : 44 - 46.
- 129- *Neaton, J.; Flack, J.M. and Grimm, R.J. (1996) :*
Blood pressure and high blood pressure : aspects of risk.
Am. J. Hypertension, 12 : 210 - 213.
- 130- *O'Brin, E.; Sheridan, J. and O'Malley, K. (1988) :*
"Dippers and non dippers".
Lancet. (2) 397-398.
- 131- *Oren, S. and Viskoper, J.R. (1991) :*
Twenty - four arterial pressure and heart rate as predictors of left ventricular hypertrophy.
Am. J. Hypertens, 302 - 307.
- 132- *Orlov, S.N.; Adragna, N.C.; Adarichev, V.A. and Hamet, P. (1999):*
Genetic and biochemical determinants of abnormal monovalent ion transport in primary hypertension.
Am. J. Physiol., 276 (3 pt 1): C511-36.
- 133- *Palatini, P.; Penzo, M. and Racioppa, A. (1992) :*
Clinical relevance of night - time blood pressure and of day - time blood pressure variability.
Arch. Intern. Med., 151 : 1855- 1860.
- 134- *Palatini, P. and Pessina, A.C. (1997):*
A new approach to define the upper normal limits of ambulatory blood pressure.
J. Hypertens., 8 : 65 - 70.
- 135- *Pannier, L.; Prisant, L.M.; Bottini, P.B. and Carr, A.A. (1989) :*
Left ventricular hypertrophy in hypertensive patients in relation to hypertension duration.
J. Clin. Pharmacol. Vol. 22, p. 260 - 266.
- 136- *Perloff, D.; Sokolow, M. and Cowan, R. (1983) :*
The prognostic value of ambulatory blood pressure.
JAMA, 249 : 2792 - 2798.

- 137- *Pickering, T.G. (1992) :*
Ambulatory blood pressure monitoring and the definition of hypertension.
J. Hypertens, 10 : 401 - 409.
- 138- *Pickering, T.G. (1991):*
Ambulatory blood pressure monitoring in clinical practice.
Clinic Cardiol., 14: 557-562.
- 139- *Pickering, T.G. (1990):*
The clinical significance of diurnal blood pressure variation :
Dippers and nondippers".
Am. J. Hypertens, 6 : 19 -28.
- 140- *Pickering, T.G.; Harshfield, G.A.; Kleinert, H.D.; Blank, S. and Laragh, J.H. (1986):*
Blood pressure during normal daily activities, sleep and exercise.
JAMA, 278 : 991 - 998.
- 141- *Raftery, E.B. (1983) :*
Zirkadine blutdruckschwankungen (circadian blood pressure variability) in : Megawitschnigg, D.; Hitznberger, G. (Eds)
Blutdruck Variabilität. (Uhlen - Verlagsges MbH : Wien) 53 - 66.
- 142- *Rasooly, Y.; Sasson, Z. and Gupta, R. (1993) :*
Relation between body fat distribution and left ventricular mass in men without structural heart disease or systemic hypertension.
Am. J. Cardiol., 71 : 1477 - 1479.
- 143- *Reavin, L.M.; Devereux, R.B. and Camber, J. (1994) :*
Pathogenesis of left ventricular hypertrophy.
Am. J. Cardiol. 83 : 1302 - 1308.
- 144- *Reeves, R.A.; Shapiro, A.P. and Thompson, M.E. (1986) :*
"Loss of nocturnal decline of blood pressure after cardiac transplantation".
Circulation, 73 : 401 - 408.

- 145- Reicher, N. and Devereux, R.B. (1981) :**
Left ventricular hypertrophy relationship of anatomic, echocardiographic and electrocardiographic finding.
Circulation, 639 : 1391 - 1939
- 146- Reichek, N. and Devereux, R.B. (1982) :**
Reliable estimation of peak left ventricular systolic pressure by M. mode echographic determined end - diastolic relative wall thickness : identification of severe valvular aortic stenosis in adult patients.
Am. Heart. J. 103 : 202 - 209.
- 147- Reichek, N.; Pomidossi, G.; Albini, F. and Malaspina, D. (1984) :**
M. Mode echocardiographic detection of left ventricular hypertrophy in hypertension and hypertrophic cardiomyopathy.
Am. Heart. J. 121 : 307 - 309.
- 148- Reik, S.F.; Savage, D.D.; Drayer, JIM.; Henry, W.L. and Mathews, E.C. (1984) :**
Echocardiographic detection of left ventricular hypertrophy.
Circulation, 63 : 29 - 34.
- 149- Reinberg, A.; Smolensky, M.H. (1983) :**
Investigate methodology of chronobiology in: Reinberg A.; Somolensky, M.H. (Eds) Biological rhythms and medicines (Springer Verlag Berlin/Heidlber/New York, 23 - 43.
- 150- Rizzoni, D. and Agabiti-Roseit (1996) :**
Relationship of cardiac function and structure to blood pressure rhythms.
Ann. N-Y Acad. Sci. 783 : 159 - 171.
- 151- Rosner, B. and Polk, B.F. (1979) :**
The implications of blood pressure variability of clinical and screening purposes.
J. Chronic. Dis., 32 : 451 - 461.
- 152- Rowlands, D.B.; Glover, D.R. and Reland, M.A. (1982) :**
Assessment of left ventricular mass and ® its response to antihypertensive treatment.
Lancet, 1 : 467 - 470.

- 153- *Sahn, D.J.; De Maria, A.; Kisslo, J. and Weyman, A. (1978) :*
The committee on M. Mode standardization of the American Society of Echocardiography.
Recommendation regarding quantitation in M. mode echocardiography : results of a survey of echocardiographic measurements.
Circulation, 58 : 1072-1082.
- 154- *Savage, D.D.; Drayer, JIM and Henry, W.L. (1983) :*
Assessment of left ventricular mass by echocardiogram.
Am. Heart, J. 109 : 73 - 77.
- 155- *Savage, D.D.; Drayer, JIM.; Henery, W.L.; Mathews, E.C.; Ware, J.H. and Gardin, J.M. (1979) :*
Echocardiographic assessment of cardiac anatomy and function in hypertensive subjects.
Circulation, 59 : 623 - 632.
- 156- *Savage, D.D.; Garrison, R.J. and Kannel, W.B. (1987) :*
The spectrum of left ventricular hypertrophy in a general population sample : the Framingham study.
Circulation, 75 : 26 - 33.
- 157- *Schillaci, G.; Verdecchia, P. and Borgioni, P. (1994):*
Improved electrocardiographic diagnosis of left ventricular hypertrophy.
Am. J. Cardiol, 74 : 714 - 719.
- 158- *Schmieder, R.E. (1994) :*
Reversal of left ventricular hypertrophy : analysis of 412 published studies.
Am. J. Hypertens. 7 : 25 - 33.
- 159- *Schmieder, R.E. and Messerli, F.H. (1993) :*
Does obesity influence early target organ damage in hypertensive patients?
Circulation, 87 : 1482-1488.
- 160- *Schmieder, R.E.; Messerli, F.H.; Caravaglia, G.E. and Nunez, B.D. (1988) :*
Dietary salt intake : a determinant of cardiac involvement in essential hypertension.
Circulation, 78 : 951 - 956.

- 161- *Sechtem, S.C.; Devereux, R.B.; Lutas, E.M. and Casale, P.N. (1987) :*
Detection of left ventricular mass by magnetic response imaging a recent approach.
Br. Heart. J. 48 : 398 - 401.
- 162- *Sen, S. and Youn, D.R. (1986) :*
Role of sodium in modulation of myocardial hypertrophy in renal hypertensive subjects.
Hypertension, 8 : 918 - 924.
- 163- *Shapiro, L.M. and Gibson, D.G. (1988) :*
Patterns of diastolic dysfunction in left ventricular hypertrophy.
Br. Heart. J. 59 : 438 - 445.
- 164- *Siegel, W.C.; Blumenthal, J.A. and Divine, G.W. (1990) :*
Physiological, psychological and behavioural factors and white coat hypertension.
Hypertension, 16 : 140 - 146.
- 165- *Smith, V.E.; White, W.B.; Karimeddini, M.K.; McCabe, E.J. and Katz, A.M. (1986) :*
Lowest not highest blood pressure may determine left ventricular filling.
Circulation, 74 : 29 - 36.
- 166- *Smith, V.E.; White, W.B.; Karimeddini, M.K. and McCabe, E.J. (1987):*
Echocardiographic assessment of left ventricular diastolic performance in hypertensive subjects. Correlation with changes in left ventricular mass.
Hypertension, 9 : 81 - 84.
- 167- *Souchek, J.; Stamler, J.; Dyer, A.R.; Paul, O. and Lapper, M.H. (1979):*
The value of two or three versus a single reading of blood pressure at a first visit.
J. Chronic. Dis. 32 : 197 - 210.
- 168- *Staessen, J.; Fagard, R.; Lijmen, P.; Thijs, L.; Amery, A. (1990):*
Reference values for ambulatory blood pressure: a meta analysis.
J. Hypertens., 8: 57-64.

- 169- *Staessen, J.; Fagard, R.; Lijnen, P.; Thijs, L.; Vanhoof, R. and Amery, A. (1991) :*
Ambulatory blood pressure in clinical trials.
J. Hypertens. 9 : 13 - 20.
- 170- *Stella, P.; Soldati, L.; Ciurlino, D. and Bianchi, G. (1998) :*
Diagnostic evaluation of patient with systemic hypertension.
J. Hypertens. 15 : 1775 - 1778.
- 171- *Strauer, B.E. (1984) :*
Ventricular function and coronary hemodynamics in hypertensive heart disease.
Am. J. Cardiol. 74 : 106 - 113.
- 172- *Strauer, B.E. (1987) :*
Comparative analysis of cardiac function, geometry, energetics and coronary reserve in hypertensive heart disease.
Nephron, 47 : 76 - 86.
- 173- *Strauer, B.E. (1994) :*
Left ventricular wall stress and hypertrophy in : Messerli, F.H. (Ed.) the heart in hypertension.
New York : York Medical Books, 173 - 185.
- 174- *Sullivan, J.M. and Ratts, T.E. (1983) :*
Hemodynamic mechanisms of adaptation to chronic high sodium intake in normal humans.
Hypertension, 5 : 814 - 820.
- 175- *Sullivan, J.M.; Ratts, T.E. and Taylor, J.C. (1980) :*
Hemodynamic effects of dietary sodium in man a preliminary report.
Hypertension, 2 : 506 - 514.
- 176- *Susic, D.; Frohlich, E.D. (1993) :*
Left ventricular hypertrophy : a pathophysiological and molecular biological perspective.
Hypertens. Res. 16 : 163 - 177.
- 177- *Swarwicz, B. (1986) :*
Electrocardiographic diagnosis of chamber enlargement.
J. Am. Coll. Cardiol. 8 : 711 - 713.

- 178- *Szlachcic, J.; Tubau, J.F.; O'Kelly, B.; Ammon, S.; Daissk and Massie, B.M. (1992) :*
What is the role of silent coronary and left ventricular hypertrophy in the genesis of ventricular arrhythmias in men with essential hypertension.
J. Am. Coll. Cardiol., 19 : 803 - 808.
- 179- *Tanaka, T.; Natsume, T. and Shibata, H. (1983) ;*
Circadian rhythm of blood pressure in primary aldosteronism and renovascular hypertension.
J. Pn. Circ. J. 43 : 788 - 794.
- 180- *Tellier, P. (1995) :*
Epidemiology of left ventricular hypertrophy in hypertension.
Mediographia, Vol. 17, No. 1, p. 17 - 21.
- 181- *Turek, F.W. (1985) :*
Circadian neural rhythms in mammals.
Annual Review of Physiology, 47 : 49 - 64.
- 182- *Verdecchia, P.; Scillaci, G. and Guerrieri, M. (1990) :*
"Circadian blood pressure changes and left ventricular hypertrophy in essential hypertension".
Circulation, 81 : 528 - 536.
- 183- *Verdecchia, P.; Schillaci, G.; Borgioni, C.; Ciucci, A. and Porcellati, C. (1998):*
Ambulatory blood pressure monitoring in comparison to casual blood pressure readings in clinical practice.
Am. J. Hypertens., 8: 790-798.
- 184- *Verdecchia- Schillaci, G. and Guerrieri, M. (1990):*
Circadian blood pressure changes and left ventricular hypertrophy in essential hypertension.
Circulation, 81: 528-536.
- 185- *Watson, R.D.S.; Lumb, R.; Young, M.A.; Stallard, T.J.; Devies, P. and Littler, W.A. (1987) :*
Variation in cuff blood pressure in untreated outpatients with mild hypertension implications for initiating antihypertensive treatment.
J. Hypertens; 5 : 207 - 211.

- 186- *Weber, K.T. (1989) :*
Cardiac interstition in health and disease : the fibrillar collagen network.
J. Am. Coll. Ardiol., 13 : 1637 - 1652.
- 187- *Weber, K.T. and Brilla, C.G. (1992) :*
Left ventricular hypertrophy and structural remodellings as a determinants of myocardial failure. From left ventricular hypertrophy and its regression.
Science Press Ltd. London, 34 - 42.
- 188- *Weber, K.T.; Janiki, J.S. and Shroff, S.G. (1988) :*
Collagen compartment and remodeling in the pressure overloaded left ventricle.
J. Appl. Cardiol. 3 : 37 - 46.
- 189- *White, W.B. (1986) :*
Assessment of patients with office hypertension by 24-hour non invasive ambulatory blood pressure monitoring. Arch. Intern. Med. 146 : 2196 - 2199.
- 190- *White, W.B.; Schulman, P. and McCabe, E.J. (1989) :*
Average daily blood pressure, not office blood pressure, determines cardiac function in patients with hypertension, JAMA, 261 : 873 - 877.
- 191- *WHO Expert Committee on Hypertension Control (1996):*
Hypertension control: Report of a WHO Expert Committee, WHO Technical Report Series no. 862. Geneva, Switzerland: World Health Organization.
- 192- *Widimsky, J. (1995) :*
Should echocardiographic evidence of left ventricular hypertrophy lead to modification of treatment of hypertension? Medicographia, 17 : 38 - 40.
- 193- *Zanchetti, A. (1998) :*
The role of ambulatory blood pressure monitoring in clinical practice.
Am. J. Hypertens. 4 : 1069 - 1080.

- 194- *Zhang, L.; Yang, Y.; Wang, J.M. and Wang, N. (1999):*
Effect of chromosome 19 transfer on blood pressure in the
spontaneously hypertensive rat.
Hypertension, 33 (1 pt 2): 256-60.
- 195- *Zusman, R.M.; Christensen, D.M.; Higgins, J. and Boucher, C.A.*
(1991) :
Nifedipine improves left ventricular function in patients with
hypertension.
J. Cardiovasc. Pharmacol, 18 : 843 - 848.