

REFERENCES

- Abi-Samra, F.; Fouad, F.M.; Tarazi, R.C. (1983):**
Determinants of left ventricular hypertrophy and function in hypertensive patients. An echocardiographic study.
The American Journal of Medicine, 75 (Suppl 3 A) 26-33.
- Agabiti, Rosei, E.; Mulesan, M.L. and Muiesan, G. (1989):**
Regression of structural alterations in hypertension.
Am. J. Hypertens. 2: 705.
- Bachemberg, T.C.; Hauck, A.J.; Shub, C.; Edward, W.D. (1991):**
Can anatomical left ventricular mass estimated reliably by M-Mode. Echocardiography ?
Journal of Echocardiography Vol. 8 No. 1: 9-15.
- Badeer, H.S. (1964):**
Biological significance of Cardiac hypertrophy.
Am. J. cardiology 14-33.
- Balanescu, S.; Galinier, M.; Faurcade, J.; Doroban, M.; Albenque, J.F.; Massabuau, P.; Fauvel, J.M.; Boum, Houre, J.P.(1996):**
Arch. Mal. Coeur. Vaiss-aug 89(8): 987-90.
- Barlaw, J. and Kincaid, Smith, P. (1960):**
The auscultatory finding in hypertension.
Br. H.J. 22: 505.
- Barr, C.S.; Naas, a.; Freeman, M.; Lang, C.C.; Struthers, A.D. (1994):**
QT dispersion and sudden unexpected death in chronic. HF.
Lancet 343: 327-329.
- Bazett, H.C. (1920):**
Analysis of the time relation of electrocardiograms.
Am. Heart, 7, 353-370.
- Beach, C.; Kenmure, A.C.F. and Shor, T.D. (1981):**
Electrocardiogram of pure left ventricular hypertrophy and its differentiation from lateral ischaemia.
Br. H.J. 46: 285-99.

- Bhuller, H.K.; Forthergill, J.C.; Gadder, W.P. and De-Bono, D.P. (1983):**
Automatic measurement of Q.T. interval dispersion from hard copy ECGs.
J. Electrocardiol, 26 (4): 321-331.
- Black, H.R.; Weltin, G. and Jalle, C.C. (1991):**
The limited echocardiogram.: A modification of standard echocardiography for use in the routine evaluation of patients with systemic hypertension.
Am. J. Cardiol 67: 1027.
- Borer, J.S.; Jason, M.; Devereux, R.B.; Fisher, J. (1989):**
Left ventricular performance in the hypertensive patients.
Exercise mediated non invasive separation of loading influence.
Circ. 89 (suppl I): 319.
- Borrow, K.M.; Green, L.H. Grossman, W. (1992):**
Left ventricular end systolic stress shortening and stress length relations in humans.
The American Journal of Cardiology 4: 999-1006.
- Brandes, A.; Bethge, K.P. and Buerschaper, M. (1997):**
Q. T. dispersion in healthy subjects.
J. Am. Coll. Cardiol 29 (2) Suppl. A.: 508 A.
- Brentano, C.F. and Jaillon, P. (1993):**
Rate corrected Q.T. interval Techniques and Limitation and Am. J. Cardiol 72: 17 B-22 B.
- Buja, G.; Miorelli, M.; Turrini, P.; Melacini, P. and Novo, A. (1993):**
Comparison of Q.T. dispersion in hypertrophic cardiomyopathy between patients with and without ventricular arrhythmias and sudden death.
Am. J. Cardiology 72, 973-976.
- Burchell, H.B. (1974):**
Clinical recognition of cardiac hypertrophy
Circ. Res., 35 (suppl. 11): 116.

- Cameron, J.S.; Myerburg, R.J.; Wong, S.S.; Gaide, M.S.; Epstein, A.L.; Alvarez, T.R.; Gelband, H.; Guse, P.A. and Bassett, A.L. (1983):**
Electrophysiologic consequences of chronic experimentally induced left ventricular pressure overload.
J. Am. Coll. Cardiol 2: 481.
- Casale, P.N.; Devereux, R.B. and Milner, M. (1986):**
Value of echocardiographic measurements of left ventricular mass in predicting cardiovascular morbid events in hypertensive men. Ann. Intern. Med. 105: 173.
- Castelli, W.P. (1984):**
The epidemiology of coronary heart disease.
The framingham study. Am. J. med. 7 (2): 4-12
- Cavallini, B.; Perri, V.; Sali, M. (1996):**
Dispersion of Q.T. interval in systemic. Hypertension with left ventricular hypertrophy.
Minerva, cardioangiol. Jan Feb 44 (1-2) 45-8.
- Challapallis, Ligamnei, R. and Ehlert, F. (1997):**
Gender difference in Q.T. dispersion.
J. Am. Coll. Cardiol, 29 (2) Suppl. A.: 148.
- Charles, Fisch, (1992):**
Electrocardiographic and vector cardiography Quoted from Braunwald text. Book of cardiovascular medicine, Chapter 5 page 122. Philadelphia.
- Clarkson, P.B.; Naas, A.A. and MC. Leod, C. (1995):**
Q.T. Dispersion in essential hypertension.
Q.J.M. 88 (5): 327-32. (Abstr.)
- Conn, R.D. and Cole, J.S. (1971):**
The cardiac apex impulse, clinical and angiographic correlations Ann of intern. Med. 75 : 185.
- Daniarov, B.; Hilbel, T. and Brachman, J. (1997):**
Shortening of prolonged Q.T. dispersion after coronary angioplasty.
J. Am. Coll. Cardiol. 29 (2) : suppl. A: 510 A.

Davie, J.; Langley, J.; Dodson, W. (1962):

Clinical and kinetocardiographic studies of apardoxical pericordial motion.
Amer. Heart.J. 63: 775.

Day, C.P.; Mccomb, J.M. and Compbell, R.W. (1992):

Q.T. dispersion in sinus beats and vent extra systoles in normal hearts. Br. Heart. 67: 139-41.

Day, C.P.; mc. Comb, J.M. and Mathews, J.(1991):

Reduction in Q.T. dispersion by sotalol following myacardiol infarction.
Eur. Heart., J., 12 : 1423-427.

DeLuna, A.B. and Vinolas, X. (1996):

Q.T. dispersion and heart rate varibaility
Eur. Heart. J. 17: 165-166.

Devereux, R.B. (1987):

Cardiac involvement in essential hypertension. Prevalence, pathophysiology and prognostic implications.
Medical clinical of North america. 71-5 : 819.

Devereux, R.B.; Alonso, D.R.; Lutrass, E.M. (1986):

Echocardiography assesement of left ventricular hypertrophy comparison. To necorpsy findings.
The Amircan Journal of cardiology: 57: 450.

Devereux, R. B.; Carale, P.N. ; Kiligfield, P.; Eisenber, R.R.; Miller, D.; Campo, E. and Alonso, D.R. (1986):

Performance of primary and derived M. mode echocardiogrphic. Measurements for detections of left ventricular hypertrophy in necropsied subjects and in patients with systemic hypertension mitral regurgitation and dilated cardiomyopathy.
Am. J. Cardiol 57: 1388.

Deverexu, R.B.;Lutas, E.M.; Casale, P.N. (1984):

Standardization of M. Mode. Echocardiogrphic left ventircular anatomic measurements.
J. Am. Coll, Cardiol. 4: 1222.

- Devereux, R.B. and Reichek, N. (1977):**
 Echocardiographic determination of left ventricular mass in man. Anatomic validation of the method.
 Circulation 55: 613-618.
- Donohue, T.J.; Dworkin, I.D.; Longo, M.N.; Fliegner, K.; Lango, R.P.; Benstein, J.A.; Salter, W.R. and Catamese, V.M. (1994):**
 Induction of myocardial insuline like growth factor 1 gene expression in left ventricular hypertrophy.
 Circulation 89: 799.
- Douglas, P. Zipes (1992):**
 Specific arrhythmias, diagnosis and treatment. Brounwald text book of cardiovascular Medicine.
 Chapter 5. Philadilphia.
- Drayer, J.I.M.; Weber, M.A. DeYoung (1989):**
 Blood pressure as adeterminant of cardiac left ventricular mass.
 Archive of internal medicine. 193: 90.
- Dreseler, W. (1957):**
 Pulsations of the wall of the chest.
 Arch. Intern. Med. 60: 225.
- Dreslinski, G.R.; Frolich, E.D.; Dunn, F.G. (1989):**
 Echocardiogrpahic diastolic ventricular abnormality in hypertensive heart disease: atrial emptying index.
 The american. Journal of cardiology 47: (1087).
- Dunn, F.G.; Mclenochan, J. and Isies, C.G.; Brown, I.; Dargie, H.J.; Lever, A.F.; Lorimer, A.R.; Murray, G.D.; Pringle, S.D. and Robertson, J.W. (1990):**
 Left ventricular hypertrophy and mortality in hypertension: Am. Analysis of data from the glasgow blood pressure. Clinic.
 J. Hypertnes. 8: 75.
- Eilen, S.D.; Crawford, M.H. and O'Roukee, R.A. (1983):**
 Accuracy of pericordial palpation for detecting increased left ventircular volume.
 Ann. Intern. Med.99: 628.

Falase, A.O.; Cole, T.O. and Osuntokun, B.O. (1982):

Myocardial infarction in Nigerians.

Tropi. Geogr. Med. 25: 147.

Fang, Q.; Bourke, J.P. and Simeomidou, E. (1996):

Increased Q.T. dispersion; a new marker predicting induction of post myocardial infarction ventricular hypertrophy. J. eur. Soc. Cardiol., 17: 30.

Feil, Goldman, J.H. and Prasad, K. (1996):

Q.T. dispersion and R.R. variations on 12 leads EcG in patients with congestive heart Failure Secondary to idiopathic dilated cardiomyopathy.

Eur. Heart. J., 17: 258-263.

Fouad, F.M.; Slominski, F.J.M.; Tarazi, R.C. (1984):

Left ventricular hypertrophy. Relation of structure to diastolic function in hypertension.

Journal of American college of cardiology; 3: 1500.

Francis, G.S. and McDaland, K.M. (1992):

Left ventricular hypertrophy, an initial response to myocardial injury.

Am. J. Cardiol. 65: 85.

Franz, M.R.; Bargheer, K.; Rollenbeul, W.; Hoverich, a. and Lichtlen, R.R. (1978):

Monophasic action potential mapping in human subjects with normal electrographic action potential mapping in human. Subjects with normal electrographic action potential mapping in human subjects with normal electrocardiograms. Direct evidence for the genesis of the T wave circulation 75: 329-386.

Frohlich, E.D. and tarazi, R.C. (1989):

Is arterial pressure the sole factor responsible for hypertensive cardiac hypertrophy ?

Am. J. Cardiol 44: 959.

Frohlich, E.D.; Tarazi, R.C. and Dustan, H.P. (1981):

Clinical physiological correlations in the development of hypertensive heart disease.

Circulation 44: 446.

Furchgott, R.F. and Zawadski, J.V. (1993):

The obligator role of endocelial celsl in the relaxation of arterial smooth muscle by acetylcholine nature 288: 373. (1980). Quoted by; treasure C.B.; Lein, J>L.; V.Ta, J.A. Manoukian, S.V.; Renwick, G.H.; Selwyn, A.P. Ganz, P. and Alexander. W.

Hypertension and left ventricular hypertrophy are associated with impaired endothelial mediated relaxaiton in human coronary resistance vessels. Circulation 87: 86.

Galinier, M.; Balanescu, S.; Fourcade, J.; Dorabantu, M.; Massobua, P.; Dongay, B.; Carbrol, P.; Fauvel, J.M.; Bounboure, P. (1997):

Prognostic value of ventricular arrythmia in hypertensive patients Arch. Mal. Coeur. Vaiss, Aug. 40 (8), 1049-53.

Ganou, A.; Devereux, R.B.; Roman, M.J.; Desimane, G.; Pickering, T.J.; Saba, P.S.; Vargiu, P.; Simongini, I. And Laragh, J.H. (1992):

Patterns of left ventricular hypertrophy and geometric remodeling in essential hypertesnion. J. Am. Coll. Cardiol 19: 1550.

Garson, A. (1993):

How to measure the Q.T. interval what's normal. Am. J. Cardiol. 72: 14 B-16 B.

Ghali, J.K.; Kadokia, S.; Cooper, R.S. and Liao, Y. (1991):

Impact of left ventricular hypertrophy on ventricular arrhythmias in the absence of coronary artery disease. J. Am. Coll. Cardiol 17: 277.

Glancy, J. M.; Garret, C.J. and Woods, K. L. (1995):

Q.T. dispersion and mortality after myocardial infarction. Lancet, 345: 945-948.

Glancy, J.M.; Garret, C.J. and Woods, K.L. (1995):

Use of lead adjustment fomulas for Q.T. dispersion after myocardial infarction. Br. Heart J., 74: 676-679.

- Glancy, J.M.; Garrett, C.J. and Woods, K.L. (1995):**
Three lead measurement of Q.T. dispersion.
J. Cardiovasc. Electrophysiol 6 (11): 987-992. (abst).
- Glancy, J.M. Weston, P.J. and Bhullar, H.K. (1996):**
Reproducibility and automatic measurement of Q.T.
dispersion Eur. Heart, J.; 17: 1035-1039.
- Grossman, W. (1980):**
Cardiac hypertrophy: useful adaptation or pathological
process. Am. J. med. 69: 576.
- Hammond, I.W.; Devereux, R.B.; Alderman, M.H.; Lutas, E.M.;
Spitzer, M.C.; Crawely, J.S. and Laragh, J.H. (1986):**
The prevalence and correlates of echocardiographic left
ventricular hypertrophy among employed patients with
uncomplicated hypertension. J. Am. Cardiol. 7: 639.
- Hanrath, P.; Mathey, D.G.; Seigert, R. (1990):**
Left ventricular relaxation and filling pattern in different
forms of left ventricular hypertrophy. an echocardiographic
study.
The American Journal of Cardiology. 45: 15.
- Hanrath, P.; Mathey, D.G.; Siegert, R. and Bleifeld, W. (1980):**
Left ventricular relaxation and filling pattern in different
form of left ventricular hypertrophy.
Echocardiographic study. Am. J. cardiol. 45: 51.
- Hantkova, K.; Malik, M. and Kautzner, J. (1994):**
Adjustment of Q.T. dispersion assessed from 12 lead ECGs
for different numbers of analysed electrocardiographic
leads, comparison of stability of different methods.
Br. Heart, J., 72: 390-396.
- HDFP (1979):**
(Hypertension detection and follow up program)
cooperative group (1979).
Reduction of mortality of person with high blood pressure
including mild hypertension.
J. A. M. A. 241 (23) : 2561-2571.

- Higham, P.D.; Furmiss, S.S. and Campbell (1995):**
 Q.T. dispersion and components of the Q.T. interval in
 ischemia and infarction.
 Br. Heart. j., 73: 32-36.
- Hii, T.I.; Wyse, G. and Gillis, A.M. (1992):**
 Precordial Q.T. interval dispersion as a marker of trasade de
 pointes, disparate, effects of class 1 antiarrythmic drugs and
 amiodrone.
 Circ, 86 : 1376-1382.
- Hollander, W. (1976):**
 Role of hypertension in atherosclerosis and cardiovascular
 disease. Am. J. Cardiol. 38 (23) 786-800.
- Houghton, J.L.; Carr, A.A.; Prisant, L.M.; Rogers, W.B.; Vondohlen,
 T.W.; Flowers, N.C. and Frank, M.J. (1992):**
 Hemodynamic and coronary perfusion characteristics in
 severe left ventricular hypertrophy secondary to systemic
 hypertension and evidence for non atherosclerotic
 myocardial ischemia.
 Am. J. Cardiol 69: 219.
- Huwez, F.U.; Pringle, S.D. and Macferlane, G. W.; (1992):**
 Variable patterns of st. T abnormalities in patients with left
 ventricular hypertrophy and normal caronaries.
 Br. Heart., J. 67: 304.
- Ichkhan, K.; molnar, J.; Somberg, J. (1997):**
 Relation of left ventricular mass and Q.T. dispersion in
 patients with systemic hypertension.
 Am. J. Cardiol., Feb. 15, 79 (4): 508-11.
- Ito, H.; Hiroe, M.; Hirata, Y.; Fujisaki, H.; Adachis, Akimoto, H.;
 Ohto, Y. and Marumo, F. (1994):**
 Endothelin, ETA receptor antagonist blocks cardiac
 hypertrophy provoked by hemodynamic overloads.
 Circulation 14: 590.

- Ito, H.; Hirata, Y.; Hirae, M.; Tsuino, M.; Adachi, S.; Takamotot, Nito, M.; Tniquchi, K. and Narumo, F. (1991):**
Endothelin-1 induces hypertrophy with enhanced expression of muscle specific genes in cultured neonatal rat cardiomyocytes.
Circ. Res. 69: 209.
- James, M.; Mclenachan, Esther, Henderson, Karen, I.; Magris and Henery, J. (1987):**
Ventricular arrhythmias in patients with hypertensive left ventricular hypertrophy. New Eng. J. of medicine. 317: 787-92.
- Joint, National Committee (1988):**
The report of the joint national committee and detection, evaluation and treatment of high blood pressure, archintern, med. 148: 1023.
- Kannel, W.B. (1983):**
Prevalence and natural history of electrocardiographic left ventricular hypertrophy.
Am. J. Med. 75 (suppl. 3 A): 4.
- Kannel, W.B.; Doyle, J.T.; M.C. Namare, P.M.; Qukikenton, P. and Gordon, T. (1975):**
Precursors of sudden coronary death; Factors related to the incidence of sudden death.
Circulation. 51: 606.
- Kannel, W.B.; Gordon, T.; Costelli, W.P. and Magolis, J.R. (1980):**
Electrocardiographic left ventricular hypertrophy and risk of coronary heart disease.
The framingham study. Ann. Intern. Med. 72: 813.
- Kannel, W.B.; Schatzkin (1985):**
Sudden death; lessons from subsets in population studies
Acc, 5, B. 141-149.
- Karen, M.I.; Devereux, R.B.; Carae, P.N.; Savage, D.D. and Laragh, J.H. (1991):**
Relation of left ventricular mass and geometry to morbidity and mortality in uncomplicated essential hypertension.
Ann. Intern. Med. 114: 345.

- Katoh, Y.; Kamuro, I.; Shibasaki, Yamaguchi, H. and Yazaki, Y. (1989):**
 Angiotensin II. Induces hypertrophy and oncogene expression in cultured cardiac Myocytes.
 Circulation 80 (Suppl 11): 1145-Abstract.
- Kaul, V.; Mohan, J.C. and Bhatai, M.L. (1984):**
 Effect of labetalol on left ventricular mass and function in hypertension, an assesment by serial echocardiography.
 Int. J. Cardiol 5: 461.
- Khachig, Kchkhan, Janos, Monnar and John, J. (1997):**
 Relation of left ventricular mass and Q.T. dispersion in patients with systemic hypertension.
 Excerpto, Medica-September. Vol. (23) 508-11.
- Khairallah, P.A. and Kamabus, J. (1987):**
 Angiotensin and myocardial protein synthesis, cardiac-hypertrophy in hypertension.
 New York, raven Press, 1983.
- Kleiman, R.B. and Houser, S.R. (1989):**
 Outward currents in normal and hypertrophied feline ventricular myocytes.
 Am. J. Physical 256 H1 450.
- Rleiman, R.B. and Houser, S.R. (1988):**
 Calcium currents in normal in hypertrophied isolated feline ventricular myocytes.
 Am. J. Physiol 255: H. 1434.
- Kojinam, M.; Shiojima, I.; Yamazaki, T.; Komuro, I.; Yunzeng, Z.; Ying, W.; Mizuno, T.; Uekik, Tobe, K.; Kadowaki, T.; Nagi, R. and yazaki, Y. (1994):**
 Angiotensin II. Receptor antagonist, TCV-116 induces regression of hypertensive left vnetircualr hypertrophy in vivo and inhibits the intracellualr signaling pathways of stretch mediated cardiomyocyte hypertrophy in vitro.
 Circulation 89: 2204.

- Komuro, I.; Katoh, Y.; Kaida, T.; Shibasaki, Y.; Kawogushi, M.; Hoh, E.; Tsakaku, F. and Yazaki, Y. (1991):**
Mechanical loading stimulates cell hypertrophy specific gene expression in cultured rat cardiac myocytes.
J. Biochem. 266: 1265.
- Komuro, I.; Kurabyashi, M.; Takaku, F. and Yazki, Y. (1988):**
Expression of cellular oncogene in the myocardium during the developmental stage and pressure. Over load hypertrophy of rats.
Circ. Res. 62: 1075.
- Kuulan, K.; Kansuaglu, B. and Tuncerc, C. (1996):**
Significance of Q.T. dispersion in ventricular tachyarrhythmias. In mitral valve. Prolapse. Int. J. Cardiol, 54 (3): 251-257.
- Kuo, C.S.; munakata, K. and Reddy, C.P. (1983):**
Characteristics and possible mechanisms of ventricular arrhythmia depend on dispersion of action potential duration.
circ. 67 (6): 1356: 1367.
- Kupari, M.; Kasknen and Viralamen, J. (1994):**
Correlates of left ventricular mass in a population sample aged 36 to 37 years. Focus on life style and salt intake.
Circulation 8: 1041.
- Lavie, C.J.; Ventura, H.O. and Messerli, F.H. (1993):**
Hypertension, obesity, left ventricular hypertrophy, complex ventricular ectopic activity and increased risk for sudden death review of ohasner studies and the literature.
J. Cardiopulm rehabil 13: 264.
- Lavie, C.J.; Ventura, H.O. and Messerli, F.H. (1991):**
Regression of increased left ventricular mass by antihypertensives.
Drugs. 42 : 945.
- Leibson, P.R. and Savage, D.D. (1986):**
Echocardiography in hypertension.
Echocardiography, Vol. 3: No. 3: 181-218.

- Levy, D. (1990):**
Stratifying the patient at risk from coronary heart disease.
Am. Heart, J. 119: 712.
- Levy, D. (1988):**
Left Ventricular hypertrophy: epidemiologic insights from the framingham heart study.
Drugs. 35 (Suppl. 5) 1.
- Levy, D.; Anerson, M.; Savage, D.D.; balkus, S.A.; Kammel, W.B. and Kastelli, W.P. (1987):**
Risk of ventricular arrhythmias in left ventricular hypertrophy, the framing ham study.
Am. J. Cardiol. 60: 560.
- Levy, D.; Labib, S.B.; Anderson, K.M.; Christiansen, J.C.; Kannel, W.B. and Castelli, W.P. (1990):**
Determinants of sensetivity and speceificity of eleanorcardiographic criteris for left ventricular hypertrophy.
Circulation 81: 815.
- Linker, N.J.; Colonna, P. andd Kekwick, C.A. (1992):**
Assesement of Q.T. dispersion in symptomatic patients with congenital long Q.T. syndroms.
Am. J. Cardiol, 69: 634-638.
- Lo, S.S.; Mathras, C.J. and John, S.M. (1996):**
Q.T. Interval dispersion in primary autonomic failure.
Heart.,75: 498-501.
- Lynch, J.J.; Parkewitz, D.A.; Gimbel, K.S. and Thomas, S.A. (1987):**
Physiological aspectsof cardiac arrhythmias.
Am. Heart., J. 93: 645.
- Macfarlane, P.w.; Veitch, Lawrie, T.D. eds (1989):**
The normal electrocardiogram and echocardiogram in comperehensive eleanorcardiology.
Vol. 1 New York Pergamon press. 451-52.

- Marcus, M.L.; Harrison, D.G.; Chilian, W.M.; Koyanogic, S.; inou, I. Tomanek, R.J.; martens, J.B.; Easthma, C.L. and Hirotzka, L.F. (1987):*
 Alternations in the coronary circulation in hypertrophied ventricles circualtion (suppl. 1) 1-19.
- Maron, B.J. and Roberts, W.S. (1979):*
 Quantitative analysis of cardiac muscle cell disorganization in ventricular septum of patients with hypertrophic cardiomyopathy. *Circulation* 590: 689.
- Mayet, J.; Shahi, M.; Mc. Grath, K.; Poulter, N.R.; Sever, P.S.; Foale, R.A.; Thom. S. A. 91996):*
 Left ventricular hypertrophy and Q.T. dispersion in hypertension *Hypertension*, no. 28 (5), 791-6.
- McLaughlin, N.B.; Compbell, R.W. and Murray (1995):*
 Comparison of automatic Q.T. measurement techniques in the normal 12 leads ECG.
Br. Heart, J., 74: 84-89.
- Mclenachan, J. M.; Henderson, E.; Morris, K.I. and Dargie, H.J. (1987):*
 Ventricular arrhythmias in patients with hypertensive left ventricular hypertrophy.
N. Engl. J. Med. 317: 787.
- Mclenachan, J.M. and Dargie, H.J. (1990):*
 Ventricualr arrhythmias in hypertensive left ventricular hypertrophy; relationship to coronary disease, left ventricular dysfunctions and myocardial fibrosis.
Am. J. Hypertens, 8: 735.
- Meerson, F.Z. and Breger, A.M. (1987):*
 The common mechanism of heart to adaptation and deadaptation, hypertrophy and atrophy of the heart muscle.
Basic, res cardiol 72-228.
- Messerli, F.H. (1983):*
 Clinical determinants and consequences of left ventricular hypertrophy. *An. J. med.* 75: 51.

- Messerli, F.H.; Numez, B.D.; Ventue, H.O. and Synder, D.W. (1987):**
Overweight and sudden death; increased ventricular ectopy in cardiomyopathy of obesity.
Arch. Intern. Med. 147: 1725.
- Messerli, F.H.; Ventura, H.O.; Elizardi, D.J.; Dunn, F.G. and Frohlich, E.D. (1984):**
Hypertension and sudden death, increased ventricular ectopic activity in left ventricular hypertrophy.
Am. J. med. 77: 18.
- Miller, J.T.; O'Rourke, R.A. and Grawford, M.H. (1988):**
Left atrial enlargement an early sign. Of hypertensive heart disease. Am. Heart. J. 116: 1048.
- Milliken, J.A.; Macfarlane, P.W. and Lawrie, T.O.V. (1989):**
Enlargement and hypertrophy an comprehensive electrocardiography oxford: pergamon, 631.
- Misserli, F.H. (1983):**
Clinical determinants and consequences of left ventricular hypertrophy. Am. J. Med. (75)-(51-56).
- Miorelli, M.; Bujo, g. and Melacini, P. (1994):**
Q.T. interval variability in hypertrophic cardiomyopathy. Patients with cardiac arrests.
Int. J. Cardiol. 45 (2): 121-127.
- Mirvis, D.M. (1985):**
Spatial variation of Q.T. intervals in normal persons and in patients with acute myocardial infarction.
J. Am. Coll. Cardiol, 3: 625-631.
- Mohamed, R.; Forsey, P.R. and Davis, M.K. (1996):**
Effect of liver transplantation on Q.T. interval prolongation and autonomic dysfunction in end stage. Liver disease.
Hepatology, 23 (5) : 1128-1134.
- Monlar, J.; Weiss, J.S.; Posinthal, J.E. and Somberg, J.C. (1996):**
Q.T. interval dispersion in survivors of sudden Cardiac Death. A 24 hour assesement.
J. Eur. Soc. Cardiol 17-29.

- Moore, J.W.; Hutchins, J.M.; Bulkely, B.M. (1984):**
 Constituents of Human ventricular myocardium, tissue hyperplasia accompanying muscular hypertrophy.
 Am. Heart. J. 100: 61.
- Moss, A.J. (1993):**
 Measurement of interval and the risk associated with Q.T. interval prolongation, a review Am. J. Card. 72/23-25.
- Murray, A.; mclaughlin, N. and Bourke, J.P. (1994):**
 Britich, Heart. Journal, 71: 386-390.
- Murray, J.A.; Johnston, W. and Ried, J.M. (1972):**
 Echocardiographic determination of left ventricular dimension, volumes. And performance.
 Am. J. Cardiol 30: 252.
- Nicolas, J.; Paolo, Colonna, Christopher, A. (1992):**
 Assesment of Q.T. dispersion syndrome in patients with congenital prolonged Q.T. syndrome.
 Am. J. Cardiol 69: 634-38.
- Niwa, H.; meeda, T. and Shimoyama, N. (1993):**
 Serial changes in Q.T. interval during acute myocardial infarction; comparison of Q.T. interval of reprefused and non reprefused cases.
 J. Cardiol, 23 (1): 14-50 (Abstr).
- Nunez, B.D.; lavie, C.J.; Messerli, F.H.; Schmieder, R.E.; Garavaglia, G.E. and Nunez, M. (1994):**
 Comparison of diastolic left ventricular filling and cardiac dysrhythmias in hypertensive patients with and without isolated septal hypertrophy.
 Am. J. Cardiol. 74 : 585.
- Oberg, K.C. and Bauman, J.L. (1995):**
 Q.T. Interval prolongation and trasades de pointes due to erythromycin. Lacto bionates pharmacotherapy.
 15: (6): 687-692.
- Oxford, M. (1986):**
 Hypertension, a major risk factor for coroanry heart disease.
 Am. J. Card. 50: 370-75.

- Perkiomaki, J.S.; Ikaheimo, M.J.; Pikkujamsa, S.M.; Rantalo, A. Lilja, M.; Kesaniemi, Y.A.; Huikuri, H.V. (1996):**
Dispersion of the Q.T. interval and automatic modulation of heart rate variability in hypertensives with and without left ventricular hypertrophy.
Hypertension Jul. 28 (1): 16-21.
- Pichard, A.D.; Gohim, R.; Smith, H.; Ambrase, J. and Meller, J. (1981):**
Studies in patients with left ventricular hypertrophy of the hypertensive type. Evidence for an impaired vascular reserve.
Am. J. Cardiol 47: 547.
- Priori, S.G.; Napolitano, C. and Diehl, L. (1994):**
Dispersion of the Q.T. interval, a Marker of therapeutic efficacy in the idiopathic long Q.T. syndrome.
Circ., 89 (4): 168-1689.
- Pye, M.; Quin, A.G. and Cabbe, S.M. (1994):**
Non invasive marker of susceptibility to arrhythmia in patients with sustained ventricular arrhythmias.
Br. Heart, J., 71 (6): 511.
- Rackley, C.E.; Dodge, H.T.; Cable, Y.D. and Hay, R.E. (1984):**
A method for determining left ventricular mass in man.
Am. J. Cardiol 69: 913.
- Ren, J.F.; Hakki, A.H.; Kotler, M.N.; Iskandrian, A.S. (1985):**
Exercise systolic blood pressure, a powerful determinant of increased left ventricular mass in patients with hypertension.
Journal of American College, of cardiology, 5: 1129.
- Rolands, D.B.; Ireland, M.A.; Stallard, M. (1982):**
Assessment of left ventricular mass and responses to antihypertensive.
Treatment Lancet, 1: 487.
- Romhilt, J.C.; Kleinman, L.H.; Fedor, J.M. (1978):**
Myocardial blood flow distribution in concentric left ventricular hypertrophy.
J. Clin. Invest. 62: 369.

- Ross, A.M.; Pisarczyk, M.J.; Calabresi, M. (1988):**
 Echocardiographic and clinical correlation in systemic hypertension journal of clinical ultrasound.; 6: 95.
- Safar, M.E.; Lehmer, J.P. Vincent, M.I. (1989):**
 Echocardiographic dimensions in borderline and sustained. Hypertension.
 The American. Journal of Cardiology 94-930.
- Sahn, D.J.; Demaria, a.; Kisslo, J.; Weymna, A. (1978):**
 Recommendations regarding quantitation in M mode.
 Echocardiograph. Results of a survey of echocardiographic measurement. Circulation 58, 1072-1083.
- Sausa, M.J.; Armaha, J. and Cotrin, C. (1995):**
 The QTC interval and its dispersion in hypertrophic cardiomyopathy, its relation to complex ventricular arrhythmias and the effect of antiarrhythmic agents.
 Rev. Port. Cardiol, 14 (6): 461-7. 477-8 (Abst).
- Savage, D.D.; Devereux, P.D.; Garrison, M.S. et al. (1987):**
 The spectrum of left ventricular hypertrophy in a general population, Sample the framingham study.
 Circulation 75 (suppl. 1) 1-26 & 1-40.
- Savage, D.D.; Drayer, J.I.M. Hanry, W.L. (1983):**
 Echocardiographic assesment of cardiac anatomy and function in hypertensive subjects.
 Circulations. 84. 21.
- Savage, D.D.; Drayer, J.I.M. and Honny, W.L. (1979):**
 Echocardiographic assesment of cardiac anatomy and function in hypertensive subjects.
 Circulation 59: 623.
- Schaller, W.A.; Cobba, L.A. (1975):**
 Recurrent vent fibrillation and modes of death in survivors of out of hospital ventricular fibrillation.
 N. Nj. J-Med. 293, 259-262.
- Schmider, R.E. and Messerli, F.H. (1992):**
 Determinants of ventricular ectopy in hypertensive cardiac hypertrophy.
 Am. Heart, J. 123: 89.

- Schmider, R.E.; Messerli, H.F.; Garavaglia G.E. and Nunez, B.D. (1988):**
Dietary salt intake. Adeterminant of cardiac involvement in essential hypertension.
Circulation 78: 951.
- Schwartz, P.J. and Wolf, S. (1978):**
Q.T. interval prolongation as predictor of sudden death in patients with myocardial infarction circulation 57, 1074-1072.
- Seedat, Y.K. Pillay, N. and Fojas, H.M. (1987):**
Myocardial infarction in the African hypertensive patients.
Am. Heart J. 94 (3): 398-399.
- Sen, S.; tanazi, R.C. and Rampus, F.M. (1989):**
Cardiac effect of angiotensin antagonist in normotensive rates Clin. Sci. 56: 439.
- Sharpiro, L.M. and Mckemna, W.J. (1984):**
Left ventricular hypertrophy. Relation of structure to diastolic function.
The British heart journal 51: 637.
- Smith, V.E.; Schulmna, P. Karimedдини, M.K. (1985):**
Rapid ventricular filling in left ventricular hypertrophy.
Journal of the American collage of Cardiology 5: 869.
- Simpson, P. (1983):**
Noradrenaline, Stimulated hypertrophy of cultural rat myocardial cells, in an alpha. 1 adrenergic response.
J. Clin. Invest. 72: 732.
- Statters, D.J.; Malik, M. and Ward, D.E. (1999):**
Q.T. dispersion, Problem. Of methodology and clinical significance/ J. cardiovasc. Electrophysiol, 5 (8) : 627-85. (Abstr.).
- Sterns, L.D.; Hilbel, T. and Horacek, M. (1997):**
Measures of Q.T. dispersion from signal averaged 12 leads ECGs are better discriminators than those measured from individual beats. J. Am. Coll. Cardiol, 29 (2) Suppl. A148-A.

Surawicz, B. (1996):

Will Q.T. dispersion play a role in clinical decision making
J. Cardiovasc. Electrophysio., 7 (8): 777-84.

Swan, L.; Birnie, D. and Connell, J. (1997):

Familial influences on Q.T. and Q.T. dispersion.
J. Am. Coll. Cardiol, 29 (2) A suppl: 82 a.

Unverferth, D.V.; Majorien, R.D.; Lewis, R.P. and Leier, C.V. (1983):

The role of subendocardial ischemia in perpetuating
myocardial failure. In patients with non ischaemic
cardiomyopathy.
A.M. Heart, J. 105: 176.

Van, De, Loo, Klinghenheben, T. and Hohnlaser, S.H. (1994):

Amiodarone therapy after sotalol induced Torsade
depointes prolonged Q.T. interval and Q.T. dispersion in
differentiation of proarrhythmic. Effect.
Z. Cardiol, 83 (12) : 887-90 (Abst.).

Viklas, Boas, F.; Lima, A.A.; Torreaaj, Feitosa, G.S.(1997):

Q.T. dispersion in patients with systemic hypertension.
Arq. Bras, Cardiol, may. 58 (5) 343-6.

Warner, J.P.; Barnes, T.R. and Henry, J.A. (1996):

Electorcardiographic changes in patients receiving
neuroleptic medications.
Acta. Psychiatr. Secand 93 (4) 311-3 (Abstr.).

**Weber, K.T.; Janicki, J.S.; Shroff, S.G.; Pick, R.; Chen, R.M. and
Bashey, R.I. (1988):**

Collagan remodeling of the pressure. Over load.
Hypertophied non human. Primate myocardium.
Circ. Res. 62: 757.

Wit, A.L. and Coromilas, J. (1993):

Role of alternaiton sin refractoriness and conduction in the
genesis of rentront arrhythmias, in plication for
antiarrhythmemic effects of class III. Drugs.
Am. J. Cardiol. 72: 3 F.

Yamari, Y.; Tarzai, R.C. and Ooshima, A. (1981):

Effect of beta-receptro blocking agents on cardioovascular structural changes. In spontaneous and noradrenalme-induced hypertension in rats.
Clin. Sci. 59:457.

Yanagisawa, M.; Kurithara, H.; Kimura, S.; tomobek, Y.; Kobayashi, M.; Mitsui, Y.; Yazaki, Y.; Gata, K.; Masaki, I. (1988):

A noval potent vasoconstrictor peptide prodced by vascular endothelialc ells. Nature 332: 411.

Yelamanchi, V.; Ranado, V. and Cao, W. (1997):

Influecne of electorlyte abnormalities on inteerlead variability of ventricular repolarization times. In 12 leads Ecg. J. Am. Col. Cardiol. 29 (2) suppl. A: 149 A.

Yotsukura, M.; Toamno, Y. and Yamauchi, T. (1997):

Q.T. dispersion and vnetircular arrythmias in patients. With progressive muscular dystophy of duchenne type J. Am. Coll. Cardiol 29 (2) suppl. A: 149 A-1997.

Zareba, W. and Maass, A.J. (1997):

Dispersion of repolarization duration, relation to heart rate and repolarization duration.
J. Electorcardial, 28 Suppl. 202-6.