

REFERENCES

American Heart Association (1990):

Heart Facts. Dallas, American heart Association National center P. 1.

Apple, F.S.; Voss, E. and Lund, L. et al., (1995):

Cardiac troponin, CK-MB and myoglobin for the early detection of acute myocardial infarction and monitoring of reperfusion following thrombolytic therapy.
Clinical Chemistry Acta., 237: 59-66.

Assoian, R.K.; Grotendorst, G.R.; Miller, D.M.; miller, D.M. and Sporn, M.B. (1984):

Cellular transformation by coordinated action of three peptide growth factors from human platelets.
Nature 309: 804.

Babior, B.M. (1978):

Oxygen dependent microbial killing by phagocytes.
N. Engl. J. Med. 288: 668, 721-725.

Bagly, G.C. and Segal, G.M. (1995):

Growth factors and the cotnrol of haematopoiesis in: Haematology basic principle & Practice. Eds. (Hoffmann. R.; Benz, E.J.; Shattil, S.J.; Furie, B., Cohen H.J. and Siberstein L.E.) 2nd edition. Churchill fiving stone P: 207.

Bartelmesz, S.H.; Bradley, T.R. and Bertoncello, I. (1989):

IL-1 plus IL-3 plus CSF.1 are essential for clonal proliferation of primitive myeloid bone marrow cells.
Exp. Hematology, 17: 240.

Bednar, M.; Smith, B.; Pinto, A. and Mullane, K.M. (1985):

Nafazatrom-induced salvage of ischemic myocardium in anesthetized dogs is mediated through inhibition of neutrophil function.
Circ. Res. 57: 131-141.

- Bell, D.; Jackson, M.; Nicoll, J.J.; Millar, A.; Dawes, J. and Muir, A.L. (1990):**
Inflammatory response, neutrophil activation and free radical production after acute myocardial infarction. Effect of thrombolytic treatment.
Br. Heart J. 63: 82-87.
- Bertagolli, M.; Lin, B. and Young, D. (1992):**
IL-12 augment antigen dependent proliferation of activated T-lymphocytes.
J. Immunol., 140: 3778.
- Bevilacqua, M.P.; Pober, J.S.; Cotran, R.S. and Gimbrone, M.A. (1985):**
Interleukin-1 acts upon vascular endothelium to stimulate procoagulant activity and leukocyte adhesion.
J. coll. Biochem. 1A: 148.
- Bomsztyk, K.; Sims, J.E. and Stanton, T.H. (1989):**
Evidence for different interleukin. 1 receptors in murine B and T-cell lines.
Proc. Natl. Acad. Sci. U.S.A. 86: 8034.
- Boom, W.; Liano, D. and Alloas, A. (1988):**
Heterogeneity of helper./ inducer T-lymphocytes.
J. Exp. Med., 167: 1352.
- Bot, F.J.; Dorssers, L. and Wagemaker, G. (1988):**
Stimulating spectrum of human recombinant multi CSF (IL-3) on human marrow precursors: importance of accessory cells.
Blood, 71: 1609.
- Brownwold, E. (1996):**
Pathogenesis of atherosclerosis. A textbook of cardiovascular medicine by Brownwold, E., Fifth edition chapter 34 page 1110.
- Brownwold, E. and Ganz, P. (1996):**
Reperfusion injury, A textbook of Cardiovascular Medicine by Braunwold, E., Fifth edition Chapter 36 page 1179.

- Braunwold, E. and Kloner, R.A. (1985):**
Myocardial reperfusion, A double edged sword.
J. Clin invest 76: 1713-1719.
- Brounwold, E. and Sobel, E. (1992):**
Coronary blood flow and myocardial ischemia. A textbook of cardiovascular medicine by Brounwould, E., forth edition chapter 38, P. 1187.
- Campbell, M.J. and Mackin, D. (1993):**
Medical statistics A common sense approach 2nd ed.
John Wiley and sons New York.
- Campbell, H.D.; Tucker, W.Q.J.; Hort, Y. et al., (1987):**
Molecular cloning and expression of the gene encoding human eosionphil differentiation factor.
Proc. Natl. Acad. Sci., U.S.A., 84: 6629.
- Cathcart, M.K.; Morel, D.W. and Chisolm, G.M. (1985):**
Monocytes and neutrophils oxidize low density lipoprotein making it cytotoxic.
J. leukoc. Biol. 38: 341.
- Chan, S.; Kolyashi, M. and Santoli, D. (1992):**
Mechanism of IFN and induction of IL-12 role of transcription and mRNA stability in the synergistic interaction between NKSF (IL-12) and IL-2.
J. Immunol., 148: 92.
- Chen, W.F. and Zlotnick, A. (1991):**
A novel cytotoxic T cell differentiation factor.
J. Immunol., 147: 528.
- Clinton, S.K.; Fleet, J.C.; Loppnow, H.; salomon, R.N; Clark, B.D.; Cannon, J.G.; Shaw, A.R.; Dinarello, C.A. and Libby, P. (1991):**
Interleukin-1 gene expression in rabbit vascular tissue in vivo.
Am. J. Pathol. 138: 1005-10014.
- Cooper, R.J.; Gordon, M.S. and Weber, J. (1993):**
The in vivo adminstration of recombinant human IL-6 to patients with advanced malignancies has a profound effect on haemopoiesis.
Exp. Hematol., 21: 1117.

Daveenport, R.D.; Strieter, R.M.; standiford, T.J. and Kunkel, S.L. (1991):

IL-8 production in red blood cell in compatibility.
Blood, 76: 2439.

Detmers, P.A.; Los, K. and Cohn, Z.A. (1990):

Neutrophil-activating protein: IL-8 Stimulates the bunding activity of the leukocyte adhesion receptor CD11, CD8 on human neutrophils.
J. Exp. Med, 171: 1155.

Dinareello, C.A. (1988):

Biology of IL-1.
FASEB. J., 2: 108.

Dinerman, J.L.; Mehra, J.L.; Saldeen, T.G.; Emerson, S.; Wallin, R. and davda, R. (1990):

Increased neutrophil elastrase release in unstable angina pectoris and acute myocardial infarction.
J. Am. Coll. Cardiol. 15: 1559-63.

Donahue, R.E.; Yang, Y.C. and Clark, S.C. (1990):

Human P 40 T-cell growth factor (IL-9) supports erythroid colony formation.
Blood, 75: 2271.

Dreyer, W.J.; Michael, L.H.; Nguyen, T.; Smith, C.W.; Anderson, D.C.; Entman, M.L. And Rossen, R.P. (1992):

Kinetics of C5a release in cardiac lymph of dogs experiencing coronary artery ischemia reperfusion injury.
Circ. Res. 71: 1518-1524.

Dreyer, W.j.; Smith, C.W.; Michael, L.H.; Rossen, R.D.; Hughes, B.J.; Entman, M.L. and Anderson, D.C. (1989):

Canine neutrophil activation by cardiac lymph obtained during reperfusion of ischemic myocardium.
Circ. Res. 65: 1751-1762.

Du, X.X.; Neben, T.; Goldman, S. and Williams, D.A. (1993):

Effects of IL-11 on hematopoietic reconstitution in transplant mice: Acceleration of recovery of peripheral blood neutrophil and platelets. Blood, 81: 27.

- Elner, V.M.; einer, S.G. and Kunkel, S.L. (1990):***
Neutrophil chemotactic factor (IL-8) gene expression by cytokine treated retinal pigment epithelial cells.
Am. DJ. Path., 135: 745.
- Engler, R.E. (1987):***
Granulocytes and oxidative injury in myocardial ischemia and reperfusion.
Fed. Proc. 46: 2395-2396.
- Engler, R.I.; Dahlgren, M.D.; Morris, D.; Peterson, M.A. and Schmial-Schoenbein, G. (1986):***
Role of leukocytes in the response to acute myocardial ischemia and reflow in dogs.
Am. J. Physiol. 251: H 314-H 323.
- Ensrud, K.; Richard, H.; Grimm, J.R.; Minneapolis, Minn (1992):***
The white blood cell count and risk for coronary heart disease.
Am. Heart. J. July 207: 213.
- Enteman, L.M.; Youker, K.; Kukielka, L.G. and Ballantyne, M.C. (1993):***
Reaction to injury in cardiovascular disease: A therapeutic challenge.
Accrrent Journal September /October, P. 13:15.
- Fallon, J.T. (1996):***
Pathology of myocardial infarction and reperfusion. In faster, V. Ross, R. and Topol, E.J. (eds): Atherosclerosis and coronary Artery Disease. Philadelphia, lippincott- Roven, PP. 791-796.
- Fanone, J. C. and Ward, P.A. (1984):***
Role of oxygen free radicals and metabolites in leukocyte dependent inflammatory reaction.
Am. J. Pathol. Vol. 107 No. 3: 397-418.
- Feldman, M. (1996):***
Cytokines. Immunol. Today June (3) 11.8-11.10.

- Fiorentino, D.; Zlotnik, A. and Viera, P. (1991):**
IL-10 acts on the antigen presenting cells to inhibit cytokine production by Th₁ cells.
J. Immunol., 146: 3444.
- Garabedion, H.D.; Gold, H.K. and Yasuda, T. (1988):**
Detection of coronary artery reperfusion with creatine kinase-MB determination during thrombolytic therapy: correlation with acute angiography.
J. Am. Coll. Cardiol., 11: 729-734.
- Gately, M.; Desai, B. and Quinn, R. (1991):**
Regulation of human lymphocyte proliferation by a heterodimeric cytokine IL-12 (CLME).
J. Immunol., 147: 874.
- Gown, A.M.; Tuskada, T. and Ross, R. (1986):**
Human atherosclerosis II. Immunocytochemical analysis of the cellular composition of human atherosclerotic lesions.
Am. J. Pathol. 125:191-207.
- Gubler, U.; Chua, A.O. and Schoenhout, D.S. (1991):**
Coexpression of two distinct genes is required to generate secreted bioactive cytotoxic lymphocyte maturation factor.
Proc. Natl. Acad. Sci. U.S.A., 88:4143.
- Hansson, G.K.; Jonasson, L.; Holm, J.; Clowes, M.M. and Clowes, M.M. and Clowes, A.W. (1993):**
Gamma-interferon regulates vascular smooth muscle proliferation and Ia antigen expression in vivo and in vitro.
Circ. Res., 63: 712-719.
- Hansson, G.K.; Holm, J. and Janasson, L. (1989):**
Detection of activated T-lymphocytes in the human atherosclerotic plaque.
Am. J. Pathol. 135: 169.
- Hansson, G.K.; Seifert, P.S.; Olsson, G. and Bondjers, G. (1991):**
Immunohistochemical detection of macrophages and T lymphocytes in atherosclerotic lesions of cholesterol fed rabbits.
Arterioscl. Thromb. 11: 745-750.

- Haple, A.J.; Osborn, J.M. and Fung, M.C. (1985):**
Expression of 20- hydroxsteroid dehydrogenase in mouse macrophages, hemopoietic cells and cell lines and its induction by CSF.
J. Immunol., 134: 92.
- Hearse, D.J. and Bolli, R. (1992):**
Reperfusion-induced injury: Manifestations, mechanisms and clinical relevance.
Cardiovasc. Res. 26: 101.
- Hearse, D.J. (1977):**
Reperfusion of the ischaemic myocardium.
J. Mol cell cardiol. 9: 605-615.
- Hébert, C.A.; Lusinskas, F.W.; Kiely, J.M.; Luis, C.C.; Obin, M.S.; Gimbrone, M.A. and Baker, J.B. (1990):**
Endothelial and leukocyte forms of IL-8 conversion by thrombin and interaction with neutrophils.
J. Immunol., 145: 3033.
- Hill, A.B. (1979):**
In "A short text book of medical statistics 10th edition". The English language Book Society and Hodder and Stoughton.
- Hoang, T.; Haman, A. and Gancalves, O. (1988):**
IL-6 enhances growth factor dependent proliferation of the blast cells of acute myeloblastic leukemia.
Blood, 72: 823.
- Holbrook, S.T.; Ohls, R.K. and Yang, Y.C. (1991):**
Effect of IL-9 on clonogenic maturation and cell cycle status of fetal and adult hematopoietic progenitors.
Blood 77: 2129.
- Hoshida, S.; Kuzuya, T. and Kin, Y. (1993):**
Transcardiac alteration of neutrophil function relates to myocardial ischemia reperfusion injury.
Cardiovascular research, 27: 377-383.
- Huber, A.R.; Kunkel, S.T.; Todd, R.F. and Weiss, S.J. (1991):**
Regulation of transendothelial neutrophil migration by endogenous IL-8.
Science 254: 99-102.

Ida, N.; Sakurai, S.; Hosoi, K. and Kunitomo, T. (1992):

A highly sensitive enzyme- Linked immunosorbent assay for measurement of interleukin-8 in biological fluids.
J. Immunol. Meth. 156: 27-38.

Ihle, J.N. (1983):

Biochemical and biological properties of IL-3 A lymphokine mediating the differentiation of a lineage of cells which includes prothymocytes and mast like cells.
J. Immunol., 131: 282.

Jabara, H.H.; Ackerman, S.J. and Vercelli D. (1988):

Induction of IL-4 dependent IGE synthesis and IL-5 dependent eosinophil differentiation by supernatants of a human helper T-cell clone.
J. Clin. Immunol., 8: 437.

Jakubowski, A.A. and Golde, D.W. (1995):

Therapeutic use of cytokines. In William's. J. Beutler, E.; Arsleve, A.; Lichtman, M.A. (eds) Hematology.
Mc Grow-hill Book Company P. 155.

Jenings, R.B.; Schaper, J.; Hill, M.L., et al. (1985):

Effect of reperfusion late in the phase of reversible ischemic injury: Changes in cell volume, electrolytes, metabolites and ultrastructure.
Circ. Res. 56: 262.

Jolly, S.R.; Kane, W.J.; Hook, B.G.; Abrams, G.D.; Kunkel, S.L. and Lucchesi, B.R. (1986):

Reduction of myocardial infarct size by neutrophil depletion: Effect of duration of occlusion.
Am. Heart. J. 112: 682-690.

Jonasson, L.; Holm, J.; Skalli, O. et al., (1986):

Regional accumulations of T-cells, macrophages and smooth muscle cells in the human atherosclerotic plaque.
Arteriosclerosis. 6: 131-138.

Kimberly, J.V.Z.; Fischer, E.; Howes, A.S. et al.(1992):

Effect of intravenous IL-8 administration in non-human primate.
J. Immunol., 148:1746.

Klein, J. (1991):

Lymphokines and their receptors. In Roitt, I.; Brostoff, J. and Male, D. (eds).
Immunology Mosby P: 1019.

Kloner, R.A.; Ganote, C.B. and Jennings, R.B. (1974):

The "no reflow" phenomenon after temporary coronary occlusion in the dog.
J. Clin. Invest. 54: 1496-1508.

Kloner, R.A.; Hale, S.; Alker, K. et al., (1992):

The effects of acute and chronic cocaine use on the heart.
Circulation 85: 407.

Kloner, R.A.; Ellis, S.G.; Lange, R. and Braunwald, E. (1983):

Studies of experimental coronary artery reperfusion. Effects on infarct size, myocardial function, biochemistry, ultrastructure and microvascular damage.
Circulation 68 (Suppl. 1): 8.

Kloner, R.A.; Ellis, S.G.; Calson, N.V. and Braunwald, E. (1983):

Coronary reperfusion for the treatment of acute myocardial infarction: postischemic ventricular dysfunction.
Cardiology. 70: 233.

Kuijpers, T.W.; Hakkert, B.C.; Hart, M.H.C. and Ross, D. (1992):

Neutrophil migration across monolayers of cytokine-prestimulated endothelial cells.
A role of platelet-activating factor and IL-8.
J. Cell. Biol. 117: 565-572.

Kukerja, C. and Hess, L. (1992):

The oxygen free radical system: from equations through membrane-protein interactions to cardiovascular injury and protection.
Cardiovasc. Res. 26: 641-655.

Kukielka, G.L.; Smith, C.W.; Lo Rosa, G.L.; Manning, A.M.; mendoza, L.H.; Daly, T.J.; Hughes, B.J.; Youker, K.A.; Howkins, H.K.; Michael, L.H.; Rot, A. and Entman, M.L. (1995):

Interleukin-8 gene induction in the myocardium after ischemia and reperfusion in vivo.
J. Clin. Invest. 95: 98-103.

Kukielka, G.L. and Entman, M.L. (1994):

Adhesion molecule- dependent cardiovascular injury. In cellular Adhesion: Molecular Definition to therapeutic potential. Poste, G. and Metcalf, B., editors. Plenum Publishing Corp. New York. 187-212.

Lavin, F.; Kane, M. and Forde, A. et al., (1995):

Comparison of five cardiac markers in the detection of reperfusion after thrombolysis in acute myocardial infarction. Br. Heart. J., 73: 422-427.

Le Beau, M.M.; Lemons, R.S. and Espinosa, R. (1989):

IL-4 and IL-5 map to human chromosome 5 in a region encoding growth factors and receptors and are detected in myeloid leukemias with a del (5q). Blood, 73: 647.

Lee, T.H.; Rouan, G.W. and Weiberg, M.C. et al. (1987):

Sensitivity of routine clinical criteria for diagnosing myocardial infarction within 24 hours of hospitalization. Ann. Intern. Med., 106: 181-186.

Lefer, D.J.; Suresh, M.L.; Shandelya, M.L.; Serrano, C.V.; Becker, L.C.; Kuppusamy, P. and Zweier, J. L. (1993):

Cardioprotective actions of a monoclonal antibody against CD-18 in myocardial ischemia-reperfusion injury. Circulation 88: 1779-1787.

Lefer, A.M.; Tsao, P.S.; Lefer, D.J. and Ma, X.L. (1991):

Role of endothelial dysfunction in the pathogenesis of reperfusion injury after myocardial ischemia. FASEB, J. 5: 2029

Leonard, E.J.; Sheel, A.; Yoshimura, T.; Noer, K.; Kutvirt, S. and Van Epps, D. (1990):

Leukocyte specificity and binding of human neutrophil attractant. Activation protein-1. J. Immunol. 144: 1323-1330.

Lewis, D.B.; Prickett, K.S. and Lørsen, A. (1988):

Restricted production of IL-4 by activated human T-cells. Proc. Natl. Acad. Sci, U.S.A., 85: 9743.

- Lindsten, T.; June, C.H. and Ledbetter, J.A. (1989):**
Regulation of lymphokine messenger RNA stability as a surface-mediated T-cell activation pathway.
Science, 244: 339.
- Lumpkin, M.D. (1981):**
The regulation of ACTH secretion by IL-1.
Science, 238: 452.
- Mac Neil, I.; Suda, T.; Moor, K. et al. (1992):**
IL-10, a novel growth factor for mature and immature T-cells.
J. Immunol. 145: 4167.
- Mair, J.; Artner, D worzak, E. and Dienstle, A. et al. (1991):**
Early detection of acute myocardial infarction by measurement of mass concentration of creatine kinase-MB.
Am. J. Cardiol, 68: 1545-1550.
- Malech, H.L. and Gallin, J.I. (1987):**
Neutrophils in human diseases.
N. Engl. J. Med. 317: 687-694.
- Mario, J.B.; Mikael, L.A.; Christopher, G.S.; Frank, M.T.; Inge, O.; Stephen, P.E. and Fletcher, S.J. (1993):**
IL-1B induces IL-1 receptor antagonist, tumour necrosis factor binding protein in human.
Cancer Res., 53: 4010.
- Martin, T.R.; Altman, L.C.; Albert, R.K. and Henderson, W.R. (1984):**
Leukotriene B₄ production by human alveolar macrophage: a potential mechanisms for amplifying inflammation in the lung.
Am. Rev. Respir. Dis. 129: 106.
- Matsuhima, K.; Morishita, K.; Yoshimura, T.; Lavu, S.; Kobayashi, Y. and Lew, W. (1988):**
Molecular cloning of a human monocyte-derived neutrophil chemoattractant factor (MDNCF) and the induction of MDNCF mRNA by interleukin 1 and tumor necrosis factor.
J. exp. Med. 167: 1883-1893.

Mc Corcky, P.L. (1994):

Down regulation of cytokines action in: cytokines and Growth Factors.

Palliers, Clinical haematology, Palliers C indoll., 7 (1): 472.

Mc Cord, J.M. (1987):

Oxygen derived free radicals: a link between reperfusion injury and inflammation.

Fed. Proc., 46: 2402-2406.

Mc Cord, J.M. (1985):

Oxygen-derived free radicals in post-ischemic tissue injury.

N. Engl. J. Med., 312: 159-163.

Mckenize, A.N.J.; Culpepper, J.A.; De Wall Malefyt, R. et al. (1993):

IL-13 a novel T-cell derived cytokine that regulates human monocyte and B-cell function.

Proc. Natl. Acad. U.S.A. 90: 3735.

Mc-Mahan, C.J.; Slack, J.L. and Mosley, B. (1991):

A novel IL-1 receptor, cloned from B-cells by mammalian expression is expressed in many cell types.

EMB O.J., 10: 2821.

McNiece, I.K.; Bertoncello, I.; Kriegler, A.B.; et al. (1990):

Colony forming cells with high proliferative potential.

Int. J. cell cloning, 8: 146.

Metacalf, D. (1989):

The molecular control of cell division, differentiation commitment and maturation in hemopoietic cells.

Nature (Lond), 339:27.

Miller, M.D. and krangel, M.S.; (1992):

Biology and biochemistry of the chemokines: a family of chemotactic and inflammatory cytokines.

Crit. Rev. Immunol., 12: 17-46.

Minick, C.R. and Murphy, G.E. (1973):

Experimental induction of atheroarteriosclerosis. By the synergy of allergic injury to arteries and lipid-rich diet. II. Effect of repeated injections of horse serum in rabbits fed a lipid-risk, cholesterol-poor. Diet.

Am. J. Pathol. 73: 265.

- Minty, A.; Chalon, P.; Derocy, J.M.; et al. (1993):**
IL-13 is a new human lymphokine regulating inflammatory and immune response.
Nature, 362: 248.
- Mkhitarian, L.S.; Kornilina, E.M.; Dubrovshaice, G.V. and Babov, K.D., (1992):**
Autoimmune process in patients with different forms of the clinical course of ischemic heart disease.
Vrach. Delo. Mar. (3): 12-15.
- Mortensen, R.F.; Shopiro, J. and Lin, B.F. (1988):**
Interaction of recombinant IL-1 and recombinant TNF in the induction of mouse acute phase protein.
J. Immunol., 140: 2260.
- Moyer, C.F.; Sajuthi, D.; Tulli, H. and Williams, J.K (1992):**
Synthesis of IL-1A and IL-1B by arterial cells in atherosclerosis.
Am. J. Path., 138: 951-960.
- Mullane, K.M.; Salmon, J.A. and Kraemer, KR. (1987):**
Leukocyte-derived metabolites of arachidonic acid in ischemia-induced myocardial injury.
Fed. Proc. 46: 2422-1433.
- Nathan, C.F.; Murray, H.W. and Cohn, Z.A. (1980):**
Current concepts: The macrophage as an effector cells.
N. Engl. J. med. 303: 622.
- Niemeyer, C.M.; Sieff, C.A. Mathey, Prevot, B. et al. (1989):**
Expression of human IL-3 is restricted to human lymphocytes and T-cell tumor line.
Blood, 73: 945.
- Ohman, E.M.; Casey, C.; Bengtson, J.R.; Pryor, D.; Toomey, W. and Horgan, J.H. (1990):**
Early detection of Acute myocardial infarction.
Additional diagnostic infarction by serum myoglobin in patients without ST elevation.
Br. Heart. J., 63: 335-338.

- Oppenheim, J.J.; Ruscette, F.W. and Faltynek, C. (1994):**
Cytokine in: basic and clinical immunology. Eds (stites, D.P.; Ters A.T and Parslow, T.G.)
Eight edition Appleton and Lange P: 105. Paccud: J.P.
- Paccaud, J.P.; Schifferli, J.A. and baggiolini, M. (1990):**
NAP. 1/IL-8 induces up-regulation of CRI receptors in human neutrophil leukocytes.
Biochem. Biophys. Res. Comm. 166: 187-192.
- Paul, S.R.; Bennett, F.; Keller, K. et al. (1990):**
Molecular cloning of ac-DNA encoding IL-11 a stromal cell-derived lymphopoietic and hemopoietic cytokine.
Proc. Natl. Acad. Sci., U.S.A., 87: 7512.
- Pedoe-Tunstal, H.; Kuulasmaa, K.; Amouyel, P. et al. (1994):**
Myocardial infarction and coronary deaths. In the world health organization MONICA project. Circulation, 90: 583.
- Peschel, C.; Green, I.; Ohara, J. et al. (1987):**
Role of B-cell stimulatory factor I. IL-4 in clonal proliferation of B-cells.
J. Immunol., 139: 3338.
- Petit-Frere, C; Ongas, B. and Mencia-Huerta, J. (1993):**
IL-9 potentiates the IL-4 induced IgE and IgG release from murine B-lymphocytes.
J. Immunol., 79: 146.
- Petrov, D. (1990):**
The prognostic significance of high leukocytosis in acute myocardial infarction.
Vutr. Boles. 29 (1): 56.
- Peveri, P.; Waltz, A.; Dewald, B. and Baggiokini, M. (1988):**
A novel neutrophil activating factor produced by human mononuclear phagocytes.
J. Exp. Med., 167: 1547-1550.
- Quesenberry, P.J. (1995):**
Hemopoietic stem cells progenitor cells and cytokines, in william's, J.; Beutler, E. Erslev, A.J.; Lichtman, M.A. (eds).
Hematology McGraw-Hill book company P. 211.

- Ranies, E.W.; Dower, S.K. and Ross, R. (1989):**
Interleukin-1 mitogenic activity for fibroblasts and smooth muscle cells is due to PDGF-AA.
Science (Wash). 243:393-396.
- Romson, J.L.; Hook, B.G.; Kunkel, S.I.; Abrams, G.D.; Schork, M.A. and Lucchesi, B.R. (1983):**
Reduction of the extent of ischemic myocardial injury by neutrophil depletion in the dog.
Circulation, 67: 1016-1023.
- Rook, G.; Roitt, I.; Brostoff, J. et al. (1996):**
Cytokine network: cytokines form a complex communication network between cells.
Immun. 96: 212.
- Ross, R.; Raines, E.W. and Bowen-Pope, D.F. (1986):**
The biology of platelet-derived growth factor.
Cell 46: 155.
- Schall, T.J.; Bacon, K.; Comp, R.D.R. et al. (1993):**
Human macrophage inflammatory protein-1 α (MIP-1 α), MIB-1B chemokines attract distinct populations of lymphocytes.
J. Exp. Med., 177: 1821.
- Schröder, J.M. (1989):**
The monocyte -derived neutrophil activating peptide (NAP/ Interleukin-8 stimulates human neutrophil arachidonate-5-lipoxygenase, but not the release of cellular arachidonate.
J. Exp. Med., 170: 847-863.
- Schröder, J.M.; Mrowietz, U.; Morita, E. and Christophers, E. (1987):**
Purification and partial biochemical characterization of a human monocyte derived neutrophil activating peptide that lacks interleukin-1 activity.
J. Immunol., 139: 3474-3483.
- Seigel, J.; Happer, M.E. and wong stall, F. (1984):**
Gene for T-cell growth factor location on human chromosome 4q.
Science, 223: 175.

Sckido, N.; Mukaida, N.; harada, I.; Nakanishi, Y.; Watanabe, Y. and Matsushima, K. (1993):

Prevention of lung reperfusion injury in rabbits by a monoclonal antibody against interleukin-8.
Nature (Lond). 365-654-657.

Sieff, C.A.; Neimyer, C.M. and Nathan, D.G. (1987):

Stimulation of human hematopoietic colony formation by recombinant gibbon multi-colony stimulating factor or interleukin-3.
J. Clin. Invest., 80: 818.

Smith, K.A. (1988):

IL-2 inception impact and implications.
Science, 240: 1169.

Sonoda, Y.; Arai, N. and Ogawa, M. (1989):

Humoral regulation of eosinophilopoiesis in vitro, analysis of the targets of IL-3, granulocyte/macrophage colony stimulating factor (GM-CSF) and IL-5.
Leukemia, 3:14.

Springer, T.A. (1994):

Traffic signals for lymphocyte recirculation and leukocyte emigration: the multistep paradigm.
Cell. 76: 301-314.

Stemme, S.; Holm, J. and Hansson, G.K. (1992):

T-lymphocytes in human atherosclerotic plaques are memory cells expressing CD45 RO and the integrin VIA-1.
Arterioscl. Thromb., 12: 206-211.

Sutherland, G.R.; Baker, E. and Callen, D.F. (1988):

IL-5 is at 5q 31 and is detected in the sq-syndrome.
Blood, 71: 1150.

Swain, S.L. and Dutton, R.W. (1982):

Production of a b cell growth promoting activity from a cloned T-cell line and its assay.
J. Exp. Med., 156: 1821.

- Takatsu, K.; Tominaga, A. and Mamaoka, T. (1980):**
Antigen induced T-cell replacing factor (TRF).
J. Immunol., 124: 2414.
- Thompson-Sniger, L.A.; Dhar, V.; bond, M. et al. (1991):**
IL-10: A novel stimulatory factor for mast cells and their progenitors.
J. Exp. Med. 173: 507.
- Tovernier, J.; Devos, R. and Cornelis, S. (1991):**
A human high affinity interleukin-5 receptor (IL-5 R) is composed of an IL-5 specific α chain and B chain shared with the receptor for GM-CSF.
Cell., 66: 1175.
- Tsugiyasu, K.; Yoshio, H.; Shigeru, O.; Kazuo, Y.; Koichi, T.; Ryozo, N. and Lsao, K. (1996):**
Interleukin-8 as a sensitive marker of unstable coronary artery disease.
The Am. J. of card. 77: 304-307.
- Van der wal, A.C.; Das, P.K.; Van de Berg, D.B.; Van der loos, C.M. and Becher, A.F. (1992):**
Atherosclerotic lesion in human, in situ immunophenotypic analysis suggesting an immune mediated response.
Lab. Invest. 61: 166-170.
- VanLenten, B.J. and Fogelman, A.M. (1992):**
Lipopolysaccharide induced inhibition of scavenger receptor expression in human monocyte, macrophages is mediated through tumor necrosis factor alpha.
j. Immunol., 148: 112-116.
- Waldman, T.A.; Goldman, C.K.; Robb, R.J. et al. (1984):**
Expression of Il-2R on activated human B-cell.
J. Exp. Med., 160: 1450.
- Wasserman, S.I. (1985):**
Mediators of immediate hypersensitivity.
J. allergy clin. Immunol., 72: 101.

- Wèisman, H.F.; Barton, T.; Leppo, M.K.; Marsh, H.C.; Carson, G.R.; Concino, M.F.; Boyle, M.P.; Roux, K.H.; Weisfeldt, M.L. and Fearon, D.T. (1990):**
Soluble human complement receptor type 1: in vivo inhibitor of complement suppressing post-ischemic myocardial inflammation and necrosis.
Science (Wash, DC), 249: 146-151.
- Werns, S.W.; Shea, M.J. and Lucchesi, B.R. (1986):**
Free radicals and myocardial injury: pharmacologic implications.
Circulation, 74: 1-5.
- Whasi, S.; kaul, N.; Ganguly, N.K.; Varma, S.; Sharma, B.K. and Wahi, P.L. (1991):**
Neutrophil oxygen free radical production proportionates with the degree of myocardial ischemia.
Can. J. Cardiol. Jun.; 7 (5): 229-233.
- Yasunori, A.; Masanolu, K.; Masatoshi, K.; Tsugumichi, Y.; Mikihiisa, F.; Hisoyuki, K.; Toshio, Y.; Akiyohi, K.; Muneyosu, S. and Koyi, M.; (1993):**
Transient rise in serum interlukin-8 concentration during acute myocardial infarction
Br. Heart. J. 70: 132-134.
- Yokota, T.; Arai, N. and De vries, J. (1988):**
Molecular biology of IL-4 and IL-5 genes and biology of their products that stimulate B-cells, T-cells and hemopoietic cells.
Immunol. Rev., 102: 137.
- Yoshimura, T.; Matsushima, K.; Tanaka, S.; Roleinson, E.R.; Appella, E.; Oppenheim, J.J. et al. (1987):**
Purification of a human monocyte-derived neutrophil chemotactic factor that has peptide sequence similarity to other host defense cytokines.
Proc. Natl. Acad. Sci. U.S.A. 84: 9233-9237.
- Youker, K.; Smith, C.W.; Anderson, D.C.; Miller, D.; Michael, L.H.; Rossen, R.D. and Entman, M.L. (1992):**
Neutrophil adherence to isolated adult cardiac myocytes.
Induction by cardiac lymph collected during ischemia and reperfusion
J. Clin. Invest. 89: 602-609.

Ytehus, K.; Myklebust, R. and Mjos, O. (1986):

Influence of oxygen free radicals generated by xanthine oxidase in the isolated perfused rat heart.
Cardiovasc. Res. 20: 597-603.

Zweeier, J.L.; Flaherty, J.T. and Weisfeldt, M.L. (1987):

Direct measurement of free radical generation following reperfusion of ischemic myocardium.
Proc. Natl. Acad. Sci. U.S.A. 84: 1404-1407.