
REFERENCES

[A]

- Abi-Samara F.; Fouad F.M. and Tarazi R.C. (1983):*
Determinants of LVH and function in hypertensive patients.
Am. J. Med. 75 "Suppl. 3A": 26-33.
- Adair R.F.; Hellerstein H.K. and White L.W. (1972):*
Digoxin-induced exercise ECG changes in young men.
Circulation, 45: 1.
- Agabiti-Rosei E. and Muiesan M.L. (1993):*
Hypertension and diastolic function. Drugs 46 Suppl. 2: 61-7.
- Agabiti-Rosei E.; Mulesan M.L. and Muiesan G. (1989):*
Regression of structural alterations in hypertension. Am. J. Hypertens., 2: 705.
- Anderson M.T.; Lee G.B. and Champion B.C. (1972):*
Cardiac dysrhythmia associated with exercise stress testing.
Am. J. Cardiol., 30: 763.
- Aronow W.S. and Harris C.N. (1975):*
Treadmill exercise test in aortic stenosis and mitral stenosis.
Chest, 68: 507.
- Asmussen E.; Perry G. and Borow K. (1981):*
Similarities and dissimilarities between static and dynamic exercise.
Circ. Res., 48: 13.

[B]

- Baron D.W.; Ilseley C.; Sheiban I.; Poole-Wilson P.A. and Pickards A.F. (1980):*
R-wave amplitude during exercise, relation to left ventricular function and coronary artery disease. Br. Heart, J., 44: 512-7.

Beller G.A. (1991):

Pharmacologic stress imaging. JAMA, 265:633-638.

Blake J.; Devereux R.B.; Herrold E.M.; Jason M.; Fisher J. and Laragh J. (1988):

Relation of concentric LVH and extra cardiac target organ damage to supranormal left ventricular performance in established essential hypertension. Am. J. Cardiol., 62: 246-252.

Bloom W. and Fawcett D.W. (1994):

"A textbook of histology" 2nd edition: New York: Chapman and Hall. P., 137-142.

Bonoris P.E.; Greenberg P.S. and Christison G.W. (1978):

Ability of R-wave amplitude changes to reduce false negative and false positive responses by ST depression in treadmill stress testing. Am. J. Cardiol., 41: 378.

Brilla C.G.; Janicki J.S. and Weber K.T. (1991):

Cardioreparative effects of lisinopril in rats with genetic hypertension and LVH.
Circulation, 5: 1771-1779.

Bruce R.A. and McDonough J.R. (1969):

Stress testing increasing for cardiovascular disease. Bull. NY Acad. Med., 45: 1288.

Bruce R.A.; Kusumi F. and Hosmer D. (1973):

Maximal oxygen intake and nomographic assessment of functional aerobic impairment in cardiovascular disease. Am. Heart J., 85: 546-62.

Bruce R.A.; Fisher L.D.; Pettinger M. et al. (1988):

ST segment elevation with exercise: A marker for poor ventricular function and poor prognosis. Coronary Artery Surgery Study (CASS). Circulation, 77: 897.

Buda A.J.; Li Y.; Brant D. (1991):

Changes in left ventricular diastolic filling during the development of LVH in a unique canine model. Am. Heart J., 121: 1759-1767.

Busby M.J.; Shefrin E.A. and Fleg J.L. (1989):

Prevalence and long-term significance of exercise-induced frequent or repetitive ventricular ectopic beats in apparently healthy volunteers. *J. Am. Coll. Cardiol.*, 14: 1659.

Buttrick P.; Malhotra A. and Factor S. (1991):

Effect of aging and hypertension on myosin biochemistry and gene expression in the heart. *Circulation*, 88: 645.

[C]

Califf R.M.; Ohman E.M.; Frid D.J.; Fortin D.F.; Mark O.B. and Hlatky M.A. (1990):

"Restenosis: the clinical issues. In textbook of interventional cardiology" Philadelphia, Saunders; 363-394.

Casale P.N.; Devereux R.B. and Milner M. (1986):

Value of echocardiographic measurement of LV mass in predicting cardiovascular morbid events in hypertensive men. *Ann. Intern. Med.*, 105: 173-178.

Chevalier B.; El-Amrani F.C.; Heymes C. and Swynghedauw B. (1994):

Molecular basis of regression of cardiac hypertrophy. *Am. J. Cardiol.*, 73: 100-170.

Chiariello M. and Betocchi S. (1991):

Ventricular hypertrophy as an expression of hypertensive damage. *Cardiologia Dec.*, 36 (12 Suppl. 1): 79-84.

Chung E.K. (1983):

Principles of cardiac arrhythmias "third edition". Baltimore, Williams and Wilkins.

Cole J.P. and Ellestad M.H. (1978):

Significance of chest pain during treadmill testing: correlation with coronary events. *Am. J. Cardiol.*, 41: 227-232.

Crawford M.H.; Petru M.A. and Rabinowitz C. (1985):

Effect of isotonic exercise training on left ventricular volume during upright exercise. 72: 1237-1243.

[D]

Dalen, H.; Saetevsdal T. and Odegarden S. (1987):

Some ultrastructural features of the myocardial cells in the hypertrophied human papillary muscle. Virchows Arch. A. 410: 281.

Darnell J.; Lodish, H.; Baltimore D. (1990):

Actin, myosin, and intermediate filaments: cell movements and cell shape. Molecular cell Biology 2nd ed., New York, W.H. Freeman.

De Divitiis, O.; Celentano. A.; De Simone G. et al. (1993):

Management of the patient with LVH. Eur. Heart J., 14 "Suppl. D": 22.

Derman W.E.; Sims, R. and Noakes. T.D. (1992):

The effects of anti-hypertensive medications on the physiological response to maximal exercise testing. J. Cardiovasc. Pharmacol., 19: S122.

Devereux R.B. (1987):

Cardiac involvement in essential hypertension. Prevalence, pathophysiologic and prognostic implications. Medical Clinics of North America, 71-5: 819.

Devereux R.B. (1990):

Does increased blood pressure cause LVH or vice versa? Ann. Intern. Med., 112: 157-9.

Devereux R.B. and Reichek N. (1980):

Left ventricular hypertrophy. Cardiovascular Review Rep., 1: 55-68.

Devereux R.B.; Alonso D.R.; Lutas E.M.; Gottlieb G.J.; Campo E.; Sachs, I. And Reichek N. (1986):

Echocardiographic assessment of LVH: Comparison to necropsy findings. Am. J. Cardiol., 57: 450-458.

Devereux R.B.; Lutas E.M. and Casale P.N. (1984):

Standardization of M-mode echocardiographic left ventricular anatomic measurements. Am. J. Cardiol., 4: 1222.

Devereux R.B.; Pickering T.G.; Harshfield G.A. et al. (1983):

LVH in patients with hypertension: Importance of blood pressure response to regularly recurring stress. *Circulation*, 68: 470-476.

Dohlman H.G.; Thorner J.; Caron M.G. and Lefkowitz R.J. (1991):

Model system for the study of seven-transmembrane-segment receptors. *Am. Rev. Biochem.*, 60: 653-688.

Donhoue T.J.; Dworkin L.D.; Longo M.N. et al. (1994):

Induction of myocardial insulin-like growth factor I gene expression in left ventricular hypertrophy. *Circulation*, 89:799.

Douglas P.S.; O'Toole M.L.; Hiller W.D.; Reichel N. (1986):

Left ventricular structure and function by echocardiography in ultraendurance athletes. *Am. J. Cardiol.*, 58: 805-809.

[E]

Ehsani A.A.; Hagberg J.M. and Hickson R.C. (1978):

Rapid changes in left ventricular dimensions and mass in response to physical conditioning and deconditioning. *Am. J. Cardiol.*, 42: 521.

Ellestad M.H. and Halleday W.K. (1977):

Stress testing for the prognosis and management of ischemic heart disease. *Angiology*, 28: 149.

[F]

Factor S.M.; Flomenbaum M.; Zhao M.J.; Eng C. and Robinson T.F. (1988):

The effects of acutely increased ventricular cavity pressure on interintrinsic myocardial connective tissue. *Am. J. Cardiol.*, 12: 1582-1589.

Fagard R.; Bielen E. and Amery A. (1992):

Automated versus observer blood pressure as determinants of left ventricular structure. *Eur. Heart J.*, 13: 1373-1379.

Fagard R.; Bielen E.; Lijnen P. and Amery A. (1993):

Cardiac variables and blood pressure as determinants of left ventricular inflow velocities.
J. Hum. Hypertens., 7: 7-12.

Fagard R.; Brooke C. and Ogawa T. (1989):

Left ventricular dynamics during exercise in elite marathon runners. J. Am. Coll. Cardiol., 14: 112.

Fagard R.; Staessen J.; Thijs L. and Amery A. (1995):

Relation of left ventricular mass and filling to exercise and rest blood pressure. Am. J. Cardiol., 75: 53-57.

Faris J.V.; McHenry P.L. and Morris S.N. (1978):

Concepts and applications of treadmill exercise testing and the exercise ECG. Am. Heart J., 95: 102.

Ferrans V.J. (1989):

Pathologic anatomy of the dilated cardiomyopathies.
Am. J. Cardiol., 64: 9C.

Fletcher G.F. ed. (1988):

Exercise in the practice of medicine. 2nd ed. Mt. Kisco, NY: Futura.

Fletcher G.F.; Balady G.; Froelicher V.F. et al. (1995):

Exercise standards. A statement for health care professionals from the American Heart Association. Circulation, 91: 580.

Fouad F.M.; Tarazi R.C.; Gallagher J.H.; Macintyre W.J. and Cook S.A. (1980):

Abnormal left ventricular relaxation in hypertensive patients. Clin. Sci., 41 "Suppl. 6": 411S.

Fox M.K.; Jonathan A. and Selwyn A. (1983):

Significance of exercise induced ST segment elevation in patients with previous myocardial infarction. Br. Heart J., 49: 15-19.

Francis G.S. and McDonald K.M. (1992):

LVH: An initial response to myocardial injury. Am. J. Cardiol, 69: 3G.

Franklin B.A. (1985):

Exercise testing, training and arm ergometry. Sports Medicine, 2:100-119.

Frimm C.C.; DeMoraes A.V.; Medeiros C.; Silva H.B.; Bellotti G.; Krieger E.M. and Pileggi F. (1994):

Role of blood pressure in rest and exercise conditions and importance of clinical factors in LVH in arterial hypertension. *Arq. Bras. Cardiol.* Jul., 63 (1): 21-26.

Frohlich E.D. (1987):

Cardiac hypertrophy in hypertension. *N. Engl. J. Med.*, 317: 831-833.

Frohlich E.D. and Tarazi R.C. (1979):

Is arterial pressure the sole factor responsible for hypertensive cardiac hypertrophy? *Am. J. Cardiol.* Volume, 44, October, 22: 959-963.

Furchgott R.F. and Zawadzki J.V. (1993):

Hypertension and LVH are associated with impaired endothelium mediated relaxation in human coronary resistance vessels. *Circulation*, 87: 86.

[G]

Garber C.E.; Carleton R.A.; Camaione D.N. and Heller G.V. (1991):

The threshold for myocardial ischaemia varies in patients with coronary artery disease depending on the exercise protocol. *J. Am. Coll. Cardiol.*, 17: 1256-1262.

Geiser E.A. and Bove K.E. (1974):

Calculation of left ventricular mass and relative wall thickness. *Arch. Pathol.*, 97: 13.

Geltman E.M.; Ehsani A.A.; Campbell M.K.; Roberts R. and Sobel B.E. (1979):

The influence of location and extent of myocardial infarction on long-term ventricular dysrhythmia and mortality. *Circulation*, 60: 805-813.

Genovesi-Ebert A.; Marabotti C.; Palombo C.; Giaconi S.; Rossi G. and Ghione, S. (1994):

Echo-Doppler diastolic function and exercise tolerance. *Int. J. Cardiol.* Jan., 43 (1): 67-73.

Gordan S.; Admant P. and Hood M. (1982):

Guidelines for the functional testing. *Am. J. Cardiol.*, 43: 181.

Gosse P.; Campello G.; Aouizcrate E.; Roudant R.; Broustet J.P. and Dallochio M. (1986):

LVH in hypertension: correlation with rest, exercise and ambulatory systolic blood pressure. *J. Hypertens.* 4 (Suppl. 5): S297-S299.

Gottdiener J.S.; Brown J.; Zoltick J. and Fletcher R.D. (1990):

LVH in men with normal blood pressure: relation to exaggerated blood pressure response to exercise. *Ann. Intern. Med.*, 112: 161-166.

Granger C.B.; Karimeddini M.K.; Smith V.E.; Shapiro H.R.; Katz A.M. and Riba A.L. (1985):

Rapid ventricular filling in LVH: I. Physiologic hypertrophy. *J. Am. Coll. Cardiol.*, 5: 863.

Grant S.C.D.; Rohman A.N. and Brooks N.H. (1992):

Regression of LVH in a transplanted heart. *Br. Heart J.*, 68: 55-57.

Grassi G.; Giannathasio C.; Cleroux J.; Cuspidi G.; Sampieri L.; Bolla G.B. and Mancia G. (1988):

Cardiopulmonary reflex before and after regression of LVH in essential hypertension. *Hypertension*, 12: 227.

Grossman W. (1980):

Cardiac hypertrophy: useful adaptation or pathological process? *Am. J. Med.*, 69:576.

Grossman w.; Jons D. and McLaurin L.P. (1975):

Wall stress and patterns of hypertrophy in the human left ventricle. *J. Clin. Invest.*, 56: 64.

[H]

Hagberg M.; Chen Z. and Sagar L. (1987):

Values of functional exercise testing in cardiopulmonary diseased patients. *Circulation*, 63: 115.

Hanrath P.; Mathey D.G.; Seigert R. and Bleifeld W. (1980):

Left ventricular relaxation and filling pattern in different forms of LVH: An echocardiographic study. *Am. J. Cardiol.*, 45: 15.

Harris C.N.; Aronow W.S.; Parker D.P. and Kaplan M.A. (1973):

Treadmill stress test in LVH. *Chest*, 63: 353.

Harrison M.R.; Grigsby C.G.; Souther S.K. et al. (1991):

Midventricular obstruction associated with chronic systemic hypertension and severe LVH.

Am. J. Cardiol., 68: 761-765.

Haskell W.L. and Debusk R. (1979):

Cardiovascular responses to repeated treadmill exercise testing soon after myocardial infarction. *Circulation*, 60: 1247.

Haughton J.L.; Carr A.A.; Prisant L.M.; Rogers W.B.; Von Dohlen T.W.; Flowers N.C. and Frank M.J. (1992):

Morphologic, haemodynamic and coronary perfusion characteristics in severe LVH secondary to systemic hypertension and evidence for nonatherosclerotic myocardial ischaemia. *Am. J. Cardiol.*, 69: 219.

Hollander W. (1976):

Role of hypertension in atherosclerosis and cardiovascular disease. *Am. J. Cardiol.*, 38: 786-800.

[I]

Imai K.; Sato H.; Hori M. et al. (1994):

Vagally mediated heart rate recovery after exercise is accelerated in athletes but blunted in patients with chronic heart failure. *J. Am. Coll. Cardiol.*, 24: 1529.

Inouye I.; Massie B.; Loge D.; Topic N.; Silverstein D.; Simpson R. and Tubau J. (1984):

Abnormal left ventricular filling: Early finding in mild to moderate essential hypertension. *Am. J. Cardiol.*, 53: 120.

Irving J.B. and Bruce R.A. (1977):

Variations in and significance of systolic pressure during maximal exercise testing. Relation to severity of coronary artery disease and cardiac mortality. *Am. J. Cardiol.*, 39: 841.

Ito H.; Hirata Y.; Hiroe M.; Tsujino M.; Adachi S.; Takamoto T.; Nilata M.; Taniguchi K. and Marumo F. (1991):

Endothelin -1 induces hypertrophy with enhanced expression of muscle specific genes in cultured neonatal rat cardiomyocytes. *Circ. Res.*, 96: 209.

Ito H.; Hiroe M.; Hirata Y.; Fujisaki H.; Adachi S.; Akimoto H.; Ohta Y. and Marumo F. (1994):

Endothelin ETA receptor antagonist blocks cardiac hypertrophy provoked by haemodynamic overload. *Circulation*, 89: 2198.

Iwai N.; Ohmichi N.; Nakamura Y. and Kinoshita M. (1994):

DD genotype of the angiotensin-converting enzyme gene is a risk factor for LVH. *Circulation*, 90: 2622-2628.

[J]

Jouko K.; Matti M.; Matti V. and Urho K. (1997):

Left ventricular mass, geometry, and filling in endurance athletes: Association with exercise blood pressure. *J. Appl. Physiol.*, 82 "2": 531-537.

[K]

Kadish A.H.; Weisman H.F.; Veltri E.P. et al. (1990):

Paradoxical effects of exercise on the QT interval in patients with polymorphic ventricular tachycardia receiving type Ia antiarrhythmic agents. *Circulation*, 81: 14.

Kannel W.B. (1974):

Role of blood pressure in cardiovascular morbidity and mortality. *Prog. Cardiovasc. Dis.*, 17: 5.

Kannel W.B. and Sorlie P.D. (1979):

Some health benefits of physical activity: The Framingham study. *Arch. Intern. Med.*, 139: 857.

Kannel W.B.; Doyle J.T.; McNamara P.M.; Quikenton P. and Gordon T. (1975):

Precursors of sudden coronary death: factors related to the incidence of sudden death. *Circulation*, 51: 606.

Kaplan N.M. (1990):

Primary hypertension: pathogenesis and natural history, in clinical hypertension. 5th ed. Williams and Wilkins, P. 54-135.

Karjalainen J.; Mantysaari M.; Viitasalo M. and Kujala U. (1997):

Left ventricular mass, geometry, and filling in endurance athletes: association with exercise blood pressure. *J. Appl. Physiol.*, Feb., 82 "2": 531-7.

Kattus A.A. (1977):

Exercise electrocardiography: Recognition of the ischaemic response, false positive and false negative pattern, in exercise in cardiovascular health and disease. Edited by E.A. Amisterdam, J.H. Wilmore and A.N. Demoria, P. 161-178.

Kaul V.; Mohan J.C. and Bhatai M.L. (1984):

Effects of labetalol on left ventricular mass and function in hypertension. An assessment by serial echocardiography. *Int. J. Cardiol.*, 5: 461.

Kelly S.; Hauch A. and Pombo J. (1986):

Indication of exercise test in ischeamic heart disease. *Br. Heart J.*, 72: 151.

Kishida H.; Cole J.S. and Surawicz B. (1982):

Negative U-wave: A highly specific but poorly understood sign of heart disease.
Am. J. Cardiol., 49: 2030.

Klug D.; Robert V. and Swynghedauw B. (1993):

Role of mechanical and hormonal factors in cardiac remodeling and the biological limits of myocardial adaptation. *Am. J. Cardiol.*, 71: 1-9.

Knutsen K.M.; Michelsen S.; Otterstad J.E.; Froland G. and Stugaard M. (1991):

Correlations between left ventricular mass and systolic blood pressure at rest and during maximal exercise in moderately hypertensive men. *J. Hum. Hypertens.*, Jun., 5 (3): 149-54.

Kojima M.; Shiojima I.; Yamazaki T.; Komuro I. et al. (1994):

Angiotensin II receptor antagonist TCV-116 induces regression of hypertensive LVH in vivo and inhibits the intracellular signaling pathways of stretch mediated cardiomyocyte hypertrophy in vitro. *Circulation*, 89: 2204.

Kotler T.S. and Diamond G.A. (1990):

Exercise thallium-201 scintigraphy in the diagnosis and prognosis of coronary artery disease. *Ann. Intern. Med.*, 113: 684-702.

- Krayenbuehl H.P.; Hess O.M.; Schmieder J. and Turina M. (1983):**
Physiologic or pathologic hypertrophy. Eur. Heart. J., 4 "Suppl. A": 29.
- Kuhm H. (1982):**
The definition of ventricular hypertrophy. Eur. Heart. J. 1 "Suppl. A": 5-8.
- Kupari M.; Koshinen P. and Virolainen J. (1994):**
Correlates of left ventricular mass in a population sample aged 36 to 37 years: Focus on life style and salt intake. Circulation, 89: 1041-1050.
- Kurita A.; Chaitman B.R. and Bourassa M.G. (1977):**
Significance of exercise-induced junctional ST-depression in evaluation of coronary artery disease. Am. J. Cardiol., 40: 492.
- Kuwajima I.; Suzuki Y.; Shimosawa T.; Kanemaru A.; Hoshino S. and Kuramoto K. (1992):**
Diminished nocturnal decline in blood pressure in elderly hypertensive patients with LVH. Am. Heart J., 67: 1307.

[L]

- Lauer M.S.; Levy D.; Anderson K.M. and Plehn J.F. (1992):**
Is there a relationship between exercise systolic blood pressure response and left ventricular mass? The Framingham Heart study. Ann. Intern. Med. Feb., 1; 116 "3": 203-10.
- Lele S.S.; Scalia G.; Thomson H. et al. (1994):**
Mechanism of exercise hypotension in patients with ischemic heart disease: Role of neurocardiologically mediated vasodilation. Circulation, 90: 2701.
- Levy D.; Garrison R.J.; Savage D.D.; Kannel W.B. and Castelli W.P. (1990):**
Prognostic implications of echocardiographically determined LV mass in the Framingham heart study. N. Engl. J. Med., 322: 1561-1566.

- Levy D.; Anderson K.M.; Savage D.D.; Kannel W.B. and Castelli W.P. (1988):**
Echocardiographically detected LVH: prevalence and risk factors: The Framingham heart study. *Ann. Intern. Med.*, 108: 7-13.
- Lewis J.; Spirito P.; Pelliccia A. and Maron B.J. (1992):**
Usefulness of Doppler echocardiographic assessment of diastolic filling in distinguishing "athlete's heart" from hypertrophic cardiomyopathy. *Br. Heart J.*, 68: 296-300.
- Liebson P.R.; Grandits G.A.; Dianzumbe S. and Prineas R.J. (1995):**
For the treatment of hypertension study research group. *Circulation*, 91: 698-706.
- Limacher M.C.; Quinones M.A.; Poliner L.R.; Nelson J.G.; Winters W.L.J. and Waggoner A.D. (1983):**
Detection of coronary artery disease with exercise two-dimensional echocardiography: Description of a clinically applicable method and comparison with radionuclide ventriculography. *Circulation*, 67: 1211-1218.
- Linhart J.W. (1978):**
Stress testing in angio. Resident and staff physician, P. 79-89.
- Linzback A.J. (1960):**
Heart failure from the point of view of quantitative anatomy. *Am. J. Cardiol.*, 5: 370.
- [M]**
- Mann S.; Altman D.G.; Raftery E.B. and Bannister R. (1983):**
Circadian variation of blood pressure in autonomic failure. *Circulation*, 68: 477.
- Marcus M.L.; Harrison D.G.; Chilian W.M.; Koyanagi S.; Inou T.; Tamonek R.J.; Martens J.B.; Esthm. C.L. and Hiratzka L.F. (1987):**
Alterations in the coronary circulation in hypertrophied ventricles. *Circulation* 75 "Suppl. 1": 1-19.
- Maron B.; Kelly A. and Aubert A. (1986):**
Structural features of the athlete's heart as defined by echocardiography. *J. Am. Coll. Cardiol.*, 7: 190-203.

Martin W.; Coyle E. and Maron B. (1986):

Effects of physical deconditioning after intense endurance training in left ventricular dimensions and stroke volume. J. Am. Coll. Cardiol., 7: 982.

McHenry P.L. (1977):

The actual prevalence of false positive ST-segment responses to exercise in clinically normal subjects remains undefined. Circulation, 55: 683.

McHenry P.L. and Morris S.N. (1976):

Exercise electrocardiography, current state of the art, in advances in electrocardiography, edited by J.W. Hurst and R.C. Schlant, New York, Grune and Stratton.

McHenry P.L.; Richman H.M. and Jordan J.W. (1981):

Evaluation of abnormal exercise electrocardiogram in apparently healthy subjects: Labile repolarization (ST-T) abnormality as a cause of false positive responses. Am. J. Cardiol., 47: 1152.

Meerson F.Z. (1969):

The myocardium in hypertension, hypertrophy and heart failure. Monograph 26; New York, Am. Heart Assoc. 6-8.

Meerson F.Z. and Breger A.M. (1977):

The common mechanism of heart to adaptation and deadaptation: Hypertrophy and atrophy of the heart muscle. Basic. Res. Cardiol., 72: 228. Quoted by: Taquini CM: Cardiac function in experimental hypertension. Am. Heart J., 166: 607, 1988.

Messerli F.H.; Aristizabal D. and Soria F. (1993):

Reduction of LVH: How beneficial? Am. Heart J., 125: 1520-1540.

Messerli F.H.; Ventura H.O.; Elizardi D.J.; Dunn F.G. and Frohlich E.D. (1984):

Hypertension and sudden death: Increased ventricular ectopic activity in LVH. Am. J. Med., 77: 18.

Montgomery H.; Clarkson P.; Dollery C.; Losi M.A.; Jubb M.; McEwan J.; Humphries S. and McKenna W. (1996):

D polymorphism of the ACE gene is strongly associated with the development of physiological LVH. J. Am. Coll. Cardiol., 27: 244A.

Morgan H.E. and Baker K.M. (1991):

Cardiac hypertrophy: Mechanical, neural and endocrinal dependence. *Circulation*, 83: 13-25.

Morgan H.E.; Chua B.H.L.; Russo L. (1991):

Protein synthesis and degradation. In: Fozzard HA, Haber E, Jennings RB, Katz AM, Morgan HE (eds): *The heart and Cardiovascular system*. New York, Raven, 1991: 1505-1524.

Moriarity T.; Byrd B. and Kenl J. (1980):

The law of Laplace: Its limitations as relation for diastolic pressure, volume, or wall stress of the LV. *Circ. Res.*, 46: 321.

Mukherjee D. and Sen S. (1990):

Collagen phenotypes, in spontaneously hypertensive rats. *Circ. Res.*, 67: 1474-1480.

Muntz K.; Gonyea W. and Hagler D. (1981):

Cardiac hypertrophy in response to an isometric training program in the cat. *Circ. Res.*, 49: 1092.

[P]

Page E. and Oparil S. (1978):

Effect of peripheral sympathectomy on left ventricular ultrastructure in young spontaneously hypertensive rats. *J. Mol. Cell Cardiol.*, 10: 301.

Pearson A.C.; Pasierski T. and Labovitz A.J. (1991):

Left ventricular hypertrophy: Diagnosis, prognosis and management. *Am. Heart J.*, 121: 148-157.

Pelliccia A.; Spataro A.; Caselli G. and Moron B.J. (1993):

Absence of left ventricular wall thickening in athletes engaged in intense power training. *Am. J. Cardiol.*, 72: 1048-1054.

Pelliccia A.; Maron B. and Robert A. (1991):

The upper limit of physiologic cardiac hypertrophy in highly trained elite athletes. *N. Engl. J. Med.*, 324: 295.

Percy R.; Conetta D. and Sander S. (1990):

Echocardiographic assessment of the left ventricle of endurance athletes just before and after exercise. *Am. J. Cardiol.*, 65: 1140.

Petru M.; Bryd S. and Yamada Y. (1990):

Exercise test as a reflection of work capacity. *Circulation*, 37: 1315.

Phillips R.A.; Goldman M.E. and Ardeljan M. (1989):

Determinants of abnormal left ventricular filling in early hypertension. *J. Am. Coll. Cardiol.*, 14: 979-985.

Pipkin T.B. (1984):

Simple significance tests, In: *Medical statistics made easy*, 1st edition. Churchill Livingstone, P. 46-61.

Rapaport E. (1982):

Pathophysiological basis of ventricular hypertrophy. *Eur. Heart J.*, 3 "Suppl. A": 5-8.

Ren J.F.; Hakki A.H.; Kotler M.N.; Iskandrian A.S. (1985):

Exercise systolic blood pressure: a powerful determinant of increased left ventricular mass in patients with hypertension. *J. Am. Coll. Cardiol.*, 5: 1224-1231.

Roeske W.R.; O'Rourke R.A.; Klein A.; Leopold G.; Karliner J.S. (1976):

Noninvasive evaluation of ventricular hypertrophy in professional athletes. *Circulation* 53: 286.

[S]

Sahn D.J.; Demania A.; Kisslo J. and Weyman A. (1978):

Recommendations regarding quantitation in M-mode echocardiography: results of a survey of echocardiographic measurement. *Circulation*, 58: 1072-1083.

Savage D.D.; Levy D.; Dannenberg A.L.; Garrison R.J.; Castelli W.P. (1990):

Association of echocardiographic left ventricular mass with body size, blood pressure, and physical activity "Framingham study". *Am. J. Cardiol.*, 65: 371-6.

Schlant R.; Blomquist G.; Brandenburg R. et al. (1986):

Guidelines for exercise testing. A report of the joint American college of cardiology/ American Heart Association Task force on assessment of cardiovascular procedures. *Circulation*, 74: 653(A)-667 (A).

- Schmieder R.E.; Messerli F.H.; Garvaglia G.E. (1988):**
Dietary salt intake. A determinant of cardiac involvement in essential hypertension. *Circulation*, 78: 951.
- Schmieder R.E.; Grube E.; Ruddel H.; Schachinger H. and Schulte W. (1990):**
Relation of hemodynamic reaction during stress to LVH in essential hypertension. *Am. J. Hypertens. Apr.*, 3(4): 281-7.
- Shah P.M. (1986):**
Controversies in hypertrophic cardiomyopathy.
Curr. Prob. Cardiol., 11: 10.
- Shapiro L.M.; Kleinebenne, A. and McKenna, W.J. (1985):**
The distribution of LVH in hypertrophic cardiomyopathy: Comparison to athletes and hypertensives. *Eur. Heart J.*, 6: 967.
- Shapiro L.M. and McKenna W.J. (1983):**
Distribution of LVH in hypertrophic cardiomyopathy: A two-dimensional echocardiographic study. *J. Am. Coll. Cardiol.*, 2: 437.
- Simone G.; Lorenzo L.; Moccia D.; Costantino G. (1987):**
Haemodynamic hypertrophied left ventricular pattern in systemic hypertension. *Am. J. Cardiol.*, 60: 1317-1327.
- Simpson P. (1983):**
Norepinephrine-stimulated hypertrophy of cultured rat myocardial cells in an alpha-1 adrenergic response. *J. Clin. Invest.*, 72: 732.
- Smith D.H.; Neutel J.M.; Graettinger W.F.; Myers J.; Froelicher W.F. and Weber M.A. (1992):**
Impact of LVH on blood pressure response to exercise. *Am. J. Cardiol.* 5, 69 , 225-8.
- Smith V.E.; Schulman P.; Karimeddini M.K. et al. (1985):**
Rapid ventricular filling in LVH. *J. Am. Coll. Cardiol.*, 5: 869.

Strnaer B. (1979):

Ventricular function and coronary haemodynamics in hypertensive heart disease. *Am. J. Cardiol.*, 44: 999.

Straure B.E. (1990):

Significance of coronary circulation in hypertensive heart disease for development and prevention of heart failure. *Am. J. Cardiol.*, 65: 34.

Sugishita Y.; Roseki S. and Saltin B. (1983):

Myocardial mechanics of athletic heart in comparison with diseased heart. *Am. Heart J.*, 105: 273.

[T]

Tan L.B.; Jalil J.E.; Tanicki J.S.; Weber K.T. and Clark W.A. (1989):

Cardiotoxic effects of angiotensin II. *J. Am. Cardiol.*, 13: 2-5.

Taquini C.M. (1988):

Cardiac function in experimental hypertension. *Am. Heart J.*, 116: 607.

Thomas J.D. and Weyman A.E. (1991):

Echocardiographic Doppler evaluation of left ventricular diastolic function. *Circulation*, 84: 977-990.

Topol E.J.; Traill T.A. and Fortuin N.J. (1985):

Hypertensive hypertrophic cardiomyopathy in the elderly. *N. Engl. J. Med.*, 312: 277.

Toyoshima H.; Park Y.D.; Ishikawa Y.; Nagata S.; Hirata Y.; Sakakibara H. and Nakayama R. (1982):

Effect of ventricular hypertrophy on conduction velocity of activation front in the ventricular myocardium. *Am. J. Cardiol.*, 49: 1638.

Treasure C.B.; Klein J.L.; Vita J.A.; Manoukian S.V.; Renwick G.H.; Sellwyn A.P.; Ganz P. and Alexander W. (1993):

Hypertension and LVH are associated with impaired endothelium mediated relaxation in human coronary resistance vessels. *Circulation*, 87: 86.