

References

Agarwal JB, and Helfant RH (1983):

Functional importance of coronary collateral circulation. *Int J Cardiol*, 1983; 4: 94.

Akhras F, Upward J, and Jackson G (1985):

Reciprocal change in ST segment in acute myocardial infarction: correlation with findings on exercise electrocardiography and coronary angiography. *BMJ* 1985; 290:1931-4.

Anderson H, Willerson J (1993):

Thrombolysis in acute myocardial infarction. *N Engl J Med* 1993; 329:703-709.

ASSENT-2 investigators (1999):

Assessment of the safety and efficacy of a new thrombolytic (ASSENT-2) investigators. Single-bolus tenecteplase compared with frontloaded alteplase in acute myocardial infarction: the ASSENT-2 double-blinded randomized trial. *Lancet* 1999; 354: 716-722.

Baim DS, and Grossman W (1986):

Coronary arteriography in: cardiac catheterization and angiography edited by Grossman, W., 3rd ed., Lea and Febiger, Philadelphia, P.179-199, 1986.

Balady GJ, Weiner DA, and McCabe CH (1985):

Value of arm exercise testing in detecting coronary artery disease. *Am J Cardiol*. 1985; 55:37-39.

References

Balter S, and Members of the Laboratory and Performance Standards Committee of the Society for Cardiac Angiography and Interventions (1993):

Guidelines for personnel radiation monitoring in the cardiac catheterization laboratory. Cathet. Cardiovasc. Diagn. 30:277, 1993.

Braunwald E (1992):

Coronary blood flow and myocardial ischemia in: Braunwald E (Ed.) Heart disease. 4th ed., text book of cardiovascular medicine. Saunders WB, company: 1161, 1992.

Braunwald E, Zipes DP, and Libby P (2001):

Assessment of normal and abnormal cardiac function in: Braunwald E (Ed.) Heart disease. 6th ed., textbook of cardiovascular medicine. Saunders WB, company: 498, 2001.

Bourdillon PVD, Broderick TM, and Sawada SG (1989):

Regional wall motion index for infarct and non infarct regions after reperfusion in acute myocardial infarction: comparison with global wall motion index. J Am Soc Echocardiogr 2:398-407, 1989.

Brown KA (1996):

Prognostic value of myocardial perfusion imaging: state of the art and new developments. J Nucl Cardiol. 1996; 3:516-537.

References

Buja CN, Shah PK, Lew AS, and Hulse S (1987):

Electrocardiographic differentiation of occlusion of the left circumflex versus the right coronary artery as a cause of inferior acute myocardial infarction. *Am J Cardiol* 1987; 60:456–459.

Buja LM, and Willersone JT (1981):

Clinicopathologic correlates of acute ischemic heart disease syndromes. *Am J Cardiol.* 47:343,1981.

Chaitman BR (1997):

Exercise stress testing. In Branwald E.: *Testbook of cardiovascular medicine*. 5th ed. Philadelphia, W.B. saunders Company, 1997.

Chaitman BR, Stone PH, and Knatterud G L (1995):

Asymptomatic Cardiac Ischemia Pilot (ACIP) study: Impact of anti-ischemia therapy on 12-week rest ECG and exercise test outcomes. *J. Am. Coll. Cardiol.* 26:585, 1995.

Christian TF, Schwartz RS, and Gibbons RJ (1992):

Determinants of infarct size in reperfusion therapy for acute myocardial infarction. *Circ* 86:81,1992.

Cohen-Solal A, Zannad F, and Kayanakis JG (1991):

Multicentre study of the determination of peak oxygen uptake and ventilatory threshold during bicycle exercise in chronic heart failure: comparison of graphical methods, interobserver variability and influence of the exercise protocol: the VO₂ French Study Group. *Eur Heart J.* 1991; 12:1055–1063.

References

Cohn JN (1987):

Quantitative exercise testing for the cardiac patient: the value of monitoring gas exchange: introduction. *Circulation* 1987; 76(suppl VI): VI-1-VI-2.

Coma-Canella I (1991):

Significance of ST segment changes induced by dobutamine stress test after acute myocardial infarction. Which are reciprocal? *Eur Heart J* 1991; 12:909-16.

Davies MJ, Treasure T, and Richardson PD (1996):

The pathogenesis of spontaneous arterial dissection. *Heart* 1996; 75:434-435.

Dewood MA, Spores J, Notske R, and Mouser LT (1980):

Prevalence of total coronary occlusion during the early hours of transmural myocardial infarction. *N Engl J Med* 1980; 303:897-902.

Dewood MA, Stifter W, Simpson CS, and Spores J (1986):

Coronary arteriographic findings soon after non-Q wave myocardial infarction. *N Engl J Med*, 1986; 315:417.

Dillahunp PH, and Miller AB (1998):

Early treadmill testing after myocardial infarction: angiographic and haemodynamic correlations. *Ann Intern Med* 1998 Jun; 94(6): 727-34.

Dubach P, Froelicher VF, and Klein J (1988):

Exercise-induced hypotension in a male population: criteria, causes, and prognosis. *Circulation* 1988;78: 1380-1387.

References

Dunn RF, Freedman B, and Bailey IK (1980):

Noninvasive prediction of multivessel disease after myocardial infarction. *Circulation* 1980; 62:726-34.

Ellestad MH (1986):

Stress testing: principles and practice. 3rd ed. Philadelphia, F.A. Davis Company, 1986.

Ellestad MH (1995):

Stress Testing: Principles and Practice. 4th ed. Philadelphia, F. A. Davis Co., 1995.

Evans GL, Sumlyan H, and Eich RH (1987):

Role of peripheral resistance in the control of cardiac output. *Am J Cardiol.* 1987; 200:216.

Falk E, Shah PK, and Fuster V (1995):

Coronary plaque disruption. *Circulation* 1995; 92:657-671.

Falk E, Shah P K, and Fuster V (1996):

Pathogenesis of plaque disruption. in Fuster V., Ross R., and Topol EJ.(Eds.): *Atherosclerosis and coronary artery disease*. Philadelphia, Lippincott-Raven, 1996, pp. 492 - 510.

Falk E (1983):

Plaque rupture with severe pre-existing stenosis precipitating coronary thrombosis. Characteristics of coronary atherosclerosis plaque underlying fatal occlusive thrombi. *Br Heart J* 1983; 50:127-134.

References

Falk, E., Shah, P.K., and Fuster, V (1995):

Coronary plaque disruption. *Circulation* 92:657, 1995.

Feil H, and Siegel M L (1928):

Electrocardiographic changes during attacks of angina pectoris. *Am. J. Med. Sci.* 175:255, 1928.

Ferguson J. meeting highlights (1997):

46th annual Scientific Sessions of the American College of cardiology: in TIME trial. *Circulation* 1997; 96:367-371.

Florence H.S, Braunwald E, and Paul Canner (1987):

Effects of intravenous thrombolytic therapy on left ventricular function: a report on tissue type plasminogen activator and streptokinase from the Thrombolysis In Myocardial Infarction (TIMI) trial, *Circulation* 1987, 75; 4:817.

Franklin BA (1985):

Exercise testing, training and arm ergometry. *Sports Med.* 1985; 2:100 – 119.

Freedman SB, Dunn RF, and Bernstein L (1985):

Influence of coronary collateral blood flow on the development of exertional ischemia and Q-wave infarction in patients with severe single-vessel disease. *Circulation* 1985; 71 : 681 .

References

Freifeld AG, Schuster E H, and Bulkley BH (1983):

Nontransmural versus transmural myocardial infarction . Am. J. Med. 75: 423, 1983.

Froelicher V F, Myers J, and Labovitz AJ (1993):

Exercise and the Heart. 3rd ed. St. Louis, Mosby-YearBook, 1993.

Fuller CM, Raizner AE, and Verani MS (1981):

Early post-myocardial infarction treadmill stress testing as accurate predictor of multivessel coronary artery disease and subsequent cardiac events. Ann Intern Med 1981; 94:734-9.

Fuster V, Badimon L, and Badimon JJ (1992):

The pathogenesis of coronary artery disease and the acute coronary syndromes. N Engl J Med 1992; 326:242-250, 310-318.

Gensini GG (1975):

Anatomy of the normal coronary arteries in Gensini, G.G. (Ed.) coronary arteriography. Futura publishing co., New York, PP.163-230.

Gersh BJ, and Anderson JL (1993):

Thrombolysis and myocardial salvage: results of clinical trials and the animal paradigm-paradoxical or predictable? Circulation 1993; 88:296-306.

Gheorghiade M, Nissen H, and Horder M (1996):

Independent prognostic value of serum creatinine kinase isoenzyme MB mass, cardiac Troponin T and Myosin light chain levels in suspected acute myocardial infarction. J Am Coll Cardiol 1996; 25:574-581.

References

Gianrossi R, Detrano R, and Mulvihill D (1989):

Exercise-induced ST depression in the diagnosis of coronary artery disease: a meta-analysis. *Circulation*. 1989;80:87-98.

Gibbons RJ, Balady GJ, and John W Beasley (1997):

ACC/AHA guidelines for exercise testing: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Exercise Testing). *J Am Coll Cardiol*. Vol.30 no1 July 1997: 260-315. .

Gibson CM (1994):

TIMI frame count: A new standardization of infarct-related artery flow grade, and its relationship to clinical outcomes in the TIMI-4 trial. *Circulation* 90 (Abs.): I-220, 1994.

Gibson RS, Watson DD, Graddock GB, and Crampton RS (1983):

Prediction of cardiac events after uncomplicated myocardial infarction: a prospective study comparing predischage exercise thallium-201 scintigraphy and coronary angiography. *Circulation* 1983; 68:321-336.

Goldhaber SZ, Haire WD, and Feldstein ML (1993):

Alteplase versus heparine in acute pulmonary embolism: randomized trial assessing right-ventricular function and pulmonary reperfusion. *Lancet* 1993; 341:507-511.

Gruber A, Pal A, and Kiss RG (1993):

Generation of activated protein C during thrombolysis. *Lancet* 1993; 342:1275-1276.

References

GUSTO investigators (1993):

An international randomized trial comparing for thrombolytic strategies for acute myocardial infarction. N Engl J Med 1993; 329:673-682.

GUSTO Angiographic Investigators (1993):

The effects of tissue plasminogen activator, streptokinase, or both on coronary artery patency, ventricular function, and survival after acute myocardial infarction. N. Engl. J. Med. 329:1615, 1993.

GUSTO-III investigators (1997):

The global use of strategies to open occluded coronary arteries: A comparison of reteplase with alteplase for acute myocardial infarction. N Engl J Med 1997; 337: 1118-1123.

Guyton AC (1995):

Textbook of Medical physiology. 9th Ed. Philadelphia, W.B. Saunders company, 1995.

Hsia J, Hamilton WP, and Kleiman N (1990):

A comparison between heparin and low-dose aspirin as adjunctive therapy with tissue plasminogen activator for acute myocardial infarction. N Engl J Med 1990; 323:1433-1437.

Ileana WH, Aronow WS, and Goodman P (1995):

Five-year follow-up of maximal treadmill stress test in asymptomatic men and women. Circulation 1995;62:522-527.

ISIS-3 study group (1992):

Third international study of infarct survival: collaborative group. ISIS-3: a randomized comparison of streptokinase vs tissue plasminogen activator vs anistreplase and of aspirin plus heparin vs aspirin alone among 41,299 cases of suspected acute myocardial infarction. Lancet 1992; 339: 753-770.

Jennings K, Reid D, and Julian DG (1983):

"Reciprocal" depression of the ST segment in acute myocardial infarction. BMJ 1983; 287:634-7.

Jennings R.B., Schaper J., and Hill M.L (1990):

Effect of reperfusion late in the phase of reversible ischemic injury: changes in cell volume, electrolytes, metabolites and ultrastructure. Circ Rec., 66:1133,1990.

Juneau M, Colles P, and Theroux P (1992):

Symptom-limited versus low level exercise testing before hospital discharge after myocardial infarction. J Am Coll Cardiol. 1992; 20:927-933.

Kenneth A, Brown MD, and Gary V (1999):

Early Dipyridamole ^{99m}Tc-Sestamibi SPECT imaging 2 to 4 days after acute myocardial infarction predicts in-Hospital and postdischarge cardiac events (comparison with submaximal exercise imaging). Circulation 1999; 100:2060-2066.

References

Kligfield P, Ameisen O, and Okin PM (1989):

Heart rate adjustment of ST segment depression for improved detection of coronary artery disease. *Circulation* 1989; 79:245–255.

Kloner RA, Ellis SG, and Braunwald E (1983):

Coronary reperfusion for treatment of acute myocardial infarction. Post ischemic ventricular dysfunction. *Cardiology*, 70: 233, 1983.

Kloner RA, Ellis SG, and Lange R (1983):

Studies of experimental coronary artery reperfusion: effects on infarct size, myocardial function, biochemistry, ultrastructure and microvascular damage. *Circulation* 68: 1, 1983.

Kloner RA, and Przylenk K (1993):

Understanding the jargon a glossary of terms used in the study of ischemia and reperfusion. *Cardiovasc Res.* 27: 162, 1993.

Kono T, Morita H, and Nishina T (1996):

Circadian variations of onset of acute myocardial infarction and efficacy of thrombolytic therapy. *J Am Coll Cardiol* 1996; 27:774–778.

Kovanen PT, Kaartinen J, and Paavonen T (1995):

Infiltrates of activated mast cells at the site of coronary atheromatous erosion or rupture in myocardial infarction. *Circulation*, 92: 1084, 1995.

Krone RJ, Dwyer EM Jr, and Greenberg H (1989):

Risk stratification in patients with first non-Q wave infarction: limited value of the early low level exercise test after uncomplicated infarcts: the

References

Multicenter Post- Infarction Research Group. J Am Coll Cardiol. 1989; 14:31-37; discussion 38-39.

Krucoff MW, Croll MA, and Pope JE (1993):

For the TAMI 7 study group: continuous 12 lead ST-segment recovery analysis in the TAMI 7 study: performance of a noninvasive method for real-time detection of failed myocardial reperfusion. Circulation 1993; 88:437-446.

Lamas J, Flaker G, and Mitchell G (1995):

For the survival and ventricular enlargement investigators: Effect of infarct artery patency on prognosis after acute myocardial infarction. Circulation 1995; 92: 1101-1109.

Lauer MS, Francis GS, and Okin PM (1999):

Impaired chronotropic response to exercise stress testing as a predictor of mortality. JAMA. 1999;281: 524-529.

Lefer A.M., Tsao P.S., Lefer D.J., and Max-L (1991):

Role of endothelial dysfunction in the pathogenesis of reperfusion injury after myocardial ischemia .FASEB. J., 2029, 1991.

Lehmann EL, annd D'Abrera HJM (1995):

Nonparametrics: statistical methods based on ranks. San Francisco; Holden-Day 1995.

Leor J, Poole WK, and Kloner RA (1996):

Sudden cardiac death triggered by an earthquake. N Engl J Med 1996; 334:413-419.

References

Libby P (1995):

Molecular basis of the acute coronary syndromes. *Circulation* 1995; 91:2844–2850.

Linderer T, Schroder R, and Arntz R (1993):

Pre-hospital thrombolysis: beneficial effects of very early treatment on infarct size and left ventricular function. *J Am Coll Cardiol.* 1993; 22: 1304-1310.

Londeree BR, and Moeschberger ML (1984):

Influence of age and other factors on maximal heart rate. *J Cardiac Rehabil.* 1984; 4:44–49.

M.Gabriel Khan (1999):

Cardiac drug therapy 5th ed., W.B.Saunders, an imprint of Harcourt Brace and Co.Ltd.1999(P 188-189).

Mahmarian JJ, Mahmarian AC, and Marks GF (1995):

Role of adenosine thallium-201 tomography for defining long-term risk in patients after acute myocardial infarction. *J Am Coll Cardiol.* 1995; 14: 1609–1615.

Manyari DE, Kostuk WJ, and Purves PP (1983):

left and right ventricular function at rest and during bicycle exercise in the supine and sitting positions in normal subjects and patients with coronary artery disease. Assessment by radionuclide ventriculography. *Am J Cardiol.* 1983; 51: 35-42.

References

Mark DB, Shaw L, and Harrell FE (1991):

Prognostic value of a treadmill exercise score in outpatients with suspected coronary artery disease. *N Engl J Med.* 1991; 325:849-853.

Markiewicz W, Houston N, and DeBusk RF (1977):

Exercise testing soon after myocardial infarction. *Circulation.* 1977; 56:26-31.

Markis JE, Brewer CC, and Braunwald E (1985):

Myocardial infarction without early coronary angiographic evidence of occlusion: the NHLBI thrombolysis in myocardial infarction trial. *Circulation*, 1985; 72: 56.

Martin GV, and Kennedy JW (1994):

Choice of thrombolytic agent. In Julian D. and Braunwald E: management of acute myocardial infarction. London, W.B., Saunders Ltd., 1994, P.71.

Martin WH, Colye EF, and Bloomfield SA (1986):

Effects of physical deconditioning after intense endurance training on left ventricular dimensions and stroke volume. *J.Am.Coll Cardiol.* 1986; 7: 982.

Matsuda M, Fujiwara J, and Onodera T (1987):

Quantitative analysis of infarct size, contraction band necrosis, and coagulation necrosis in human autopsied hearts with acute myocardial infarction after treatment with selective intracoronary thrombolysis. *Circulation* 76: 981, 1987.

References

McHam SA, Marwick TH, and Pashkow FJ (1999):

Delayed systolic blood pressure recovery after graded exercise: an independent correlate of angiographic coronary disease. *J Am Coll Cardiol.* 1999; 34:754 –759.

Meier FA, Al-Hani A J, and Dear W E and Members of the Laboratory and Performance Standards Committee of the Society for Cardiac Angiography and Interventions (1987):

Guidelines for continuous quality improvement in the cardiac catheterization laboratory. *Cathet. Cardiovasc. Diagn.* 30:191, 1987.

Metz B, White H, and Granger C (1998):

For the global use of strategies to open occluded coronary arteries in acute coronary syndromes (JUSTO-IIb) investigators randomized comparison of direct thrombin inhibition versus heparin in conjunction with fibrinolytic therapy for acute myocardial infarction: results from the JUSTO IIb trial . *J Am Coll Cardiol.* 1998; 31: 1493-1498.

Mirvis DM (1988):

Physiologic basis for anterior ST segment depression in patients with acute inferior wall myocardial infarction. *Am Heart J* 1988; 116:1308-22.

Mizuno P, Ockene I, and Goldberg R J (1988):

A prospective randomized trial of outpatient versus inpatient cardiac catheterization. *N. Engl. J. Med.* 319:1252, 1988.

Moreno PR, Bernardi VH, and Lopez-Cuellar J (1996):

References

Macrophages, smooth muscle cells, and tissue factor in unstable angina: Implications for cell-mediated thrombogenicity in acute coronary syndromes. *Circulation* 1996; 94:3090–3097.

Moreno PR, Falk E, and Palacios IF (1994):

Macrophage infiltration in acute coronary syndromes: Implications for plaque rupture. *Circulation* 1994; 90:775–778.

Muller DWM, and Topol EJ (1990):

Selection of patients with acute myocardial infarction for thrombolytic therapy. *Ann Intern Med* 1990; 113:949–960.

Neri Serner G, Gensini J, and Poggesi L (1990):

Effect of heparin, aspirin, or alteplase in reduction of myocardial ischemia in refractory unstable angina. *Lancet* 1990; 335: 615-618.

Newby LK, Califf RM, and Guerci A (1996):

Early discharge in the thrombolytic era: an analysis of criteria for uncomplicated infarction from the Global Utilization of Streptokinase and t-PA for Occluded Coronary Arteries (GUSTO) trial. *J Am Coll Cardiol.* 1996; 27:625– 632.

Newman PE (1981):

The coronary collateral circulation: determinants and functional significance in ischemic heart disease. *Amr Heart J*, 1981; 102:431.

Nissen S E (1994):

Principles and applications of digital imaging in cardiac and coronary angiography. In Pepine, C. J., Hill, J. A., and Lambert, C. R. (eds.):

References

Diagnostic and Therapeutic Cardiac Catheterization. 2nd ed. Baltimore, Williams and Wilkins, 1994, p. 162.

Norrell MS, Lyons JP, and Gardener JE (1989):

Significance of "reciprocal" ST segment depression: left ventriculographic observations during left anterior descending coronary angioplasty. J Am Coll Cardiol 1989; 13:1270-4.

O'Mera JJ III, Mcn Mc Nutt RA, and Evans ET (1994):

A decision analysis of streptokinase plus heparin as compared with heparin alone for deep-vein thrombosis. N Engl J Med 1994; 330: 1864-1869.

Okin PM, and Kligfield P (1995):

Gender-specific criteria and performance of the exercise electrocardiogram. Circulation. 1995;92:1209 -1216.

Pepine CJ, Hill JA, and Lambert C (1998):

Diagnostic and therapeutic cardiac catheterization. 3rd ed. Baltimore: (Williams & Wilkins, 1998).

Pina I L, Balady G J, and Hanson P (1995):

Guidelines for clinical exercise testing laboratories: A statement for healthcare professionals from the Committee on Exercise Cardiac Rehabilitation, American Heart Association. Circulation 91:912, 1995.

Pujadas G (1980):

coronary angiography. Mc Graw-Hill book company, 1980; chapter 13, P 226-227.

References

Quyyumi AA, Crake T, and Rubens MB (1986):

Importance of "reciprocal" electrocardiographic changes during occlusion of left anterior descending coronary artery. Studies during percutaneous transluminal coronary angioplasty. *Lancet* 1986;I: 347-50.

Rapaport E (1989):

Thrombolytic agents in acute myocardial infarction. *N Engl J Med* 1989; 320: 861-864.

Reimer KA, and Jennings RB (1979):

The "wavefront phenomenon" of myocardial ischemic cell death. II. Transmural progression of necrosis within the framework of ischemic bed size (myocardium at risk) and collateral flow. *Lab Invest* 1979; 40:633-644.

Rentop KP, Cohen M, and Blake H (1985):

Changes in collateral channel filling immediately after controlled coronary artery occlusion by an angioplasty balloon in human subjects. *J Amr Coll Cardiol*, 1985 ; 5 : 587 .

Rentrop J M, and Cohen KW (1986):

The future of digital angiography. *Cathet. Cardiovasc. Diagn.* 27:14, 1986.

Ridker PM, Herbert PR, and Fuster V (1993):

Are both aspirin and heparin justified as adjuncts to thrombolytic therapy for acute myocardial infarction? *Lancet* 1993; 341: 1574-1577.

References

Rodgers GP, Ayanian JZ, and Gary Balady (2000):

ACC/AHA clinical competence: a statement on stress testing. J Am Coll Cardiol. 2000; 36:1441-53.

Ronnevik PK, and Von der Lippe G (1992):

Prognostic importance of predischARGE exercise capacity for long-term mortality and non-fatal myocardial infarction in patients admitted for suspected acute myocardial infarction and treated with metoprolol. Eur Heart J. 1992; 13:1468 -1472.

Rothe C F (1986):

Physiology of venous return: an unappreciated boost to the heart. Arch. Intern. Med. 1986; 146:977.

Rowell LB (1986):

Human Circulation: Regulation During Physical Stress. New York: Oxford University Press; 1986.

Ryan TJ, Anderson JL, and Antman EM (1996):

ACC/AHA Practice Guidelines: guidelines for the management of patients with acute myocardial infarction. Circulation. 1996; 94:2341-2350.

Ryan TJ, Anderson JL, and Antman EM (1996):

ACC/AHA guidelines for the management of patients with acute myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Management of Acute Myocardial Infarction). J Am Coll Cardiol. 1996;28:1328 -1428.

References

Sacks FM, Pfeffer MA, and Moye LA (1996):

The effect of Pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. N Engl J Med 1996; 335:1001-1009.

Schartz KM, Turner JD, and Sheffield LT (1981):

Limited exercise testing soon after myocardial infarction. Ann Intern Med 1981;94:727-34.

Schlant RC, Silverman ME, and Roberts WC (1990):

Anatomy of the heart. In Hurst (The heart) , 7th ed.,Mc Graw-Hill, New York, 1990; chapter 2: P 24-26.

Schlesinger M J, and Reiner L (1995):

Focal myocytolysis of the heart. Am J Physiol. 31 : 443,1995.

Schoen F J (1994):

The heart. In Cotran, R.s., Kumar, V., and Robbins, S. L. (eds.): pathologic Basis of disease. Philadelphia, W.B. Saunders Company, 1994, P.517.

Schwartz H, Leiboff RH, and Bren GB (1984):

Temporal evaluation of the human coronary circulation after myocardial infarction . J Amr coll Cardiol , 1984 ; 4: 1088 .

Shah PK, Cercek B, and Lew AS (1993):

Angiographic validation of bedside markers of reperfusion. J Am Coll Cardiol 1993; 21:55-61.

References

Shepherd JT (1987):

Circulatory response to exercise in healthy. *Circulation* 1987; 76: 3-10.

Sheps DS, Ernst JC, and Myerburg R.J (1978):

Diastolic blood pressure response to treadmill exercise test. *Circulation* 1978; 58:141-153.

Silber H, Khan S, and Matloff J (1993):

The St Jude valve: thrombolysis as the first line of therapy for cardiac valve thrombosis. *Circulation* 1993; 87:30-37.

Simes R, Topol E, and Holmes Jr (1995):

For the GUSTO-1 investigators. Link between the angiographic substudy and mortality outcomes in a large randomized trial of myocardial reperfusion: importance of early and complete infarct artery reperfusion. *Circulation* 1995; 91: 1923-1928.

Sivarajan ES, Snydsman A, and Smith B (1977):

Low-level treadmill testing of 41 patients with acute myocardial infarction prior to discharge from the hospital. *Heart Lung*. 1977;6: 975-980.

Stevenson RN, Umachandran V, and Ranjadayalan K (1994):

Early exercise testing after treatment with thrombolytic drugs for acute myocardial infarction: importance of reciprocal ST segment depression. *BMJ*1994;308:1189-1192(7 May).

Stewart JT, French JK, and Theroux P (1998):

References

Early noninvasive identification of failed reperfusion after intravenous thrombolytic therapy in acute myocardial infarction. J Am Coll Cardiol 1998; 31:1499-1505.

Sullivan ID, Davies DW, and Sowton E (1984):

Submaximal exercise testing early after myocardial infarction. Prognostic importance of exercise induced ST segment elevation. Br Heart J 1984; 52:147-53.

Tamesis B, Stelken A, and Byers S (1993):

Comparison of the Asymptomatic Cardiac Ischemia Pilot and modified Asymptomatic Cardiac Is Pilot versus Bruce and Cornell exercise protocols. Am. J. Cardiol. 72:715, 1993.

Terrin ML, William DO, and Braunwald E (1993):

Two- and three-year results of the Thrombolysis In Myocardial Infarction (TIMI) phase II clinical trial. J Am Coll Cardiol. 1993; 22:1763-1772.

Theroux P, Waters DD, and Halphen C (1979):

Prognostic value of exercise testing soon after myocardial infarction. N Engl J Med. 1979; 301:341-345.

TIMI Study Group (1985):

The Thrombolysis in Myocardial Infarction (TIMI) trial: Phase I findings. N. Engl. J. Med. 312:932, 1985.

Topol E (1991):

Which thrombolytic agent should one choose? Prog Cardiovasc Dis 1991; 34:165-178.

References

Topol EJ, and Califf RM (1998):

Thrombolytic therapy for elderly patients. *N Engl J Med* 1998; 327:45-46.

Topol EJ, and Nissen S (1995):

Our preoccupation with coronary luminology: The dissociation between clinical and angiographic findings in ischemic heart disease. *Circulation* 1995; 92:2333-2342.

Tubau DR, Wondrow MS, and Tulsrud P R (1981):

Radiographic techniques used in cardiac catheterization. In Pepine, C. J., Hill, J. A., and Lambert, C. R. (eds.): *Diagnostic and Therapeutic Cardiac Catheterization*. 2nd ed. Baltimore, Williams and Wilkins, 1981, p. 141.

Valsania P, Zorio HS, and Kowalschuk G (1991):

Severity of coronary artery disease in young patients with insulin dependant diabetes mellitus. *Am Heart J* 1991; 122:695.

Waters DD, Chaitman BR, and Bourassa MG (1980):

Clinical and angiographic correlates of exercise-induced ST segment elevation. Increased detection with multiple ECG leads. *Circulation* 1980;61:286- 95.

Weiner DA, McCabe C, and Klein MD (1978):

ST segment changes postinfarction: predictive value for multivessel coronary disease and left ventricular aneurysm. *Circulation* 1978; 58:887-91.

References

White H (1992):

Remodeling of the heart after myocardial infarction. Aust N Z J Med 1992; 22(5suppl): 601-606.

White HD, and Van de Werf Fj (1998):

Clinical cardiology: frontiers: thrombolysis for acute myocardial infarction . Circulation 1998; 97:1632-1646.

Williams JA, Littmann D, and Hall JH (1976):

Coronary arteriography. II. Clinical experiences with the loop-end catheter. N Engl J Med 1976; 262:328-332.

Yvorchuk KJ, Davies RA, and Chang KL (1994):

Measurment of left ventricular ejection fraction by acoustic quantification and comparison with radionuclide angiography. Am.J.Cardiol 74:1052-1056; 1994.