

References

1-Abizaid AS, Mintz GS, Abizaid A , et al., (1999) :

One-year follow-up after intravascular ultrasound assessment of moderate left main coronary artery disease in patients with ambiguous angiograms.

J Am Coll Cardiol. 34: 707-715.

2-Andersen K, Vik-Mo H and Hoff PL, (1986) :

Left main or proximal left anterior descending coronary artery disease: detection by echocardiographic evaluation of interventricular septal function during exercise.

Clin Physiol. 6(3) : 269-80.

3-Anton P, Marc A, Bos A , et al ., (1988) :

An electrocardiographic pattern, characteristic for extensive myocardial. Ischemia.

Circulation, Supplement II. Vol. 78. No. 4.

4-Atie J,Brugada J, SmeetsL R, et al.,(1991):

Clinical presentation and prognosis of left main coronary artery disease in 1980s.

European Heart Journal,12:495-502.

5-Bentivoglio LG, VanRaden MJ, Kelsey SF, et al ., (1984) r

Percutaneous transluminal coronary angioplasty (PTCA) in patients with relative contraindications : results of the National Heart, Lung, and Blood Institute PTCA Registry.

Am J Cardiol. 53: 82-8.

6-Bergelson B, Tommaso CI and Chicago,(1995):

Left main coronary artery disease: Assessment, diagnosis, and therapy.

Am H J. 129:350-359.

7- Bernard JG, Braunwald E and Rutherford J (1997):

chronic coronary artery disease in Braunwald's Heart disease. A text book of cardiovascular medicine, edited by Braunwald E, 6th Edition WB saunders Company, Philadelphia pp 1289-1366.

8-Block PC, Myler RK, Stertz S, et al., (1983) :

Morphology after transluminal angioplasty in human being.

N Engl J med. 305- 382.

9-Block P and Popp R,(1985):

Detecting and excluding significant left main coronary artery narrowing by echocardiography.
Am J Cardiol.55:937-940

10-Boehrer JD, Lange RA, Willard JE, et al ., (1992) :

Markedly increased periprocedure mortality of cardiac catheterization in patients with severe narrowing of the left main coronary artery.
Am J Cardiol. 70: 1388-90.

11-Bourdillon PD, Brodeick TM, Savada SG, et al .,(1989)

Regional wall motion index for infarct and non-infarct regions after reperfusion in acute myocardial infarction : comparison with global wall motion index.
J Am Soc Echocardiogram. 2 (6) : 388-407.

12-Ronderos R, Salcedo EE, Kramer JR, et al ., (1984) :

Value and limitations of two-dimensional echocardiography for the detection of left main coronary artery disease.
Cleve Clin. 51: 7-12.

13-Boyne T, Koplan B, Parsons W, et al.,(1997):

Predicting adverse outcome with exercise SPECT technetium 99m sestamibi imaging in patients with suspected or known coronary artery disease.
Am J Cardiol.79:270-4.

14- Bowen E, (2000) :

Systolic heart failure in manual of cardiovascular medicine edited by Tobol E, Griffen B and Marso S 1st editions lippincott williams and wilkins Philadelphi, pp 543.

15-Brown KA, (1995) :

Prognostic value of cardiac imaging in patients with known or suspected coronary artery disease : comparison of myocardial perfusion imaging, stress echocardiography, and position emission tomography.
Am J Cardiol. 75: 35D-41D.

16-Buchbinder M, Reiman M, Fenner J, et al ., (1992) :

High speed rotational ablation in complex calcified lesions.

Circulation 86 (suppl1) 1-513.

17-Bulklev BH and Roberts W C, (1976):

Atherosclerotic narrowing of the left main Coronary artery a necropsy analysis of 152 patients with fatal coronary heart disease and varying degree of left main narrowing.
Circulation.53:823-828.

18-Caracciolo EA, Davis KB, Kaiser GC, et al.,(1995):

Comparison of surgical and medical group survival in patients with left main coronary artery disease long-term CASS experience
Circulation , 91:2325-2334.

19-Califf RM, Conley MJ, Behar VS, et al.,(1984) :

Left main equivalent coronary artery disease : its clinical presentation and prognostic significance with nonsurgical therapy.
Am J Cardiol. 53: 1489-1495.

20-Centi CR, Selby JH, Christie LG, et al., (1979) :

Left main coronary artery stenosis : clinical spectrum, path physiology and management.
Progress in Cardiovascular Disease, 22: 73-106.

21-Chauhan A ,Zubaid M ,Ricci DR, et al.,(1997):

Left main intervention revisited: early and late outcome of PTCA and stenting.
Cathet Cardiovasc Diagn.41:21-29.

22-Chaitman BR, Fisher Ld, Bourassa MG, et al., (1981) :

Effect of coronary bypass surgery on survival patterns in patients with left main coronary artery disease : report of the Collaborative Study in Coronary Artery Surgery (CASS).
Am J Cardiol. 48: 755-77.

23-Chaix AF, Barragan P, Silvestri M, et al., (1995) :

Rotablator and endoprosthesis on the left main coronary trunk.
Arch mal Couter Vaiss. 88: 95-97.

24-Chandraratna PA, Aronow WS, Murdock K, et al., (1980):

left main coronary arterial patency assessed with cross- section echocardiography.
American Journal of cardiology 46:91.

- 25-Charitos CE, Nanas JN, Tsoukas A, et al., (1997):**
Total occlusion of the left main coronary artery with preserved left ventricular function.
Int J Cardiol, 19;61(2) :193-6.
- 26-Chen CC, Morganroth J, Ogawa S, et al., (1980)**
Detecting left main coronary artery disease by apical cross-sectional echocardiography.
Circulation, 62: 288-93.
- 27-Chernoff DM, Ritchie CJ and Higgins CB, (1997) :**
Evaluation of electron beam CT coronary angiography in health subjects.
Am J Radiol. 169: 93-99.
- 28-Cho ZH, Mun CW and Friedenber RM, (1991) :**
MR angiography of coronary vessels with 2-D planar image scanning.
Magn Reson Med. 20: 134-43.
- 29-Christian TFMiller TD, Bailey KL, et al., (1992) :**
Noninvasive identification of severe coronary artery disease using exercise tomographic thallium-201 imaging.
Am J Cardiol. 70: 14-20.
- 30-Cohen DJ, Kim D and Baim DS, (1991) :**
Origin of the left main coronary artery from the "non-coronary sinus of Valsalva".
Cathet Cardiovasc Diagn. 22: 150-2.
- 31-Cohen MV, Cohn PF, Herman MV, et al., (1972):**
Diagnosis and prognosis of main left coronary artery obstruction.
Circulation, 45(suppl .11) 1-57.
- 32-Colle JP, Delearche N, Bourdeaul A, et al., (1991) :**
Nondiagnosed left main ostial stenosis partly due to the use of 5 French coronary angiographic catheter.
Cathet Cardiovasc Diagn. 22: 180-3.
- 33-Compeau L, Corbars F, Crocnet L, et al., (1978) :**
Left main coronary artery stenosis : The influence of aortocoronary bypass surgery on survival.
Circulation, 57: 1111-5.

- 34-Conley MJ, Ely PL, Kisslo J, et al.,(1978) :**
The prognostic significance of left main stenosis.
Circulation, 57: 947-52.
- 35-Dacosta A, Tarduy B, Favre JP, et al., (1994)**
left main coronary artery disease.
Arch Mal Coeur Vaiss, 87(9):1225-32.
- 36-Daniel WG, Scoder E, Mugge A, et al., (1988) :**
Transesophageal echocardiography in infection endocarditis.
Am J Cardiac Imaging. 2: 78-85.
- 37-Davis KB, Chaitman BR, Killip T. et al., (1983) :**
Effect of coronary bypass surgery on operative mortality and survival patterns insubsets of patients with left main coronary artery disease. In Hammermeister KE (ed.) : Coronary Bypass Surgery : the late Results.
New York Praeger, p. 129.
- 38-Davies SW, Winterton Sj and Rothman MT,(1992):**
Intravascular ultrasound to assess left main stem coronary artery lesion .
Br H J. 68:524-526.
- 39-Deepak KP, Jacobs GS , Shahriar L, et al., (1994) :**
Intravascular ultrasound for angiographically indeterminant left main coronary artery disease.
Echocardiography. 11: 65-69.
- 40-Dessault O, Pasquire G, Etlinger P, et al., (1988) :**

Stenosis of the main trunk of the coronary artery Apropos of 54 cases.
Rev Med Interne, 9(2) 145-8.
- 41-Diderholm E,Andren B,Frostfeldt G, et al.,(2002):**
ST depression in ECG at entry indicates severe coronary lesions and large benefits of an early invasive treatment strategy in unstable coronary artery disease.
European heart J.23: 41-49.
- 42-Douglas PS, Fiolkoski J, Berko B, et al., (1988) :**
Echocardiographic visualization of coronary artery anatomy in the adult.
J Am Coll Cardiol. 11: 565-71.
-

- 43-Edelman RR, Manning WJ, Bursstein D, et al., (1991) :**
Coronary arteries breath-hold MR angiography. Radiology, 181: 641-643.
- 44-Eigler NL, Weinstock B and Douglas JS, (1993) :**
Excimer laser coronary angioplasty of aorto-ostial stenoses : results of the Excimer Laser Coronary Angioplasty (ELCA) Registry in the first 200 patients.
Circulation, 88: 2049-57.
- 45-Ellis SG, Myler RK, King SB, et al., (1997) :**
Causes and correlated cardiac death after unsupported coronary angioplasty : implication for the use of advanced support techniques.
Am J Cardiol. 1447-1451.
- 46-Engelen DJ, Gorgels AP, Cheriex EC, et al., (1999) :**
Value of the electrocardiogram in localizing the occlusion site in the left anterior descending coronary artery in acute anterior myocardial infarction.
J Am Coll Cardiol. 34: 389-95.
- 47-Feigenbaum H. (1994):**
Assessment of the left ventricular function in the ECHO manual edited by Oh Jk, Seward JB and Tajik A J, 1st edition mayo foundation, USA PP.43.
- 48-Feigenbaum H. (1994):**
Coronary artery disease: regional wall motion analysis in the ECHO manual edited by Oh Jk, Seward JB and Tajik A J, 1st edition mayo foundation, USA PP.43.
- 49-Fernandez-Ortiz A, Badimon JJ, Falk E, et al., (1994) :**
Characterization of the relative thrombogenicity of atherosclerotic plaque components : implications for consequences of plaque rupture.
J Am Coll Cardiol. 23: 1562-1569.
- 50-Gandhi MM, Lampe FC and Wood DA, (1995) :**
Incidence, Clinical characteristics and short term prognosis of angina pectoris.
Br Heart J.73:1938-42.

51-Garini A, Astorri E, Pedroni P, et al.,(1999):

left main coronary artery disease, clinical and angiographic features, retrospective study regarding two groups of patients with left main coronary artery disease and three vessel disease
Minerva cardio angiol .47:231-8

52-Gensini GG, (1975) :

Coronary arteriography in prog. Cardiovasc .dis. edited by Gensini G, 2nd edition Future Publishing Company PP. 163-232.

53-Glagov S, Weisenberg BA, Zarins CK, et al., (1987) :

Compensatory enlargement of human atherosclerotic coronary arteries.
N Engl J Med. 316: 1371.

54-Gordan P R, Abrams C, Gash AK et al., (1987) :

Pericatheterization risk factors in left main coronary artery stenosis.
Am J Cardiol. 59: 1080.

55-Gorgels AP, Vos MA, Mulleneers R, et al., (1993):

Value of the electrocardiogram in diagnosing the number of severely narrowed coronary arteries in rest angina pectoris.
Am J Cardiol. 72:999-1003.

56-Harrison DG, White CW, Hiratzka LF, et al., (1984) :

The value of lesion cross-sectional area determined by quantitative coronary angiography in assessing the physiologic significance of proximal left anterior descending coronary artery stenosis.
Circulation, 69: 169 -75.

57-Helfant RH and Banka VS, (1978) :

A clinical and angiographic approach to coronary heart disease.
Circulation .78:15-27.

58-Hermiller JB, Buller CE, Tenaglia AN, et al., (1993) :

Unrecognized left main coronary artery disease in patients undergoing interventional procedures.
Am J Cardiol. 71: 173-6.

59-Hinohara T, Row MH and Robertson GC, (1991) :

Effect of lesion characteristics on outcome of directional coronary atherectomy.

J Am Coll Cardiol. 17: 1112-20.

60-Hutter AM, (1980):

is there a left main equivalent ?

Circulation, 62:207-211.

61-Isner JM, Kishel J, Kent KM, et al., (1981) :

Accuracy of angiographic determination of the left main coronary arterial narrowing : angiographic – histologic correlation analysis in 28 patients.

Circulation, 63: 1056.

62-Issaz K and Costa A, (1998) :

Use of the continuity equation for transesophageal Doppler assessment of severity of proximal left coronary artery stenosis. A quantitative coronary angiography validation study.

JACC Vol. 32. No. 1. 42-8.

63-Janosi A and Vertes A, (1991) :

Exercise testing and left main coronary artery stenosis.

Chest. 100:227-29.

64-Jackson, Atkinson and Orans, (1997):

Improvement of myocardial metabolism in coronary artery disease by beta blockade.

Br Heart J.39:829-33.

65-Janek B Malek I, ouhrabkova R, et al., (1998):

left main coronary artery stenosis.

Am J cardiol Feb 1;81(3): 343 -6.

66- Jefferson K and Rees S, (1980) :

Clinical Cardiac Radiology in modern trends in cardiology edited by Jones M 3rd edition Butter Worths. PP. 68-75.

67-joshi J, (1999) :

Stress echocardiography in practical echocardiography edited by walsh C and wilde P, 1st edition, Greenwich Medical Media limited, London, PP 162.

68-Judkins MW, (1985):

coronary arteriography in coronary arteriography and intervention edited by judkins MM, douglas and king SD ,3rd edition New York Mc Graw-Hill PP.166-169.

69-Kaji S, Akasaka T and Yoshida K, (2001) :

Noninvasive coronary imaging.
J Cardiol 37(Suppl. 11) : 51-6.

70-Kanjwal MY, Carson DE and Schwartz JS, (1999) :

Chronic/subacute total occlusion of left main coronary artery – a case report and review of literature.
Angiology. 937: 45.

71- kapadia S, (2000):

Stable angina in manual of cardiovascular Medicine edited by tobol E, Griffen B and Marso S 1st editions, lippincott williams and wilkins Philadelphia, PP.72

72-Kasprzak JD, Drozd J, Peruga J, et al., (1999) :

Doppler detection of Proximal coronary artery stenosis using transesophageal echocardiography.
Kardiologie Polske, 50: 491-500.

73-Katritis D and Wegg-People M, (1991) :

Limitation of coronary angiography : An underestimated problem?
Clin Cardiol. 14: 20.

74-Kern MJ, (1989) :

Approach to the patient with left main coronary artery stenosis. Cathet Cardiovas Diagn. 18: 181-2.

75-Kennedy JW, Baxley WA, Bunnell IL, et al.,(1982):

mortality related to cardiac catheterization and angiography
Cathet Cardiovasc Diagn.8:323-340

76-Khaja F, Sharma SD, Easlev RM, et al.,(1974) :

Left main coronary artery lesions : Risks of catheterization. Exercise testing and Surgery.
Circulation, 49 and 50 (suppl II) : 136-40.

77-Khandheria BK, Tajik AJ and Sevard JB, (1996) :

Multiphase transesophageal echocardiography : Examination technique, anatomic correlations, and image orientation.

Crit Care Clin. 12: 203.

78-Koh KK, Howng HK, Lee, et al., (1993) :

Isolated left main coronary ostial stenosis in Oriental people :
operative histopathologic and clinical findings six patients.
J Am Cordiol.21:369-73

79-Kosuga K, Tamai H, Kyo E, et al ., (2000):

Predictors of prognosis after angioplasty in the unprotected
left main coronary artery.
J Cardiol .36 (3):149-57

80-Ko JK, Nishimura RA, Holmes DR, et al ., (1991) :

Predictors of early mortality in patients with angiographically
documented left main coronary artery disease.
Cathet Cardiovas Diagn. 24: 84-7.

81-Kovac JD and de Bono DP, (1996) :

Cardiac catheter complications related to left main stem
disease.
Heart, 76: 76-78.

82-Krzanowski M, Bodzon W, Brzostek T, et al., (2000) :

Value of transthoracic echocardiography for the detection of
high-grade coronary artery stenosis : prospective evaluation in
50 consecutive patients scheduled for coronary angiography.
J Am Soc Echo Vol. 13. No. 12.

83-Lavine P, Kinbiris D, Segal BL, et al ., (1972):

left main coronary artery disease clinical, arteriographic and
hemodynamic appraisal.
Am J Cardiol 30:791-796.

84-Lee R, Lee S, Shyu K, et al., (2001) :

Immediate and long-term outcomes of stent implantation for
unprotected left main coronary artery disease.
Int J Cardiol. 9: 173-7

85-Levitin A, (2001) :

Intravascular ultrasound.
Tech Vasc Interv Radiol Marc. 4(1) : 66-74.

86-Leung WH, Alderman EL , Lee TC, et al.,(1995):

Quantitative arteriography of apparently normal coronary segments

with nearby or distant disease suggests the presence of occult, nonvisualized atherosclerosis.

J Am Coll Cardiol. 25:311-17

87-Li D and Deshpande V, (2001) :

Magnetic resonance imaging of coronary arteries.

Top Magn Reson Imaging Oct. 12 (5) : 337-47.

88-Lim JS, Proudfit W and Sones F, (1995) :

Left main coronary arterial obstruction : long term follow-up of 141 non-surgical cases.

Am J Cardiol. 36: 131-35.

89- Lipton MJ, Pfeifer JG, Murophy ML, et al ., (1977):

dangers of left main coronary artery lesions: angiographic technique and evaluation .

Inves Radiol Sep -Oct; 12 (5) : 447-54.

90-Loop FD , Lytle BW, Cosgrove DM, et al., (1979)

Atherosclerosis of the left main coronary artery : 5 year results surgical management.

American journal of cardiology, 44:195-201.

91-Lozner E, Johnson L, Johnson S, et al., (1989):

coronary arteriography 1984-1987: areport of the registry of the society for cardiac angiography and interventions. An analysis of 218 deaths related to coronary arteriography .

Cathet Cardiovasc diagn; 17:11-14

92-Manning WJ, Li W, Boyle NG, et al., (1993) :

Fat-suppressed breath-hold magnetic resonance , coronary angiography.

Circulation, 87: 94-104.

93-Memmola C, Iliceto S and Rizzon P, (1993) :

Detection of proximal stenosis of left coronary artery by digital transesophageal echocardiography : Feasibility, sensitivity, and specificity.

J Am Soc Echocardiogr. 6: 149.

94-McNeer JF, Margolis JR, Lee KL, et al ., (1978):

The role of the exercise test in the evaluation of patients for ischemic heart disease.

Circulation. 57: 64-70.

95-Miller G. Honey M and EL – Sayed H, (1986):

Isolated coronary ostial stenosis.
cathet cerviovass Diagn. 12:30-4.

96-Moreus ML, Skorton DJ, Johnson MR, et al., (1988) :

Visual estimates of percent diameter coronary stenosis : A
battered gold standard.
J Am Coll Cardiol. 11: 882.

97-Morton E, (2001) :

Stress testing in cardiac evaluation. Current concepts with
emphasis on the ECG.
Chest 119. No. 3. 342 -350.

98-Moshage WB, Achenbach S, Sees? B, et al., (1995) :

Coronary artery stenosis : three dimensional imaging with
electrocardiographically triggered, contrast enhanced, electron
beam CT.
Radiology. 196: 707-714.

99-Mott L, Brizolara A, McCreery C, et al., (1999):

Left main coronary artery diseases.
Am J Med Sci Dec ; 318(6) :405-12.

100- Motthew D, (2000):

Stress echocardiography in manual of cardiovascular
Medicine edited by Tobol E, Griffen B and Marso S, 1st
editions, Lippincott Williams and Wilkins, Philadelphia, PP.543.

101- Motthew T, (2000):

Unstable angina and non –ST- segment elevation MI in
manual of cardiovascular Medicine edited by Tobol E, Griffen
B and Marso S 1st editions, Lippincott Williams and Wilkins
Philadelphia, P.P.27.

102-Moehara A Mintz G, Castagna M, et al., (2001) :

intravascular ultrasound Assessment of the stenoses location
and morphology in the left Main Coronary Artery in Relation
to Anatomic left Main Length.
Am J Cardiol. 88:1-4.

103-Muller DW, Ellis SG and Topol EJ, (1989) :

Atherectomy of the left main coronary artery with
percutaneous cardiopulmonary bypass support.
Am J Cardiol. 64: 114-6.

104-Nishimura RA, Edward WD, Warnes CA, et al., (1990) :

Intravascular ultrasound imaging : in vitro validation and pathological correlation .
J Am Coll Cardiol. 16: 145.

105-Nissen S, (2001) :

Coronary angiography and intravascular ultrasound.
Am J Cardiol 16 Feb. 87(4A) : 15A-20A.

106-Ogawa S, Chen CC, Hubbard FE, et al., (1980) :

A new approach to visualize the left main coronary artery using apical cross sectional echocardiography.
American Journal of Cardiology. 45: 301.

107-O'Keefe JH, Hartzler GO, Rutherford BD, et al., (1989) :

Left main coronary angioplasty : early and late results of 127 acute and elective procedures.
Am J Cardiol. 64: 144-7

108-Offili EO, Kern MJ, Vrain JA, et al., (1995) :

Differential characterization of blood flow, velocity, and vascular resistance between proximal and distal normal epicardial human coronary arteries.: analysis by intracoronary Doppler spectral flow velocity.
Am Heart J. 130: 37-46.

109-Pamela S, John F and Berko B, (1988) :

Echocardiographic visualization of coronary artery anatomy in the adult.
JACC, 11(3). 565-71.

110-Pandian NG, Hsu TL, Schwartz SL, et al., (1992) :

Multiplane transesophageal echocardiography : imaging plane. Echocardiographic anatomy, and clinical experience with a prototype phased-array amnioplane probe.
Echocardiography. 9: 649-53

111-Panovasky R, Starek Z, Meluzin J, et al., (1997):

The ergometry stress test in patients with stenosis of the left main coronary artery.
Vnitr Lek.. 43 (10) : 659-62.

112-Patterson RE, Horowitz SE, Eng C, et al., (1983) :

Can noninvasive exercise test criteria identify patients with left main or 3-vessel coronary disease after a first myocardial infarction?

Am J Cardiol. 51: 361-372.

113-Pearce FB, Sheikh KH, deBruijn NP, et al., (1989) :

Imaging of the coronary arteries by transesophageal echocardiography.

J Am Soc Echocardiogr. 2: 276.

114-Pennell DJ, Keegan J, Firmin DN, et al., (1993) :

Magnetic resonance imaging for coronary arteries : technique and preliminary results.

Br Heart J. 70: 315-26.

115-Plotnick GD, Greene HL, Carliner NH, et al., (1988) :

Clinical indicators of left main coronary artery disease in unstable angina.

Annals Of Internal Medicine, 91: 149-153.

116-Poelandt JR, Thomason IR, Vliet'et WB, et al., (1992) :

Multiplane transesophageal echocardiography. Latest evolution in an imaging.

J Am Soc Echocardiogr, 5: 361.

117-Pujadas G, (1980) :

Coronary angiography in coronary arteriography and intervention edited by Pujadas, 1st edition. Mc Graw- Hill Book Company, PP. 55-100.

118-Quinn RR, Pflugfelder PW, Kosluk WJ, et al., (2000) :

Intracoronary ultrasound imaging : methods and clinical applications.

Can J Cardiol. 01- 16(7) : 911-17.

119-Quigly RL , Milanon CA, Smith LR, et al., (1993):

Prognosis and management Of anterolateral myocardial infarction in patients with severe left Main disease and cardiogenic shock: The left main shock syndrome.

Circulation, 88:65-70.

120-Rensing B, Alfons B and Robert-Jan van G. (1998) :

Intravenous coronary angiography by electron beam computed tomography A Clinical evaluation.

Circulation, 98: 2509-2512.

- 121-Rink LD, Feigenbaum H, Godfrey RW, et al., (1982) :**
Echocardiographic detection of left main coronary artery obstruction.
Circulation, 65: 719-24.
- 122-Rogers EW, Feigenbaum H, Weyman AE, et al., (1980) :**
Possible detection of atherosclerotic coronary calcification by two-dimensional echocardiography.
Circulation, 62: 1046-53.
- 123-Rolle F, Christides C, Corun Em, et al., (1994):**
Significant stenosis of the common trunk of the left coronary artery. Retrospective study of 127 cases .
Arch Mal Coeur Vaiss, 87 (7) : 899-905.
- 124-Rosch J, Demots H, Antonvic K, et al., (1974):**
coronary arteriography in left main coronary artery disease.
Am J Roentgenography, 121:533-590.
- 125-Ryan T, Armstrong WF and Feigenbaum H, (1986) :**
Prospective evaluation of the left main coronary artery using digital two-dimensional echocardiography.
J Am Coll. Cardiol. 7: 807-12.
- 126-Schluter M, Their W, Hinrichs A, et al., (1984) :**
Klinischer einsatz der transqsophagealen echokardiographie
Dtsch Med Wschr. 109: 722-727.
- 127-Segar DS, Brown SE, Sawada SG, et al.,(1992) :**
Dobutamine stress echocardiography : correlation with coronary lesion severity as determined by quantitative angiography.
J Am Coll Cardiol. 19: 1197-1202.
- 128- Sendecor and cochran, (1980):**
Statistical methods, 7th edition. In ames. Ia. Iowa state university.
- 129-Sherman TC, Litvack F, Grundfest W, et al., (1986) :**
Coronary angiography in patients with unstable angina pectoris.
N Engl J Med. 315:913.
- 130-Showl FA, Chun PK, Mutter ML, et al., (1989) :**
-

Asymptomatic left main coronary artery disease and silent myocardial ischemia.
Am H J. 117: 537.

131- Spiecker m ,Erbel L ,Rupprecht HJ, et al.,(1994):

Emergency angioplasty Of totally occluded left main coronary in acute myocardial infarction and unstable angina pectoris institutional experience And literature review.
Eur Heart J.15:602-607.

132-Stertz SH, Myler RK, Insel H, et al ., (1985) :

Percutaneous transluminal coronary angioplasty in left main stem coronary stenosis : a five year appraisal.
Int J Cardiol. 9: 149-59.

133-Stouffer GA, Morr L, Brizolara A, et al ., (2000):

left main coronary artery disease in adults younger than 50 years : a comparison with older patients.
Catheter cardivasc interv, 51(1) :11-17.

134-Stone PH, La Follette L, and Cohn K, (1982) :

Patterns of exercise treadmill test performance in patients with left main coronary artery disease : detection dependent on left coronary dominance or coexistence dominant right coronary disease.
Am H J. 104-13.

135-Taams EJ, Cornel JH, Roelandt JR, et al., (1988) :

Detection of left coronary artery stenosis by transoesophageal echocardiography.
European Heart Journal, 9: 1162-1166.

136-Takaro T, Hultgren HN, Lipton MJ,et al ., (1976) :

The VA cooperative randomized study of surgery for coronary arterial occlusive disease : Subgroup with significant left main lesions.
Circulation, 54(suppl 11) : III. 107-16.

137-Tanaka S, Ueda K, Yung,-Sheng H, et al ., (1996) :

Initial long-term results of directional coronary atherectomy of undirected left main coronary artery.
Circulation, 94(Suppl 1) : 361-67

138-Tardif JC, Vannan MA, Taylor K, et al., (1994) :

- Delineation of extended lengths of coronary arteries by multiplane transesophageal echocardiography.
J Am Coll Cardiol. 24: 909.
- 139-Taylor HA, Deumite NJ, Chaitman BR, et al., (1989):**
Asymptomatic left main coronary artery disease in the coronary artery surgery study (CASS) registry
Circulation, 79:1171-1179.
- 140-Tenz M, Laske A, Niederhauser U, et al., (1992):**
Main coronary artery stenosis: a continuous challenge
Schw Med wschr, 122:1479-83.
- 141-Tobis JM, Mallery J, Mahon D, et al., (1991) :**
Intravascular ultrasound imaging of human coronary arteries in vivo; analysis of tissue characterization with comparison to in vitro histological specimens.
Circulation, 81: 913.
- 142-Tommaso CL, Vogel JH and Vogel RA, (1992) :**
Coronary angioplasty in high-risk patients with left main coronary stenosis : results from the national registry of elective supported angioplasty.
Cathet Cardiovasc Diag. 25: 169-73.
- 143-Tommaso CL, Applefeld MM, Singietom RT, et al., (1988) :**
Isolated left main coronary artery stenosis and mediastinal radiotherapy as an etiologic factor.
Am J Cardiol. 61:1119-20
- 144-Topaz O, Warner M and Lenter P, (1991):**
Isolated significant left main coronary artery stenosis hemodynamic and clinical finding in 16 patients.
Am Heart J. 122:1308-14.
- 145-Topaz O, Disciascio G, Cowley MJ, et al., (1991) :**
Absent left main coronary artery : angiographic findings in 83 patients with separate ostia of the left anterior descending and circumflex arteries at the left aortic sinus.
Am Heart J. 122: 447-52.
- 146-Turi ZG, Rezkella S, Campbell CA, et al., (1990) :**
Left main percutaneous transluminal coronary angioplasty with the autoperfusion catheter in an animal model.
-

Cathet Cardiovasc Diagn. 21: 45-50.

147-Vered Z, Katz M, Rath S, et al., (1986) :

Two-dimensional echocardiographic analysis of proximal left main coronary artery in humans.
Am Heart J. 112: 972-977.

148-Vrublevsky AV, Boshchenko AA and Karpov RS, (2001) :

Diagnostics of main coronary artery stenoses and occlusions :
multiplane transoesophageal Doppler echocardiographic
assessment.
Eur J Echocardiogr. 2 :170-7.

149- wallentin J, Jernberg D, Genberg E, et al., (1980):

Prospective randomized study of coronary artery bypass in
stable angina pectoris.
Lancet. 2: 49-5.

150-Wakasugi S, Shibata N, Kobayashi T , et al., (1986) :

Specific perfusion pattern in stress. Tl myocardial
scintigraphy of left main coronary artery disease.
Eur J Nucl Med. 12: 369-74.

151-Weidinger F Frick M , Alber F, et al., (2002) :

Association of wall thickness of the brachial artery measured
with high resolution ultrasound with risk factors and coronary
artery disease.
Am J Cardiol. 89: 1025-1029.

152-Weiner DA, McCabe CH and Ryan TJ, (1980) :

Identification of patients with left main and three vessel
coronary disease with clinical and exercise test variables.
Am J Cardiol. 46: 21.30

153-Weller BF,(1989):

Anatomy, histology ,and pathology of the major epicardial
coronary arteries relevant to echocardiographic imaging
techniques.
J Am soc Echocardiogr,2:232.

154-Weyman AE, Feigenbaum H, Dillon JC, et al.,(1979) :

Nonivasive visulization of left main coronary artery by cross
sectional echocardiography.
Circulation, 45: 169-174.

- 155-White C, Gobel F, Campeau L, et al.,(2001):**
Effect of an aggressive lipid lowering Strategy on progression of atherosclerosis in the left main coronary artery from patients in the post coronary artery bypass graft trial.
Circulation,104:2660-2665.
- 156-Williams K, Candace M and Schneider BS, (1999) :**
Increased stress right ventricular activity on dual isotope perfusion SPECT. A sign of multivessel and/or left main coronary artery disease.
J Am Coll Cardiol. 34: 420-7.
- 157-Williams DO, (1996) :**
Catheterization interventions for the treatment coronary artery disease : techniques, results, and patients selective in : Smith TW., ed., Cardiovascular Therapeutics : A Companion Braunwald's Heart Disease. Philadelphia. Pa : WB Saunders. 57-65.
- 158-Yamagishi M, Miyatake K, Bepp S, et al., (1988) :**
Assessment of coronary blood flow by transesophageal two-dimensional pulsed Doppler echocardiography.
Am J Cardiol. 62: 641-644.
- 159-Yamagishi M, Miyatake K, Beppu S, et al.,(1989):**
Visualization of coronary blood flow by color Doppler imaging with transesophageal approach.
Chest, 96: 972-5.
- 160-Yamagishi M, Yasu T, Ohara K, et al., (1991) :**
Detection of coronary blood flow associated with left main coronary artery stenosis by transesophageal Doppler color flow echocardiography.
J Am Coll Cardiol. 17: 87.
- 161-Yamaji H, Iwasaki K, Kusach, S, et al., (2001) :**
Prediction of acute left main coronary artery obstruction by 12-lead electrocardiography ST segment elevation in lead aVR with less ST segment elevation in lead V1.
J.ACC 38(5) : 1348-54.
- 162-Yamanaka O and Hobbs RE, (1990) :**
Coronary artery anomalies in 126, 595 patients. Undergoing coronary arteriography.
Cathet Cardiovasc Diagn. 21 : 28-40.
-

163-Yip HK, Wu CJ, Chen MC, et al. (2001) :

Effect of primary angioplasty on total or subtotal left main occlusion : analysis of incidence, clinical features, outcomes and prognostic determinants.
Chest, 120: 1212-17.

164-Yoshida K, Yoshikawa J, Hozumi T, et al., (1990) :

Detection of left main coronary artery stenosis by transesophageal color Doppler and two-dimensional echocardiography.
Circulation, 81: 1271.

165-Youn Ho-Joong, and Foster Elyse, (2000) :

Transesophageal echocardiography (TEE) in the evaluation of coronary arteries.
Cardiology Clinics, Vol 18, No 4.

166-Zwicky P, Daniel WG, Mugge A, et al., (1988) :

Imaging of coronary arteries by color-coded transesophageal echocardiography.
Circulation, 62: 639.