

REFERENCES

- 1- Alexander, J.O.D. and Pasieczny , T. (1954): Follicular lymphoma of the skin. *Dermatologica*, 109: 1.
- 2- Allen, A.C.(1948) : Persistent "insect bites" (Dermal eosinophilic granulomas) simulating lymphoblastoma as, histiocytoses and squamous cell carcinomas. *Amer.J. Path.*, 24: 367-375.
- 3- Andersen, B.L., Brandrup , F. and Betri,J. (1982): Lymphocytoma cutis: a pseudomalignancy treated with penicillin. *Acta Derm. Venereol.*, 62:83-85.
- 4- Bafverstedt, B. (1943) : Lymphadenosis benigna cutis, clinical and pathological study . *Acta Derm. Venereol.*, 24:202 .
- 5- Bafverstedt,B. (1953) : Lymphadenosis benigna cutis as a symptom of malignant tumours. *Acta Derm.Venereol.*, 33:171.
- 6- Bafverstedt, B., Lundmark, C., Mossberg, H., Stenbeck, A. (1956): Lymphadenosis benigna orbitae. *Acta ophthalmol.* 34:367.

- 7- Bafverstedt, B. (1960): Lymphadenosis benigna cutis, its nature, course and prognosis. Acta Derm. Venereol., 40:10-17.
- 8- Bafverstedt, B. (1968) : Lymphadenosis benigna cutis . Acta Derm. Venereol., 48:1-6.
- 9- Baraf, C.S. and Shapiro .L. (1969) : Arch . Derm., 99:589.
Quoted by : Beare and Cunliffe, (1979).
- 10- Beare, J.M. and Cunliffe, W.J. (1979): Benign lymphoplasias of the skin. In Textbook of Dermatology (ed. A. Rook, D.S. Wilkinson and F.J.G. Ebling), 3rd edn, 46, P.1529-1532 . Blackwell Scientific publications, Oxford.
- 11- Berde, K. and Lang, M. (1935) : Miliary lymphocytoma of the skin. Dermat. Wchnschr. 100:641.
Quoted by: Hallam and Vickers (1939).
- 12- Bernengo, M.G., Capella, G., Mascoli, I. and Zina, G. (1977): Active rosette test in cutaneous lymphoproliferative disorders. Dermatologica, 154/6 : 342-349.
Quoted in: Excerpta Medica, 32.6:401., 1978.

13-Bernstein, H., Shupack, J., and Ackerman. A.B. (1974):
Cutaneous pseudolymphoma resulting from antigen injection. Arch. Derm., 110:756-757 .

14-Bettley ,F.R. (1947) :Miliary lymphocytoma. Brit. J. Derm., 59:70

15-Bianchi, G.E. (1950) : Die Penicillinbenhandlung der lymphozytome. Dermatologica, 100:273.
Quoted by : Self et al,(1969).

16-Bielicky, T., and Trapl, J. (1963) : Nicktvernabender Chronischer Erythematodes. Arch. Klin. exp. Derm., 217:438.
Quoted by: Lever and Schaumburg-Lever, (1975) .

17-Bluefarb, S.M. (1960) : "Cutaneous manifestations of the benign inflammatory reticuloses". Springfield, III: Charles C. Thomas, Publisher: P.131.
Quoted by: Self et al ,(1969) .

18-Bluefarb, S.M. (1975) : In Current medical management 2nd edn., Ed. Maddon S. St. Louis C.V. Mosby ,:P. 274.
Quoted by : Beare and Cunliffe, (1979).

- 19- Brown , S., Lane, P.R. , and Magnus, I.A. (1969) :
Skin photosensitivity from fluorescent Lighting.
Brit . J. Derm., 81:420.
- 20- Burckhardt, J.L. (1911) : Frankf. Z.Path., 6: 352.
Quoted by : Hellier (1939) .
- 21- Burnham, T.K., Fine,G., and Neblett, T.R. (1970) :
Immunofluorescent "band" test for lupus erythematosus.
Arch. Derm., 102:42.
- 22- Cabre, J. and Steigleder G.K. (1961) : Das Krankheits-
bild der lymphozytaren Infiltration (lymphocytic
Infiltration) im Sinne von Jessner und Kanof. Arch.
Klin. exp. Derm., 212:525.
Quoted by: Lever and Schaumburg-Lever, (1975).
- 23- Calnan, C.D. (1957) : Lymphocytic infiltration .
of the skin (Jessner) . Brit . J. Derm., 69:169.
- 24- Caro, W.A., Helwig, E.B. (1969) : Cutaneous lymphoid
hyperplasia. Cancer., 24:487-502 .

- 25- David, M., Shohat ,B., Hazaz, B., Feurman, E.J.,
and Joshua, H., (1980) : Identification of T and B
lymphocytes on skin sections from patients with
lymphoproliferative disorders of the skin. J.invest.
Derm., 75:491-494 .
- 26- Domonkos, A.N., Arnold, H.L. and Odom, R.B. (1982) :
Andrews' Diseases of the skin . Clinical Dermatology.
7th edn. Ch. 16 , P. 414.W.B. Saunders company .
Philadelphia.
- 27- Duncan, S.C., Evans, H.L., Winkelmann, R.K. (1980:
Large cell lymphocytoma. Arch. Derm., 116:1142-1146.
- 28- Epstein, S. (1935): Lymphocytoma of skin with involve-
ment of bulbar conjunctiva. Arch . f. Derm. u, Syph.,
173:181
Quoted by : Mopper and Rogin, (1951).
- 29- Evans, H.L., Winkelmann, R.K., Banks, P.H. (1979) :
Differential diagnosis of malignant and benign cutan-
eous lymphoid infiltrates (A study of 57 cases in
which malignant lymphoma had been diagnosed or
suspected in the skin). Cancer., 44:699.

- 30- Everett, M.A., Crockett. W., Lamb, J.H., and Minor, D. (1961) : Light sensitive eruptions in american indians. Arch. Derm., 83:243.
- 31- Fendt, H. (1900): Beitrage zur Kenntniss der sogenannten Sarcoiden Geschwulste der Haut. Arch. Derm. Syph., 53:213 .
Quoted by: Silvers and Glickman (1962) .
- 32- Fitzpatrick, T.B., Szabo, G., Hori, Y. (1968) : White leaf shaped macules. Arch. Derm., 98:1 .
- 33- Frain-Bell, W., and Magnus, I.A. (1971) : A study of the photosensitivity factor in cutaneous lymphocytoma. Brit.J.Derm., 84:25-31.
- 34- Gaul, L.E. (1953) : Heredity of multiple cystic benign epithelioma. Arch. Derm. Syph., 68:517.
Quoted by : Lever and Schaumburg-Lever, (1975)
- 35- Gertler, W. (1955) : Reticulosarkomatose umwandlung tumorartiger lymphozytome. Dermatol. Wochensch., 132: 1035-1042 .
Quoted by : Shelley et al , (1981) .

- 36- Godtfredsen , E. and Lindgren , S. (1953) : Orbital, palpebral and epibulbar lymphomas . Acta . Ophthalmol., 31:29.
Quoted by : Bafverstedt, (1968).
- 37- Gottlieb. B., and Winkelmann, R.K. (1962) : Lymphocytic infiltration of the skin. Arch. Derm., 86:626-633.
- 38- Haim, S. and Shafrir, A. (1970) : The nature of discoid lupus erythematosus . Acta Derm. Venereol., 50:86.
- 39- Hallam,R., and Vickers, H.R. (1939) : Two cases of miliary lymphocytoma or benign lymphadenoid granuloma of the skin. Brit .J. Derm., 51:251-259.
- 40 Hargreaves , A.(1949): Two cases of lymphocytoma. Dermatologica, 99:220-227.
Quoted by : Beare and Cunliffe, (1979).
- 41- Hashimoto, K., DiBella, R.J., Borsuk, G.M., and lever, W.F. (1967) : Eruptive hidradenoma and syringoma . Arch. Derm., 96:500.

- 42- Headington ,J.T.,Koski, J., and Murphy ,P.J. (1972)
Clear cell glycogenesis in multiple syringomas.
Arch. Derm., 106:353.
- 43- Hefer, W. (1956) : Arch.Klin. exp. Derm., 203:23 .
Quoted by : Beare and Cunliffe , (1979).
- 44- Hellier, F.F. (1939) : Lymphocytoma of the face. Brit.
J. Derm., 51:260-265 .
- 45- Herzberg, J. (1952). Retikulosarcomatose Umwandlung
multipler subkutaner Lymphozytome ber Acrodermatitis
chronica atrophicans . Dermtol. Wochenschr.,125:422.
Quoted by : Duncan et al, (1980) .
- 46- Hofer,W. (1956) : Lymphadenosis benigna cutis. Arch.
Klin. exp. Derm., 203:23.
Quoted by : Lever and Schaumburg-Lever, (1975).
- 47- Jadassohn , J.(1906) Arch. f. Dermat. u. Syph. 82:297.
Quoted by: Mopper and Rogin, (1951).
- 48- Jessner, M. and Kanof, N.B. (1953) :Lymphocytic
infiltration of the skin. Arch. Derm. Syph., 68:447.
Quoted by : Bafverstedt , (1968).

- 49- Johnson , W.C., Higdon, R.S., and Helwig, E.B.(1959):
Granuloma faciale. Arch. Derm., 79:42.
- 50- Kauffmann - Wolf, M. (1921): Scrotal lymphocytoma of
children. Arch.f. Dermat. u. Syph. 130:425.
Quoted by : Hellier, (1939).
- 51- Kawada, A., Mori, S., and Hayashi, T. (1970): Lymphod-
enosis benigna cutis :Pseudomalignant form and its
imprint smear cytology. Dermatologica, 141:339-347.
- 52- Kindermann, G. and Seibicke, E. (1977): Lymphodenosis
benigna cutis (M.Bafverstedt). Geburtshilfe, Frauenh-
eilkd., 37/12 : 1024-1027.
Quoted in : Excerpta Medica, 33.2:112., (1979).
- 53- Laymon, C.W. : Lupoid rosacea. Arch. Derm., 63:409.
- 54- Lever, W.F. and Schaumburg-Lever, G. (1975) :
"Histopathology of the skin". 5th edn. ch.33, P:708.
J.B.Lippincott Company, Philadelphia, Toronto.

- 55- Lewis, G.M. (1935): Is Spiegler-Fendt sarcoid a clinical or histologic entity . Arch. Derm. Syph., 31:67.
Quoted by: Hellier, (1939).
- 56- Loveman, A.B., and Fliegelman, M.T. (1951) : Lymphocytoma cutis. Arch . Derm., 63:169-183.
- 57- Lowenstein, A. (1911) : Wissenschaftl Gesellschaft Deutscher Aerzte in Bohmen. Munchen Med. Weshr., 58: 660.
Quoted by : Self et al, (1969) .
- 58- Mach ,K.W., and Wilgram, G.F. (1966) : Characteristic histopathology of cutaneous lymphoplasia (Lymphocytoma). Arch. Derm., 94:26-32
- 59- Mach, K.W., and Wilgram, G.F. (1966): Cutaneous lymphoplasia with giant follicles. Arch. Derm., 94:749-756.
749-756
- 60- Matras, A. (1930) : Uber lymphatische Knotcheneinlagerung (lymphozytome) in der Gesichtshaut. Arch. Franc. Derm., 161:520
Quoted by : Mach and Wilgram, (1966) .

61- Matras, A. (1964) : Hautarzt, 15:657.

Quoted by : Beare and Cunliffe, (1979) .

62 Miescher, G. (1937) : Le Granulome lymphodenoide de la peau (Lymphocytome), Bull. Soc. Franc. Derm., 44:1254.

Quoted by : Mach and Wilgram (1966).

63- Montgomery, H.(1951) : Personal communication to the authors.

Quoted by: Mopper and Rogin, (1951).

64- Mopper, C., and Rogin, J.R. (1951) : Benign solitary lymphocytoma. Arch . Derm., 63:184-190.

65- Muller-Hess, S., and Delacretaz, J. (1973) : Tricho epitheliom mit strukturen eines apokrinen Adenoms. Dermatologica, 146:170.

Quoted by : Lever and Schaumburg-Lever, (1975).

66- Muller, S.A., Wolff, K., and Winkelmann, R.K. (1967): Generalized eruptive histiocytoma. Arch. Derm., 96:11.

67- Nickel, W.R., and Reed, W.B. (1962) : Tuberous sclerosis. Arch. Derm., 85:209.

- 68- Ormsby ,O.S., and Montgomery , H. (1948) : Diseases of the skin , edn.7, philadelphia, Lea and Febiger.
Quoted by : Mopper and Rogin, (1951) .
- 69- Paschoud, J.M. (1958) : Die Lymphodenosis benigna cutis als ubertrogbare Infektionskrankheir, Neue Gesichtspunkte uber Verlauf, Histologie undtherapie. Hautarzt, 8:197.
Quoted by : Mach and Wilgram, (1966).
- 70- Pegum, J.S., and Landells, J.W. (1970) : Lymphosarcoma supervening on lymphocytoma. Trans. st.Johns. Hosp. Dermatol. Soc., 56:149-155.
Quoted by: Duncan et al (1980).
- 71- Pillsbury, D.M., Shelley , W.D., and kligman, A.M. (1956) : Follicular Lymphoma in Dermatology, Philadelphia, W.B. Saunders company., P:927.
Quoted by : Silvers & Glickman, (1962)
- 72- Pomeranz, J. (1970) : In discussion of : Roenigk, H. H., Jr., and Lesbowitz, S.A. : Lymphocytoma cutis. Arch . Derm., 101:248.

- 73- Postoma , C., and Sluiter, J.T.F. (1958) : The relationship between Bafverstedt's benign lymphadenosis of the skin and Jessner's lymphocytic infiltration of the skin. Acta Derm. Venereol., 30:180 .
- 74- Rayan, T.J. and Wilkinson, D.S. (1979): cutaneous vasculitis "angiitis" In Textbook of Dermatology (ed.A.Rook,D.S Wilkinson and F.J.G. Ebling). 3rd edn. ch. 31, P:1023. Blackwell Scientific publications, Oxford.
- 75- Sanderson, K.V. and Rona- Mackie (1979): Tumours of the skin. In Textbook of Dermatology (ed. A. Rook, D.S. Wilkinson and F.J.G.Ebling) 3rd edn.,ch. 66, P: 2155. Blackwell Scientific Publications, Oxford.
- 76- Schneider, W. (1964) : Lymphadenosis benigna cutis (blaige lymphozytome) der Mundschleimhaut, Z. Haut. Geschlechtskr., 36:17-24
Quoted by : Self et al, (1969).
- 77- Self, S.J., Carter, V.H., Noojin, R.O. and Ala, B. (1969): Disseminated lymphocytoma cutis (case report of miliarial and Nodular types). Arch. Derm., 100:459-464 .

-
- 78- Shelley, W.B., Wood, M.G., Wilson, J.F. , Goodman, R. (1981): Premalignant lymphoid hyperplasia (Preceding and Coexisting with Malignant lymphoma in the skin. Arch. Derm., 117:500-503.
- 79- Silvers, S.H., and Glickman, F.S., (1962) : Lymphadenosis benigna cutis. Arch. Derm., 86:512-514 .
- 80- Slepyan, A.H., (1958) : Case presentation. Chicago Dermatological Society.
Quoted by : Frain-bell and Magnus , (1971).
- 81- Spiegler, E. (1894): Uber die Sogenannte Sarkomatosis Cutis . Arch. Derm. Syph. 27:162 .
Quoted by : Silvers and Glickman, (1962).
- 82- Thies , W. (1955) : Beitrag zur Kenntniss der miliaren lymphocytome. Hautarzt, 6:32 .
Quoted by: Mach and Wilgram, (1966) .
- 83- Tobias, N. (1949) : Tick bite granuloma . J.Invest. Derm., 12:255
- 84- Waisman, M. (1973) : Immunofluorescent studies in lichen nitidus. Arch. Derm., 107:200 .

- 85- Weiss, P.M. and Cohen, A.D. (1971) Lichen nitidus of palms and soles. Arch. Derm., 104:538 .
- 86- Wilkinson, D.S. (1979) : Sarcoidosis. In Textbook of Dermatology (ed.A. Rook, D.S. Wilkinson and F.J.G. Ebling) , 3rd edn, ch. 48, P.: 1575. Blackwell Scientific Publications, Oxford.
- 87- Winchester, R.J.S.M.Fu., Wernet, P., Kunkel, H.G., Dupont, B. & Jerslid, C. (1975) : Recognition by pregnancy serum of non HL-A alloantigens Selectively expressed on B lymphocytes. J.Exp. Med., 141 :924-929 .
Quoted by : David et al, (1980) .
- 88- Winer, L.H., and Strakosch, E.A. (1941) : Tick bites dermacenter variabilis . (say). J.Invest. Derm., 4:249.
- 89- Winkelmann, R.K., and Muller, S.A. (1963) : Generalized eruptive histiocytoma. Arch. Derm., 88:586 .
- 90- Wiskemann, A. (1957) : Miliare Lymphocytome bei gleichzeitig bestehenden chronisch polymorphen lichtexanthemen. Hautarzt. , 8:275.
Quoted by : Frain-bell and Magnus, (1971).

91- Wiskemann, A. (1963) : Lymphozytome bei Lichturtikaria.

Derm. Wschr., 147:450 .

Quoted by : Frain-bell and Magnas (1971).

92- Wislinski, M. (1967) : Limfadenoma łagodna skóry

Leczona Doogńiskowymi Wstrzykiwaniami Hydrokortyzonu,

Przegl. Derm., 54:685-687 .

Quoted by : Self et al , (1969) .