

REFERENCES

References

Abbasciano, V.; Levato, F. and Zavasil, G. (1989) :

" Specificity of tumor markers (CEA, GICA, TpA, AFP) for the diagnosis of Large bowel cancers"

Med. Oncol. Tumor Pharmacother 6 (2) : 129 - 32 .

Abbasciano, V.; Levato, F. and Zavasil, G. (1988) :

"Specificity of CEA, GICA, Fibrinopeptide A and gamma - GT in the diagnosis and Follow up of gastric Cancer" .

Oncology 45 (3) : 159 - 61 .

Abdallah, N.; M.; Nassar, M., I.; Abd El - Hameed, G. and Marcell, N., G. (1990) :

" Carcinoembryonic antigen in bilharzial polypi " .

2 nd Int. Conference of Tumor Markers Oncology, Cairo, Egypt .

Abdel - Fattah, M.; Reiner, S.; Fouad, F., M. and Baner, Co., R. (1981) :

"Kinetics of acute phase proteins in rats after tumor transplantation" .

Cancer Res. 41 : 2548 - 55 .

Abdel - Hay, M.; Moussa, L.; Khalil, R.; Hassab, A.; Fahmy, E.; Moussa, S., and Beshir, N. (1990) :

" The value of serum levels of CEA and Ca 19 - 9 in patients with col-

orectal carcinoma"

2nd Int. Conference of Tumor Markers Oncology, Cairo, Egypt .

Alexander, J , C.; Chretien , P., B.; Dellon, A., L. and Snyder, J. (1978) :

"CEA levels in patients with carcinoma of oesophagus"

Cancer 42 : 1492 - 97 .

Anderson, A. and Nordlie, R., C. (1967) :

"Inorganic pyrophosphat - glucose phosphotransferase activity associated with alkaline phosphatase of E. coli"

J. Biol. Chem. 242 : 114 - 20 .

Andren - Sandberg, A. and Isacson, S. (1975) :

"Serum tissue polypeptide antigen (TPA) in surgical trauma" .

Proc. Ann Meet., Swedish Med. Soc., Stockholm .

Anonymous (1981) :

"Carcinoembryonic antigen : its role as a marker in the management of cancer " .

National Institutes of Health Consensus Development Conference Statement .

Cancer Res. 41 : 2017 - 22.

Attiyeh, F., F. and Stearns, M., W. (1981) :

"Second - Look Laparotomy based on CEA elevation in Colorectal cancer " .

Cancer 47 : 2119 - 2125 .

Banjo, C.; Gold, P. and Freedman, S.,O. (1973) :

"Structure of human digestive system"

Fed Proc 32 : 4444 - 50 .

Banjo, C.; Gold, P.; Freedman, S., O. and krupey, J. (1972) :

"Immunologically active heterosaccharides of carcinoembryonic antigen of human digestive system "

Nature 233 : 183 - 85 .

Baskies, A., M.; Chretien, P., B.; weiss, j, F.; Catalona, W., J. and spiegel, H.,E. (1980)

"Serum glycoproteins in cancer pts. First report of correlation, with in vitro and in vivo parameters, of cellular immunity"

Cancer 45 : 3050 - 60

Bastable , A. (1979) :

"Acute phase reactant proteins in gastrointestinal malignancies .

Arch. Biochem Biophys. 118 : 350 - 75 .

Nucl - Med - Commun 7 (2) : 121 - 27 .

Begent, R., H. (1987) :

"The value of carcinoembryonic antigen in clinical practice"

Br. J. Hosp. Med. 37 (4) : 335 - 38 .

Behring , C. (1985) :

Z. Klin. Chem. 6 : 113

Birkmayer, G.D. (1988) :

Tumor markers in diagnosis and therapy of cancer : Recent developments. Proceeding of the 1st Int. Conference of Tumour Markers Oncology (Feb 15-17, 1988) Cairo , Egypt .

Birkmayer, G.D; and Klavins J, V. (1990) :

Classification of tumour markers. Proceeding of the 2nd Int. Conference of Tumour Markers Oncology (Jan 16-19, 1990) Cairo, Egypt .

Birkmayer, G.D. and Klavins, J.V. (1990) :

Standardization and quality assurance of tumour markers tests. Proceeding of the 2nd Int. Conference of Tumour Markers Oncology (Jan 16-19, 1990) Cairo , Egypt .

Bjorklund, B. (1981) :

"Tissue polypeptide antigen (TPA), review and recent progress : critical evaluation of tumour markers"

Contr. Oncol. 7 : 73-80.

Bjorklund, V. and Bjorkland, B. (1979) :

Localizaion of synthesis of TPA in normal and malignant human tissues by imonurohistological techniques. In peeters, H.(ed.), protides of the biological fluids (Pergamon press, oxford), P 229-39.

**Bjorklund, B; Bjorklund, V.; Wiklund, B., Lundstrom, R.; Ek-
dahl, P.H., Hagbarg, L.; Kaijser, K., Ekland, G. and Lunning,
B. (1973) :**

"A human tissue polypeptide related to cancer and placenta". I. Prepar-
ation and properties. II. Assay techniques III. Clinical studies of 1.483
individuals with cancer and other conditions" in Bjorkland B. (ed.),
Immunological techniques for detection of cancer, P 133, (Bronniers,
Stockholm).

Bjorklund, B. and Bjorklund, V. (1957) :

"antigenicity of pooled human malignant and normal tissues by cytoim-
munological technique : presence of an insoluble heat - labile tumor an-
tigen".

Int. Arch. Allergy 10 : 153-84.

Braasch, J.W. and Denbo, H., E. (1964) :

"Tumors of the small intestine".

Surg. Clin. North. Am. 44 : 791-809.

Buda, F.; Aragona, P.; Giani, G.; Sisto, R.; Di-Nardo, G.; Binotto, F., Ciangemi, P.; De-Zerbi, T., Gentile, G. and Papadia, a. (1989) :

"Pretreatment evaluation of Ca 72-4 in patients with carcinoma of the stomach (I-IV stages) versus CEA, TPA, Ca 19-9".

G. Ital. Oncol. 9 : 67-72.

Carlson, R. W.; Wada, H., C. and Sussman, H., H.(1976) :

" The plasma memberane of human placenta, isolation of microvillus membrane and characterization of protein and glycoprotein subunits".

I. Biol. Chem. 251 : 4139-45.

Carter, A., C.; Goldenberg, D., M.; and Nerille, M. (1981) :

CEA: Its role as a marker in management of cancer"

Ann. Inter . Med. 94 : 402 - 9.

Chiandussi, L.; Greene, s., F. and Sherlock, S. (1962) :

"Serum alkaline phosphatase fraction in hepatobiliary and bone diseases".

Clin. Sci. 22 : 425-34.

Chin, K., M.; Morensen, R., F. and Cewurz, A.,P. (1977) :

"Interaction of alpha-I-acidglycoprotein with the immune system. Puri-

fication and effects upon lymphocytes responsiveness".

Immunology 32 : 97---102.

Ciaccino, E.; Capeni, A.; Cappetti, P.; Fanciullacci, A.; Bartoli, P. and Rerti, P. (1985) :

"Changes in carcinoembryonic antigen and tissue polypeptide antigen in monitoring the post-operative course of some tumor types".

Quad. Schivo. Diagn. 21 (2) : 199-205.

Coligan, J.E. and Todd, C., U. (1975) :

"Structural studies on carcinoembryonic antigen periodate oxidation".

Biochemistry 14 : 805-10.

Coligan, J.E.; Egan, M., L. and Guyer, R., L. (1975) :

"Structural studies on the carcinoembryonic antigen".

Ann Ny Acad Sci 259 : 355-65.

Constanza, H., E.; Saro, J, D.; Nathanson, L.; Rule, A. and Schwartz, R., S. (1973) :

"Carcinoembryonic antigen : Report of a screening study".

Cancer 30 : 583 : 90.

Coomb's, E., C. and Powles, T., J. (1982) :

"Tumor markers in the management of human cancer".

In : Dealey T., J. (ed.), Topical review in radiotherapy and oncology"
P 39. Wright P.S.G., Bristol.

Cooper, E., M. and Milford-Word, A. (1979) :

"Acute phase reactant proteins as aids to monitoring disease".

Invest. Cell Pathol. 2 : 293-96.

Cox, R., P.; Gilbert, P., J. and Griffin, M., J. (1967) :

"Alkaline phosphatase activity of mammalian cell alkaline phosphatase".

Biochem J. 105 : 155-60.

Crombach, G. and Warz, H. (1987) :

"Tumor marker system CEA-TPA" In Luthgens, M.E and Schlegel, G. (eds).

Tumor Diagnostik Leonberg, FRG.

Culter, S., J and Devesa, S., S. (1973) :

"Trends in cancer incidence and mortality in the USA "In Doll, R., Vodopija (eds) :

Host environment interaction in the etiology of cancer in man. Lyon, France, WHO. International agency for cancer research.

Davis, T., E.; Kahan, L.; Tormey, D., C.; Larson, F., C.; Crowley, J.J and Caney, R., N. (1982) :

"Clinical studies of a fast homoarginine-sensitive alkaline phosphatase in patients with cancer".

Cancer Res 41 (3) : 1110-3.

Davis, T., E.; Kahan, L.; Tormey, D.; Larson, F.; Entine, J; Metzenberg, R.; Ansfield, F. and Moon, M. (1979) :

"A fast homoarginine-sensitive alkaline phosphatase isoenzyme (FHAP) in cancer".

Proc. Am. Assoc. Cancer Res. 20 : 122-28.

De-Mello, J; Struthers, L.; Turner, R.; Cooper, E., H. and Giles, G., R. (1983) :

"Multivariate analysis as aids to diagnosis and assessment prognosis in gastrointestinal cancer".

Br. J. Surg. 48 (3) : 341-8.

Denk, H.; Tappeiner, G. and Exerstorfer, R. (1972) :

"CEA in gastrointestinal and extra gastrointestinal tumors and its relationship to tumor cell differentiation".

Int. J. Cancer 10 : 262-72.

D

P 39. Wright P.S.G., Bristol.

Cooper, E., M. and Milford-Word, A. (1979) :

"Acute phase reactant proteins as aids to monitoring disease".

Invest. Cell Pathol. 2 : 293-96.

Cox, R., P.; Gilbert, P., J. and Griffin, M., J. (1967) :

"Alkaline phosphatase activity of mammalian cell alkaline phosphatase".

Biochem J. 105 : 155-60.

Crombach, G. and Warz, H. (1987) :

"Tumor marker system CEA-TPA" In Luthgens, M.E and Schlegel, G. (eds).

Tumor Diagnostik Leonberg, FRG.

Culter, S., J and Devesa, S., S. (1973) :

"Trends in cancer incidence and mortality in the USA "In Doll, R., Vodopija (eds) :

Host environment interaction in the etiology of cancer in man. Lyon, France, WHO. International agency for cancer research.

D

avis, T., E.; Kahan, L.; Tormey, D., C.; Larson, F., C.; Crowley, J.J and Caney, R., N. (1982) :

"Clinical studies of a fast homoarginine-sensitive alkaline phosphatase in patients with cancer".

Cancer Res 41 (3) : 1110-3.

Davis, T., E.; Kahan, L.; Tormey, D.; Larson, F.; Entine, J; Metzenberg, R.; Ansfield, F. and Moon, M. (1979) :

"A fast homoarginine-sensitive alkaline phosphatase isoenzyme (FHAP) in cancer"

Proc. Am. Assoc. Cancer Res. 20 : 122-28.

De-Mello, J; Struthers, L.; Turner, R.; Cooper, E., H. and Giles, G., R. (1983) :

"Multivariate analysis as aids to diagnosis and assessment prognosis in gastrointestinal cancer".

Br. J. Surg. 48 (3) : 341-8.

Denk, H.; Tappeiner, G. and Exerstorfer, R. (1972) :

"CEA in gastrointestinal and extra gastrointestinal tumors and its relationship to tumor cell differentiation".

Int. J. Cancer 10 : 262-72.

Donegan, W.,L. and Decosse, J,J. (1978) :

"Pitfalls and Controversies in the staging of colorectal cancer". In Enker W.E. : Carcinoma of the colon and rectum. P 60, Chicago, Year Book Medical.

Dukes, C., E. and Bussey, H.,J., R. (1958) :

"The spread of rectal cancer and its effect on prognosis".
Br J Cancer 12 : 309-20.

Durdey, P., Copper, J, C.; Switala, S.; King, R.,F.; and Williams, N., S. (1985) :

"The role of peptidases in cancer of the rectum and sigmoid colon".
Br. J. Surg. 72 (3) : 378-81.

Durdey, P.; Williams, N., S. and Brown, D., A. (1984) :

"Serum carcinoembryonic antigen and acute phase reactant proteins in the preoperative detection of fixation of colorectal tumors".
Br J Surg 71 : 881-84.

Edward, D., L.; Gena, F. and Slaughter, R., L. (1982) :

"Alpha-1-acidglycoprotein concentration and protein binding in trauma".
Clin. Pharm. Therap. 31 : 63.

Ehrmeyer, S., L.; Joiner, B., L.; Kahan, L; Larson, F., C. and Metzenberg, R. (1978) :

"A cancer, associated, Fast homoarginine - sensitive electrophoretic form of serum alkaline phosphatase".

Cancer Res 38 : 599-601.

El-Asfhani, A., M., A.; Abou El-Makarem; Abddel-aziz, S.; Badawi, S.; Moustafa, M.,M.; Hammouda, E. and El-Ahmady, O. (1990) :

"Evaluation of CEA, TPA, Ferritin and CRP as tumor markers in breast cancer patients in Egypt".

2nd Int. Conference of Tumor Markers Oncology, Cairo, Egypt.

El-Asfahani, A., M., A.; Abou El-Makarem, M.; Badawy, S.A. and Essam, F., A. (1989) :

"Study of certain biochemical parameters in patients with breast cancer".

M.D. Thesis, El-Azhar Faculty of medicine, El-Azhar university.

Elliot, A. (1978) :

"The immunochemical complexity of CEA : A golden dream or molecular night mare".

Cancer 42 : 1585-88.

El- Sebai, E. (1961) :

"gastrointestinal malignancies in Egypt .

١٢
/١١

Engstrom, L. (1961) :

"I. Studies on intestinal alkaline phosphatase. II. Incorporation of inorganic phosphate into a highly purified enzyme preparation".

Biochem Biophys Acta 52 : 49 - 59.

**Fahmy, E.; Abaza, H.; El-Sawy, M.; Moussa, L.; Moussa, S.,
M. and Salim, M. (1990) :**

"Fibronectin, alpha-fetoprotein and carcinoembryonic antigen in hepatitis and malignant liver disease".

2nd Int. Conference of Tumour Marker Oncology, Cairo, Egypt.

**Falcone, F.; Sabbatani, S.; Fini, A.; Turba, E.; Magri, P. and
Ales, D. (1985) :**

"Combination of tissue polypeptide antigen (TPA) and carcinoembryonic antigen (CEA) in different types of cancer".

Nucl. Med. Commun. 6 (3) 299-304.

Fernley, H., N. and Walker, P., G. (1969) :

"Studies on alkaline phosphatase transient - state and steady - state kinetics of E. Coli alkaline phosphatase".

Biochem J 111 : 187-92.

Fishman, W., H. (1974) :

"Perspectives on alkaline phosphatase isoenzymes".

Am J Med 56 : 617-48.

Fishman, W., H. (1971) :

"Organ-Specific inhibition of human alkaline phosphatase isoenzymes of liver, bone, intestine and placenta; L-phenyl alanine, L-tryptophan and L-homoarginine".

Ehzymologia 41 : 141-50.

Fishman, W., H. and Gosh, N., K. (1967) :

"Isoenzymes of human alkaline phosphatase".

Adv Clin Chem 10 : 255 - 65.

Fletcher, R., H. (1986) :

"Carcinoembryonic antigen".

Annal of Int. Med. 104 : 66-73.

Frank, J., L.; Von Laufen, J., L., D. and Bishop, M., L. (1987) :

"Tumor Markers. In : M.L. Bishop, J.L., D., Von Laufen and Fody, E.P. (eds) Clinical Chemistry : principles, procedures, correlations.

Chap 25 p 495 J. B. Lippincott Company, Philadelphia.

Fucini, C., Tommasi, S., M.; Malatantis, G.; Cardana, G., Panichi, S. and Bettin, U. (1987) :

"Follow-up of colorectal cancer resected for cure.

An experience with CEA, TPA, Ca 19-9 analysis and second-look surgery".

Dis-Colon-Rectum 30 (4) : 272-7.

Fujimoto, S.; Kitsukawa, Y. and Itoh, K. (1979) :

"CEA in gastric juice and faeces as an aid in the diagnosis of gastrointestinal cancer".

Ann Surg 189 : 34-38.

Gabril, A. (1952) :

" Colorectal cancer " .

Ann. Surg. 30 : 250 - 60 .

Gahmberg, C., G. and Anderson, L., C. (1978) :

"Leukocyte surface origin of alpha 1-acid glycoprotein".

Am. J. Clin. Path. 66 : 331- 40.

Gamal El-Din, A.; El-Ahmady, O.; Khalifa, A. and Hamza, A. (1990) :

"Certain tumor markers in apparently helthy Egyptian individuals"

2nd Int. Conference of Tumour Markers Oncology, Cairo, Egypt.

Ganz, P., A.; Shell, W., E. and Tokes, Z.,A. (1983) :

"Evaluation of radioimmunoassay for acid glycoprotein to monitor therapy of cancer patients".

JNCI 71 (.) : 25-30.

Gion, M.; Mione, R.; Dittadi, R.; Fasan, S. and Bruscagnin, G. (1986) :

"Carcinoembryonic antigen, Ferritin and tissue polypeptide antigen in serum and tissue".

Cancer 57 : 917-22.

Gneiting, G. and Haum, A. (1990) :

"Value of CEA for second look operation decision making in colorectal cancer follow up".

2nd Int. Conference of Tumor Markers Oncology, Cairo, Egypt.

Go, V., L. and Zamcheck, N. (1982) :

"The role of tumor makers in management of colorectal cancer".

Cancer 50 : 2618 - 23 .

Go, V., L.; Ammon, H., V.; Holfermuller; K., H.; Krag, E. and Phillips, S., F. (1975) :

"Guantification of carcinoembryonic antigen - like activities in normal

human gastrointestinal secretions".

Cancer 36 : 2346-2350.

Gold, P. (1972) :

"Tumor-specific antigen in GI cancer".

Hosp. Practice 7 : 79-88.

Gold, P. and Freedman, S., O. (1965) :

:"Determination of tumor specific antigen in human colonic carcinoma by immunologic tolerance and absorption techniques".

J Exp Med 121 : 349-62.

Gosh, N., K. and Fishman, W., H. (1968) :

"Purification and properties of molecular weight variants of human placental alkaline phosphatase".

Biochem J 108 : 779-83.

Goslin, R., O.; O'Brien, M., J; and Zamcheck, N. (1981) :

" Correlation of plasma CEA and CEA tissue staining in poorly differentiated colorectal cancer " .

Am. J. Med. 71 : 246 - 50 .

Gowenlock, A., H. (1986) :

"Plasma proteins". In "Varley's Practical Clinical Biochemistry" Chapter 19 P 401-435.

Heinemann Medical Books, London.

Griffiths, J. (1989) :

"Alkaline phosphatases. Newer concepts in isoenzymes and clinical applications".

Clinics in laboratory medicine 9 (4) : 717-30.

Guarino, A. and Taccone, W. (1985) :

"Clinical application of TPA and CEA assay".

Giorn. Ital. Path. Clin. 2 : 132-40.

Hagbard, L. and Sorb, B. (1978) :

"Preliminary experiences of TPA in cancer of the ovary".

Lokortidningen 75 : 3433-42.

**Hamazoe, R.; Koga, S.; Maeta, M.; Shimizu, N.; Inoue, Y.,
Ishiguro, M. and Murakami, A. (1989) :**

"Levels of tissue polypeptide antigen in serum and the progression of gastric cancer".

J. Surg. Oncol. 41 (2) : 128-33.

**Hammarstrom, S.; Engvall, E.; Johansson, B.,G. Svensson, S.;
Sundblad, G. Goldstein, I., J. (1975) :**

"Nature of the tumor-associated determinant (s) of carcinoembryonic antigen".

Proc. Nat. Acad. Sci. USA 72 : 1528-32.

**Hansen, H., J; Synder, J.J; Miller, E.; Vandevoorde, J, P.;
Miller, O., N., Hinus, L., R. and Burns, J., J. (1974) :**

"CEA assay : a laboratory adjunct in the diagnosis and management of cancer".

Human Path. 5 : 139-47.

Hansen, H., J; Lance, K., P. and Krupey, J. (1971) :

"Determination of an iron sensitive site on carcinoembryonic antigen using zirconyl phosphate".

Clin. Res. 19 : 143 - 55.

Harkness, D., R. (1968) :

"Studies on human placental alkaline phosphatase. II. Kinetic properties and studies on the apoenzyme".

Arch Biochem Biophys 126 : 513.

**Harmenberg, U; Frodin, J, E.; Ljungdahl, S., E.; Mellstedt, H.
and Stigbrand, T. (1989) :**

"Significance of ALP isoenzymes in the monitoring of patients with colorectal carcinoma".

Tumour-Etiol. 10 (5) : 225-31.

Harold,J; Wanebo, A.; Maus,S. and Schwartz,M.,K.(1978):

"Use of CEA as indicator of early recurrence and as a guide to a select-

ed 2nd look procedure in patients with colorectal cancer".

Ann. Surg. Oct. 481 : 93.

Herbsman, H.; Wetstein, L. and et al., (1980) :

"Tumor of the small intestine".

Surgery 17 : 131-184.

Herrera, M., A.; chu, T., M. and Holyoke, E. (1976) :

"CEA as a prognostic and monitoring test in clinically complete resection of colorectal carcinoma".

Ann. Surg. 183 : 5-9.

Higashino, K.; Hashinostume, M.; Kang, K., Y; Yakahashi, Y. and Yamamura, Y. (1972) :

"Studies on a variant alkaline phosphatase in sera of patients with hepatocellular carcinoma".

Clin. Chim. Acta. 40 : 67 - 77.

Hill, A., E. (1979) :

"A short text book of medical statistics". Hill, A., B. (ed) PP 86-96, 132. Published by the English Language Book Society, London, U.K.

Holland, F., J. and Emil, F. (1982) :

"Carcinoembryonic antigen".

Cancer Medicine, 4 : 15 - 20 .

Holyoke, E., D.; Chu, M. and Murphy, G., P. (1975) :

"CEA as a monitor of gastrointestinal malignancy".

Cancer 35 : 830-36.

Holyoke, E., D.; Reynoso, G. and Chu, T. (1972) :

"Carcinoembryonic antigen in patients with carcinoma of digestive tract". Proceeding of second conference on embryonic fetal antigens in cancer.

National Technical Information Service, USA Departement of commerce, Springfield 215-219.

Ikenaka, T.; Ishiguro, M.; Isemura, S.; Kaufmann, H.; Bauen, W. and Schmid, K. (1971) :

"Biochemical properties of human glycoprotein".

Bioch. Biophys. Res. Commun. 142 : 1142 - 52.

Inglis, N., R.; Kirley, S., Stolbach, L., L. and Fishman, W., H. (1973) :

"Phenotypes of the regan isoenzyme and identity between the placental D-variant and the Nagao isoenzyme".

Cancer Res. 33 : 1657 - 67.

Iwatomo, I.; Furukawa, M.; Ito, S.; Tanigawa, M.; Koga, Y. and Mogari, Y. (1988) :

"Carcinoembryonic antigen, alpha-fetoprotein and hepatoma alkaline

phosphatase in gastric carcinoma".

Gan. No. Kinsho. 34 (2) : 208-12.

Jacoby, B. and Bagshawe, K., D. (1972) :

"Placental-type alkaline phosphatase from human tumor tissue".

Clin. Chim. Acta. 35 : 473.

Janssen, C., W. (1987) :

"Immunoglobulins, complement and carcinoembryonic antigen in patients with gastric carcinoma".

Acta. Chir. Scand (Supp) 535 : 1-60.

Jeppson, A.; Wahren, B.; Stiglvard, T.; Edsmyr, F.; T. and Anderson, L. (1983) :

"A clinical evaluation of serum placental alkaline phosphatase in seminoma patient.

Eur. J. Surg. Oncol. 13 (4) : 285-95.

Johnson, R., B.(1969) :

"A new fluorometric method for the estimation or detection of total and fractionated alkaline phosphatase.

Clin. chem. 15 : 108-123.

Johnson, A., M.; Schmid, K. and Alper, C., A. (1969) :

"Inheritance of human AAG variants".

J. Clin. Invest. 48 : 2293-98.

Jubert, A., V.; Talbott, T., M. and Maycorft, A. (1978) :

"Characteristics of adenocarcinomas of the colorectum with low levels of preoperative plasma CEA".

Cancer 42 : 623-63.

Kanauchi, M.; Sawai, F.; Hirayama, T.; Yumura, F.; Kanauchi, K.; Hamaguchi, T.; Hara, Y.; Yoshimura, K.; Nishiura, K. and Dohi, K. (1985) :

"Clinical studies on tissue polypeptide antigen (TPA) in gastric cancer".

Gan - No - Rinsho 13 (11) : 1367-71.

Khalifa, A. (1991) :

"Tumor Markers In : Applied Biochemistry and Clinical Chemical Pathology" Khalifa, A. (ed.) Published by Scientific Book Center, Cairo, Egypt.

Khalifa, A. ; El-Ahmady, O, Hamza, R., and Salah , N (1990) :

" Tumor markers in cancer lung" .

M. D. Thesis Ain Shams Faculty of Medicine .

Khoo, S., K. and Mackay, E., V. (1973) :

"Carcinoembryonic antigen in cancer of the female reproductive sys-

tem : Its detection in whole serum by microradioimmunoassay".

J. Clin. Path 26 : 470-75.

Klavins, J. V. and Birkmayer, G., D. (1990) :

"Classification of Tumor Markers".

2nd Int. Conference of Tumor Markers Oncology, Cairo, Egypt.

Kleinman, M., S.; Turner, M., D. and Thayer, W. (1972) :

"Serum CEA in patients with inflammatory bowel disease".

Gastroenterology 62 : 772 - 82.

Kralavanszky, J; Szentirmay, Z.; Deszyak, I. and Eckhardt, S. (1984) :

"Placental type alkaline phosphatase in possibly premalignant alterations of human gastric mucosa".

Oncology 41 (3) : 189-94.

Kupchik, H., Z. and Zamcheck, N. (1972) :

"Carcinoembryonic antigen (CEA) in liver diseases II. Isolation from human cirrhotic liver and serum, and from normal liver".

Gastroenterology 63 : 95-101.

Lazdunski, M.; Petitelerc, C.; Chappelet, D. and Lazdunski, C. (1971) :

"Flip-Flop mechanism in enzymology. A model : the alkaline phosphatase".

tase of E. Coli.

Eur J Biochem 20 : 124-34.

Lewis, H; Blumgart, L., Carter, D., C.; Gillis, C., R., Hole, D.; Ratcliffe, Wood, C., B. and Mc Ardle, C., S. (1984) :

"Preoperative carcinoembryonic antigen and survival in patients with colorectal cancer".

Br. J. Surg. 71 : 206-8.

Loewenstein, M., S. and Zamcheck, N. (1978) :

"Carcinoembryonic antigen (CEA) levels in benign gastrointestinal disease states".

Cancer 42 : 1412-1418.

Loewenstein, M., S. and Zamcheck, N. (1977) :

"Carcinoembryonic antigen and the liver".

Gastroenterology 72 : 161-166.

LoGerfo, P. and Herter, F., P. (1975) :

"Carcinoembryonic antigen and prognosis in patients with colon cancer".

Ann. Surg. 181 : 81-84.

LoGerfo, P.; Krupey, J. and Hansen, H. (1971) :

"Demonstration of an antigen common to several varieties of neo-

plasia".

N. Eng. J. Med. 285 : 138-41.

Lowenfil, A., B. (1973) :

"Why are small bowel tumors are rare?".

Lancet 1 : 24-25.

Lundstrom, R.; Bjorklund, B. and Eklund, G. (1973) :

"A tissue-derived polypeptide antigen. Its relation to cancer and its temporary occurrence in certain infectious disease. In : Bjorklund, B. (ed.), Immunological techniques for detection of cancer".

P 243 (Bonniers, Stockholm).

Luning, B.; Wikland, B.; Redelius, P. and Bjorklund, B. (1981) :

"Biochemical properties of tissue polypeptide antigen".

Biochem. Biophys. Acta 3 (2) : 65.

Lurie, B., B.; Loewenstein, M., S. and Zamcheck, N. (1975) :

"Elevated circulating CEA levels in benign extrahepatic biliary tract obstruction and inflammation".

JAMA 233 : 326-330.

Luthgens, M. and Schlegel, G. (1987) :

"The clinical value of tissue polypeptide antigen".

Journal of Tumor Marker Oncology 2 (4) : 261.

Luthgens, M. and Schlegel, G. (1984) :

Evaluation of TPA, CEA in tumors of head and neck.

Verh. Dtsch. Kretsges 5 : 271-85.

Mahdy, Z; Khalifa A., Hassan, A. and Tonsy, H. (1990) :

" Tumor markers in breast cancer " .

M. D. Thesis, Benha Faculty of Medicine .

**Martin, E., M.; James, K., K.; Hurtubise, P., E.; Cataland, P.
and Minton, J, P. (1977) :**

"The use of CEA as an early indicator for gastrointestinal tumor recurrence and second-look procedure".

Cancer 39 : 440-46.

Mattsson, W., and Borgstrom, S. (1975) :

"Tissue polypeptide antigen as a guide in antineoplastic treatment of advanced mammary carcinoma".

Folksam symp. TPA (Folksam, Stockholm).

Meeker, W., R. and Kashniri, R., L. (1973) :

"Clinical evaluation of carcinoembryonic antigen test".

Arch. Surg. 107 : 266-74.

Menendez-Etort, M., C.; Oettgen, H., F.; Pinsky, C., M. and Schwartz, M., K. (1978) :

"A preliminary evaluation of tissue polypeptide antigen in serum or urine of patients with cancer or benign neoplasms".

Clin. Chem. 24 : 848-72.

Mercer, D., W. and Talamo, T., S. (1985) :

"Multiple markers of malignancy in sera of patients with colorectal carcinoma : preliminary clinical studies".

Clin. Chem. 31 (11) : 1824-38.

Millan, J., L. and Manes, T. (1988) :

"Seminoma-derived Nagao isoenzyme is encoded by a unique alkaline phosphatase gene".

Proc. Natl. Acad. Sci 85 : 3024-45.

Montinari, F.; Beretta, G., D.; Pancera, G. and Petrini, C. (1989) :

"CEA, TFA and GICA during long-term follow-up and therapy monitoring of gastrointestinal tumors".

J. Tumor Marker Oncology 3 (2) : 201-20.

Moore, T.; Dhar, P.; Zamcheck, N.; Keeley, A.; Gottlieb, L. and Kupchik, H., 2. (1971) :

"Carcinoembryonic antigen (S) in liver disease. Clinical and morpho-

ligical studies".

Gastroenterology 63 : 88-94.

Moore, T.; Dhar, P.; Zamcheck, N.; L. and Kupchik, H., Z. (1972):

"CEA in benign conditions"

Gastroenterology 73 : 120 - 5.

Moss, D., W. (1982) :

"Alkaline phosphatase isoenzymes".

Clin. Chem. 28 : 2007.

Munck-Wikland, E.; Kuylensstierna, R., Lindholm, J. and Wahren, B. (1990) :

"CEA, Ca 19-9 and Ca 50 in monitoring human squamous cell carcinoma of the oesophagus".

2nd Int. conference of Tumour Markers Oncology, Cairo, Egypt.

Nakayama, T., Yoshida, M. and Kitmura, M. (1970) :

"L-leucine sensitive heat-stable alkaline phosphatase isoenzyme detected in patients with pleuritis carcinomatosa".

Clin. Chem. Acta 30 : 546-60.

Nishida, O.; Shirato, H.; Satoh, N.; Nakajima, F.; Manabe, K.; Kondoh, Y.; Sano, F. and Uchino, J. (1988) :

"Clinical evaluation of a combination of assay of CEA, Ca 19-9 and

TPA in patients with colorectal cancer".

Gan-No-Rinsho 34 (9) : 1096-100.

Nugent, F., W. and Hansen, E., R. (1971) :

"Radioimmunoassay of carcinoembryonic antigen as a diagnostic test for cancer of colon".

Lahey Clin. Found. Bull. 20 : 85-88.

Oehr, P.; Schwabe, H.; Loos, U. and Utzat, H. (1989) :

"NSE, TPA and CEA compared to conventional clinical diagnostic methods for patients with small cell cancer of the lung". Proceeding of the 6th International conference of human tumor markers.

May 23-25, Tokyo, Japan.

Omar, S.; Elisa, S. and Nagieb, S. (1989) :

"Studies on the aetiological factors of stomach cancer in Egypt".

A thesis submitted for partial fulfillment of the M.D. degree in surgery, National Cancer Institute, Cairo University.

Peter et al., 1977.

Peter, A, Oehr, B. and Loose, U. (1977) :

"Significance of serum proteins in invasive bladder cancer " .

Eur. J. cancer 17 : 251 - 60 .

Petit, J; Poupon, R.; Vasseur, J.; Kelemen, N.; Milhaud, G. and Darnis, F. (1980):

"Mise en evidence et valeur diagnostique d'une isoenzyme anormale des phosphatases alcaline au cours des cancers du foie".

Gastroenterol. Clin. Biol. 4 : 86-95.

Piafsky, K., M.; Borga, O.; Odar, C., L.; Johansson, C. and Sjoquist, F. (1978) :

"Increased plasma protein binding to propranolol and chlorpromazine mediated by disease-induced elevation of plasma acid glycoprotein".

N. Eng. J. Med. 299 : 435-42.

Piantino, F.; Furaso, A.; Pecchio, F.; Arosio, E. and Rapellino, M. (1979) :

"TPA in gastrointestinal diseases". Proceeding of the 6th Int. Conference on human tumor markers. May 23-26, Tokyo, Japan.

Piantino, F.; Tappero, G.; Cerchier, A.; Pecchio, F.; Tridico, F. and Cattaneo, U. (1986) :

"Tissue polypeptide antigen as a tumor marker in disorders of the digestive tract : Comparison with CEA".

Minerva Med. 77 (16) : 617-20.

Plow, E., F. and Edgington, T., S. (1975) :

"Isolation and characterization of a homogenous isometric species of carcinoembryonic antigen".

Int. J. Cancer 15 : 748-61.

Primus, F.,J; Goldenberg, D.,M. and Hansen, H.,J.(1973) :

"Metabolism of carcinoembryonic antigen (CEA) in a human tumor-hamster host model".

Fed. Proc. 32 : 834-45.

Pritchard, D. and Todd, C. (1979) :

"Purification of carcinoembryonic antigen by removal of contaminating mucopolysaccharides".

Cancer Res. 36 : 4699-4701.

Puntum, F., W. (1975) :

"The role of plasma proteins. In the plasma protein structures, function and genetic control".

2nd edition, Vol. 1 : 58-125.

Putzki, H.; Ledwoch, J., Student, A.; Jablonski, M. and Heymann, H. (1988) :

"Tumor markers : carcinoembryonic antigen, tissue polypeptide anti-

gen and carbohydrate antigen 19-9 in liver diseases".

J. Surg. Oncol. 37 (2) : 133-5.

Rashid, S., A.; Quigley, O.; Axon, A.,T. and Cooper, E., H.
(1982) :

"Plasma proteins and prognosis in gastric cancer"

Br. j. Cancer 45 (3) : 390 - 4 .

Redelius, F. (1980) :

"Studies, on the primary structure of TPA"

Chem. Commun. Unir., Stockholm, No. 2.

Reiners, C., H. (1982) :

"Clinical application of TPA and CEA assay in different tumors "

Tumor Diagnostik 2 : 109 - / 5 .

Ross, M., H.; and Archer, j,G. (1955) :

"Alkaline phosphatase activity and pH optimun" .

J Biol Chem 192 : 561 - 80 .

Roy, S., K. (1990) :

" Upper gastrointestinal tract malignencies in jubail, kingdom of Saudia Arabia" .

2nd Int. Conference of Tumor Markers Oncology, Cairo, Egypt .

Samak, R.; Edelstein, R and Israel, R. (1982) :

"Immunosuppressive effect of acute phase reactant proteins in vitro and its relevance to cancer " .

Cancer Immunol. Immunoth. 12:38 - 48 .

Sarcioni, E., H. (1963) :

"Clinical and immunological studies on the products of human plasma fractionation" .

Arch. Biochem. Biophys. 100 : 516 - 25 .

Sasaki, M., and Fishman, W., H. (1973) :

"Ultrastructural studies on Regan and non - Regan isoenzymes of alkaline phosphatase in human ovarian cancer cells .

Cancer Res. 33 : 3008 - 20 .

Schmidt, K.; Kaugman, H.; jsemura, S.; Bauer, F., Enuta, J.; Motoyama, T., Ishiguro, M. and Nanno, S . (1973) :

"Determination of human serum glycoprotein"

Biochemistry 12 : 2711 - 18 .

Shaarawy, M. and Omar, S. (1988) :

"The diagnostic values of TPA, CEA, B2- microglobulin in breast can-

cer " .

Proceeding of 1st Int. Conference of Tumor Marker, Oncology. Cairo, Egypt .

Sharma, M., P., Gregg, j, A. and Mc. Cobe, R., P. (1974) :

"CEA - Live activity in pancreatic juice of patients with pancreatic carcinoma and pancreatitis "

Gastroenterology . 66 : 776 - 82 .

Sharp, H., L. (1972) :

"The current status of antitrypsin in a protease inhibitor in gastrointestinal diseases "

Gastroenterology 4 : 611 - 631 .

Sherif. M., Tawfik, N. and Badawi, S. (1989) :

"Gastrointestinal malignancies in Egyptian patients. A Clinicopathological study with a special reference to carcinoembryonic antigen"

M. D. Thesis, National Cancer Institute, Cairo University .

Shibata, , G., Anderson, A. and et al. (1977) :

"Plasma proteins : Chemistry and metabolism"

Clin. Chera. Acta 50 : 40 - 50 .

Shimano, T.; Okuda, H.; Moden, T.; Inaji, H. and Mari; H. (1987) :

"Usefulness of carcinoembryonic antigen measurement in faeces of patients with colorectal cancer" .

Dis - Colon - Rectum 30 (8) : 607 - 10 .

Shimizu, N.; Wakatsuki, T.; Murakami, A.; Yoshioka, H.; Hamazoe, R.; Kanayama, H.; Maeta, M. and Koga, S. (1987) :

"Carcinoembryonic antigen in gastric cancer patients"

Oncology 44 (4) : 240 - 4 .

Shuster, j; Silverman, M. and Gold, P. (1973):

"Metabolism of human carcinoembryonic antigen in xenogenic animals"

Cancer Res 33 : 65 - 68 .

Skinner, j, M. and Whitehead, P. (1981):

"Carcinoplacental alkaline phosphatase in malignant and premalignant conditions of human digestive tract"

Path - Anat. 394 : 109 - 18 .

Staab, H., j; Ahlemann, L., M.; Anderer, F., A.; Zwirner, M. and Schindler, A., E. (1985) :

"Optimizing tumor markers in breast cancer, monitoring, prognosis

patocellular, bile duct and pancreatic carcinoma" .

Gan. No. Rinsho. 31 (6swpp.) : 623 - 30 .

Talamo, R., C. (1975) :

"Basic and clinical aspects of AAT"

Pediatrics 56 (1) : 91 - 95 .

Thibeault, D., W.; Lee, S., K. and Heiner, D., C. (1978) :

"AAT and AAG levels in cord blood and amniotic fluid of infant with respiratory distress syndrome "

pediatric Res. 12 : 775 - 80 .

Thomas, P. and Hems, D., A. (1975) :

"The hepatic clearance of circulating native and a sialo carcinoem-tryonic antigen by the rat"

Biochem. Biophys. Res. Comm. 67 : 1205 - 9 .

Tietz, N., W. (1987) :

In " Fundamentals of clinical chemistry"

Chapter 3 P : 330, W. B. Saunders, Company, Philadilphia .

Torriani, A. : (1968) :

"Alkaline phosphatase subunits and their dimerization in vivo"

J . Bacter. 96 : 1200 - 10

Turnbull, R., B.; Kgle, K.; Watso, F.,R. and et al.(1967) :

"Cancer of the colon. The influence of the no - touch isolation tectnique on survival rates "

Ann, Surg. 166 : 420 - 27 .

Twining, S., S. and Brecher, A., S. (1977) :

"Identification of AAG, alpha 2-MG and antithrombin III as component of normal and malignant human tissues " .

Clin. Chem. Acta. 76 : 143 - 50 .

Van Andel, j, G.; Dees, j. and et al. (1979) :

"Carcinoma of the oseophagus. Results of treatment" .

Ann. Surg. 190 : 684 - 93 .

Van - Eden, P.,J., Bezuidenhout, D.,J; Kock, j ; Weidemann, A; Rossouw, M.,A. and Carthy , N.,A. (1987) :

" Gastric juice carcinoembryonic antigen etimation .

Avseful addition test in the diagnosis of gastric carcinoma" .

S. Afr. Med. j. 71 (4) : 241 - 3 .

Vayliang, G., O. and Zamcheck, N. (1982) :

"The role of tumor markers in the management of colorectal cancer" .

Cancer 50 : 2618 - 23 .

**Viot, M.; Thyss, A.; Schneider, M.; Ramaiali, A.; Cambon, P.
and Lalanne, M., C. (1983) :**

" α 1 isoenzyme of alkaline phosphatases"

Cancer 52 : 140 - 45 .

Warnes, T., W. (1972) :

"Alkaline Phosphatase "

Gut 13 : 926 - 40 .

Warnock, M., L. and Reisman, R., (1969) :

"Variant alkaline phosphatases in human hepatocellular cancers " .

Clin. Chim. Acta. 24 : 5 - 10 .

**Weber, K.; Osborn, M.; Moll, R.; Wiklund, B. and Luning, B.
(1984) :**

**"TPA is related to the non - epidermal Keratins 8, 18 and 19. Re -
evaluation of a human tumor markers"**

EMBOJ 3 : 2707 - 14 .

Wick, M., R.; Swanson, P., E. and Manivel, j, C. (1987) :

"Placental - like alkaline phosphatase reactivity in human tumors "

Human Pathol 18 : 946 - 52 .

Wiernik, P., H. and Serpick, A., A. (1973) :

"Plasma carcinoembryonic antigen in patients with advanced cancer " .

Proc. Am. Assoc. Cancer Res 14 : 14 - 20 .

Wiklund, B. (1980) :

"Studies on TPA and its antigenic determinant "

Chem. Comm. Univ., Stocholm, No. 1 ; Diss. Stokholm .

**Winter, S.; Schmidt, R., P., Schulz, K., D. and Sturm, G.
(1985) :**

"The prognostic significance of consecutive determination of serum
TPA in human breast cancer "

Verh. Dtsch. Krebses 5 : 552 - 62 .

**Wolf, G., T.; Chretien, P., B., Elias, E., G.; Baskias, A., M.;
and weiss, J.F. (1979) :**

" Serum glycoprotein in heat & neck squamous carcinoma "

Am. J. Surg . 138 : 489 - 500 .

**Wood, C., B.; Ratcliffe, j, G.; Burt, R., W.; Malcolm, j, H.
and Blumgart, L., H. (1980) :**

" The clinical significance of the pattern of elevated CEA levels in re-
current colorectal cancer " .

Br j Surg 67 : 46 - 48 .

Wychulis, A., R.; Woolam, G., L.; Anderson, H., A. and Ellis, F., H. (1971) :

"Achalasia and carcinoma of oesophagus"

JAMA 215 : 1638 - 48 .

Zamcheck, N.; Moore, T., L.; Dhar, P. and Kupchick, T., L. (1972) :

"Immunologic diagnosis and prognosis of human digestive tract cancers"

New Eng. J. Med. 13 : 83 - 87 .