REFERENCES

- Abbott RD, Donahue RP, Kannel WB, and Wilson PWF (1988):

 The impact of diabetes on survival following myocardial infarction in men vs women: the Framingham study.

 JAMA. 260:3456-3460.
- Airaksinen KEJ, Heimo MJ, Linnaluoto M, et al. (1998) Gender difference in autonomic and hemodynamic reactions to abrupt coronary occlusion. J Am Coll Cardiol. 31:301-306.
- Alexander AP, Shaw LJ, DeLong ER, et al. (1998) Value of exercise treadmil testing in women. J Am Coll Cardiol. 32 L1657-64.
- Ali AS, Rybicki BA, Alam M, et al (1999): Clinical predictors of heart failure in patients with first acute myocardial infarction. Am Heart J. 138: 1133-1139.
- Andreotti F; Conti E, Sanna T. Crea F, Zecchi P, and Maseri A

 (1999). Acute coronary syndromes in women:
 pathophysiology and therapentic options. Cardiologia.

 Dec; 44 Suppl (pt2): 5114.
- Antman EM, McCabe CH, Gurfinkel EP, et al (1999). Enoxaparin prevents death and cardiac ischemic events in unstable angina/non-Q- wave myocardial infarction. results of the thrombolysis in myocardial infarction(TIMI). IIB trial. Circulation . 100: 1593-601.

- Antman EM, Guigliano RP, Gibson CM, et al (1999). Abciximab facilitates the rate and extent of thrombolysis: results of the Thrombolysis In Myocardial Infarction (TIMI) 14 trial. Circulation. 99: 2720 –2732.
- Ball D and Bowman L (1999). Underutilization of abciximab and coronary stents in women, minorities, and the elderly undergoing coronary angioplasty. Paper presented at:

 American Public Helath Association Annual Meeting; November; Chicago, 111.
- Barrett-Conner E, Grady D, Sashegyi A et al (2002). Raloxifene and cardiovascular events in osteoporotic postmenopasal women four year results from the MORE (Multiple Outcomes of Raloxifene Evaluation) randomized trial. JAMA . 287:847-57.
- Becker RC, Terrin M, Ross R, et al (1994). Comparison of clinical outcomes for women and men after acute myocardial infarction. Ann Intern Med. 120:638-645.
- Bell MR, Grill DE, Garratt KN, et al (1995). Long- term outcome of women compared with men after successful coronary angioplasty. Circulation. 91:2876-2881.
- Berkowitz SD, Granger CB, PieperKS, et al (1997). Incidence and predictors of bleeding after contemporary thrombolytic therapy for myocardial infarction. Circulation .95:2508-2516.

- Birdwell BG. Herbers JE, and Kroenke K (1993). Evaluating chest pain. Arch Intern Med. 153: 1991.
- Bodi vv, Facila L, Sanchis J, Liacer A, Ferrero JA, and chorro FJ(2003). Gender differences in patients with suspected Non St- segment Elevation Acute coronary Syndromes. Implications for Invasive management. Rev ESP cardiol; Apr: 56 (4): 412-416.
- Bonow RO(1999). Cardiovascular Disease in Women: Viva la difference ? XXIst Congress of the European society of Cardiology.
- Boresma E, Peiperks, Steyeberg EW, et al (2000). Predictors of outcome in patients with acute coronary syndromes without persistent ST- segment elevation: results from an international trial of 9461 patients. Circulation. 101: 2557-67.
- Bosch x, Casanovas N, Miranda- Guardiola F, Diez Ajas, Sitges M, Anguera I, Sanz G, and Betriu A(2002): Long Term prognosis of Women with Non- ST segment Elevation Acute Coronary Syndromes, a case contral study. Rev ESP Cardiol. Dec: 55 (12): 1235- 42.

- Braunwald E, Antman EM, Beasley JW, et al (2000). ACC/ AHA
 Guideline for the management of patients with Unstable
 Angina .A Report of the American college of
 cardiology/ American Heart Association Task Force on
 Practice Guidelines (Committee on the Management of
 patients with Unstable Angina). J Am Coll Cardiol. 36:
 970-1062.
- Bueno H, Vidan MT, Almazan A, Lopez-SendonJL, and Delcan JL (1995). Influence of sex on the short term outcome of elderly patients with a first acute myocardial infarction. Circulation.;92:1133.
- Bugiardini R, Pozzati A, Ottani F, et al (1993): Vasotonic angina:

 A spectrum of ischemic syndromes involving functional abnormalities of the epicardial and microvascular coronary circulation. J Am Cardiol. 220: 417.
- Burt VL, Whelton P, Roccella EJ, et al (1995). Prevalence of hypertension in the US adult population. Results from the Third National Health and Nutrition Examination Survey, 1988-1991. Hypertension. 25: 305.
- Cannon RO and Epstein SE (1988). "Microvascular angina" as a cause of chest pain wit angiographically normally coronary arteries. Am J Cardiol.; 61: 1338.
- Cannon RO, Camici PG, and Epstein SE(1992). Pathophysiological dilemma of syndrome X. Circulation.; 85:883.

- Cannon RO (1994). Imipramine in patients with chest pain despite normal coronary angiograms. N Engl J Med. 330:1411.
- Cannon RO (1995). The sensitive heart. A syndrome of abnormal cardiac pain perception JAMA. 273:883.
- Castelli WP (1984). Epidemiology of coronary heart disease: the Framingham study. Am J Med. 76:4-12.
- Chardra NC, Ziegelstein RC, Rogers WJ, et al (1998):

 Observations of the treatment of women in the United

 States with myocardial infarction. Arch Intern

 Med.158: 981.
- Chasan-Taberl, and Stampfer MJ (1998): Epidemiology of oral contraceptives and cardiovascular disease. Ann Intern Med. 128: 467.
- Christina C, Demosthenes B. panagiotakos, Christos p, kokkinos P, et al (2003). Gender Differences on the risk Evaluation of Acute coronary syndromes: the CARDIO 2000 study. Preventive cardiology. spring: 6 (2): 71-7.
- Chua TP, Saia F, Bhardwaj V, et al (2000). Are there gender differences in patients presenting with unstable angina?

 Intern J Cariol. 72:281.
- Clarke KW, Gray D, Keating NA, and Hampton JR (1994). Do women with acute mycoardial infarction receive the same treatment as men? BMJ. 309: 563-566.

- Colditz GA, and Coakley E(1997). Weight, weight gain, activity, and major illnesses: the Nurses' Health Study. Int J Sports Med 18 (Suppl 3): \$162-70.
- Collins, L.J.; and Douglas P.S (1999): Acute coronary syndromes.

 In charney P. (ed): coronary artery disease in women: prevention, diagnosis and management. Philadelphia, American college of physicians. P. 401.
- Compell MJ, and Machin D (1993). Medical statistics: a common source approach. 2nd edition. John wiley and Sonf chiechester, New york, pris bone, Toronta and thing apore.
- Cotter G, Cannon CP, Mc Cabe CH, et al., (2003). Prior peripheral arterial disease and cerebrovascular disease are independent predictors of adverse outcome in patients with acute coronary syndromes: are we doing enough? Results from the Orofiban in Patients with Unstable Coronary Syndromes— Thrombolysis IM Myocardial Infarction (OPUS-TIMI) 16 study. Am Heart J. 145 (4): 622-7.
- Cunningham MA, Lee TH, Cook EF, et al (1989). The effect of gender on the probability of myocardial infarction among emergency department patients with acute chest pain: a report from the Multicenter Chest Pain Study Group. J Gen Intern Med. 4: 392-398.

- Cushman M, Legault C, Barrett-Connor E, Stefanick ML, Kessler C, Judd HL, et al (1999). effect of postmenopausal hormones on inflammation senstitive proteins: the Postmenopausal Estrogen/Progestin Interventions (PEPI) Study. Circulation. 100: 717-22.
- Davis KB, Chaitman B, Ryan T, Bittner V, and Kennedy JW (1995). Comparison of 15-year survival for men and women after initial medical or surgical treatment for coronary artery disease: a CASS Registry study: Coronary Artery Surgery study. J Am Coll Cardiol. 25:1000-1009.
- Davies MJ (2000): The pathophysiology of acute coronary syndromes. Heart. 83:361-6.
- Davar JI, Brill Dj, Bulugahipitiya S, et al (1999). Prognostic value of negative dobutamine stress echo in women with intermediate probability of coronary artery disease. Am J Cardiol. 83: 100.
- De Bacquer D, De Backer G, Kornitzer M, et al(1998). Prognostic value of ischaemic electrocardiographic findings for cardiovascular mortality in men and women. J Am Coll Cardiol. 32:680.

- De Sanctis RW(1993). Clinical manifestations of coronary artery disease: chest pain in women. In: Cardiovascular Health and Disease in Women, Wenger NK, Speroff L, Packard B. Eds), Le Jacq Communications, Connecticut. p. 67.
- Diabetes Control and Complications Trial Research Group (1995).

 Effect of intensive diabetes management on macrovascular events and risk factors in diabetes control and complications trial, Am J cardiol 75: 894.
- Dittrich H, Gilpin E, Nicod P, Cali G, Henning H, and Ross J Jr (1988). Acute myocardial infarction in women: influence of gender on mortality and prognostic variables. Am J Cardiol. 62:1-7.
- Donahue RP, Goldberg RJ, Chen Z, Gore JM, and Alpert JS (1992). The influence of sex and diabetes mellitus on survival following acute myocardial infarction: a community-wide perspective. J Clin Epidemiol. 46:245-252.
- Douglas PS and Ginsburg GS(1996). The evaluation of chest pain in women. N Eng J Med. 334: 1311.
- Dunn N, Thorogood M, Faragher B, et al (1999). Oral contraceptives and myocardial infarction: Results of the MICA case control study. BMJ. 318: 1579.

- Elhendy A, van Domburg RT, Bax JJ, et al (1999). Gender differences in the relation between ST-T-wave abnormalities at baseline electrocardiogram and stress myocardial perfusion abnormalities in patients with suspected coronary artery disease Am J cardiol. 84. 865.
- Elliot PM, Kryzowska DK, Calvino R. et al (1997): Effect of oral aminophylline in patients with angina and normal coronary arteriogram (cardiac syndrome X). heart. 77:523.
- Eriksson BE, Jansson E, Kaijser L, and Sylven C (1999). Impaired exercise performance but normal skeletal muscle characteristics in female syndrome X patients. Am J Cardiol. 48(2):176-80.
- Ever DG and David RR (2003). Acute coronary syndrome: unstable angina and non- ST segment elevation myocardial infarction: ABC of interventional cardiology. BMJ. 326; 1259-1261.
- Feldman R (2000). ACE inhibitors versus ATI blockers in the treatment of hypertension and syndrome X. cand J Cariol. 16 suppl E: 41 E- 44E.
- Feldman T, and Silver R (2000). Gender differences and outcome of interventions for acute coronary syndromes. Caridol Rev. Jul-Aug: 9 (4) -240-7.

- Fetters JK, peterson ED, Shaw LJ, et al (1996): Sex specific differences in coronary artery disease risk factors, evaluation, and treatment: Have they been adequately evaluated Am Heart J. 131:796.
- Florence AT, Jean- Michel P, Collette C, and Claude R (1999).

 Coronary Heart Disease Risk Factors and Menopause: a study in 1684 French women. Atherosclerosis. 142:415.
- Fox KA (2000). Acute coronary syndromes: presentation, clinical spectrum and management Heart. 84:93.
- Fox KA, Poole- Wilson PA, Henderson R A, et al (2002). The Randomized Intervention Trial of unstable Angina (RITA) investigators. Interventional versus conservative treatment for patients with unstable angina or non- ST-elevation myocardial infarction: the British Heart Foundation RITA 3 randomized trial. Lancet. 360 (9335): 743.
- Fragasso g, Chierchia SL, Lu C, Dabrowski P, Pognotta P, and Rosano GM (1999). Left ventricula dysfumction during dobutamine stres echocardiography in patients with syndrome X and positive myocardial perfusion scintigraphy. G Ital Cardiol.29 (4): 383-90.
- Glaser R, Herrmann HC, Murphy SA, et al(2002). Benefit of an early invasive Management strategy in women with Acute coronary syndromes. JAMA. 288:3124-3129.

- Gillum RF (1989). Sudden coronary death in the United States 1980-1985. Circulation. 79:756.
- wide perspective of sex differences and temporal trends in the incidence and survival rates after acute myocardial infarction and out of hospital deaths caused by coronary heart disease. Circulation. 87:1947-1953.
- Goldschmidt- clermont PJ, Schulman Sp, Bray PF, et al (1996).

 Refining the treatment of women with unstable angina- a randomized, double-blind, comparative safety and efficacy evaluation of integralin versus aspirin in the management of unstable angina. Clin cardiol. 19: 869-874.
- Gottliebs, Harpaz D, Shotan A, Baykov, Leorj, Cohen M, et al. (2000), sex differences in management and outcome after acute myocardial infarction in the 1990s circulation. 102: 2484-2490.
- Greenland P, Reicher- Reiss H, golbourt U, and Behar S (1991). In

 -hospital and 1-year mortality in 1524 women after

 myocardial infarction: comparison with 4315men.

 Circulation. 83:484-491.
- Grover- McKay M (1996). gender related imaging issues in assessment of coronary artery disease by nuclear techniques. Am J Card Imaging. 10:54

- Grundy SM (1994). Second Report of the Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel II). Circulation. 89:1329.
- Hanes DS, Weir MR, and Sowers JR (1996). Gender considerations in hypertension pathophysiology and treatment. Am J Med. 101 (Suppl 3A): 10 S.
- Harpaz D,. Benderyl M, Gold bourt U, et al (1996): Effect of aspirin on mortality in women with symptomatic or silent myocardial ischemia. Am J cardiol. 78: 1215.
- Hayashi T, Yamada K, Esaki T, et al (1995). Estrogen increases endothelial nitric oxide by a receptor mediated system. Biochem Biophys Res Commun. 214:847.
- Hayes SN, and Taler SJ (1998): Hypertension in women: current understanding of gender differences. Mayo Clin proc 73:157.
- He J, Klag MJ, Whelton PK, Zhoa Y, and Weng X (1994). short and long-term prognosis after acute myocardial infarction in Chinese men and women. Am J Epidemiol. 139: 693-703.
- Heinsimer JA and Dewitt CM (1989). Exercise testing in women. J Am Coll Cardiol. 14:1448.

- Herlitz J, Hansson E, et al (2002). Characteristics and outcome among women and men transported by ambulance due to symptoms arousing suspicion of acute coronary syndrome. med Sci Monit. 4:251-6.
- Hitoshi S, Hajime W, Kiyotaka K, et al (1998). Does Passive smoking Impair Endothelium- dependent coronary Artery Dilatation in Women? J Am Coll Cardiol. 31: 811.
- Hochman JS, Mc Cabe CH, Stone PH, et al (1997). For the TIMI Investigators: Thrombolysis In Myocardial Infarction Outcome and profile in women and men presenting with acute coronary syndromes: a report from TIMI IIIB trial.

 J AM Coll Cardiol. 30:141-8.
- Hochman JS, Tamis JE, Thompson TD, et al (1999). For the Global Use of Strategies to Open Occluded coronary Arteries in Acute Coronary Syndromes IIb investigators. Sex, clinical presentation, and outcome in patients with acute coronary syndromes. N Engl J Med 341:226-32.
- Holmvang L, Clemmensen P, Wagner G, and Grand P (1999).

 Admission standard electrocardiogram for early risk stratification in patients with unstable coronary disease not eligible for acute revascularization therapy: a TRIM substudy. Am Heart J. 137: 24-33.

- Horwitz KB, and Horwitz LD (1982). Canine vascular tissues are targets for androgens, estrogens, progestins, and glucocorticoids. J Clin Invest. 69: 750.
- HRT Update 2001. National Institutes of Health NH, Lung, and Blood Institute. Women's Health Intitiative. Accessed March 2000 at http://www.nhlbi.nih.gov/whi/hrt.htm.
- Hulley S, Grady D, Bush T, Furberg C, Herrington D, Riggs B, et al (1998). Randomized trial of estrogen plus progestin for secondary prevention of coronary heart disease in postmenopausal women. Heart and estrogen / progestin Replacement Study (HERS) Research Group. JAMA. 280: 605-13.
- Humphrey LL, Chan BK, and Sox HC (2002). Postmenopausal hormone replacement therapy and the primary prevention of cardiovascular disease. Ann Intern Med. Aug 20;137 (4):273-84, Review.
- Inoue S, Hoshino S, Miyoshi H, et al (1996). Identification of a novel isoform of estrogen receptor, a potential inhibitor of estrogen action, in vascular smooth muscle cells.

 Biochem Biophys Res Commun. 219: 766-772
- Jacobs AK, Kelsey SF, Brooks MM, et al (1998). Better outcome for women compared with men undergoing coronary revascularization: a report from the Bypass Angioplasty Revascularization Investigation (BARI). Circulation. 98:1279-85.

- Jenkins JS, Flaker GC, Nolte B, et al (1994). Causes of higher in hospital mortality in women than in men after acute myocardial infarction Am J cardiol. 73: 319.
- Joan B, Loren L, and Sabina D. (2001). Coronary artery disease prevention: what's different for women? Am Fam physician. 63:1393-400, 1405-6.
- Jong P, Mohammed S, and sternberg L (1996). Sex differences in the features of coronary artery disease of patients undergoing coronary angiography. Can J cardiol. 12: 671.
- Jousilahti P, Tuomilehto J, Vartiainen E, et al (1996). Body weight, cardiovascular risk factors, and coronary mortality. 15 year follow -up of middle aged men and women in eastern finland. Circulation. 93: 1372.
- Jousilhati P, Puska P, Vartiainen E, et al(1996). Parental history of pre mature coronary heart disease an independent risk factor for myocardial infarction. J Clin Epidemiol. 1996; 49: 49%-503.
- Kannel WB, Wilson WF, D' Agostino RB, and Janet Cobb (1998).

 Sudden Coronary Death in women Am Heart J. 136:

 305.
- Kaski JC (1995). Cardiac syndrome X: Clinical characteristics and left ventricular function. Long-term follow-up study. J Am Coll Cardiol 25: 807.

- Kaski JC (1999). Chest Pain with Normal Coronary Angiograms:

 Pathogenesis, Diagnosis and Management. Kluwer

 Academic Publishers, Boston.
- Kaski JC and Russo G (2000). Microvascular angina in patients with syndrome X Kardiol.89 Suppl 9: IX/121-5.
- Kaski JC (2002). Overview of gender aspects of cardiac syndrome X. cardiovasc Res. 53 (3): 620-6.
- Kelsey SF, James M, Holubkov AL, et al (1993). Results of percutaneous transluminal coronary angioplasty in women: 1985-1986 National Heart, Lung and Blood Institute's Coronary Angioplasty Registry. Circulation. 87:720-7.
- Kemp HG, Kronmal RA, Vlietstra RE, and Frye RL (1986). Sevenyear survival of patients with normal or near normal coronary arteriograms: a CASS registry study. J Am Coll Cardiol. Mar:7 (3): 479-83.
- Kereiakes D, Lincoff A, Simoons M, et al (1999). Abciximab benefit in coronary intervention by gender and device: results of randomized trials (abstract). Eur Heart J. 20: 471.

- Kereiakes D, Giardina E. Lansky AJ, et al (2001). Managing cardio vascular disease in women: Role of clinical and interventional cardiology. Women's Health, CME circle 2001 NCM publishers, INC, New York, New York (http://www.ncmp.Com).
- Killip T, and Kimball J (1976). Treatment of myocardial infarction in coronary care unit. A two year experience with 250 patients. Am J cardiol. 20: 457.
- Kim C, Kwok YS, Saha S, et al (1999). Diagnosis of suspected coronary artery disease in women: a cost -effectiveness analysis. Am Heart J. 137:1019.
- Kober L, Torp- Pedersen C, Ottesen M, et al (1996). On behalf of the TRACE study group influence of gender on short-and long- term mortality after acute myocardial infarction. Am J cardiol. 77:1052.
- Kokkinos PF, Holland JC, Pittaras AE, Narayan P, Dotson CO, and Papademetriou V (1995). Cardiorespiratory fitness and coronary heart diesase risk factor association in women. Cardiovasc Rev Rep. 11: 49.
- Kostis JB, Wilson AC, O'Dowd K, et al (1994). sex differences in the management and long-term outcome of acute myocardial infarction: a state wide study. Circulation. 90: 1715-1730.

- Kronowski R, Lansky AJ, Mintz GS, et al (1997). Comparison of men versus women in cross- sectional area luminal narrowing, quantity of plaque, presence of calcium in plaque, and lumen location in coronary arteries by intravscular ultrasound in patients with stable angina pectoris. Am J Cardiol. 79: 1601-1605.
- Kudenchuk PJ, Maynard C, Martin JS, Wirkus M, and Weaver WD (1997). Comparison of presentation, treatment, and outcome of acute myocardial infarction in men versus women (the Myocardial Infarction Triage and Intervention Registry). Am J Cardiol. Mar 15; 79 (6): 844.
- Kurrelmeyer KM (2002). Non invasive evaluation of women with coronary artery disease. Current Opinion in Cardiology. 17:464-469.
- Kwok Y, Kim C, Grady D, et al (1999). Meta-analysis of exercise testing to detect coronary artery disease in women. Am J Cariol. 83:660.
- Lansky AJ, Peterson ED, Popma JJ, et al (1998). Gender differences in device selection and procedural outcome: a National Cardiovascular Network (NCN) report. J Am Coll Cardiol. 31:234A. Abstract 1086-82.

- Lansky AJ, Popma JJ, Mehran R, et al (1999). Tubular slotted stents: a "breakthrough therapy" for women undergoing coronary interventions. Pooled results from the STARS, ACSENT, SMART, and NIRVANA randomized clinical trials. J Am Coll Cardiol. 33: 58A. Abstract 833-5
- Lanza GA, Andreotti F. Sestito A, Sciahbasi A, Crea F, and Masei A. (2001) Platelet aggregability in cardiac syndrome X. Eur Heart J. 22 (20): 1844-5.
- Lewis JF, Lin L, McGorray S, et al (1999). Dobutamine stress echocardiography in women with chest pain. pilot phase data from the National Heart, Lung and Blood Institute Women's Ischemia Syndrome Evaluation (WISE).

 J AM Coll Cardiol. 33:1462-8.
- Lewis SJ, Sacks Efm Mitchell JS, et al (1998). Effect of pravastation on cardiovascular events in women after myocardial infarction: the cholesterol and Recurrent Events (CARE) trial. J Am Coll Cardiol. 32:140-146.
- Liao Y, Cooper RS, Ghali JK, Lansky D, Gao G, and Lee J (1993).

 Sex differences in the impact of coexistent diabetes with coronary heart disease. Diabetes care. 16:708-13.
- Lieberman EH, Gerhard MD, Uehata A, et al (1994). Estrogen improves endothelium- dependent, flow mediated vasodilation in postmenopausal women. Ann Intern Med. 121: 936.

- Lincoff AM, Califf RM, Ellis SG, et al (1993). for the Thrombolysis and Angioplasty in Myocardial infarction study Group.

 Thrombolytic therapy for women with myocardial infarction: is there a gender gap? J Am Coll Cardiol.

 22:1780 -1787.
- Linden W, Stossel C, and Mourice J (1996). Psychosocial interventions for patients with coronary artery disease. A meta-analysis. Arch intern Med. 156: 745.
- Losordo DW, Kearney M, Kim EA, Jekanowsky J, and Isner JM (1994). Variable expression of the estrogen receptor in normal and atherosclerotic coronary arteries of premenopausal women. Circulation. 89: 1501-1510.
- Mac Donald S, Joffres M, and Stachenko S (1992). Multiple cardiovascular risk factors in canadian adults. Can Med Assoc J. 146: 2021.
- Malacrida R, Genoni M, Maggioni Ap, et al (1998). For the Third International Study of Infarct Survival Collaborative Group (ISIS-3). A comparison of the early outcome of acute myocardial infarction in women and men. N Engl J Med. 338:8-14.
- Manson JE, Willet WC, Stampfer MJ, et al (1995). Body weight and mortality among women. N Engl J Med. 333; 676.

- Marrugat J, Sala J, Masia R, et al (1998). Mortality differences between men and women following first myocardial infarction. JAMA. 280: 1405-1409.
- Marrugat J, Masia R, Sala J, Elosua R, JAM Anto, and the REGICOR Investigators (2001). Role of age and sex in short-term and long term mortality after a first Q wave myocardial infarction. J Epidemiol Community Health. 55: 487-493 (July).
- Mark DB, Shaw LK, DeLong ER, et al(1994). Absence of sex bias in the referral of patients for cardiac catheterization. N Engl J Med. 330:1101.
- Mautner SL, Lin F, Mautner GC, Roberts we (1992): Comparison in women versus men of composition of atherosclerotic plaques in native coronary arteries and in saphenous veins used as aorto-coronary conduits J Am coll Cardiol. 21: 1312.
- Maynard C, and Weaver WD (1992). Treatment of women with acute MI: New findings from the MITI registry. J Myocardial Ischemia. 4; 27.
- Maynard C, Litwin PE, Martin JS, et al (1992). Gender differences in the treatment and outcome of acute myocardial infarction: results from the Myocardial Infarction Triage and Intervention Registry. Arch Intern Med. 152: 972-976.

- Maynard C, Litwin PE, Martin JS, and Weaver WD (1993).

 Treatment and outcome of acute myocardial infarction in women 75 years of age and older: findings from the Myocardial Infarction Triage and Intervention Registry.

 Cardiol Elderly. 1: 121-5.
- MC Guire DK, Emanulesson H, Charnwood A et al (1998).

 Diabetes mellitus is associated with worse clinical outcomes across the spectrum of acute coronary syndromes Results from Gusto-II b (abstract) Circulation. (suppl I): 1-259, 1998.
- Miller VT (1994). Lipids, liporpoteins, women and cardiovascular disease. Atheroselerosis. 108: S73.
- Milner KA, Funk M: Arnold A, and Vaccarino V (2002). Typical symptoms are predictive of acute coronary syndromes in women. Am heart J Feb: 143 (2): 383-8.
- Moller J, Hallqvist J, Diderichsen F, Theorell T, Reuterwall C, and Ahlbom A (1999). Do episodes of anger trigger myocardial infarction? A case—cross over analysis in the Stockholm Heart Epidemiology program (SHEEP). Psychosom Med. 61: 842.
- Mosca, L.; Manson, J.E.' Sutherland, S.E. et al (1996).

 Cardiovascular disease in women: A statement for health
 care professionals from AHA. Circulation. 2468,1997.

- Mosca L, Grundy S, Judelson D, et al (1999). Guide to preventive cardiology in women J Am Coll Cardiol. 33:1751-1755.
- Muller C, Neumann FJ, Roskamm H, Buser P, Hodgson JM, Perruchoud AP, and Buettner H (2002). Women do have an improved long term outcome after non- ST-elevation acute coronary syndromes treated very early and predominatly with percutanous coronary interventions: a prospective study in 1,450 consecutive patients. J Am Coll Cardio. Jul 17: 40 (2): 245-50.
- Njistad I, Arnesen E, and Lund- Larsen PG (1996). Smoking, Serum lipids, blood pressure, and sex differences in myocardial infarction: a 12-year follow up of the finnmark study. Circulation. 93:450 -456.
- Oe K, Shimizu M, Yamaguchi M, et al (2002). Effects of gender on the number of diseased vesses and clinical outcome in Japanese patients with acute coronary syndrome.

 Circulation. 66 (5): 435-40.
- Onaka H, Hirota Y, Shimada S, Kita Y, Sakai Y, Kawakami Y, et al (1996). Clinical observation of spontaneous anginal attacks and multivessel spasm in variant angina pectoris with normal coronaries: Evaluation by 24-hour 12 lead electrocardiography with computer analysis. J Am Coll Cardiol. 26: 38.

- Opherk D, Schuler G, Wetterauer K, Manthey J, Schwarz F, and Kubler W (1989). Four-year follow-up study in patients with angina pectoris and normal coronary arteriograms ("syndromeX"). Circulation. Dec; 80 (6): 1610-6.
- Orth-Gomer K, Mittleman MA, Schenck- Gustafsson K, et al (1997). Lipoprotein-a as a determinant of coronary heart disease in young women. Circulation. 95:329.
- Ozaki Y, Keane D, and Serruys P (1995). Fluctuation of spastic location in patients with vasospastic angina: A quantitative angiographic study. J Am coll cardiol. 26: 1606.
- Panagiotakos DB, Pitsavos C, Chrysohoou C, et al (2001). The effect of short term depressive episodes in the risk stratification of Acute coronary syndromes: a case control study in Greece (CARDIO 2000). Acta Cardiol. 56 (6): 357-365.
- Pepine CJ, El- Tamini H, and Labert CR (1992). Prinzmetal's angina (variant angina). Heart Dis stroke. 1: 281.
- Pfeffer MA, Brounwald E, Moye LA, et al (1992). On behalf of the SAVE investigators: Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction: Results of the survival and ventricular enlargement trial. N Eng J Med. 327:669.

- Phibbs B, Marcus F, Marriott HJC, et al (1999). Q wave versus non. Q wave myocardial Infarction: A meaning less distinction. J Am cardiol. 33: 576-582.
- Pinto RJ, Bhagwat AR, Loya Ys, and sharma S (1992). Coronary artery disease in premenopausal indian women: risk factors and angiographic findings Indian Heart J. 44: 99.
- Pramparo P (2000). Diangosis, Treatment, and Access to Care., The first international conference on Women, Heart Disease and Stroke: Science and Policy in Action. Conference summary.
- PURSUIT Trial Investigators (1998). Inhibition of platelet glycoprotein IIb/ IIIa with eptifibatide in patients with acute coronary syndromes. N Engl J Med. 339:436-443.
- Rexrode KM, Carey VJ, Henneckens CH, et al (1998). Abdominal Adiposity and Coronary Heart Diease in Women. JAMA. 280; 1843.
- Rich Edwards JW, Manson Je, Hennekens CH, and Buring JE (1995). The primary prevention of coronary heart disease in women. N Engl Med. 332:1758.
- Ridker PM, Buring JE, Shih J, et al (1998): Prospective study of C-reactive protein and the risk of future cardiovascular events among apparently healthy women. Circulation. 98:731-733.

- Ridker PM, Hennekens CH, Buring JE, and Rifai NC (2000). Creactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. N Eng J Med 342: 836-843.
- Robinson S (1995). Coronary heart disease in women. Urban Cardiol. 17:23.
- Robinson K, Conroy RM, Mulcahy R, and Hickey N (1988). Risk facotrs and in-hospital course of first episode of myocardial infarction or acute coronary insufficiency in women. J Am Coll Cardiol. 11: 932 -936.
- Roger VI, Jacobsen SJ, Pellikka PA, et al (1998). Gender differeces in use of stress testing and coronary heart disease mortality: a population-based study in Olmasted County, Minnesota. J Am Coll Cardiol. 32; 345.
- Rosen SD, Paulesu E, Wise RJ, and Camici PG (2002). Central neural contribution to the perception of chest pain in cardiac syndrome X. Heart. 88 (4): 328-30.
- Rosenberg L, Palmer JR, and Shapiro S (1990). Decline in the risk of myocardial infarction among women who stop smoking. N Engl J Med. 322:213.

- Rossouw JE, Anderson GL, Prentice RL et al (2002). Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal resuilts from the Women's Health Initiative randomized controlled trial. JAMA. 288 (3):31-33.
- Rude RE, Poole WK, Muller JE, et al (1983). Electrocardiographic and clinical criteria for recognition of acute myocardial infarction based on analysis of 3, 697 patients. Am J Cardiol. 52: 936.
- Safstrom K, Nielsen NE, Bjorkholm A, et al (1998). For the IRIS Study Group. Unstable coronary artery disease in postmenopausal women: identifying patients with significant coronary artery disease by basic clinical parameters and exercise test. Eur Heart J. 19.899-907.
- Schrrott H, Bittner B, Vittinghoff E, Herriington D, and Hulley S
 (1997). For the HERS Research Group. Adherence to
 National Cholesterol Education program treatment goals
 in postmenopausal women with heart disease. JAMA.
 227: 1218.
- Schwartz LM, Fisher ES, Tosteson NA, et al (1997). Treatment and health outcomes of women and men in a cohort with coronary artery disease. Arch Intern Med. 157:1545-51.

- Shah PK, Cercek B, and Lew As (1993). Angiographic validation of bedside markers of reperfusion. J Am Coll Cardio. 21:55-61.
- Sheldon WC (1993). Indications for coronary arteriography. Heart Dis Stroke.2:192.
- Shlipak MG, Simon JA, Vittinghoff E, Lin F, Barrett Connor E, Knopp RH, et al (2000). Estrogen and progestin, lipoprotein(a), and the risk of recurrent coronary heart disease events after menopause. JAMA. 283:1845-52.
- Sowers JR (1998). Diabetes Mellitus and Cardiovascular Disease in Women. Arch Intren Med. 158: 617.
- Stampfer MJ, Hu FB, Manson JE, Rimm EB, and Willett WC (2000). Primary Prevention of coronary heart disease in women through diet and lifestyle. N Engl J Med; 343: 16-22.
- Stokes J III, Kannel WB, Wolf PA, Cupples LA, and Agostino RB (1987). The relative importance of selected risk factors for various manifestations of cardiovascular disease among women and men from 35-64 years: 30 years of follow- up in the Framingham Study. Circulation.;75 (Suppl V) V65-73.
- Stone GW, Grines CL, Browne KF, et al (1995). Comparison of inhospital outcome in men versus women treated by either thrombolytic therapy or primary coronary angioplasty for acute myocardial infarction. Am J Cardiol. 75:987-992.

- Stone PH, Thompson B, Anderson HV, et al (1996). Influence of race, sex, and age on management of unstable angina and non-Q-wave myocardial infarction: the TIMI III registry. JAMA. 275: 1104-12.
- Stone GW, Marcovitz P, Lansky AJ, et al (1999). Differential effects of stenting and angioplasty in women versus men undergoing a primary mechanical reperfusion strategy in acute myocardial infarction. the PAMI Stent Randomized Trial. J Am Coll Cardiol. 33: 357 A. Abstract 832-6.
- Sullivan AK, Holdright DR, Wright CA Sparrow JL, Cunningham D, and Fox KM (1994). Ches pain in women: Clinical, investigative, and prognostic features. BMJ. 308:883-6.
- Suzanne O (1998). Pathophysiology of Sudden Coronary Death in Women: Implications for prevention. Circulation. 97:2103-2105.
- Taillefer R, DePuey EG, Udelson JE, et al (1997). Comparative diagnosis accuracy of TI201 and Tc-99m sestamibi SPECT imaging (perfusion and ECG gated SPECT) in dectecting co-onary artery disease in women. J Am Coll Cardiol. 29:69.
- The TIMI IIIB Investigators (1994). Effects of tissue plasminogen activator and a comparison of early invasive and conservative stratigies in unstable angina and non- Q- wave myocarial infarction: Results of the TIMI III B trial. Circulation. 89: 1545-1556.

- Thompson PM, and Wolf JL (1999). The sexual revolution in science: what gender -based research is telling us. J investig Med. 47:106-113.
- Timoteo AT, Ferreira J, Aguiar C, Almeida Mde S, Ribeiro MA, Cavaco DM, Trabulo M, and Seabra-Gomes R (2003).

 Revascularization and prognosis in female patients with non-ST-segment elevation acute coronary syndromes.

 Rev Port Cardiol. Mar; 22 (3): 367 -74.
- Tofler GH, Stone PH, Muller JE, et al (1987). Effects of gender and race on prognosis after myocardial infarction: adverse prognosis for women, particularly black women.

 J Am Coll Caridol. 9:473-482.
- Tyni-Lenne R, Stryjan S, Eriksson B, Berglund M, and Sylven C (2002). Benificial therapeutic effects of physical trining and relaxation therapy in women with coronary syndrome X. Physiother res Int. 7(1):35-43.
- Vaccarino V, Krumholz Hm, Berkman LF, and Horwitz RI (1995).

 Sex differences in mortality after myocardial infarction.

 Circulation. 91 1861.
- Vaccarino V, Parsons L, Every NR, Barron HV, and Krumbholz HM. (1999). For the National Registry of Myocardial Infarction 2 Participants. Sex-based differences in early mortality after myocardial infarction. N Engl J Med. 341:217-225.

- Vaccarino V, Abramson J, Veledar E, and Weintraub WS (2002).

 Sex differences in Hospital mortality After Coronary

 Artery By pass surgery Evidence for higher mortality in
 younger women: Circulation. 105: 1176.
- Vaziri Sm, Evans JC, Larson MG, and Wilson PW (1993). The impact of female hormone usage on the lipid profile: the framingham Offspring study. Arch Intern Med. 153:2200-6.
- Vermeer F, Bosl L, Meyer J, Bar F, Charbonnier B, Windeler J, and Barth H (1999). Saruplase is a safe and effective thrombolytic agent; observations in 1698 patients; results of PASS study. J thromb thrombolysis. 8:1-8.
- Vermeer. F (2001). Thrombolytic therapy in patients of female gender. Thrombosis Research. 103: 101-104.
- VeroniqueRL, Farkouh ME, Weston SA, et al (2002). Sex Differences in Evaluation and outcome of Unstable Angina. JAMA. 283 (5): 646-652.
- Viscoli CM, Brass LM, Kernan WN, Sarrel PM, Suissa S, and Horwitz RI (2001). A clinical trial of estrogen-replacement therapy after ischemic stroke. N Engl J Med. 345:1243-9.

- Wallentin L, Lagerqvist B, Husted S, et al (2000). Outcome at 1 year after an invasive compared with a non- invasive strategy in unstable coronary-artery disease: the FRISC II invasive randomized trial. FRISC II investigators. Fast Revascularisation during instability in Coronary artery disease. Lancet. July: 356 (9223): 9-16.
- Walsh BW, Schiff 1, Rosner B, Grienberg L, Ravinkar V, and Sacks F (1991). Effects of postmenopasual estrogen replacement on the concentrations and metabolism of plasma lipoproteins. N Engl J Med. 325: 1196.
- Walsh JME, and Grady D (1995). The treatment of hyperlipidemia in women. JAMA. 274: 1152.
- Waters DD, Alderman EL, Hsia J, et al (2002). Effects of a hormone replacement therapy and antioxidant vitamin supplements on coronary atherosclerosis in postmenopausal women: a randomized controlled trial JAMA. 288 (19): 2432-40.
- Weaver WD, White HD, Wilcox RG, et al (1996). Comparisons of characteristics and outcomes among women and men with acute myocardial infarction treated with thrombolytic therapy: GUSTO- I investigators. JAMA. 275:777-782.

- Weksler BB(1993) Hemostasis and thrombosis. In Douglas PS (ed):
 Cardiovascular Health Disease in women. Philadelphia,
 WB sounders. P 231.
- Wenger NK (1994). coronary heart disease in women: gender differences in diagnostic evaluation JAMA. 49: 181.
- Wenger NK (1996). Coronary heart disease in older women: prevention, dianosis, management, and prognosis. Clin Geriatr. 4: 53.
- Welexer LF (1999). Studies of acute coronary syndromes in women, lessons for everyone. N Engl J Med. 341: 275.
- White HD, Barbash GI, Modan M, et al (1993), for the Investigators of the International Tissue Plasminogen Activator / Streptokinase Mortality Study. After correcting for worse baseline characteristics. Women treated with thrombolytic therapy for acute myocardial infarction have the same mortality and morbidity as men except for a higher incidence of hemorrhagic stroke. Circulation. 88 (pt I): 2097-2103.
- Wilkinson P. Laji K, Panjadoyalon K, et al (1994). Acute myocardial infarction in women: Survival analysis in first six months. BMJ. 309-566.
- Willet W, Green A, Stampfer M, et al (1987). Relative and absolute excess risks of coronary heart disease among women who smoke cigarettes. N Engl J Med. 317: 1303.

- Williams JK, Adams MR, and Klopfenstein HS (1990). Estrogen modulates responses of atherosclerotic coronary arteries.

 Circulation. 81: 1680.
- Wong SC, Sleeper LA, Monrad ES, Menegus MA, Palazzo A, Dzavik V, Jacobs A, Jiang X, and Hochman JS (2001).

 SHOCK Investigators. Absence of gender differences in clinical outcomes in patients with cardiogenic shock complicating acute myocardial infarction. A report from the SHOCK Trial Registry. J Am Coll Cardiol. Nov 1, 38(5): 1395-401.
- Yarzebski J, Col N, Pagley P, et al (1996). Gender differences and factors associated with the receipt of thrombolytic therapy in patients with acute myocardial infarction: a community-wide perspective. Am Heart J. 131:43-50.
- Yusuf S, Peto R, Lewis J, et al (1985) Beta blockade during and after myocardial infarction: An over view of the randomized trials. Progr Cardiovasc Dis. 27:335.
- Zhang F, Ram JL, Standley PR, and Sowers JR (1994). 17 B estradiol attenuates voltage- dependent Ca2+ currents in vascular smooth muscle. Am J physiol. 193: C975.
- Zuanetti G, Latini R, Maggioni AP, et al (1993). Influence of diabetes on mortality in acute myocardial infarction:

 Data from the GISSI-2 study. J Am Coll Caridol. 22:

 1788.