

VII-REFERENCES

VII- REFERENCES

- Abbassy, M.A. (1974):**
Toxicity of some plant constituents to insects. Ph. D. Thesis.,
Fac., Agric. Alex. Univ., PP. 180.
- Abbassy, M.A.; Zein, A. Ashray M. and Abou-Shamish S. (1981):**
Naturally occurring chemicals for pest control. Insecticidal
and synergistic alkaloids isolated from black pepper, *piper*
nigrum (L.) Proc. 4th Arab Pesticides Conf. Tanta Univ. Vol.
(111A): 339-407.
- Abbott, W.S. (1925):**
A method of computing the effectiveness of an insecticide. J.
Econ. Entomol., 18: 265-267.
- Abd El-Naby, L.M.; Farrag R.M. and El-Sheikh F.M. (1988):**
Relative potency of certain insecticides against the green
peach aphid. (*Myzus persica*: Homoptera: aphididae). J.
Agric. Sci. Mansoura Univ., 13 (1): 334-338.
- Abd El-Samea, S.A. (1990):**
Biological studies on some lepidopterous insects by using
extractions of some plants. M.Sc. Thesis, Fac. of Sci.
Menoufia Univ., Egypt.
- Abdel-Hafez, M.M. (1978):**
Some biochemical studies on cotton Leafworm *Spodoptera*
littoralis Ph. D. Thesis, Fac. Agric., Cairo Univ., Cairo,
Egypt.
- Abul Dahab F. F. (1998):**
Toxic action of some plant extracts on *Musca domestica*. J.
Egypt. Ger. Soc. Zool. Vol. 27 (E), Entomology, 33 - 4.1.
- Afifi, F.A.; Salem, M. and Hekal, A.M. (1989):**
Insecticidal properties of the extracts of lupin seed and
caraway fruits against some stored product insects. Ann.
Agric. Sc., Fac. Agric. Ain Shams Univ., Cairo, Egypt, 34:
401-404.

- Agrowal, H.S.; Gupta, N.K.; Prasad, V.K. and Vishwanama, S. (1979):**
Chemical control of yellow mosaic of "moong". Pesticides 13 (5): 44-47.
- Ahmed, A.H.M.; Elhag, E.A. and Bashir, N.H.H. (1987):**
Insecticides resistance in the cotton whitefly (*Bemisia tabaci* Genn.) in the Sudan Gizira. Tropical Pest management, 33 (1) 67-72.
- Ahmed M.S. and Mahgoub S. M. (1996):**
Insecticidal properties of the extracts of *Ricinus communis* seed against the Cowpea Beetle *Callosobruchus maculatus*. Egypt J. Agric. Res., 74 (3).
- Al Shayji Y., Shaheen N., Saleem M., and Ibrahim M. (1998):**
Susceptibility of aphid, *Aphis gossypii* (Aphididae, Homoptera) to certain *Bacillus thuringiensis* formulations in kuwaite Egypt. J. Agric. Res. 76(2) 599-606.
- Amir, D., Thomas O., Leigh F. and Granett J. (1992):**
Insecticide resistance in field populations of green house whitefly (Homoptera: Aleyrodidae) in the San Joaquin Valley (California) cotton cropping system. J. Econ. Entomol., 85 (1): 21-27.
- Anderson, T.E. Babu, J.R. Dybas R.A. (1986):**
Avermectin, B, ingestion and contact toxicity against *Spodoptera cridania* and *Heliothis virescens* (Lepidoptera : Noctuidae) and potentiation by oil and piperonyl butoxide. J. Econ. Entomol, 79 : 197 - 201.
- Areekul, S., Sinchairsri, P. and Tigvatanon, S. (1987):**
Effect of the plant extracts on the oriental fruit fly. Kasetstart. Journal Natural. Sciences, 21: 395-407.
- Ascher, K.R.S., Eliyahu, M., Nemny, N.E. and Ishaaya, I. (1986):**
The toxicity of synthetic pyrethroids and other insecticides and pesticides to *Spodoptera littoralis* (Boisd), *Drosophila melanogaster* (Meis) and *Bemisia tabaci* (Genn.). Int. Pest. Control, V (3): 68-71.

Atha, F.B. and Hamilton, J. T. (1978):

Insecticide resistance in *Myzus persicae* in Australia. J. Econ. Entomol. 7 - 851 - 853 (1978).

Ayad, F.A., Keddis M.E., Ghoneium, Y.F. Mourad, E.I. Allam, A. M. and El-Guindy M.A. (1992):

Toxicity of some insecticides. Against *Aphis Gossypii* Glover collected from different governorates of Egypt during 1990 cotton season. Bull. Ent. Soc. Egypt, Econ. Ser. 19, (101).

Ayoub S.M. A. (1994):

Field and laboratory studies on the susceptibility of the whitefly *Bemisia tabaci* (Genn) to insecticides. M.sc. thesis (pesticides). Fac. Agric. Cairo univ.

Badr N.A., Hindy, M.A.; Negm, M.F. Abd El-Halem S.M. and EL-Sisi A.G (1999) :

Performance of ground low volume spraying machines for controlling cotton leafworm larvae (Glover) plants. Egypt J. Agric. Res 77 (3) 1035 – 1043.

Balasubramaian, G. and Chelliah, S. (1974):

Chemical control of pests of sunflower. Pesticides, 19 (4): 21-22.

Bashir, S.E. (1974):

Effect of some insecticides on immature stage of the cotton whitefly. Cotton Growing Review, 51 (1): 62-69.

Berlinger Menachem J.; Sara Lebiush. Mordechi, Richard Dahan, Robin A.J. Taylor (1996):

Rapid Method for Screening insecticides in the laboratory pesticides. Sci. 46, 345-353.

Belal M.H. and, Othman S.A. K. (1992):

The susceptibility and resistance status of the Egyptian cotton leaf worm *Spodoptera littoralis* (Boisd) to some currently used organophosphorous insecticides. Bull. Ent. Soc. Egypt. Econ. Ser., 19 : (25).

- Bhattachar Jee, N.S. (1976):**
Control of the spread of the yellow mosaic virus of "moong" in soybean. Entomologist. Nevsietter 6 (11/12): 64-65.
- Bodhade, S.N. Norkhede P.W. and Borle, M.N. (1992):**
Residual toxicity of different synthetic insecticides against aphid on Bhindi Abel Monschus. Esculentus Moench. Indian. J. Ent., 54 (1) : 20 - 26.
- Butler, G.D.; Coudriet, D.L. and Henneberry, T.J. (1988a):**
Toxicity and repellency of soybean and cotton seed oils to the sweetpotato whitefly and the aphid on cotton in green house studies. Southwest. Entomol, 13 (2): 81-86.
- Butler, G.D.; Hutchison, W.D. and Broza, M. (1988b):**
Effect of aldicarb treatments to cotton on *Bemisia Tabaci* and *Aphis gossypii* populations in Israel. Southwest Entomol., 1-13 (2): 87-93.
- Cahill M., Gorman K., Day S., Denholm I., Elbert A., Nauen R. (1996):**
Baseline determination and detection of resistance to imidacloprid in *Bemisia tabaci* (Homoptera : Aleyrodidae). Bulletin of Entomological Research, 86, 343 - 349.
- Chauhan, S.P.; Kumar, A.S.; Singh, C.L. and Pandey, U.K.(1989):**
Toxicity of some plant extracts against rice moth *Corcyra cephalonica* strain lepidoptera. India J. Entomol., 49: 532-534.
- Chaufaux J, Cohn J. M., Buisson C., Sanchis V., Lreclus D. and Pasteur N. (1997):**
Inheritance of resistance to *Bacillus thuringiensis* crylc toxin in *Spodoptera littoralis* (Lepidoptera : Noctuidae) J.Econ. Entomol 90(4) 873-878.
- Chavan, S.R. and Nikam S.T. (1982):**
Mosquito larvicidal activity of *Ocimum basilicum* linn. Indian J. Med. Res, Vol, 75: PP. 220-2.
- Chao S. L, Dennehy T.J. and Casida J. E.,(1997):**
Whitefly (Hemiptera:Aleyrodidae) site. for imidacloprid and related insecticides : aputative nicolinic acetylcholine receptor.J.Econ. Entomol (4).879-882.

Cochran, D.G. (1994):

Abamectin resistance potential in the German Cockroach (Dictyoptera: Blattellidae). J. Econ. Entomol., 87 (4): 899-903.

Cohen, S. and Veierov, D. (1989):

Effect of various insect control agents on the landing behaviour of *Bemisia tabaci*. Phytoparasitica, 17(3): 229-30.

Collins, P.J. Rose, H.A. and Wege Csany, M. (1992):

Enzyme activity in strains of the sawtoothed grain beetle (Coleoptera: Cucujidae) differentially resist to fenitrothion malathion, and chlorpyrifos methyl. J. Econ. Entom. 85-1571-1575.

DeCock, A. Ishaaya, I.; Degheele, D. and Veiero, V. (1990):

Vapor toxicity and concentration depends persistence of buprofezin applied to cotton foliage for controlling the sweetpotato whitefly (Homoptera: Aleyroididae). J. Econ. Entomol., 83, 1254-1260.

Dittrich, V. and Ernst, G.H. (1983):

The resistance pattern in whitefly of sudanese cotton. Entomologie, 4 (1/3): 96-97.

Dittrich, V.; Hassan, S.O. and Ernst, G.H. (1985):

Sudanese cotton and whitefly: a case study of the emergence of a new primary pest. Crop prot., 4 (2): 161-176.

Dittrich, V.; Ernst, G.H.; Ruesch, O. and Uk, S. (1990):

Resistance mechanisms in sweetpotato whitefly (Homoptera: Alyrodidae), populations from Sudan/Turkey/ Guatemala and Nicaragua. J. Econ. Entomol., 83 (5): 1665-1670.

El-Dahan A.A. (1991):

Resistance status of *Spodoptera littoralis* (Boisd), to insecticides in Kafr El-Sheikh Governorate during, 1984-1988 cotton growing seasons. Egypt. J. Agric. Res., 69 (1).

El Dahan A.A. and Hallawa L.A. (1991):

The joint action of pyrethroids with acephate and Amitraz on the cotton leafworm *Spodoptera littoralis* (Boisd). Egypt J. Agric. Res., 69 (1).

El Defrawi, M.E., Toppozada A., Mansour N., and Zeid M. (1964):
Toxicological studies on the Egyptian cotton leafworm *Prodenia littura* (F). Susceptibility of different larval instars of *Prodenia* sp. to insecticides. J. Econ. Entomol., 57: 591-593.

El Guindy, M.A. Elsayed G.N. and madi S.M. (1975):
Distribution of insecticide resistant strains of cotton leafworm, *Spodoptera littoralis* (Boid) in two governorates of Egypt. Bull-Ent Soc Egypt, Econser 191-195.

El Guindy, M.A., Madi, S. M., El Sayed, G.N., Farrag, A. M. I. Dogheim, S. M.A. Keddis, M.E. and Issa ,Y. H. (1981):
Further studies on resistance spectrum and subgrouping of certain insecticides in field populations of the Egyptian cotton leaf worm, *Spodoptera littoralis* (Boisd) during the 1980 cotton season. Bul, Ent. Egypt. Econ. Ser., 11 : (243).

El Guindy, M.A. Keddis, M. E. Abdel Sattar M.M. and Ghoneim Y.F. (1989):
Status of resistance to pesticides in the cotton leafworm *Spodoptera littoralis* (Boisd), under the present Egyptian cotton pest control programme. Proc. 1st Int. Conf. Econ. Ent. Vol. 11 1989 (453).

El Hag, F.A. and Horn, D.G. (1983):
Resistance of greenhouse whitefly (Homoptera: Aleyroididae) to insecticides in selected Ohio greenhouse. J. Econ. Entomol., 76: 945-948.

El Lakwah, F.A. and Abd El- latif A.M. (1998):
Effect of surface treatment of grain bags with the botanical insecticide Neemazol-T and certain plant extracts on weight loss of stored wheat and Maize grains due to infestation with *Sitophilus oryzae*(L).Egypt. J.Agric. Res. 76(3) 1003-1016.

EL Lakwah, F. A., Abdel Latif A., and EL-Sayed F.M. (1999):
Toxicity and joint action of the botanical insecticide neemazal-w with malathion to the cowpea weevil, *Callosobruchus maculalus* (F). Egypt. J. Agric. Res. 77 (3)1023 –1034.

El Meniawi F. A. and Hashem, M. (1997):

Insecticidal activity of detergents against the adult and immature stages of the cotton whitefly *Bemisia tabaci*. Genn on tomato. Alex J. Agric Res .42.(3), 75-87

El Sayed, E.I. (1982):

Effect of different insecticides in combination with neem (*Azadirachta indica* A. Juss) seed suspension on the Egyptian Cotton leafworm *Spodoptera littoralis*, Zagazig J. Agric. Res. (Egypt), 9: 457-469.

Elsebae A. A, Nousier N. I., El-Deeb Y. A.A., and Taman F. A. (1997):

Antifeeding and ovicidal effects of 1-5` x plant extracts against the larvae and egg mass. of *Spodoptera littoralis* (Boisduval). Alex. Exchange, Vol. 18- 321-334.

Ernst, G.H., Ruesch, O., Dittrick .N. and Uk, S. (1987):

Unpublished data cited from whiteflies their "bionomics", pest status and management" (D.Gerling.Ed.), chapll. Intercept, 1990.

Evans, R. and Kaleysa Raj (1988):

Extracts of Indian plants as mosquito larvicides. Indian J. Med. Res. 88, July 1988, PP. 38, 41.

Evans, R. and Kaleysa, R. (1991):

Larvicidal efficacy of Quassin against *Culex quinquefasciatus*. Indian J. Med. Res. [A] : 93, September 1991, PP. 324 - 327.

Finney, D.J. (1971):

Propit analysis. Cambridge university press, cambridge, London, 333 pp.

Freedman, L.J., Kwolek, W.F. Berry, E.C. and Guthrie, W.D. (1979):

A bioassay for plant derived pest control agents using the European corn borer. J. Econ. Entomol, 72 : 541 - 545.

Furk, C, and Hines,C.M.(1993):

Aspects of insecticide resistance in the melon and cotton aphid, *Aphis gossypii* (Hemiptera:Aphididae). Ann. Appl. Biol. 123(1):9-17.

Gameel, O.I. (1974):

Field evaluation of insecticides for jassid, *Empoasca lybica* De Berg and Whitefly, *Bemisia tabaci* (Gennaduis) control on cotton (Hemiptera : Jassidae and Aleyroididae). Bull. Soc. Ent. Egypt., 7 : 113 - 122.

Goel S.C. and Kumar V. (1992):

Food-energy, budget and secondary productivity of noctuidae larva *Plusia orichalceafabr.* Lepidoptera. Indian J. Ent. 54 (I) : 27 - 33.

Gopalan, M.; Madhusudhan R. and Balasubramanian G. (1987):

Studies on biological activity of plants on *Dysdercus cingulatus* Madros - Agric. J. 47 : 61 - 69.

Gravena, S. and Nakano, O. (1975):

Experimental test with some insecticides for the control of the whitefly *Bemisia tabaci* (Genn) (Homoptera : Aleyrodidae) an autumn sown beans in the north of Parana. Cientifica. 3 (1) : 187 - 193.

Gubran, E.M.E; Delorme, R; Auge, D; Moreau-J.P.(1993):

Pyrethroids and organochlorines resistance in cotton aphid *Aphis gossypii* (Glov.) (Homaptera: Aphididae) in the Sudan Gezira. Int. J. Pest Mana. 39 (2):197-200.

Guy, B. and David, W. (1994):

Methidathion Resistance in the sweetpotato whitefly (Aleyrodidae : Homoptera) in Israel : Selection, heritability, and correlated changes of esterase activity. J. Econ. Entomol. 87 (5) : 1147 - 1156.

Habsah, A.K., Zakeria, M.B. Kechil A.A. and Azirun M.S. (1989):

Toxicity and electrophysiological effects of *Spilanthes acmella* Murr. Extracts on *Periplaneta americana* (L). Pesticide. Sci, 25 : 329 - 335.

Hardee, D.D. and Answorth, J.M. (1993):

Cotton aphid (Homoptera-Aphididae effect of In-Furrow Insecticides on pesticide resistance. J. Econ. Entomol., 86 (4): 1026-1029.

- Hollingsworth R.G; Tabashnik B.E, Ullman. D.E; Johnson, M.w; Messing, R. (1994):**
Resistance of *Aphis gossypii* (Homoptera" Aphididae_ to insecticides in Hawaii spatial patterns and relation to insecticide use. J. Econ. Entomol 87(2) 243-300.
- Holtkamp R.H. Edge, V.E. Dominiak, B.C. and Walters, P.J. (1992):**
Insecticide resistance in *Therioaphis trifollii* (Homoptera : Aphididae) in Australia. J. Econ. Entomol., 85 (5) : 1576 - 1582.
- Horowitz, A.R.; Toscano, N.C.; Youngman, R.R.; Kido, K.; Kaneko, J.J. and Georgiou, G.P. (1988):**
Synergism : Potential new approach to whitefly control. Calif. Agric. 42 (1) : 21 - 22.
- Horowitz, A.R. and Ishaaya, I. (1992):**
Susceptibility of the sweet potato whitefly (Homoptera : Aleyrodidae) to buprofezin during the cotton season. J. Econ. Entomol. 85 (2) : 318 - 324.
- Ishaaya, I.; Austerweil, M. and Frenkel, H. (1986):**
Effect of the petroleum oil virol on toxicity and chemical residue of fenpropathrin applied against adult of *Bemisia tabaci* (Homoptera : Aleyrodidae) as high and low volume sprays. J. Econ. Entomol, 79 : 596 - 599.
- Ishaaya, I.; Mendelson, Z. and Madjar, M. (1988):**
Effect of buprofezin on embryogenesis and progeny formation of sweet potato whitefly (Homoptera : Aleyroididae) J. Econ. Entomol, 81 (3) : 780, 784.
- Issa, Y.H. Keddis, M.E. Abdel Sattar, M.A., Ayad F. A., and EL-Guindy M. A. (1985 a):**
Survey of resistance to pyrethroids in field strains of the cotton leafworm *Spodoptera littoralis* (Boisd) during 1980 - 1984 cotton growing season. Bull. Ent. Soc. Egypt. Econ. Ser., 14, (405).
- Issa, Y.H.; Keddis, M.E.; Ayad, F.A. Abdel Sattar, M. M.; and El-Guindy, M. A. (1985 b):**
Survey of resistance to organophosphorous insecticides in field strains of the cotton leafworm *Spodoptera littoralis* (Boisd), during 1980 - 1984 cotton growing seasons. Bull - Ent. Soc. Egypt. Econ. Ser. 14 (399).

- Issa, Y.S. Farag, A.A. Keddis M. E. and El-Hamaky M.A. (1993):**
Toxicity of certain plant extracts to *Spodoptera littoralis* (Boisd) and *Aphis durantae* (Theoald). Egypt. J. Appl. Sci., 8 (10).
- Jagdish, R.K.,; Krupadanam, G.L.D and Srimamnaryana, G. (1989) :**
Isoflavans from *Milleta racemosa* phytochemistry 28 : 913 - 916.
- James, M.C. and Ubina, C.D. (1990):**
Evaluation of some botanicals against tobacco insect pest national review and planning workshop on batacias pesticides. Los Banos, Lagunal Philippines), 19 - 20.
- Katundu M.J. and Aliniaze, E. (1990):**
Variable resistance of filbert aphid (Homoptera : Aphididae to insecticides in the willamette valley, Oregon - J. Econ. Entomol, 83 (1) : 41 - 47.
- Kalsansundaram, M. and Das P.K. (1985):**
Larvicidal synergistic activity of plant extracts for mosquito control. Indian J. med. Res., 82 : 19 - 23.
- Kandil, M.A, Elkabbany, G.H. and Abdalah, M.D(1991):**
Efficacy of some insecticides against the cotton whitefly *Bemisia tabaci* (Genn) with special regard to their side effect on predator Bull. Soc. Ent .Egypt 19-9-17.
- Kayitare, J. and Ntezarubanza, L. (1989):**
Evaluation of the toxicity and repellent effect of certain plants from Rwanda against the bean bruchids. *Acanthoscelides obtectus* Sary et *Zabrates subfaciantus* Boheman. Inset. Science and its application, 12, 695 - 697.
- Keddis, M.E., Ayad, F. A. Abdel Fattah M. S. and El-Guindy M. A. (1986):**
Studies of resistance to urea derivatives and their mixtures with insecticides in field strains of the cotton leaf worm *Spodoptera littoralis* (Boisd) during the cotton seasons 1983, 1984 and 1985. Bull. Ent. Soc. Egypt. Ser., 15 (229).

Keddis, M.E.; Omar M.E. and El-Guindy M.A.(1988):

Monitoring insecticides resistance in field strains of the cotton leafworm, *Spodoptera littoralis* (Boisd), during 1986 cotton season. Agric. Res. Revi., 66 (1) : 25 - 31 (1988).

Kerns, V. and Gaylor M.J. (1992):

Insecticide resistance in field populations of the cotton Aphid (Homoptera-Aphididae) J. Econ. Entomol., 85 (1): 1-8.

Khan, M.A.J. and Khan R.G. (1984):

Insecticidal effects of indigenous vegetable oils (Tarmaria and Artemisia) on some fruit tephritids in Pakistan. Proceeding of the Entomological, Society of Karachi, 14/15, 113 - 118.

Kisha, J.S.A. (1986):

Comparison of electrodynamic spraying with use of knapsack sprayer for control of whitefly on tomato in the sudan Gezira. Ann. Appl. Biol. 108, Suppl., 36 - 37.

Madi, S.A.; El Guindy, M.A. El-Sayed, G. N.; Keddis, M.E.; Issa Y.H.; and Abd El-Sattar, M.M. (1979):

Evaluation of certain insecticides against field strains of the Egyptian cotton leaf worm, *Spodoptera littoralis* (Boisd) during 1977, 1978 and 1979), cotton seasons in Egypt. Bull - Ent. Soc. Egypt. Econ. Ser., 11[201].

Madjar, M.V.; Hameir, J. and Chen, M. (1987):

Influence of aldicarb at different stages of cotton plants growth on the development of *Bemisia tabaci* populations. Phytoparasitica 15 (3) : 263.

Matter, M.M. Marei, S.S.; Moqwad, S.M and El Gengaihi, S. (1993):

The reaction of *Aphis gossypii* and its predator *Coccinella undecimpunctata*. Bull. Fac. Agric. Cairo Univ., 417 - 432.

Mckenzie C.L; Cartwright, B. (1994):

Susceptibility of *Aphis gossypii* (Glover) to insecticides as affected by host plant using a rapid bioassay. J. Entomol. Sci. 29 (3):289-301

Misra, A.K.; Misra D.N. and Tripathi N.N. (1988):

Mycotoxic evaluation of some higher plants. National - Academy - Science letters, India, 11, (1) 5-6.

Mourad, E.I.; Keddis M.E. and Ghoneim Y.F. (1991) :

Potency of certain insecticides against the cotton whitefly, *Bemisia tabaci* (Genn.). J. Agric. Sci. Mansoura Univ. 16 (1) : 172 - 175.

Mourad, E.I.; Keddis, M.E.; Osma, M.S. Ayad, F.A. and El-Gunindy M.A. (1994):

Suitability of some natural and mineral oils as pesticidal synergists against *Aphis gossypii* Glover. Egypt J. Agric. Res., 72 (3).

Moustafa, F.I.; Shawira M.S. and Mourad M.A. (1992):

Studies on resistance to insecticides 11- comparative susceptibility of insect populations of *Spodoptera littoralis* from several locations of the country to several insecticides from different groups - Alex - J. Agric. Res., 37 (1) : 505 - 521.

Moustafa, O.K. and El Attal, Z.M. (1984) :

The joint action of triflumuron and some insecticides against the cotton leafworm *Spodoptera littoralis* (Boisd). Bull Ent. Soc . Egypt. Econ. sev 14 (161).

Nassef. M. A. and Watson W.M. (1999):

Sequential spray schedules of insecticides to control boll worms as target pests in addition to certain sap suckers as non target pests in cotton fields. Egypt Agric. Res. 77(3): 155 – 1162.

Nauen R., Koob B. and Elbert A. (1998):

Antifeedant effects of sublethal dosages of imidacloprid on *Bemisia tabaci*. Entomologia Experimentalis et Applicata 88: 287-293.

O'Brien, P.J. Abdel Aaol, Y.A. Ottea, J. A. and Graves, J. B. (1992):

Relationship of insecticide resistance to carboxylesterases in *Aphis gossypii* (Homoptera : Aphididae) from Mid South cotton. J. Econ. Entomol. 85 (3) : 651 - 657.

Oetting, R.D. and Anderson, A.L. (1990):

Imidacloprid for control of whiteflies *Trialeurodes vaporariorum* and *Bemisia tabaci*, on greenhouse grown poinsettias. Brighton crop protection conf., Pests and Diseases, 367 - 372.

Omar, H.I.; Hegab, M.F.A. and El Hamaky, M.A. (1988):

Combined action of some insecticides and fertilizers on pests infesting sweet melon. Agricultural Research review 66 (1) : 85 - 90.

Owusu. E.O; Kim-C.S; Horiike, M.; and Hirano, C. (1995):

Some dichlorvos induced plant enzymes and their possible influence on field control of cotton aphid (Homoptera Aphididae). Journal of Horticultural Science 70(3) 461-467.

Owusu .E.O; Horiike, M; and Hirano, C.(1996):

Inhibition by insecticides of partial purified carboxylesterase from *Aphis gossypii* (Homoptera : aphididae). J. Econ. Entomol. 89(2):307:310.

Pandy, N.D., Mathur, K.K. Pandey S., and Tripathi, R.A. (1986):

Effects of some plant extracts against pulse beetle, *Callosobruchus chinensis* Linnaeus. Ind. J. Ent. 48 : 85 - 90.

Pandy, N.D., Singh, L. Singh Y.P. and Tripathi, R.A. (1987):

Effect of certain plant extracts against *Lipaphis erysimi* Kalt under laboratory conditions. Ind. J. Ent. 49 : 238 - 242.

Patterson, S.K., Betty, D. Wahbakhalil, S.K. Schermeister L.J. and Quaishi M. S. (1975):

Plant insect interactions. I. Biological and phytochemical evaluation of selected plants. Lloychia, 38 : 391 - 403.

Peterson, G.S., Kandil, M.A. Abdallah, M.D. and Farag A.A.A. (1989):

Isolation and characterisation of biologically - active compounds from some plant extracts. Pesticide. Sci. 25 : 337 - 342.

Prabhaker, N.; Toscano, N.C. and Coudriet, D.L. (1989):

Susceptibility of the immature and adult stages of the sweetpotato whitefly (Homoptera : Aleyrodidae) to selected insecticides. J. Econ. Entomol. 82 (4) : 983 - 988.

- Prabhaker, N.; Toscano, N.C.; perring, T.M.; Nuessly, G.; Kido, K. and Youngman, R.R. (1992):**
Resistance monitoring of the sweetpotato whitefly (Homoptera : Aleyrodidae) in the Imperial Valley of California. J. Econ. Entomol. 84 (4) : 1063 - 1068.
- Quarles, W. (1992):**
Botanical pesticides from chinopodium IPM. Practitioner, 14 : 1-11.
- Radwan, H.S.A.; Ammar, I.M.A.; Eisa, A.a.; Assal, O.M. and Omer, H.I.H. (1985):**
Development retardation and inhibition of adult emergence in cotton white fly *Bemisia tabaci* (Genn.) following immature stages treatment with two molt inhibitors. Bull. Soc. Ent. Egypt (18): 175- 181.
- Rae, N.V.; Reddy, A.S. and Reddy, D.D.R. (1990):**
Relative toxicity of some insecticides to cotton whitefly, *Bemisia tabaci*. Indian Journal of plant protection 18 (1) : 97 - 100.
- Raman, K.V.; Booth, R.H. and Milalacios (1987):**
Control of potato tuber moth *Phthoremaea operculella* (Zeller) potato stores. Tropical. Science, 27 : 157 - 194.
- Rashwan, M.H. El-Bermawy, Z.A. El-Sheikh A.E. and Radwan, H.S.A. (1992):**
The onset of organophosphate and carbamate resistance among lower Egypt population soil. The cotton leaf worm, *Spodoptera littoralis* (Boisd). Bull. Ent. Soc. Egypt. Econ. Ser., 19, 1991 - 1992 (211).
- Rataul, H.S.; Harbans, S. R. H. and Singh, L. (1976) :**
Field studies on the control of soybean yield - mosaic virus, transmitted by whitefly *Bemisia tabaci* (Genn.) by using different granular systemic insecticides. Journal of Research. Punjab Agricultural University 13 (3): 298 - 302.
- Ravensberg, W.J.; Malais, M. and Schaaf, D.A. (1990):**
Verticillium lecanii as a microbial insecticide against glasshouse whitefly. Brighton crop protection Conf. Pests and Diseases, 265 - 268.

Reddy, P.V.R.; K.C. Chitra, R.C. and Rao P.K. (1990):

Field evaluation of certain plant extracts for the control of brinjal spotted leafbeetle, *Henosepilachna rigintioctopunctat*. Journal of Insect. Science, 3 : 194 - 195.

Robert G. Sworth H., Bruce E., Ink T., Johnson M., Russell Vc H. Messing, Z. and Diane E. M. (1997):

Relationship Between susceptibilty to insecticides and fecundity across populations of cotton aphid *Aphis gossypii* (Homoptera Aphididae). J. Econ. Entomol. 90(1) 55-58.

Roush R.T. and Miller G.L. (1986):

Considerations for design of insecticide resistance monitoring programs. J. Econ. Entomol., 79 : 293 - 298.

Saxena, J.D. Banerjee S.K. and Sin H.A. (1992):

Pyrethroid resistance in field populations of pink Bollworm *Pectinophora gossypiella* Saunders in India. Indian. J. Ent. 54 (3) : 347 - 350.

Sawicki R. M.S. and Rice A.D. (1978):

Response of susceptible and resistant peach potato aphids *Myzus persicae* (Sul.) to insecticides in leaf dip bioassays. Pestic Sci., 1978, 9, 513 - 516.

Saxena, R.C. (1989):

Insecticides of plant origin insecticides from neem. Entomology department, international rice research institute, P.O. Box 933, Manila, philippines American Chemistery Society chapter (9): 11 - 131.

Scott, W.P. and Mckibben G.H. (1978):

Toxicity of black pepper extract to boll weevils. J. Econ. Entomol.; 331 : 343.

Selander, J.; Markkula, M. and Tittanen, K. (1972):

Resistance of the Aphids *Myzus persicae* (Sulz) *Aulacorthumsolani* (Kalt.) and *Aphis gossypii* Golv. to insecticides and the influence of the host plant on this resistance. Ann. Agric. Fenn. 11 : 141 - 145.

Sharaf, N.S. and Allawi, T.F. (1980):

Studies on whiteflies on tomato in Jordan Valley III. Laboratory and field experiments on the control of whitefly (*Bemisia tabaci*) Genn. (Homoptera : Aleyroididae). Populations with organophosphorous insecticides and the incidence of the tomato leaf curl virus. Zeitschrift fur pflanzenkrankheiten und Pflanzenschutz, 83 (3) : 176 - 184.

Sharma, S.K.; Mathur, S.D.; Khan, B.W. and Mathur, B.N. (1971):

Evaluation of some insecticides for the control of insect pests of cotton by means of aerial spraying and their effect on parasites and predators. Zeitschrift fur pflanzenkrankheiten und Pflanzen. Schutz, 78 (5) : 286 - 295.

Shawir, M.S. Moustafa F., and Mourad M.A. (1991):

Studies on resistance to insecticides. Resistance levels of six field strains. of the Egyptian cotton leafworm *Spodoptera littoralis* (Boisd) to certain insecticides in two cotton seasons. Alex. J. Agric. Res., 36 (3) : 327 - 340.

Sherif, A. and Hall, R.G. (1985):

Plant ingredients detrimental to mosquito development. Atlantic city. New Jersey, 249 - 255.

Sherif, A. and Hall R.G. (1986):

The utilization of plant ingredients as aids in mosquito control proceedings and papers of the Annual, Conference of the California Mosquito and vector control Association, 53 : 113 - 116.

Sherif, A.; Hall R.G. and El- Amamy, M. (1987):

Drugs, insecticides and other agents from Artemisia, Medical Hypothesis 23 : 182 - 193.

Shires, S.W.; Inglesfield, C.; Tipton, J.D. and Sittingbourne, U.K. (1987):

Effects of chlorfenvinphos on *Bemisia tabaci* Genn. and its parasites on cotton in the Sudan Gezira. Crop. Prot. 2(2) : 109 - 116.

Sidhu, A.S. and Dhawan, A.K. (1981):

Evaluation of some new insecticides against foliage feeding pests of cotton. Pesticides, 15 (6) : 20 -22.

- Silver A., R.J., Helmut, F., Emden V. and Battersby M. (1995):**
A biochemical mechanism of resistance to pirimicarb two glass house clones of *Aphis gossypii*. Pestic. Sci., 43: 21-29.
- Singh, D.S. Dhingra, S., Srivastava, V.S., Sircar, P. and Rattanlal (1980):**
Relative susceptibility of *Aphis craccivora* Koch, to pesticides in relative to different hosts. Indian. J. Ent. 42(4): 746-756.
- Smith J.E. and Yu S.J. (1991):**
Insecticide resistance in the fall armyworm. *Spodoptera frugiperda*. Pesticide. Biochemistry and physiology, 39, 84 - 91.
- Soedomom, P. and Sunaryono, H. (1988):**
Seed storage of bushita with several kinds of plant pesticide treatments, Buletin - penelitan. Hort Kultura, 18 : 24 - 28.
- Stein, U. and Klingauf F. (1990):**
Insecticidal effect of plant extracts from tropical and subtropical species. Traditional methods are good as long as they are effective. J. Appl. Ent. 110 - 160.
- Stribley, M.F. Moores, G.D.; Devonshire A.L.; and Sawicki R.M. (1983):**
Application of the FAO - recommended method for detecting insecticide resistance in *Aphis fabae* Scopoli, *Sitobin avenae* (F). *Metopolophium dirhodum* (walker) and *Rhopalosiphum padi* (U) (Hemiptera : Aphididae). Common health Agricultural Bureaux.
- Su, H. C.F. (1972):**
Toxicity of citrus oils to several stored products insects : Laboratory evaluation. J. Econ. Entomol., 65 : 1438.
- Su, H.C.F. (1991):**
Toxicity and repellency of chenopodium oil four species of stored product insects. J. Ent. Sci., 26 : 178 - 182.
- Su, H., and Horvat R. (1977):**
Insecticidal properticidal properties of black pepper to rice weevils and cowpea weevils. J. Econ. Entomol., 70 : 19 - 21.

Thapliya, P.N.; Dubey, K.S. and Bhadula, H.K. (1987):

Control of yellow mosaic of soybean with granular insecticides. Indian phytopathol, 40 (1) : 10 - 11.

Thurston, R. and Webster J. A. (1962):

Toxicity of *Nicotiana gossei* (Domin) to *Myzus persicae* (Sulzer). Ent. Expand Appl., 5 : 233 - 238.

Uk, S. and Dittrich, V. (1986):

The behaviour modifying effect of the adult whitefly *Bemisia tabaci* (Genn.) which attack cotton in Sudan crop prot 5 (5) : 341 - 347.

Vidyasekhar, S.M. Reddy, G.P.V., Murthy, M.M.K. and Nagalingam, B. (1989):

Efficacy of certain newer insecticides against the sucking pests of Radifed cotton. Pesticides 23 (8) : 45 - 48.

Wada, J. and Munakata L. (1968):

Naturally occurring insect control chemical inolboldine a feeding inhabitant conceculolike insecticide in leaves of *cocculus tribolus* D.C. J. Agric. Fd. Chem., 16 : 471 - 474.

Wang, Y. Toyota M., Krause F., Hamburger M. and Hostettmann, K. (1990):

Antifungal and larvicidal polyacetylenes from *Artemisia borealis* planta Medica 56.

Wardlow, L.R, Ludlam, F.A.B and Hammcn, R.P (1975):

Comparison of the effectiveness of insecticides against glasshouse whitefly (*Trialeurodes vaporariorum*) Ann. Appl. Biol 81 433-435.

Yassin, K.M.; Bashir, N.H.H. and Gadalla, B.H. (1990):

Effect of endosulfan, chlorpyrifos and their mixtures on *Bemisia tabaci* of Sudan Gezira. Tropical Pest management 36 (3) : 230 - 233.

Yee W., Toscano N., Hendrix D., and Eberry T. H.

(1998):

Effects of insecticide applications on *Bemisia argentifolii* (Homoptera:Aleyrodidae) densities and honeydew production. Environ Entomol 27 [1] 22-32.