

VIII. LITERATURE CITED

Aitken, T. H. G.

- (1977) An in vitro feeding technique for artificially demonstrating virus transmission by mosquitoes.
Mosq. News, 37 (1): 130-133.

Al-Mashhadani, H. M., Davidson, G. and Curtis C. F.

- (1980) A genetic study of the susceptibility of Anopheles gambiae to Plasmodium berghei.
Trans. Roy. Soc. Trop. Med. Hyg., 74 (5): 585-594.

Backhouse, T. C., and Woodhill, A. R.

- (1954) Mosquito-born diseases, Filariasis, Wuchereria bancrofti from New Caledonia in relation to certain scutellaris group mosquitoes, Noumea.
South Pacific Commission Technical Information Circular, No. 11.

Bianchi, U.

- (1969) Genetical aspect of an esterase in the larva of Anopheles stephensi.
Atti. Ass. Genet. Ital., 14: 139-140.

Boyed, M. F., and Russell, J. C.

- (1943) Preliminary observations on the inheritance of susceptibility to malaria infection as a character of Anopheles quadrimaculatus.
Say. Am. J. Trop. Med., 23, 451-457.

Bryan, J. H. and Southgate, B. A.

- (1976) Some observations in filariasis in Western Samoa after mass administration of diethylcarbazine.
Trans. Roy. Soc. Trop. Med. Hyg., 70: 39-48.

Chen, C. C. and Laurence, B. R.

- (1987) Selection of a strain of Anopheles quadrimaculatus with high filaria encapsulation rate.

- J. Parasitol., 73 (2): 418-419.
- Collins, F. H., Sakai, R. K. and Vernick, K. D.
(1986) Genetic selection of a Plasmodium-refractory strain of the malaria vector Anopheles gambiae.
Science, USA, 234 (Oct. 31) 607-610.
- Coluzzi, M. and Cancrini, G.
(1974) Genetica della suscettibilita di Aedes aegypti a Dirofilaria repens.
Parassitologia 16: 239-256.
- Commission on Biochemical Nomenclature.
(1972) ENZYME NOMENCLATURE.
American Elsevier Publ. Co., New York, U.S.A.
- Craig, G. B., Jr.
(1964) Applications of genetic technology to mosquito rearing.
Bull. Wld Hlth Org., 31, 469-473.
- Craig, G. B., Jr and Hickey, W. A.
(1967) Genetics of Aedes aegypti. In "Genetics of Insect Vectors of Disease" (J. W. Wright and R. Pal, eds.), pp. 67-131. Elsevier Publishing Co., Amsterdam.
- Curtis, C. F. and Graves, P. M.
(1983) Genetic variation in the ability of insects to transmit filariae, trypanosomes and malarial parasites.
Advances to parasitology, 1, 31-62.
- Curtis, C. F., Ellis, D. S., Doyle, P. E., Ramji, B. D.,
(1983) Hill, N., Irungu, L. W., and Townson, H.
Susceptibility of aposymbiotic Culex quinquefasciatus to Wuchereria bancrofti.
J. Invert. Path., 41: 214-223.

- Curtis, C. F., Kihamia, C. M. and Ramji, B. D.
(1981) Tests of susceptibility of Liberian Culex quinquefasciatus to Tanzanian Wuchereria bancrofti.
Trans. Roy. Soc. Trop. Med. Hyg. 75: 736-739.
- De Meillon, B., Hayashi, S. and Sebastian, A.
(1967) Infection and reinfection of Culex pipiens fatigans with Wuchereria bancrofti and the loss of mature larvae in blood-feeding.
Bull. Wld Hlth Org., 36, 81-90.
- Desowitz, R. S., and Chellappah, W. T.
(1962) The transmission of *Brugia* sp. through Culex pipiens fatigans: the effect of age and prior non-infective blood meals on the infection rate.
Trans. Roy. Soc. Trop. Med. Hyg., 56, 121-125.
- Duhrkopf, R. E. and Trpis, M.
(1980) The degree of susceptibility and levels of infection in ten different strains of Aedes polynesiensis marks infected with subperiodic Brugia malayi and Brugia pahangi.
Am. J. Trop. Med. Hyg., 29 (5): 815-819.
- El-Dine, K. Z.
(1966) Preliminary studies on autogeny in Culex pipiens complex.
J. Egypt. Pub. Hlth. Assoc., 41: 115-126.
- Ewart, A.
(1967) Studies on the transfer of infective Brugia pahangi larvae from vector mosquitoes to the mammalian host.
Trans. Roy. Soc. Trop. Med. Hyg., 61: 110-113.
- Ford, E. B.
(1965) Genetic Polymorphism.
Lond.: Faber and Faber.

- Frizzi, G., Rinaldi, A. and Bianchi, U.
(1975) Genetic studies on mechanisms influencing the susceptibility of Anopheline mosquitoes to Plasmodial infection.
Mosq. News, 35 (4): 505-508.
- Gad, A. M.
(1963) Insects of medical importance "notes in arabic"
Institute of medical entomology, Ministry of Public Health, Dokki, Cairo, Egypt.
- Gelfand, H. M.
(1955 a) Studies on the vectors of Wuchereria bancrofti in Liberia.
Am. J. Trop. Med. Hyg., 4, 52-60.
- Gelfand, H. M.
(1955 b) Anopheles gambiae Giles and Anopheles melas Theobald in a coastal area of Liberia, West Africa.
Trans. Roy. Soc. Trop. Med. Hyg., 49, 508-527.
- Gubler, D. J. and Bhattacharya, N. C.
(1974) A quantitative approach to the study of bancroftian filariasis.
Am. J. Trop. Med. Hyg., 23, 1027-1036.
- Hardy, J. L., Apperson, G., Asman, S. M. and Reeves, W. C.
(1978) Selection of a strain of Culex tarsalis highly resistant to infection following ingestion of western equine encephalomyelitis virus.
Am. J. Trop. Med. Hyg., 27 (2): 313-321.
- Hariston, N. G. and De Meillon, B.
(1968) On the inefficiency of transmission of Wuchereria bancrofti from mosquito to human host.
Bull. Wld Hlth Org., 38, 935-941.

Hertig, A. T. and Hertig, M.

- (1927) A technique for the artificial feeding of sandflies (*Phlebotomus*) and mosquitoes.
Science, 65: 328-329.

Hertig, M. and Mc Connell, E.

- (1963) Experimental infection of Panamanian *Phlebotomus* sandflies with *Leishmania*.
Exp. Parasitol., 14: 92-106.

Hovanitz, W.

- (1947) Physiological factors which influence the infection of *Aedes aegypti* with *Plasmodium gallinaceum*
Am. J. Hyg., 45, 67-81.

Huff, C. G.

- (1929) The effects of selection upon susceptibility to bird malaria in *Culex pipiens* Linn.
Ann. Trop. Med. Parasitol., 23, 427-442.

Huff, C. G.

- (1931) The inheritance of natural immunity to *Plasmodium cathemerium* in two species of *Culex*.
J. Prev. Med., (Baltimore), 5: 249-259.

Ibrahim, M. S. and Trpis, M.

- (1987) The effect of *Brugia paphangi* infection on survival of susceptible and refractory species of the *Aedes scutellaris* complex.
Med. Vetr. Entomol. 1: 329-337.

Jeffery, G. M.

- (1944) Investigation of the factors influencing the mosquito transmission of *Plasmodium lophurae*.
Doctorate dissertation: Johns Hopkins Univ., Baltimore.

Kartman, L.

- (1953) Factors influencing infection of the mosquito with Dirofilaria immitis (Leidy, 1856).
Exp. Parasitol., 2: 27-78.

Khalil, M.

- (1936) Filariasis and elephantiasis in Rosetta and the means of their effective control.
J. Egypt. Med. Ass., 19: 701-711.

Khalil, M., Halawani, A. and Hilmy, I. S.

- (1932) On the transmission of filariasis bancrofti in Egypt.
J. Egypt. Med. Assoc., 15:317.

Khalil, H. M., Rifaat, M. A., Gad, A. M. and Sadek, S.

- (1974) Filarial infectivity rate of Culex pipiens molestus subjected to sub-lethal concentrations of insecticides abote and sevin and distribution of infective filaria larvae in mosquito body regions.
J. Egypt. Publ. Hlth. Assoc., 4-9(4/5): 221-224.

Kilama, W. L. and Craig, G. B., Jr.

- (1969) Monofactorial inheritance of susceptibility to Plasmodium gallinaceum in Aedes aegypti.
Ann. Trop. Med. Parasitol., 63 (4): 419-432.

Loose, A.

- (1916) Mense Handbuck der Tropenkrankheiten I eipxig.

Macdonald, W. W.

- (1962 a) The selection of a strain of Aedes aegypti susceptible to infection with semi-periodic Brugia malayi.
Ibid., 56, 368.

Macdonald, W. W.

- (1962 b) The genetic basis of susceptibility to infection with semi-periodic Brugia malayi in Aedes aegypti. Ann. Trop. Med. Parasitol., 56: 373-382.

Macdonald, W. W.

- (1963 a) Further studies on a strain of Aedes aegypti susceptible to infection with sub-periodic Brugia malayi. Ann. Trop. Med. Parasitol., 57: 452-460.

Macdonald, W. W.

- (1963 b) A preliminary cross-over value between the gene f^m (filarial susceptibility, Brugia malayi) and the sex locus in Aedes aegypti. Ann. Trop. Med. Parasitol., 57: 461-465.

Macdonald, W. W.

- (1965) Factors affecting the transmission of filariae: the effect of parasitization by Brugia malayi in laboratory populations of Aedes aegypti. In premier Congres International de Parasitologie, Rome, Paris, ORSTOM (Cahiers ORSTOM Entomologie medicale, 3, 111-113).

Macdonald, W. W.

- (1967) The influence of genetic and other factors on vector susceptibility to parasites. In "Genetics of Insect Vectors of Disease" (J. W. Wright and R. Pal, eds.), pp. 567-584. Elsevier Publishing Co., Amsterdam.

Macdonald, W. W.

- (1972) Sub-Department of Entomology. In: The Incorporated Liverpool school of Tropical Medicine. 72. Ann. Rep. 1970-71, 78-80.

- Macdonald, W. W., and Ramachandran, C. P.
(1965) The influence of the gene f^m (filarial susceptibility, Brugia malayi) on the susceptibility of Aedes aegypti to seven strains of Brugia, Wuchereria and Dirofilaria.
Ann. Trop. Med. Parasitol., 59: 64-73.
- Macdonald, W. W., and Sheppard, P. M.
(1965) Cross-over values in the sex chromosomes of the mosquito Aedes aegypti and evidence of the presence of inversions.
Ann. Trop. Med. Parasitol., 59: 74-87.
- Magayuka, S. A. and White, G. B.
(1971) Studies on the mosquito Culex pipiens fatigans Wiedemann in East Africa: Hybrid compatibilities and susceptibility to Wuchereria bancrofti (Cobbold) of six populations.
Bull. W.H.O., 46: 801-808.
- Mahdi, A. H., Wassif, S. F. and Gad, A. M.
(1969) Biological studies of Culex pipiens in the Nile Delta. Part I. Screening to filarial infection.
J. Egypt. Pub. Hlth. Ass., 44, 189-192.
- McGreevy, P. B., McClelland, G. A. H. and Lavoipierre, M. M. J.
(1974 a) Inheritance of susceptibility to Dirofilaria immitis infection in Aedes aegypti.
Ann. Trop. Med. Parasitol., 68, 97-109.
- McGreevy, P. B., Theis, J. H., Lavoipierre, M. M. J. and Clark, J.
(1974 b) Studies on filariasis. III. Dirofilaria immitis: emergence of infective larvae from the mouth parts of Aedes aegypti.
J. Helminthol., 48, 221-228.

Micks, D. W.

- (1949) Investigations on the mosquito transmission of Plasmodium elongatum Huff, 1930.
J. nat. Malar. Soc., 8: 206-218.

Miles, M. A., Lanham, S. M., De Souza, A. A. and Povia, M.

- (1980) Further enzymic characters of Trypanosoma cruzi and their evaluation for strain identification.
Trans. Roy. Soc. Med. Hyg., 74 : 221-237.

Milton, J. S. and Tsokos, J. O.

- (1983) Statistical methods in the biological and health sciences.
Kosaido printing Co., LTD. Tokyo, JAPAN.

Obiamiwe, B. A. and Macdonald, W. W.

- (1973) Evidence of a sex-linked gene, sb, controlling susceptibility of Culex pipiens to Brugia pahangi.
Trans. Roy. Soc. Trop. Med. Hyg. 67: 32-33.

Partono, F. and Oemijati, S.

- (1970) Susceptibility of Culex pipiens fatigans to Wuchereria bancrofti in Djakarta, Indonesia.
Southeast Asian J. Trop. Med. Public Health 1: 516-518.

Raghavan, N. G. S., Das, M., Mammen, M. L., Singh, N. N. and Wattal, B. L.

- (1967) Genetic basis of differential susceptibility of Aedes aegypti to dirofilarial infection.
Part 1: Preliminary observations on selection of Aedes aegypti strains susceptible and refractory to Dirofilaria immitis infection.
Bulletin of the Indian Society for Malaria and other Communicable Diseases, 4, 318-323.

- Ramachandran, C. P., Edeson, J. F. B., and Kershaw, W.
(1960) E.
Aedes aegypti as an experimental vector of Brugia malayi.
Ann. Trop. Med. Parasitol., 54: 371-375.
- Rifaat, M. A., Hassan, Z. A., Sidky, H. S. A. and Gaafar,
(1978) S. A.
Studies on filariasis in Egypt, epidemiological
and entomological studies in an endemic village.
J. Egypt. Soc. Para., 8 (1): 95-101.
- Rockefeller Foundation.
(1948) International Health Division Annual Report,
pp. 6-7.
- Rockefeller Foundation.
(1950) International Health Division Annual Report,
pp. 29-31.
- Rodriguez, P. H.
(1973) Differential development of Brugia pahangi in
laboratory strains of Aedes aegypti.
J. Med. Ent., 10 (2): 194-197.
- Rodriguez, P. H. and Craig, G. B.
(1973) Susceptibility to Brugia pahangi in geographic
strains of Aedes aegypti.
Am. J. Trop. Med. Hyg. 22: 53-61.
- Rosen, L.
(1955) Observations on the epidemiology of human filar-
iasis in French Oceania.
Am. J. Hyg., 61:219.
- Roubaud, E.
(1937) Nouvelles recherches sur l'infection du moustique
de la fièvre jaune par Dirofilaria immitis Leidy:
Les races biologiques d' Aedes aegypti et l'infe-

- ction filarienne.
Bull. Soc. Path. exot., 30: 511.
- Roubaud, E., Colas-Belcour, J., Toumanoff, C., and Treillard, M.
(1936) Recherches sur la transmission de Dirofilaria immitis Leidy, Bull. Soc. Path. exot., 29: 1111-1120.
- Sadek, S., Gad, A. M., Rifaat, M. A. and Khalil, H. M.
(1975) Effect of diethylcarbamazine given to filarial patients on the filaria cycle in Culex pipiens molestus mosquitoes.
J. Egypt. Publ. Hlth. Assoc., L (1): 43-48.
- Samarawickrema, W. A., Spears, G. F. S., Sone, F., Ichimori, K. and Cummings, R. F.
(1985) Filariasis transmission in Samoa. 1. Relation between density of microfilariae and larval density in laboratory - bred and wild - caught Aedes (Stegomyia) Polynesiensis (Marks) and wild - caught Aedes (Finlaya) samoanus (Grueenberg).
Ann. Trop. Med. Parasitol., 79, 89-100.
- Sasa, M.
(1976) Human filariasis
A global survey of epidemiological and control.
Univ. Park Press, pp. 819.
- Shawarby, A. A., Mahdi, A. H., Naguib, K. and Moharram, A.
(1965) Incidence of filariasis in Egypt.
J. Egypt. Pub. Hlth. Assoc., 40 (4): 267-282.
- Singh, K. R. P. and Curtis, C. F.
(1974) Attempt to select a strain of Culex pipiens fatigans non-susceptible to infection with periodic Wuchereria bancrofti.

J. Commun. Dis. 6: 88-90.

Smithies, O.

- (1955) Zone electrophoresis in starch gels: group variations in the serum protein of normal human adults. Biochem. J., 61: 629-641.

Soliman, B. A.

- (1985) Field studies on the transmission of bancroftian filariasis by mosquitoes.
M. Sc. thesis, Ain Shams Univ. Cairo.

Southgate, B. A.

- (1979) Bancroftian filariasis in Egypt.
Tropical diseases bulletin, 76 (12): 1045-1068.

Spielman, A.

- (1979) Autogeny in Culex pipiens populations in nature: effects of inbreeding.
Ann. Entomol. Soc. Amer., 72: 826-828.

Steiner, W. M. M. and Joslyn, D. J.

- (1979) Electrophoretic techniques for the genetic study of mosquitoes.
Mosq. News, 39: 35-54.

Terwedow, H. A. and Craig, G. B., Jr.

- (1977) Waltonella flexicauda: Development controlled by a genetic factor in Aedes aegypti.
Exp. Parasitol. 41: 272-282.

Thomas, V. and Ramachandran, C.P.

- (1970) Selection of Culex pipiens fatigans for vector ability to the rural strain of Wuchereria bancrofti - A preliminary report.
Med. J. Malaya, 24: 196-199.

Townson, H.

- (1971) Mortality of various genotypes of the mosquito

Aedes aegypti following the uptake of microfilariae of Brugia pahangi.

Ann. Trop. Med. Parasitol., 65: 93-106.

Townson, H.

(1974) The development of Brugia pahangi in male Aedes aegypti of 'refractory' genotype.

Ann. Trop. Med. Parasitol., 68 (2) : 239-240.

Townson, H., Sulaiman, I. and Matthews, H. A.

(1981) Genetic aspects of the susceptibility of Aedes aegypti to Dirofilaria and Brugia.

Trans. Roy. Soc. Trop. Med. Hyg. 75: 175-176.

Trager, W.

(1942) A strain of the mosquito Aedes aegypti selected for susceptibility of the avian malaria parasite Plasmodium lophurae.

J. Parasitol., 28: 457-465.

Van der Kaay, H. J. and Boorsma, L.

(1977) A susceptible and refractive strain of Anopheles atroparvus to infection with Plasmodium berghei berghei.

Acta Leidensia, 45, 13-20.

Wallis, G. P., Aitken, T. H. G., Beaty, B. J., Lorenz, L., Amato, G. D. and Tabachnick, W. J.

(1985) Selection for susceptibility and refractoriness of Aedes aegypti to oral infection with yellow fever virus.

Ann. J. Trop. Med. Hyg., 34 (6): 1225-1231.

Ward, R. A.

(1963) Genetic aspects of the susceptibility of mosquitoes to malarial infection.

Exp. Parasitol., 13, 328-341.

Wassif, S. F.

- (1969) Survey of the Egyptian Culicines in the Nile Delta with special reference to filaria transmission.
M. D. thesis, Ain Shams Univ. Cairo.

Weber, K. and Osborn, M.

- (1969) The reliability of molecular weight determinations by dodecyl sulfate polyacrylamide gel electrophoresis.
J. Biol. Chem., 224: 4406-4412.

Weinstein, P. P.

- (1986) Filariasis: problems and challenges.
Am. J. Trop. Med. Hyg., 35 (2): 221-233.

Wharton, R. H.

- (1960 a) Studies on filariasis in Malaya: Field and laboratory investigations of the vectors of a rural strain of Wuchereria bancrofti.
Ann. Trop. Med. Parasitol., 54: 78.

Wharton, R. H.

- (1962) The biology of *Mansonia* mosquitoes in relation to the transmission of filariasis in Malaya.
Bull. Inst. med. Res. Malaya, No. 11.

Wharton, R. H., Laing, A. B. G., and Cheong, W. H.

- (1963) Studies on the distribution and transmission of malaria and filariasis among aborigines in Malaya.
Ann. Trop. Med. Parasitol., 57: 235-254.

W. H. O.

- (1965) Standardized strains of insects of public health importance. In "Genetics of Insect Vectors of Disease" (J. W. Wright and R. Pal, eds.),

pp. 735-766. Elsevier Publishing Co., Amsterdam.

WHO

- (1974) WHO expert committee on filariasis.
WHO, Geneva, pp. 50-51.

WHO

- (1975) Manual on practical entomology in malaria, Part II.
WHO, Geneva, pp. 55-66.

Wright, S.

- (1952) The genetics of quantitative variability.
London, Agr. C. Res. Council. H. M. S. O., 151.

Zielke, E.

- (1973 a) Untersuchungen zum Mechanismus der Filarienuber-
tragung bei Stechmücken.
Zeitschrift für Tropenmedizin und Parasitol.,
24: 32-35.

Zielke, E.

- (1973 b) Untersuchungen zur verebung der Empfanglichkeit
gegenüber der hundefilarie Dirofilaria immitis
bei Culex pipiens fatigans and Aedes aegypti.
Zeitschrift für Tropenmedizin und Parasitol.,
24: 36-44.

Zielke, E. and Kuhlow, F.

- (1977) On the inheritance of susceptibility for infec-
tion with Wuchereria bancrofti in Culex pipiens
fatigans.
Tropenmed. Parasitol. 28: 68-70.