

REFERENCES

REFERENCES

- Abdel-Salam, E., Hafez, A.A., Abdel-Azim, S., and
El-Saphori, L. (1982):
Alpha-1-antitrypsin in Bilharzial fibrosis
and normal controls.
Egyp. J. of Biochem.
- Aisen P. and Laitowsky, I., (1980):
Iron transport and storage proteins.
Annual Reviews of Biochemistry, 49: 357.
- Anderson, H.C., and Mc Carty, M. (1950):
Determination of c-reactive protein in the
blood as a measure of the activity of the
disease process in acute rheumatic fever.
Am. J. Med., 8: 445.
- Areekul, S., Chantachum, Y., and Matrakul, D. (1974):
Serum haptoglobin levels in patient with liver
diseases.
J. Tropical Med. Public Health, 5 (4): 601.

Arima, T., Motoyama, Y., Yamamoto, t., Nagata, K., and
Kondo, T. (1977):

Serum glycoprotein in liver diseases. Alpha-
1-acid glycoprotein in liver cirrhosis.
Gastroenterol. Jpn., 12 (2): 43.

Aronsen, K.F., Ekelund, G., Kindmark, C.O., Laurell,
C.B. (1972):

Sequential changes of plasma proteins after
surgical trauma. Scand. J. Clin. Lab. Invest.
, 29 : 127.

Balldin, G., Laurell, C.B. and Ohlsson, K. (1978) :

Increased Catabolism of alpha-2-macroglobulin
after intravenous infusion of trypsin-alpha-1
-antitrypsin complexes in dogs. Hoppe-Seyler
Z. physiol. Chem., 359 : 699.

Barrett, A. J. and Starkey, P.M. (1973):

Inhibition by alpha-2-macroglobulin and other
serum proteins. Bioch. J., 131 : 823

Benakappa, D.G., (1983):

XV II th int. Congr. of ped.
Philippine. Etract of free papers,
Vol. 1, P 105.

Bodmer, B., and Siboo, R. (1977):

Isolation of mouse c-reaction protein from
liver and serum.

J. of Immunology, 118 (3): 1086.

Bohm, N. (1980):

Alpha-1-antitrypsin and its deficiency states
Path. Res. Pract., 168: 1.

Bourrillon, R., Razafinahaleo, E.A and Parraudeau, M.

(1968) : Etudes sur la structure de l'alpha-2-
macroglobuline humaine (α 2 - M). II.

Preparation et proprietes des glycopeptides
isoles d'un hydrolysat enzymatique d'alpha-2-
M. Biochem.

Biophys. Acta, 154 : 405.

Calhoun, D.B., and Englander, S.W. (1979):

Measurement of haptoglobin concentration in
plasma and other fluids by a simple spectro-
metric procedure.

Analytical Biochem., 99: 421.

Carlson, J., Eriksson, S., and Hagerstrand, I. (1981):

Intra- and extra-cellular alpha-1-antitrypsin
in liver disease with special reference to Pi
phenotype. J. Clin. Pathol., 34: 1020.

Chapelle, J.P., Albert, A., Smeets, J.P., et al.
(1981):

The prognostic significance of serum alpha-1-
acid glycoprotein changes in acute myocardial
infarction.

Clin. Chim. Acta (Netherlands), 115 (2): 199.

Charles, W.G., Craig, L.F. and Carolinda, L.H. (1971):

Rapid method for protein quantitation by
electro immunodiffusion. Clin. Chem.,
17/6 : 501.

Chiu, K.M., Mortensen, R.F., Osmand, A.P. and Gewurz, H.

(1977) : Kinetics of association of serine
proteinases with native and oxidized alpha-1-
proteinase inhibitor and alpha-1
antichymotrypsin.

J. Biol. Chem., 256 : 3931.

Clarence, A.G. & Martin, H.W. (1977):

Chronic Respiratory Tract infection, pulmonary
medicine, P. 315.

Claus, D.R. Osmand, A.P., and Gewurz, H. (1976):

Radioimmunoassay of human c-reactive protein
and levels in normal sera.

J. Lab. Clin. Med., 87 (1): 120.

Connell, G.E., and Smithies, O. (1959):

Human haptoglobins : estimation and
purification.

Biochem. J., 72: 115.

Courtoy, P.J., Lombart, C., Feldmann, G., Mouguilevsky,
N. and Rogier, E. (1981) :

Synchronous increase of four acute-phase
proteins synthesized by the same hepatocytes
during the inflammatory reactions. Int. Acad.
Path., 44/2 : 105.

Craig, L., Fischer, M., Gill, C.W. (1980) :

C-reactive protein (CRP). Am. J. Clin. Path.,
75/1 : 22.

Crockson, R.A., Payne, C.J., Ratcliff, A.P., et al.
(1966):

Time sequence of acute phase reactive proteins
following surgical trauma.
Clin. Chim. Acta., 14 : 435.

Crofton, J. & Douglas, A., (1981) :

Respiratory Diseases, 3rd Ed.

Blackwell scientific publications, London.

Daniel, T.M., Balestrino, E.A., Balestrino, O.C.,

Davidson, P.T., Debanne, S.M., Kataria, S.,

Kataria, Y.P. & Scolozza, J.P., (1982) :

The tuberculin specificity in humans of

Mycobacterium T.B. Antigens.

Am. Rev. chest. Dis. Vol. 126, No. 4.

P. 600-606.

David, R., Claus, A.P., Osmand, A.P. and Gewurz, H.

(1976) :

Radio-immunoassay of human C-reactive protein

and levels in normal sera. J. lab. Clin. Med.,

87/1 : 120.

De Beer, F.C., Nel, A.E. and Gie, R.P., (1984) :

Serum amyloid-A-protein and C-reactive protein

levels in pulmonary tuberculosis :

Relationship to amyloidosis. Thorex, 39/3:196.

De Klerk, E. and Anderson, R. (1982) :

Accurate laboratory measurement of serum

C-reactive protein levels. J. SA Med.,

62 : 1030.

Denko, C.W., and Gabriel, P. (1979) :

Serum proteins-transferrin, ceruloplasmin,
alpha-1-acid glycoprotein, alpha-1-antitrypsin
in rheumatic diseases.

J. Rheumatol. (Canada), 6 (6) : 664.

Dermer, G.B., Silverman, L.M., Gendler, S.J. and Tokes,

Z.A. (1980) : Incidence of a split alpha-2-
glycoprotein band in the electrophoretic
pattern for serum of adenocarcinoma patients.
Clin. Chem., 26 : 392.

Deshmukh, M.D., (1972) :

T.B. In children, pulmonary T.B. & common
chest diseases, 1st ed., P. 9.

Unichem Laboratories, LTD.

Bombay - India.

De-Vonne, T.L. and Mouray, H., (1978) :

Human alpha-macroglobulin and its antitrypsin
and antithrombin activities in serum and
plasma Clin. Chim. Acta., 90 : 83.

Doolittle, R.F., (1973) :

Structural aspects of the fibrinogen to fibrin conversion. In : Anfinsen C.B., Edsall J.T., Richards F.M. (eds). Advances in protein chemistry.

Academic Press, New York, vol. 27, P. 1 - 109.

Edwards, D.I., Lalka, D., Cerra, F., and Slaughter, R.L.

(1982) :

Alpha-1-acid glycoprotein concentration and protein binding in trauma.

Clin. Pharmacol. Ther., 31 (1) : 62.

El-Hefny, A., Samuel, S. and Guirguis, I. (1982) :

2- T.U. PPD : A comparison test between a group of normal BCG vaccinated infant and children and a group of tuberculous children. Egypt. J. Chest Dis. and Tuber.

Elson, E.C., (1974) :

Quantitative determination of serum haptoglobin.

A simple and rapid method.

Am. J. Clin. Pathol., 62 : 655.

Eriksson, S., (1965) :

Studies in alpha-1-antitrypsin deficiency.

Acta. Med. Scand, 177/432 : 1.

Felix, N.S., Nakajima, H., and Kagan, B.M. (1966) :

Serum c-reactive protein in infections during
the first 6 months of life.

Pediatrics, 37 (2) : 270.

Fischer, C.L., and Gill, C.W., (1975) :

Acute phase proteins. Serum protein
abnormalities. Determination and clinical
aspects. Am. J. Clin. Path., 66 (5) : 331.

Fisher, R.L., Taylor, L. and Sherlock, S. (1976) :

Alpha-1-antitrypsin deficiency in liver
disease : The extent of the problem.

Gastroenterology, 71 : 646.

Fishman, A.P., (1980) :

Pulmonary diseases & disorders. vol. 11,

P. 1264 - 1323.

Mc-Crow Hill Book Co.

New York - Toron'to - Sydney - London.

For Far, J. & Gawin A., (1984) :

Diseases due to infection.

Textbook of pediatries

3rd ed., P. 1391 : 1406.

Churchill Living Stone, - London.

Fredic, B. & Donald, B., (1978) :

Tuberculosis, children in health & disease.

P. 685 - 701.

Bailliere Tindall - London.

Friedrich, K., and Black, L.F., (1974) :

Alpha-1-antitrypsin and its deficiency.

Am. Review of Resp. Disease, 110 : 176.

Gahmberg, C.G., and Andersson, L.C., (1978) :

Leukocyte surface origin of human alpha-1-acid
glycoprotein (orosomuroid).

J. Exp. Med., 148 (2) : 507.

Ganrot, P.O. and Schersten, B., (1967) :

Serum alpha-2-macroglobulin concentration and
its variation with age and sex. Clin. Chim.

Acta, 15 : 113.

Ganrot, P.O. and Bjerre, B., (1967) :

Alpha-1-antitrypsin and alpha-2-macroglobulin
concentration in serum during pregnancy. Acta.
Obstet. Gynecol. Scand., 46 : 126.

Ganrot, P.O. and Bjerre, B., (1978) :

Alpha-1-antitrypsin and alpha-2-macroglobulin
concentration in serum during pregnancy, Acta.
Obstet. Gynecol. Scand., 46 : 126.

Gatner, E.M. and Anderson, R., (1980) :

An in-vitro assesement of cellular and
humoral immune function in pulmonary
tuberculosis :

Correction of defective neutrophil motility by
ascorbate, levimasole, metoprolol and
propranolol.

Clin. Exp. Immunol., 40 : 327.

Gatner, E.M., (1982) :

Circulating Immunoglobulin levels in patients
with pulmonary tuberculosis with special
reference to IgE. Tubercle 63/2 : 113.

Gerbeaux, J., Back Lard, A. & Couvereur, J., (1965) :

Am. J. Dis. child., Vol. 110 No. 507.

Ghanem, M.H., Guirgis, F.K., and El-Sawy, M., (1973) :

serum glycoproteins in schistosomal hepatic fibrosis.

J. Trop. Med. Hyg., 76 : 13.

Gobbi, P.G., Merlini, G., Parrunello, G.A., Cavalli, P., and Ascari, E., (1982) :

Serum alpha-1-acid glycoprotein, haptoglobin and C3 in Hodgkin's disease. A comparison with other acute phase indicators.

Acta Haematol. (Basel) (Switzerland), 67 (4) : 255.

Gordon, T., (1978) :

The acute-phase plasma proteins. In plasma protein turnover. Feder. Proc. 37 : 2754.

Gottschalk, A. and Beddecke, E., (1972) :

Occurance and distribution of glycoproteins
Elsevier Publishing Company, New York, London.

Grange, J.M., Kardjito, T., (1980) :

A study of antibodies to mycobacterium tuberculosis in the IgG, IgA & IgM. Classes in tuberculosis, Sarcoidosis and Crohn's disease.
Tubercle, 61/3 : 145.

Grange, J.M., Kardjito, T. and Setiabudi, I., (1984) :

A study of acute phase reactant proteins in
Indonesian patients with pulmonary
tuberculosis.

Tubercle, 65 : 23.

Gray, C.H., and Howorth, P.J.N., (1980) :

Plasma protein and immunoglobulin.

Clinical Chemical Patholo., 66.

Guilbert, L.J. and Iscove, N.N., (1976) :

Partial replacement of serum by selenite,
transferrin and lecithin in haemopoietic cell
cultures.

Nature, 263 : 594.

Gyton, A.C., (1981) :

Textbook of Medical physiology, 6th edition.

W.B. Saunders Company, P. 98.

Hahn, D. and Ganzoni, A.M., (1975) :

Functional heterogeneity of the transport iron
compartement. Acta Haematol., 53 : 321.

Hall, P.K. and Roberts, R.C., (1978) :

Physical and chemical properties of human
plasma alpha-2-macroglobulin. Biochem. J.,
171 : 27.

Hamaguchi, H., Isomoto, A., Miyake, Y., and Nakajima,
H., (1971) :

Some spectral properties of the human
hemoglobin - haptoglobin complex.
Biochemistry, 10 (10) : 1741.

Harpel, P.C., (1973) :

Studies on human plasma alpha-2-macroglobulin-
enzyme interactions : Evidence for proteolytic
modification of the subunit chain structure.
J. Exp. Med., 138 : 508.

Harris, D.C. and Aisen, P., (1975) :

Functional equivalence of the two iron binding
sites of human transferrin. Nature, 257 : 821.

Harris, R.I., Stone, P.C., Hudson, A.G., and Stuart, J.
(1984) :

C-reactive protein rapid assay techniques for
monitoring resolution of infection in immuno-
suppressed patients. J. Clin. Path., 37 : 821.

Heide, K., Heimbürger, N. and Haupt, H. (1965) :
Bausteinanalysen von. Humanserumproteinen.
Clin. Chim. Acta, 11 : 82.

Heimbürger, N., (1974) :
Biochemistry of proteinase inhibitors from
human plasma : A review of recent development.
In : Fritz H., Teschesche H., Greene L.J.,
Truscheit E., eds. Bayer symposium V.
Proteinase inhibitors.
New York, Springer-Verlag, 14 : 22.

Hemmaplårdh, D. and Morgan E.H. (1974) :
The mechanism of iron exchange between
synthetic iron helators and rabbit
reticulocytes.
Biochem. Biophys. Acta, 373 : 84.

Henriksen, H.J., Petersen, M.U., and Pedersen, F.B.,
(1982) :
Serum alpha 1-acid glycoprotein (orosomucoid)
in uremic patients on hemodialysis.
Nephron (Switzerland), 31 (1) : 24

Hinshow, H.C. & Murry, J.F., (1980) :

Tuberculosis, Diseases of the chest. P.
298 - 349.

W.B. Saunders Co.

Philadelphia - London, Toronto.

Hoeprich, P.D., (1977) :

Infectious Diseases.

2nd ed. P. 343.

Harper & Row Publishers.

New York, San Francisco, London.

Horne, C.H., Brigys, T.P., Howie, P.W. and Kennedy, A.C.

(1972) :

Sequential changes of the plasma protein
pattern in inoculation hepatitis. Scand J.
Lab. Clin.

Invest., 29 : 105.

Hously, J., (1968) :

Evidence for alpha-1-acid glycoprotein
populations of different pI values after
concavalin-A affinity chromatography.

Biochim. Biophys. Acta, 668 : 235.

Huebers, H., Bauer, W., Huebers, E. and Finch, C.

(1981) :

The behaviour of transferrin iron in the rat.

Blood, 57 : 218.

Hurlimann, J., Thorbecke, G.J., and Hochwald, G.M.

(1966) :

The liver as the site of c-reactive protein
formation.

J. Exp. Med., 123 : 365.

Hutchings, S.E. and Sato, G.H., (1978) :

Growth and maintainance of hela cells in serum
free medium supplemented with hormones. Proc.
Nat. Acad. Scie., 75 : 901.

Ikenaka, T., Ishiguro, M., Isemura, S., Kaufmann, H.,

Bauer, W., and Schmid, K. (1971) :

The amino acid sequence of the carboxyl-
terminal CNBr-fragment of alpha 1-acid
glycoprotein.

Biochemical and Biophysical Research Communica
., 42 (6) : 1142.

Jackson, P.R., Tucker, G.T., and Woods, H.F. (1982) :

Altered plasma drug binding in cancer role of
alpha 1-acid glycoprotein and albumin.

Clin. Pharmacol. Ther., 32 (3) : 295.

Jones, M.C. and Thomas, G.O. (1971) :

Alpha-1-antitrypsin deficiency and pulmonary
emphysema. Thorax, 26 : 652.

John, J.M. (1972) :

Biochem. J., 127 : 187.

Kaplan, M.H. and Volkanakis, J.E. (1974) :

Interaction of C-reactive protein complexes
with complement system-1-consumption of human
complement associated with the reaction of CRP
with pneumococcal C-polysaccharide and with the
choline phosphatides. Lecithin and sphingomy-
elin. J. Immunol., 112 : 2135.

Kardjito, T. and Grange, J.M. (1980) :

Immunological and clinical features of smear-
positive pulmonary tuberculosis in East Java.
Tubercle, 61 : 231.

Katz, J.H. (1965) :

The delivery of iron to the immature. red cell
Seminars of Haematology, 6 : 15.

Khalil, I.F. (1984) :

Tuberculosis, communicable disease, synopsis
of public health.

Cairo University, Egypt.

Kluthe, R., Hagemann, U. and Kleine, N. (1967) :

U : Bestimmung des alpha-2-macroglobulinmstas
bein. Menschen. Vox Sang (Basel), 12 : 308.

Koj, A. (1974) :

Acute-phase reactants. Their synthesis, turn-
over and biological significance. In Structure
and Function of Plasma Proteins. Vol. 1, Ch. 4
, P. 73. Edited by A.C. Allison, Plenum Press,
London.

Krugman, S. & Katz, S. (1981) :

Infectious Disease in children.

7th ed. P. 455.

The C.V. Mosby Co. - Saint Louis.

Kueppers, F., Dickson, E.R., and Summerskill, W.H.J.

(1976) :

Alpha-1-antitrypsin phenotypes in chronic active liver disease and primary biliary cirrhosis.

Mayo Clin. Proc., 51 : 286.

Kushner, I., Volonakis, J., Gewurz, H. (1982) :

C-reactive protein and the plasma protein response to tissue injury. Ann. N.Y. Acad. Sci., 389 : 1.

Kuyas, C., Haeberli, A. and Straub, W. (1982) :

A subfraction of human fibrinogen with high sialic acid content and elongated o-Chains. J. Biol. Chemist., 257 : 1107.

Labib, M.F. (1976) :

Principles of Public Health Part II, 3rd Ed. Sherif's Bookshop, Cairo, P. 273.

Laurell, C.B. (1966) :

Quantitative estimation of proteins by electrophoresis in agarose gel containing antibodies Ann. Biochem., 15 : 45.

Laurell, C.B. and Jeppsson, J.O. (1975) :

"The plasma protein structure, function and genetic control". 2nd Ed. Edited by F.W. Patnam, Vol. I, P. 229, New York Academic Press.

Laurell, C.B. (1985) :

Acute-phase proteins- a group of protective proteins. Recent Advances In Clinical Chemistry, Ch. 4, P. 103, Edited by C.P. Price and K.G.H.M. Alberti
Churchill Livingstone.

Macleod, C.M., and Avery, O.T. (1941) :

The occurrence during acute infections of a protein not normally present in the blood. Isolation and properties of the reactive protein.

J. Exp. Med., 73 : 183.

Makie, P.H., Crockson, R.A. and Stuart, J. (1979) :

C-reactive protein for rapid diagnosis of infection in leukaemia. J. Clin. Pathol., 32 : 1253.

Mancini, G., Carbonara, A.O. and Heremans J.F. (1965) :
Immunochemical quantitation of antigens by
single radial immunodiffusion. Immunochem.,
2 : 235.

Maurice, K., Felicity, K. & Soebagyo, G. (1978) :
Primary child care.
1st ed. P. 159.
Oxford University Press - London.

Mc Conkey, B., Crockson, R.A., Crockson A.P., and
Wilkinson, A.R., (1973) :
The effects of some anti-inflammatory
drugs on the acute phase proteins in
rheumatoid arthritis.
Quarterly J. of Medicine, XL11 (168) : 785.

Miller, F.J.W. (1984) :
T.B. in childhood, Part I, Post Graduate
Doct., Middle East Vol. 7, No. 10
P. 90 : 95.

- Morelis, P., Ambrosioni, J.C., Got, R. and Fontages, R.
(1969) :
Observation au microscope electronique due
complexe forme par l'alpha-2-macroglobuline
de serum de lapin avec la trypsin. C.R. Acad.
Sci. Scr. D., 289 : 1453.
- Morrel, A.G., Bergoriordias, G.T., Scheinberg, I.H.,
Hickman, J. and Ashwell, G. (1971) :
J. Biol. Chem., 246 : 1461.
- Mortenson, R.F. and Gewurz, H. (1976) :
Effects of C-reactive protein on the lymphoid
system. J. Immunol., 116 : 1244.
- Mosher, D.F. and Wing, D.A. (1976) :
Synthesis and secretion of alpha-2-macroglo-
bulin by cultured human fibroblasts. J. Exp.
Med., 143 : 462.
- Nagel, R.L., and Gibson, Q.H. (1971) :
The binding of hemoglobin to haptoglobin and
its relation to subunit dissociation of hemo-
globin. J. of Biolog. Chem., 246 (1) : 691.

Nandi, M., Lewis, G.P., Jick, H., Slone, D., Shapiro, S., and Siskind, V. (1970) :
Evaluation of haptoglobin phenotype O-O
in cirrhotic and non-cirrhotic hospital
populations.
J. Clin. Path., 23 : 695.

Nelson, W.E., Vaugh, V.C. & Behrman, R.E. (1983) :
Infectious Diseases, Nelson Textbook of
Pediatrics, 12th ed., P. 708 : 726. W.B.
Saunders Co. Philadelphia, London, Toronto,
Mexicocity, Riodejaneiro, Sydney, Tokyo.

Nemir, R.L., Roberts, P.H., and Barry, L.S. (1957) :
Observation of antistreptolysin O, c-reactive
protein and electrophoretic patterns in
maternal and neonatal sera.
J. Pediatr., 51 : 493.

Norberg, R., Birke, G., Hodfors, E. and Plantin, L.O.
(1970) :
Plasma Protein Metabolism. (M.A. Roth-Schild
& T. Waldman, eds.), P. 427 - 436.
Acad. Press, New York.

Norum, R. A., Bearn, A. G., Briscoe, W.A., and Briscoe, A. (1977) :
Alpha-1-antitrypsin and disease.
The Mount Sinai J. of Medicine, 44 (6) : 821.

Oliveira, E. B., Gotschlich, E. C., and Liu, T. (1977) :
Primary structure of human c-reactive protein.
Proc. Natl. Acad. Sci., 74 (8) : 3148.

Osmand, A.P., Mortensens, R.F., Siegel, J., and Gewurz, H. (1975) :
Interactions of c - reactive protein with the complement system.
J. Exp. Med., 142 : 1065.

Pagel, W., Simonds, F.A.H., Mc Donald, N. & Nassan, E., (1964) :
Pulmonary T.B., 4th ed. London, Oxford Univer. Press.

Palmblad, J., Cantell, K., Holm, G., Norberg, R., Strander, H., and Sundblad, L. (1977) :
Acute energy deprivation in man, effect on serum immunoglobulins, antibody response, complement factors 3 & 4 acute phase reactants and interferon producing capacity of blood lymphocytes. Clin. Exp. Immunol., 30 : 50.

Patwardhan, V.N., Maghrabi, R.H., Mousa, W., Gabr, M.K., and El-Maraphy, S. (1971) :
Serum glycoprotein in protein - calorie deficiency disease. The Amer. J. of Clinic. Nutrition, 24 : 906.

Paul, L.W. & Juhl J.H. (1972) :
Pulm. T.B., the essentials of Roentgen Interpretation 3rd, ed. P. 808.
Harper & Row Publishers, New York, Evanston, San Francisco, London.

Perry, K.M.A. & Sellors, T.H., (1963) :
Pulm. T.B., Chest Diseases, Vol. 2, London Butter Worths.

Phelan, P. D., Landan, I. I. & Olinsky, A. (1982) :
T.B. in childhood, Respiratory Illness in children.
2nd ed. P. 316.
Blackwell Scientific Publications. Oxford - London - Edinburgh - Melbourne.

Piasky, K.M., and Woolner, E.A. (1982) :
The binding of basic drugs to alpha 1-acid glycoprotein in cord serum.
J. Paediat., 100 (5) : 820.

Rajendran, G., Dhara, R., and Srinivasan, N.G. (1975) :

Serum haptoglobin in health and disease.

Indian J. Med. Res., 63 (4) : 525.

Ralph, D. & James, D.C. (1981) :

Bacterial Infections, Textbook of Pediatric

Infectious Diseases. P. 1016 : 54.

W.B. Saunders Co.

Philadelphia - London - Toronto - Sydney.

Richard, W.B. & Ross, G.M. (1968) :

Tuberculosis, Disease In Infancy & Childhood

E. & S. Livingstone.

Edinburgh & London.

Rifikin, D.B., Loeb, J.N., Moore, G. and Reich, H.

(1974) :

Properties of plasminogen activators formed by

neoplastic human cell cultures. J. Exp. Med.,

139 : 1317.

Ritchie, D.G., and Fuller, G.M. (1981) :

An in-vitro bioassay for leukocytic endogenous

mediators using cultured rat hepatocytes in

flamm., 5 : 287.

Rose, P.E., Johnson, S.A., Meakin, M., MacKie, P.H.

and Stuart, J. (1981) :

Serial study of C-reactive protein during
infection in leukaemia. J. Clin. Path.,
34 : 263.

Rosenthal, S.R. (1961) :

The Disk - Tine tuberculin test, J. Am. Med.
Ass., 177 : 452.

Ross, J.D. & Horne, N.W. (1976) :

Modern Drug treatment in T.B., Tavistock
House North - London.

Roudolph, A.M., Barnett, H.L., & Einhorn, A.H., (1977) :

Bacterial & Viral infections, pediatrics,
Appleton - Century - Crafts - New York.

Roy, R.B., Shaw, R.W., and Connell, G.E. (1969) :

A simple method for the quantitative
determination of serum haptoglobin.
J. Lab. Clin. Med., 74 (4) : 698.

Sakr, R., Samuel, S., El-Heneidy, F., and Atallah, Al.
(1981) :

Clinico - radiological versus pathological
diagnosis in tuberculous cervical lymphadenop-
athy. G.E.P.A., 29, No. 152, P. 151 - 161.

Sakr, R., Samuel, S., and Helmy, N.T. (1984) :

Studies on extrapulmonary tuberculosis in
Egypt, accepted in the Egyptian Journal of
chest Diseases and Tuberculosis.
Vol. 27, No. I - January.

Saksela, E. (1981) :

In Interferon-3. Ed. I., Gresser Academic
Press. London, P. 45 : 63.

Schmid, K., Burgi, W., Collins, J.H., and Nanno, S.
(1974) :

The disulfide bonds of alpha 1-acid glyco-
proteins.
Biochemistry, 13 (13) : 2694.

Schmid, K. (1975) :

The plasma proteins : structure, function
and genetic control. 2nd ed. Edited by F.W.
Putnam, Vol. I, P. 183 : 223. Acad. Press,
New York.

Schofield, K.P., Voulgari, F., Gozzard, D.I., Leyland
 , M.J. Beeching, N.J. and Stuart, J. (1982) :
C-reactive protein concentration as a guide
to antibiotic therapy in acute leukaemia. J.
Clin. Pathol., 35 : 366.

Schreiber, G., Dryburgh, H., Weigand, K., Schreiber, M.
 , Witt, I., Seydewitz, H., and Howlett, G.,
(1981) :
Intracellular precursor forms of transferrin,
alpha 1-acid glycoprotein and alpha-1-anti_
trypsin in human liver.
Acta Biochem. Biophys., 212 (2) : 319.

Schultze, H.E. (1960) :
Gamma-x-globulin and immuno-electrophoretic
demonsterable gamma-globulin-complement with
relation to C-reactive protein. Verhandl.
Deutsch. Ges. Inn. Med., 1 : 255.

Schultze, H.E. (1962) :
Arch Biochem Biophys., 1 : 290.

Schultze, H.E., Heremans, J.F. (1966) :
Synthesis of the plasma proteins. In molecular
Biology of Human Proteins, Vol. I. Elsevier,
Amestrdam, P. 321 : 449.

Shelter, M.R. (1961) :

Ann. N.Y. Acad. Sci., 94 : 44.

Shibata, K., Okubo, H., Ishibashi, H. and Tsuda, K.

(1977) :

Rat alpha-1-acid glycoprotein. Purification
and immunological estimation of its serum
concentration.

Biochem. Bioph. Acta, 495/1 :37.

Shibata, K., Okubo, H., Ishibashi, H., Tsuda-Kawamura,

K., and Yanase, T. (1978) :

Rat alpha 1-acid glycoprotein, uptake by
inflammatory and tumour tissues.

Br. J. Exp. Path., 59 (6) : 601.

Sipe, J.D. and Rosentrich, D.L. (1981) :

Serum factors associated with inflammation.
In Cellular Functions in Immunity and Infl-
ammations. P. 411. Edited by J.J. Oppenheim
, D. L. Rosentrich and M. Potter. Edward
Arnold, London.

Siy, D.A., Grohlich, D. and Bezkorovainy, A. (1975) :

Transferrin in the reticulocyte cytosol.

Biochem. Biophys. Acta, 3885 : 36.

Stollerman, G.H, Glock, S., Patel, D.J., et al.

(1953) :

Determination of c-reactive protein in serum
as a guide to the treatment and management
of rheumatic fever.

Am. J. Med., 25 : 645.

Strobel, S., Bender, S.W., Posselt, H.G., and Hubner,

K. (1980) :

Alpha-1-antitrypsin deficiency : fulminant
course in early infancy.

Helv. Paedia. Acta, 35 : 75.

Sutton, David (1975) :

Pulm. infection & infestation.

Textbook of Radiology.

Churchill Livingstone, Edinburgh, London,

Melbourne, New York.

Swenson, R.P. and Howard, J.B. (1979) :

Structural characterization of human alpha-2-
macroglobulin subunits. J. Biol. Chem.,

254 : 4462.

Swingler, G.H., Miller, M.S.H. & Beatty, D.W. (1983) :

S. AF. Med. J. Vol. 63, No. 6.

Talamo, R.C. (1975) :

Basic and clinical aspects of the alpha-1-
antitrypsin. J. Pediatric, 56 : 91.

Tillet, W.S. and Francis, T. (1930) :

Serological reaction in pneumonia with a non-
protein fraction from pneumococcus. J. Exp.
Med., 52 : 561.

Twining, S.S., and Brecher, A.S. (1977) :

Identification of alpha 1-acid glycoprotein,
alpha-2-macroglobulin and antithrombin III as
components of normal and malignant human
tissues.

Clin. Chimica. Acta., 75 (1) : 143.

Udani, P.M. & Maddocks, I. (1978) :

Diseases of children in the tropics and
subtropics. 3rd ed. P. 308 : 18.

Edward Arnold publisher - London.

Van Lewen, F., Cassiman, J. J. and Van Den Berghe, H.

(1979) :

Demonstration of an alpha-2-macroglobulin
receptor in human fibroblasts, absent in
tumour-derived cell lines. J. Biol. Chem.,
254 : 5155.

Van Rizk, P.P., and Van Den Hamer, C.J.A. (1976) :

I^{125} -asialo- α_2 -acid glycoprotein.

Investigation of its use for liver function tests. Metabolism in the rat.

J. Lab. Clin. Med., 88 : 142.

Voclkkel, E.F., Levine, L., Alper, C.A. and Tashjian, A.H.
(1978) :

Acute-phase reactants ceruloplasmin and haptoglobin and their relationship to plasma prostaglandins in rabbits bearing the VX2 carcinoma. J. Exp. Med., 147 : 1078.

Voulgari, F., Cummins, P., Gardecki, T.I., Beeching, N.J., Stone, P.C. and Stuart, J. (1982) :

Serum level of acute - phase and cardiac proteins after myocardial infarction. Surgery, and infection. Br. Heart. J., 48 : 352.

Wada, T., Ohara, H., Watanabe, K., Kinoshita, H., and Nishio, H. (1970) :

Studies on haptoglobin synthesis in reticulo-endothelial tissues.

J. of Reticuloendothelial Society, 8 : 195.

Wagnerova, M., Wagner, V., Kriz, J., Wokounova, D.,
Madlo, Z., and Slesingerova, B. (1980) :
The morbidity of children with decreased serum
levels of alpha -1- antitrypsin in an air
polluted area.
Czechoslovak Medicine, 3 (4) : 280.

Wandall, J.H. (1974) :
Concentrations of serum proteins during and
immediately after surgical trauma. Acta. Chir.
Scand., 140 : 171.

Welch, M.J., Reineke, M.E., Hammarsten, J.F. and Guenter
, C.A. (1969) :
Antitrypsin deficiency in pulmonary disease.
The significance of intermediate levels. Ann.
Int. Med., 71 : 533.

Williams, R.C., Quie, P.G. (1968) :
Studies of human C-reactive protein in an in-
vitro phagocytic system. J. Immunol.,
101 : 426.

Williams, H.E. & Phelan, P.D. (1975) :

T.B. and pulmonary complications, Respiratory
Illness in children., Blackwell scientific
publications, Oxford - London - Edinburgh,
Melbourne.

Wilding, P., Adham, N.F., Mehl, J.W. and Haverbac, B.T.,
(1967) :

Alpha-2-macroglobulin concentration in human
serum. Nature, 217 : 1220.

Wintroub, B.U., Goetzi, E.J. and Austen, K.F. (1974) :

A neutrophil - dependent pathway for the
generation of a neutral peptide mediator. J.
Exp. Med., 140 : 812.

Wintrobe (1982) :

A Textbook of Haematology.

Wood, H.F., Mc Carty, M., and Slater, R.J. (1954) :

The occurrence during acute infections of a
protein not normally present in the blood.
Physical-chemical properties of the c-reactive
protein crystallized by a modified technique.
J. Exp. Med., 100 : 71.

Youmans, G.P., Brennan, P.J., Doub, L. and Goren, M.B.

(1979) :

Tuberculosis, 2nd ed.

W.B. Saunders Co.

Philadelphia, London, Toronto.

Zapolski, E.J. and Princiotto, J.V. (1980) :

Transferrin acquisition of catabolized erythrocyte iron in the rat. Life Sci., 27 : 739.

Zeinet, R.A., and Kukral, J.C. (1970) :

The turnover rate of orosomucoid in burned patients.

J. Trauma, 10 : 493.