

REFERENCES

REFERENCES

- Abbas, A.k. (1994):
Cellular and Molecular Immunology
Philadelphia , WB saunders .
- Abd El-Rahman A.M. and Rosenberg, A.A. (1994):
Prevention of intraventricular hemorrhage in the preterm infant.
Clin. Perinatol., 21 : 505 –21.
- Alpert, G.; Campos, M.; Preblud, S.R. and Plotkin, S.A. (1984):
Vancomycin dosage in pediatrics.
Am. J. pharmacol., 63 : 190.
- Amara, Et Al. (1995):
A study of calcium homeostasis in septicemic neonates.
Thesis, submitted for partial fulfillment of M.D. degree in
pediatrics. Benha Faculty of Medicine.
- Amato, M.; Howald, H. and Von-Moralt, G.(1988):
Quantitative changes of white blood cells and perinatal diagnosis
of infection in high risk preterm infants.
Pediater., 23 (2): 129-34.
- Amin, M.B. (1994):
Adrenal function in neonatal septicemia.
Thesis, M.D. degree in pediatrics, Benha Faculty of Medicine.
- Apps, D.K.; Cohen, B.B. and Stell, C.M (1996):
Common terminal pathways of metabolism; Oxygen toxicity In:
Biochemistry, a concise text for medical students: 212.
- Asada; Kozi and Yoshikawa (1993):
Frontiers of reactive oxygen species in biology and medicine.

Proceedings of the sixth International Conference on superoxide and superoxide dismutase.

Kyoto, Japan. October 11-15, 1993.

Experta Medica, Amsterdam, 1994.

Austgulen, R.; Liabakk, N. and Lein, E (1993):

Increased levels of soluble tumor necrosis factor alpha receptors in serum from pregnant women and serum and urine samples from newborn.

Pediatr., Res; 33:82-6.

Avery, G. B. (1994):

Bacterial and viral infection of the newborn.

Arch. Dis. Child., 71: 112

✓Avery, M. and First, R. (1994):

Disorders of respiratory, hematology and neurology ,neonatology, In: Pediatric Medicine seicond Edition, Waverly Company,4:199-209

Babior, B.M.; Kipnes, R.S. and Curnette, J.T. (1973):

Biological defense mechanisms. The production by leukocytes of superoxide. A potential bacteriological agent.

J. Clin. Invent, 52 : 741 - 4.

Ballard J.K., Khoury, J.C. and Wedigk (1991):

New Ballard score, expanded to include extremely prmatue infants.

J.Pediatr. 119 : 417.

Barbara J. Stoll and Robert M.Kliegman (2000):

Prematurity and intrauterine growth retardation In:- Nelson text book 16th Ed., W. B. Saunders, Philadelphia, London, Toronto, Montreal, Sydney, Tokyo, W. B. Saunders, Philadelphia,London,Toronto, Montreal, Sydney, Tokyo:451-538.

- Barbra, J. Stoll; Tavia G. and Sheldon, B.(1996):
Late onset sepsis in very low birth weight.
Journal of Pediatrics July, 43: 71.
- Bast,A. ; Haenen,G.R. and Doelman C.J. (1991):
Oxidant and antioxidant state of the art.
Am. J. Med. ;91 (3c):2s-13s.
- Bates, C.J (1995):
Vitamin A .
Lancet, 345, Jan. 7: 31-5.
- Battisti, O.; Mitchinson, R. and Davies, P.A. (1991):
Changing blood culture isolates in a referral neonatal intensive
care .
Arch. Dis. Child., 86: 775.
- Beaton, G.H.; Martorell, R; Aronson, K.J. et al. (1993):
Effectiveness of vitamin A supplementation in control of young
child morbidity and mortality in developing contraries.
ACC / SNC State of Art Series Neutrution Policy Discussion
Paper, No. 13, December.
- Bechman, JSA; Bechman, T.W., Chen J, et al. (1990):
Apparent hydroxyl radical production by peroxynitrite:
Impaction for endothelial injury from nitric oxide and superoxide.
Proc. Natl Acad Sci USA 87: 1620-1624.
- Bendich, A. and Langseth, L. (1995):
The health effect of vitamin C supplementation. A Review,
Journal of the American college of Nutrition, 14;2,: 124-36.

- Bennet, R.; Bergdahl, S.; Erikson M. and Zetterstrom, R.(1989):
The outcome of neonatal septicemia during fifteen years.
Acta. Pediatr. Scand. 78:40.
- Bennett, F.C. (1994):
Developmental outcome In :
Avery GB , Fletcher MA, Mac Donald MG (eds) :
Neonatology, pathophysiology and management of the
Newborn (4th ed), J B Lippincott Company, Philadelphia: 1367.
- Bertil, E. and Michael, D. Amylon. (1994):
Hemostatic disorders in the Newborn Diseases In :
Schaffer and Avery 6th (ed) vol. 2 , p777 W.B. Saunders
Company . Philadelphia , Lodon , Toronto , Montreal ,
Sydney and Tokyo.
- Bifano, E.M. and Ehrenkranz, Z.(1995):
Perinatal hematology
Clin. Perinatal., 23 : 3.
- Bont, E.S.; Martens, A. and Van, J. (1994):
Diagnostic value of plasma level of TNF α and IL -6 in newborn
with sepsis.
Acta. Pediatr., 83 : 696 – 9.
- Boyer, K.M.; Samuel, P. and Gotoff, S. (1986):
Prevention of early onset neonatal group B streptococcal
disease with selective intrapartum chemoprophylaxis.
N. Engl. J. Med., 314 : 1665 .
- Cairo, M.S., Rucker, R.; Bennetts, G. A.; Hicks, D.;
Katz, J. and Johnson, S. (1984):
Improved survival of newborns recieving leukocyte transfusions for
sepsis.
Pediatr., 74 (5) : 887

- Cairo, M.S. (1989):
Neonatal neutrophil host defence.
AJDC. , 143 : 40.
- Caplan MS; Sun X.M. and Hsueh, W. (1990):
Role of platelet activating factor and tumor necrosis factor-
alpha in neonatal nectrozing enterocolitis.
J Pediatr., Gastroenterol. Nutr., 11 : 317.
- Caplan, E.A. and Oed, L.J. and Kassel, R.L. (1990):
Cytokines serum level in neonatal sepsis.
Ann. Surg., 112-210.
- Carrol (1987):
Oxygen radicals and human nutrition In:
Davis Conference ann. of Int. Med., 107: 526 – 545.
- Chance, B. ; Sies, H and Boveris, A.(1989):
Hydrogen peroxide metabolism in mammalian organs.
Physiological review, 59: 577 – 605.
- Cheaseman, K. H. and Slater, T. F. (1993):
An introduction to free radical in medicine In:
Biochemistry Churchill. Livingstone; 481 –92.
- Christen, J.R. (1994):
Antioxidant and Eye Diseases.
American Journal of Medicine. 97, 3A : 175.
- Christensen, R. D.; Bradly, R. and Rothstein, G.(1981):
The leukocyte left shift in clinical and experimental
neonatal sepsis.
J. Pediatr., 98:101.

- Christensen, R. D.; Rothstein, G.; Harold, B.; Anstall, B. and Bybee, B.(1982):
Granulocyte transfusion in neonates with bacterial infection, neutropenia and depletion of mature marrow neutrophils.
J. Pediatr., 70 (1): 1.
- Christensen, R.D.; Mark, S.C.; Harry, R. and Hill, A.(1991):
Effect on neutrophil kinetics and serum opsonic capacity of intravenous administration of immunoglobulin to neonates with clinical signs of early onset sepsis.
J. Pediatr., 118 : 606.
- Clapp, D.W. ; Kligman, R.M. ; Baley, J.E; Shenker, N. ; Fanaroff, A.A. and Berger, M.(1989):
Use of intravenous administer immunoglobulin to prevent nosocomial sepsis in low birth weight infants : Report of a pilot study .
J.pediatr. 115 (6) : 973.
- Clemens, M.R. and Waller, H.D.(1987):
Lipid peroxidation in erythrocytes.
Chem. Phys. Lipids, 42 : 251.
- Collee , J.G. ; Duguid , J.P. ; Fraser , A.G. and Marmion, B.B. (1989):
Blood culture In : Practical Medical Microbiology 13 (ed.)
Churchil Livingstone .
- Coovadia, Y. M.; Mayosi, B. and Adhikari, M. Z.(1989):
Hospital acquired neonatal bacterial meningitis: the impact of cefotaxime usage on mortality and of Amikacin usage on incidence.
Ann. Trop. Pediatr., 9 (4):233.

Corrigan, J. J.(1984):

Thrombocytopenia : A laboratory sign of septicemia in infant and children.

J. Pediatr., 105:219.

Cosmi, E.V.;Torre, R.L. and Oiazze, J.L.(1996):

New Treatment of neonatal RDS17th international Symposium on Neonatal, intensive care, Sanremo, Italy, June:13-6.

Courser RJ, Ferrara T.B. and Wright G.B. (1996):

Prophylactic indomethacin therapy in the first 24 hr. of life the prevention of PDA.

J Pediatr., 128 : 631 – 7.

Crissinger K.D. and Granger, D.N. (1989):

Mucosal injury induced by ischemia and reperfusion in piglet intestine : Influences of age and feeding.

Gastroentrology, 97 : 920 –6.

Crissinger, K.D.(1994):

Regulation of hemodynamic and oxygenation in developing intestine: insight into pathogenesis of necrotizing enterocolitis.

Acta Pediatre Suppl., 396 : 8 – 10.

Crissingre, K.D.; Grisham, M.B.; and Granger, D.N.(1989):

Developmental biology of oxidant – producing enzymes and antioxidants in the piglet intestine.

Pediatric Res. 25: 612-616.

Crouch, S.P.M.; Salter, K. J. and Fletcher, J(1992):

Regulation of cytokine releasing from mononuclear cells by the iron-binding protein lactoferrin.

Blood, 80 (1): 235.

Crystal, R.C.(1991):

Oxidants and antioxidants ' Introduction'.
Am. J. Med.; 91:3c-1s.

Dalvi, R.; Rao, S. and Rangnoker, J.(1991):

Exchange transfusion in neonatal sepsis.
Indian J. Pediatr., 28 (1) : 39.

David, J.(1999):

Left to right shunt lesions.
Pediatric Clinic of North America vol. 46 N.2: 355-68.

Davison, A.M.; Cameron, J.S. and grunfeld, J.P., (1998):

Mechanism of glomerular injury In:
Oxford text book of clinical nephrology (2nd ed.)
Oxford University Press, 3 : 420.

De Bont ESGM; Matens, A and Van Rann, J. (1994):

Diagnostic value of plasma level of tumor necrosis factor alpha
and interleukin-6 in newborns with sepsis.
Acta Pediatr., 83: 696-9 .

De Tejada, J.S. ; Goldstein , J. and Cotten, R.P.(1989):

Impaired neurogenic and endothelial mediated relaxation of
penile smooth muscle from diabetic men with impotence.
N. End. J. Med., 320: 1025 - 30.

De Wayne, M.P.; and John, P.Cloherty (1998):

Identifying the high risk newborn and evaluating gestational age
prematurity and postmaturity In :
Mannual of neonatal Care, Harvard Medical School , Brigham
and Women's Hospital . The Children's Hospital . Boston.
Editiors John, P. and Ann, R. Lippicott – Raven. fourth Edition,
:40.

- Denuk, M. and david, S. (1993):
Diagnostic Value of leucocyte indices in neonatal infection.
Trop. Geogr. Med., 55 (4) : 360.
- Dietzman, D.E. ; Frischer, G.W. and Schoen, K.D. (1994):
Neonatal Escherichia coli septicemia.
J. Pediatr., 185 : 128.
- Diguiseppi and Fridovich (1985):
The toxicology of molecular oxygen.
CRC critical review in toxicology, 12 : 315 – 42.
- Dormandy, T.L. (1990):
Free radical in biological systems.
Ann. Roy. Coll. Surge. Eng.; 62 : 188 - 94.
- Dormandy., T.L. (1983):
An approach to free radicals.
Lancet ; 2 : 1010 –14.
- Dowton, S. B. and Calton, H. R.(1988):
Acute phase reactants in inflammation and infection.
Hematol., 25:84.
- Doyle, J.; Verman, H.J.and Stevenson, D.K. (1997):
Dose vitamin C cause hemolysis in premature newborn infants?
Results of a multicenter double-blind, randomized controlled trial.
J. of Pediatrics, January.
- Edward, J.O' Rourke (1994):
Antibiotics and Infectious diseases In :
Mannul of Pediatric Therapeutics 1st (Ed.): 362 – 437.

- El – Beshlawy, A. (1993):
✓ Recombinant human erythropoietin in the anemia of prematurity.
Annual Scientific Meeting of Egyptian Society of Hematology
Cairo University.
- ✓ El-Dahr S.S. and Chevalier R.L. (1990):
Special needs of the newborn infant in fluid therapy.
Pediatr. Clin. North Am., 37 (2) : 323.
- EL - Maraghi, S.; EL - Saifi, A. and Abou - Alam., M. (1984):
Diagnosis of neonatal sepsis by gastric aspirate examination.
J. Egypt Soc. Ob. / Gyn., 10 (1): 43
- El-Maraghi, S.; Mostafa, A. and Salem, F. (2000):
Diagnostic value of TNF α in newborn with sepsis.
Thesis, M.D., Cairo University.
- EL - Naggar, A. et al. (1995):
Role of complement and fibrinectin deficiencies in neonatal
bacterial infection.
Thesis, submitted for partial fulfillment of M. D. degree in
pediatrics. Cairo Faculty of Medicine.
- El Salab, S; Abdel Aal, A.; El-Tombary, Y. and El-Morsy, Z. (1988):
Some aspects of neonatal septicemia in Mansoura special baby
care unit.
The Egyptian J. of Pediatrics, 5 No. 3-4: 269-78.
- El - Tayeb, E. M. (1997):
Glutathione peroxidase in peripheral neuropathy of Egyptian
diabetics.
Thesis, M.D. Ain-Shams University, Faculty of medicine :6
- Eshali, H.; Ringertz, S. and Nystrom, S. (1989):
Septicemia with coagulase negative Staphylococci in a

neonatal intensive care unit.
Acta, Pediatr. Scand., 360 : 127

Faris, (1996):
Antioxidant in COPD
Thesis, M.D. Faculty of medicine. Cairo University.

Feigin, R. D. ; Abramson, S. L. and Edwards, M. S.(1987):
Infection in the fetus and neonate In : Franaroff, A. A. and
Martin, R. J. (ed.) Neonatal and perinatal Medicine.
4th. (ed.) Mosby Co., Baltimore, Boston.: 763.

Freedman, R. M.; Ingram, D. L.; Gross, L.; Warshav, J. B. and
Baltimore, M.(1981):
A half century of neonatal sepsis at Yale (1928 - 1978).
Am. J. Dis. Child., 135 : 140

Freeman, B.A. and Crapo, J.D. (1982):
Biology of disease. Free radicals and tissue injury
Lab. Invest, 47 : 412 -16.

Fridovich, J.(1986):
Biological effect of super-oxide radical
Archives of Biochem. and Biophys. 247: 1-11.

Gamberzade, S. ; Toppare, M. F. and Senses, D. A.(1996):
Neonatal sepsis and meningitis
Pak. Med. Assoc., 46 (2): 43

Gerdes, J. S. (1991):
Clinicopathologic approach to diagnosis of neonatal sepsis.
Clin. Perinatology, 18 (2): 361.

Gertner, J.M. (1990):
Disorders of calcium and phosphorus homeostasis

Pediatr. Clin. North Am., 37 (6) : 1441.

Gilbert, D.L. (1985):

Perspective on the history of oxygen and life.

Gilbert D.L."ed" Oxygen and living process Springer Verlag, New York : 1- 43.

Gilstrab, L.G. ; leveno, K.J. ; Cox, S.M. ; Burris, J.S. and Rosenfeld, C.R. (1988):

Intrapartum treatment of acute chorio-amnionitis : Impact on neonatal sepsis.

Am. J. Obstet. Gyn. , 159 : 579.

Goldie, A.S.; Fearon, C.H. and Ross, J.A. (1995):

Natural cytokines antagonists and endogenous antiendotoxin

Core antibodies in sepsis syndrome.

JAMA 274 :172-7

Goldstone, I.M.; Ehrenkranz, R.A.; Adberg, S.C. and Baltimore, R.S.(1990):

A ten years review of neonatal sepsis and comparison with the previous fifty years experience.

Pediatr. Infect. Dis.; J.: 819 – 25.

Gomella (1994):

Nutrition during infancy.

Neonatology,:514.

Gotoffi, S. P.(1996):

Sepsis of newborn

Behrman, R. E.; Kligman, R. M. and Arvin, A. M. (eds):

Nelson Text book of Pediatrics W.B. Saunders Company:528.

Grael, EL.; Halle, E.; Bollmann, R. and Buch, H.:1989

Neonatal septicemia - incidence, etiology and outcome.

Acta. Pediatr. Scand., 360 : 113.

Grisham, M.B. (1994):

Oxidants and free radicals in inflammatory bowel diseases.
Lanset, 344: 859-61.

Guillois, B.; Berthan, C.A. and Wad, H. (1989):

The importance of C3 estimation in the diagnosis of generalized bacterial infections in the newborn infants.
Acta. Pediatr. Scand., 78 : 369.

Gureina, N.G. (1993):

Neonatal Nosocomial Infections: Prevention and Management, In: Kaplan (Ed.) Current Therapy Decker (Mosby - Year Book).
Pediatric Infectious Disease (3rd Ed.) St. Louis: B.C.: 406.

Guthrie, H.A and Picciano, M.F. (1995):

Fat soluble vitamins In:
Human Nutrition Chllp. 240 Mobsy Company New York: 420 – 60.

Gutteridge, J.M.C. (1992):

Aging and free radicals.
Med. Lab. Sci.; 49 : 313 – 18.

Guttridge, J.M.C. and Halliwell, B.(1994):

Antioxidant in nutrition , health and disease.
Oxford University Press.

Hall, R.T. ; Shigheoka, A.O. and Hill, H.R. (1983):

Serum opsonic activity and peripheral neutrophil count before and after exchange transfusion in infants with early onset group B Streptococcal septicemia .
Pediatric. Infect. Dis., 2(5) ; 356.

- Halliday, H. L.(1989):
When to do lumbar puncture in a neonate.
Arch. Dis. Child, 64: 313.
- Halliwell, B. and Gutteridge J. M. C. (1990):
Free radicals in Biology and Medicine.
London. Oxford University Press: 456.
- Halliwell, B. and Gutteridge, J. M. C. (1986):
Oxygen free radicals and iron in relation to biology and
medicine, some problems and concepts.
Arch. Biochem Biophs. ; 246: 501 - 14.
- Halliwell, B. and Gutteridge, J.M.C. (1991):
Free radicals in biology and medicine.
Oxford, (2nd Ed.) Clarendon Press : 415.
- Halliwell, B. (1991):
Reactive oxygen species in living system .
Am. Med., 91 (suppl): 14-22.
- Halliwell, B. (1994):
Free radicals, antioxidants and human disease.
Lancet, 344: 721-24.
- Haque, K.N.; Ziadi, M.H. and Bahakin, H. (1988):
Ig M enriched intravenous immunoglobulin therapy in neonatal
sepsis.
Am. J. Dis. Child., 142: 1293.
- Haris M.C.; Andrew T; Costarino J; John, S et al. (1994):
Cytokin elevation in critically ill infants with sepsis and

necrotizing enterocolitis.

J. of Pediatr., January 124: 105 .

Hartz, W.; Funakoshi, S. and Deutsch (1993):

The level of super-oxide dismutase and catalase in human tissues determined immunologically.

Clin. Chem. Acta., 46 : 125.

Hauth, J.C. ; Gilstrap, L.C. ; Hankins, G.D.V. and Canner, K. (1983):

Neonatal and maternal complication of acute chorioamnionitis.

J. Obst. Gyn., 66:59.

Hawell, E.A.; Fiers, W. and Rober, J. North (1988):

“Antitumour function of TNF “

J. Exp. Med. Vol 167 : 1067–85.

Helen, A. Christou and David, H. Rowitch (1998):

Anemia of Prematurity.

Manual of Neonatal care, Harvard Medical School , Brigham and Women's Hospital . The Children's Hospital . Boston.

Editors John , P. and Ann , R . Lippicott – Raven . fourth Edition, 4th (ed.): 453.

Helen, G. and Ann, R. stark (1998):

Respiratory Distress Syndrome

Manual of neonatal care . Harvard Medical School , Brigham and Women's Hospital . The Children's Hospital . Boston.

Editors John , P. and Ann , R . Lippicott – Raven.

Fourth Edition : 329.

Hindocha, P.; Campbell, C.A. and Gould, J.D.(1984):

Serial study of C-reactive protein in neonatal septicemia.

Arch. Dis. Child, 59 : 435.

- Hocker, J.R.; Simposn, P.M.; Rabaliais, G and Stewart, L. (1992):
Extracorporeal membrane oxygenation and early onset
group B streptococcal sepsis.
Pediatr., 89 (1): 3.
- Horowitz, B.; Change, M.D. and lippen, A. (1984):
Preparation of antihemophilic factor and fibronectin from
human plasma cryoprecipitate.
Cline Bioch., 24: 357.
- Houston, J. B.(1983):
Pus cells and bacterial counts in the diagnosis of urinary
tract infections.
Arch. Dis. Child, 38: 600.
- Hume, R.(1992):
Neonatal Metabolic Disorder In: Forfar and Arneils
Text book of Pediatrics (4th ed) , Churchill livingstone,: 322
- Humphry J.H., Ago estina, T. et al. (1996):
Impact of neonatal vitamin A supplementation of infant mortality
and morbidity .
J. Pediatr., 128 : 489 – 96.
- Ichikawa, I.; Kiyama, S. and Uoshioka, T.(1994):
Renl antioxidant enzymes, their regulation on function .
Kidney Int., 45: 1 - 9.
- Ingelfinjer, J.A., Mosteller, F.M., Thibodeau, L.A. and Ware, J.H.
(1994):
Biostatistics in Clinical Medicine. McG raw-Hill – Inc., New
York.
- Iraj, R (2000):
Glutamic acid . Metabolic diseases In :

Nelson Textbook of Pediatrics, (16 Ed.) W.B.Saunders
Company, Philadelehia, London, Toronto, Monteral, Sydney, Tokyo:
366.

Jafari, H.S. and McCraKen, G.H. (1992):

Sepsis and septic shock .

A review for clinician .

Pediatr. Infect. Dis. J., 11:739-48.

Jan Kallman; Jens Schollin and Cleas Schalen.(1998):

Impaired phagocytosis and opsonization towards group B
streptococci in preterm neonates.

Arch. Dis. child 78, 46 – 50.

Jean H.; Muhammad, N. and James, T.(1996):

Impact of Neonatal vitamin A supplementation on infant
morbidity and mortality.

J. of Pediatrics, April,:489-96.

Jean, M; didier, E and Lionel D. (1996):

Elevation of I.L.-6 and TNF alpha for early diagnosis of neonatal
Infection.

J. of Pediatrics, 129, 4 : 574 – 8.

John, H. Menkes (1991):

Intracranial hemorrhage : pathogenesis

Diseases of the new born. Taeusch, H. Ballard, A. and Avery, M
six edition vol. 1 p 422 – 425 , W. B. Saunders Company.

Philadalphia, London, Toronto, Montreal, Sydney and Tokyo.

Johonson, L.; Graham, E. and A bbasi, S. (1995):

Sever retinopathy of prematurity in infants with birth weight less
than 1250 grams.

J. Pediatr., 127 : 632 – 9.

- Kafetzis, D. A.; Brater, A. and Kapiki, A. N.(1982):
Treatment of severe neonatal infection with cefotaxime.
Efficacy and pharmacokinetics.
J. Pediatr., 100:483.
- Kalra, J. ; Mantha S.V. and Prasad, K.(1994):
Oxygen free radicals : key factors in clinical diseases.
Lab Medical International. X1: 16.
- Kanner, J.; German, J.B. and kinsella, J.E. (1987):
Initiation of lipid peroxidation in biological system.
CRC. Crit. Rev. food sci. Nut . 2, 25 : 317.
- Karl, C.K. Kuban (1998):
Intracranial hemorrhage
Manual Neonatal Care, Harvard Medical School, Brigham .
Women's and Children Hospital, Boston. Editors John P. and
Ann, R. Lippicott- Raven. 4th (ed.) : 505.
- Katalin, G; Tamara, B. and Richard, J.(1996):
Vitamin E and vitamin C supplement use and risk
of all-causes and coronary heart disease mortality.
Am. J. Cline Nutr; 64: 190-6.
- Kayo, I.S. (1996):
Neonatal sepsis and meningitis.
J. Bac. Med. Assoc., 44: 63.
- Kim, J.Y.; Gonzalez-Scarano, F. and Zeichner, S.L.; (1993):
Replication of type 1 Human immunodeficiency viruses
containing linker substitution mutations in the-201 to
130 region of the long terminal repeat.
J. Virol., 67: 1658 – 62.

Kimberlee, C.; Michael, E. and John, P. (1998):
Temperature control In:
Manual of Neonatal Care. Joint Program in Neonatology, Harvard
Medical School. Lippincott-Raven 4th (Ed.): 139-40.

Klebanoff, S.J. (1992):
Oxygen metabolites from phagocytes In:
Gallin JI, Goldstein IM, Synderman R (eds):
New York, Raven Press, 541-88.

Klein, J.O. ; Dashafesky, B.; Norken, C.R. and Mayer, A. (1983):
Selection of antimicrobial agents for treatment of neonatal
sepsis.
Rev. Infect. Dis. , 5:55.

Klein, J.O. and Remington, J.S. (1990):
Current concepts of infection of the fetus and newborn infant.
Ann. Trop. Med., 11:9.

Klein, J.O.; Dashafesky, B.; Norken, C.R. and Mayer, A. (1983):
Selection of antimicrobial agent for treatment of neonatal sepsis.
Rev. Infect. Dis., 5:55.

Kliegman, RM (1996):
Fetal and neonatal medicine
In: Behrman RE, Kliegman RM (eds):
Nelson Essentials of pediatrics,
3rd ed, WBS Saunders : 167.

Knapp, R.G. and Miller, M.C. (1992):
Clinical Epidemiology and Biostatistics.
Harwal Publishing Co., Pennsylvania.

Koo, W.K. (1992):
Parenteral nutrition-related bone disease. J PEP

J. Parenter. Enteral. Nutr., 16 : 386.

Krinsky (1992):

Human biological antioxidants
J.Am. Chem. 90 ; 136.

Kurlat, L.; Stoll, B.J. and McGowan, J.E.(1989):

Time to positivity for detection of bacteremia in neonates.
J.Clin.Microbiol.,27(5):1068.

Kyllonen, K.S.; Clapp, W.; Robert, M.; Nancy, S.
and Fanaroff, A .(1989):

Dosage of intravenous administered immunoglobulin
and dosing interval required to maintain target levels of
Immunoglobulin G in low birth weight infants.
J. Pediatr., 155 (6): 1013.

Label, M. H. and Mc Craken, G. H.(1989):

Delayed cerebrospinal sterilization and adverse outcome
of bacterial meningitis in infants.
Pediatrics, 83: 161.

Laucy, J.F.and Dangman, B.(1984):

A reexamination of the role of oxygen in retrolental fibroplasia.
Pediatrics, 73 : 82 – 96.

Laurence, A. Boxer (2000):

Phagocytic system In:
Nelson text Book of Pediatric 16th (Ed.) W.B. Saunders Company
Philadelphia, London, Toronto, Montreal, Sedney, Tokyo :
606 – 12.

Leening, J.P; Sutton, T.M. and Fleming, P.Y.(1995):

Neonatal skin as a reservoir of infection in neonatal sepsis.

Pediatr.Ind.Dis.J.,14(8):719

Lerman, SJ.(1986):

**Infections of the newborn infant.
Biol. Neonates , 49 : 58.**

Leslie, R. and Allen, C.C.(1989):

**Developmental disabilities: Problem of fullterm infants with
asphyxia. Edited by Lea and Febriger, Philadelphia,: 79.**

Littlewood, J. M. (1982):

**Sixty six infants with urinary tract infection in the first
month of life.
Arch. Dis. Child., 47:218.**

**Liu, C. H.; Lehan, C.; Speer, M. E.; Smith, E. O.; Fernbach,
D. L. and Rudolph, A. J.(1985):**

**Early detection of bacteremia in an outpatient clinic.
Pediatr., 75 (5):827.**

Maberry, M.C. and Gilstrap, L.C.(1991):

**Intrapartum antibiotic therapy for suspected intra -
amniotic infection : Impact on the fetus and neonate.
Clin. Obstet . Gyn., 34 (2) : 345 .**

Macintosh, K. (1984):

**Bacterial infections of the newborn.
Biol. Neonates, 38:114.**

Mackendrick, W. and Caplan, M. (1993):

**Necrotizing enterocolitis: New thoughts about pathogenesis
and potential treatments.
Pediatr., Clin. North Am. 40: 1047-59.**

Main, D.M. (1991):

**Prevention of preterm birth .In:
Taeusch, Ballard RA,Avery ME(eds):Diseases of the newborn
(6th ed.) ,WB Samnders company:87.**

Manroe, B.L.; Weinberg, A.G.; Rosenfeld, C.B. and Browre, R.(1979):

The neonatel blood count in health and disease.

J. Pediatr., 95 : 89.

Mark, A (2000):

Hypoglycemia In :

**Nelson Textbook of Pediatric 16th (Ed.) W.B. Saundersd
Company. Philadalphia, Toronto, Montreal, Sydney, Tokyo.:
439 – 50.**

Marodi, L.; Kalmar, A. and Szabo, I.(1989):

**Opsonic activity in serum from septic infants treated with
intravenous immunoglobulin.**

Arch. Dis. Child., 64; 530 .

Marsden, Hall and Brenner, B.N.(1996):

**Relative nitrogen and oxygen intermediates and the kidney In :
The Kidney (5th Ed.) edited by Brenner, B.M: Philadelphia, W.B.
Saunders Company: 713- 53.**

Marshall, W.J. and Bongert, S.K, (1995):

**Clinical, Biochemistry Metabolic and Clinical aspects 1st ed.
Churchill Livingstone; 767 – 75.**

Martin, U.(2000):

Malabsorption disorders In:

**Nelson Text Book 16th (Ed.) W.B. Saundersd Company.
Philadalphia, Toronto, Montreal, Sydney, Tokyo.: 1089-93 .**

- Mary, CH; Andrew, T; Costarino, J et al.(1994):
Cytokine elevations in critically ill infants with sepsis and
necrotizing enterocolitis.
J. Pediatr; 124 : 105 – 11.
- McCracken, G. H. and Freij, B. J.(1987):
Perinatal bacterial disease. In :
Freij, R. D. and Chervy, J. D. (eds) : Text book of Pediatric Infectious
Disease. [1st Ed.] W. B. Saunders Co. Philadelphia,: 940.
- McCracken, G.H.(1991):
Perinatal bacterial diseases In:
Freij, R.D. and Chervy, J.D. (ed,) : Text book of Pediatric infections
Diseases (ed. 2) W.B. Saunders Co. Philadelphia .,:747.
- Meignier, M.; Renaud, P.; Robert, R. ; Roze, J.C. ; Rigal, E.
and Mouzard, A (1989):
Cephalhematoma infection in neonatal septicemia.
Pediatr., 44 (1) : 27.
- Michael R. and Jeffrey L. (1997):
Advances in therapy for sepsis in children.
Ped. Clin. of North Am., vol. 44 Number 1, Feb. 179. 205.
- Michael, L.S.; John L.S.; Paul, F. and Roy, R (1995):
Soluble interleukin-2 receptor as a predictor of neonatal sepsis.
J. of Pediatrics., June: 982 – 85.
- Minoti, G and Aust, S.D. (1987):
The requirement for iron (III) in the initiation of lipid
peroxidation by iron (II) and hydrogen peroxide.
J.Biol. Chem.; 262 : 1098 – 104.

Odio, C.M. (1995):

Cefotaxime for treatment of neonatal sepsis and meningitis.
Diagn. Microbial. Infect. Dis., 22: 111

O'Donnell, J.P. and Merenstein, G.B.(1993):

Infection in neonate. In:
Merenstein and Gardner (eds.) : Handbook of Neonatal
Intensive Care. Ed. 3 Mosby Year Bood, Boston, Chicagho,
London, Philadelphia, Sedney,:287.

Ohlsson, A.; Bailey, T. and Takiddine, F.(1986):

Changing etiology and outcome of neonatal septicemia.
Riyadh, Saudia Arabia.
Acta. Pediatr. Scand., 75: 540.

✓ Oski, F.A.(1991):

Physiologic jaundice In:
Teavsch, H.W.; Ballard, R.A; Avery, M.E.(eds) Shaffer and
Avery's diseases of the Newborn (6th ed.), Saunders
company:754.

Overall, J.C.(1987):

Infection of the newborn. In:
Behrman, R.F. and Vaugan, V.C. (eds):
Nelson Text bood of pediatrics. ed. 13 W.B. Saunders,
Philadelphia, London, Toronto, Tokyo,: 422

Overturf G, Scherman. M. and Scheifele, D.W.(1990):

✓ Neonatal necrotizing enterocolitis associated with delta toxin-
producing methicillin-resistant staphylococcus aureus Pediatr
Infect Dis. J 9 : 88.

Padmanbhan, J.; Risembery, H.M. and Rowe, D(1993):

Howell - jolly bodies in the peripheral blood of fullterm and
premature neonates.

John. Hopkins. Med. J., 132:146.

Palmer, R. M. ; Ashton, D.S. and Moncadas (1988):
Vascular endothelial cell synthesize nitric oxide.
J. Argonatur. 333: 664 - 666.

Pamela, C. Champe and Richard, A (1994):
Hexose Monophosphate pathway In :-
Illustrated Reviews, Biochemistry (2nd ed.); 111:17
J.B. Lippincott Company, Philadelphia.

Park, M.K. (1996):
Patent ductus arteriosus In:
Pediatric Cardiology 3rd (Ed.): 142 – 145.

Parola, M.; Pinzan M. and Casini, A. (1993):
Stimulation of lipid peroxidation.
Biochem. Biophys. Res. Comm. 194:1044-50.

Paul, W.B. and Seder, R.A. (1994):
Lymphocyte responses and cytokines .
Cell, 76: 241 – 51.

Peter, G.B. and Johnston M B. (1994):
Preterm infants. The Newborn child (17th ed.):93.

Pharaoh, P.O.; Cooke T. and Cooke, R.W.(1990):
Birth weight specific trends in cerebral palsy
Arch. Dis. child, 65: 602.

Philip, A. G.; and Hewitt, J. R.(1980):
Early diagnosis of neonatal sepsis
Pediatr. 65 (5): 1036.

- Philip, A. G. (1984):**
Acute phase protein in neonatal infection.
Pediatr., 105:904.
- Philip, A.G. (1981):**
Decreased use of antibiotics using neonatal sepsis screening technique.
J. Pediatr., 98: 795.
- Pitkanen, O.M; Hall-man, M. and Anderson, S.M. (1990):**
Correlation of free radical induced lipid peroxidation outcome in very low birth weight infants.
J.Pediatr., 116 : 760 - 4.
- Polk, DH.(1991):**
Disorders of carbohydrate metabolism In:
Taecisch H W, Ballard R.n, and Avery M E (eds):
Diseases of the Newborn (6th ed.): 967.
- Pommier, C.G.; O'Shea,J. and Chused, T. (1984):**
Studies on fibronectin receptors of human peripheral blood leucocytes .
J. Exp. Med., 159: 137.
- Prasad, K; Kolra, J. and Chan, W.P.; (1989):**
Effect of oxygen free radicals on cardiovascular function at organ and cellular levels.
Am. Heart J., 117:19.
- Pryor, W.A. and Godber, S.S.(1991):**
Non invasive measures of oxidative stress in human Free. Rad.
Biol. Med.; 10 : 177 - 84.
- Pyati, S.P. ; Jacobes, M.M. ; Raval, D.S ; Amma, P. and lillen,**

I. (1983):

Penicillin in infants weighing two kilograms or less with early onset group B streptococcal disease.

N. Engl. J. med. , 308 : 1383 .

Ralph, D. ; Stuart, L. and Morven, S. (1987):

Infection in the fetus and neonate in neonatal perinatal medicine.

Ind. J. Ped., 54 : 217.

Rangel - Frusto MS; Pittle, D. and Costigan, M. (1995):

The natural history of the systemic inflammatory response syndrome (SLRS).

JAMA 273: 117 – 123.

Reid, T.I.; Murthy, N.R.N.; Tanaka, N.; Rossman, M. and Musick, W.D.L. (1981):

Structure and heme environment of beef liver catalase.

Proc. Nat. Acad. Sci. USA; 78:47-67.

Reiter, R.J. and Robinson, G.D.(1995):

Where do free radicals come from?

Melatonin, Bantam Book, USA: Oxford University Press 24-34.

Rice Evans CA and Diplock, AT. (1993):

Current status of antioxidant therapy.

Free Rad Bio. Med. 15: 77 - 96

Copyright 1993 by El - sevier Science, Inc.

Richard, B. Parade and Thomas, M. Berger (1998):

Chronic lung disease.

Mannual of neonatal care , The Children and Women Hospital.

Boston. Editors, John P. and Ann , R. Lippicott- Raven. 4th

(ED) Ch. 24 : 378.

Robert, A. Ronzio (1997):

Antioxidant In:
The cyclopedia of nutrition and good health
Houston, Texas : 26-8.

Robert, K.; Daryl, K.; Peter, A. and Victor, W.(1996):
Vitamin E In :
Harpers Biochemistry 4th (ed.): 619 – 20.

Robert, M. kliegman (1996):
The fetus and neonatal infants In :
Nelson Text book of pediatric 15th ed. W.B. Saunders Company.
Philadelphie , London, Toronto, Montreal, Sydney, Tokyo P 94.

Roberton, N. R. (1992):
Neonatal infections.
Ped. Infect. Dis. J., 13:65.

Rodwell, L. R.; Anton, L. L. and David, T.(1988):
Early diagnosis of neonatal sepsis using hematological scoring
system.
J. Pediatr., 112:761.

Rogers, S; Gisela, W; Mujahid, A and Mark, H.(2000):
Antioxidant capacity and oxygen radical diseases in the preterm
newborn.
JAMA Pediatr., M.E., December: 30 – 4.

Rohrmoser, M. M. and Maver, G. (1996):
Reactive oxygen species and glomerular injury. Mechanism
of glomerular diseases.
Kidney Blood Press Res. 19: 263 - 269.

Roman, J.; Fernandez, F.; Velasco, F.; Rojas, R.; Roldan, M.R. and Torres, A. (1993):

Serum TNF levels in neonatal sepsis and shock.
Acta. Pediatr., 82 : 352 – 4.

Ronald, J. Sokol and Edward J. Hoffenberg (1996):

Antioxidant in Padiatric Gastrointestinal Diseases
Pediatric Clinics of North America, volume 43, No.2 461: 88.

Rouse, D.J.(1994):

Strategies for the prevention of early onset
neonatal group B streptococcal sepsis; A decision analysis.
Obstet. Gynecol. 83:483.

Rozychi, H.; Stahl, G. E. and Baumgarts, S.(1987):

Impaired sensitivity of a single early leucocytic count in
screening for neonatal sepsis.
Pediatr. Inf. Dis. J, 6: 440.

Rush, M.G. and Frank, C. (1994):

Intramuscular versus enteral vitamin A supplementation in very
Low birth weight neonates.
J. of Pediatrics, September : 458 – 62.

Safaa, B.A.; Mona, H. El-S. and Tahany, A. El-K.(1995):

Serum tumour necrosis factor levels in neonatal sepsis.
Thesis for Partial Fulfilment of Master Degree in Pediatrics, Ain
Shams University.

Salem, F.A.; Mostafa, A. and Abdel Ghany, H. (1999):

Diagnostic value of plasma level of TNF α in newborn with sepsis.
The GAZETTE of Egyptian Pediatric Association vol. 47 No.3
July : 121 – 32.

Samuel, P. Gotoff (2000):

Infection of neonatal infant In:

Nelson Text book 16th Ed. W.B. Saunders Company.

Philadelphia , London, Toronto, Montreal, Sydney, Tokyo:

538-52.

Saugstad, O.D.(1992):

Neonatal oxygen radical disease In:

Recent Advances in Pediatrics, 10:173. }

Saugstad, O.D. (1996):

Mechanism of tissue injury by oxygen implications for neonatal disease.

Acta Paediatr., 85:1-4.

Saugstad, O. (1997):

Bronchopulmonary dysplasia and oxidative stress.

Acta Paediatr., 86 : 1277 – 82.

Saugstad, O.D. and Asen, A.O.(1980):

Plasma Hypoxanthine levels as a prognostic aid of tissue hypoxia

Eur. Surg. Res., 12: 123 – 9.

Saugstad, O.D.(1988):

High postmortem levels of hypoxanthine in the vitreous humor of premature babies with RDS.

Pediatrics, 81: 395 – 8.

Scheuch, C; Bernatt, C. and Gross, J.(1989):

Influence of hypoxanthine- xanthine oxidase system on striatal (H) dopamine up-take.

Biochim. Acta, 48 : 212 – 6.

Schreck, R; Albermann, k and Baeuerle, P.A.: (1992):

An oxidant stress-responsive transcription factors of eukaryotic cells.
Free Radic Res commun; 17: 221-37.

Shatla, H.M.; Basiem, H.M. and Shehata, M.H., (1997):
Clinical value of tumour necrosis factor alpha and
Dexamethasone in neonatal sepsis. The Gazette of the Egyptian
Pediatric Association, 45 (4): 1-17.

Shenai, J; Rush, M. B. and Stahlman, M. T. (1990):
Plasma retinol binding protein response to vitamin A
administration in infant susceptible to bronchopulmonary
dysplasia.
J. Pediatr., 116: 27.

Sherman, M. P.; Chance, K. H. and Goetzman, B. W.(1989):
Gram's stains of tracheal secretions predict neonatal bacteremia.
Am. J. Dis. Child., 138:848.

Shi, Y.; Shen, C.; Wang, J., et al.(1994):
Role of TNF in neonatal sepsis.
Clin. Med. Sci. H., 1 : 45-8.

Shohaib, S.; EL-Far, N. and Zoair, A.(1989):
Blood and Throat culture in neonates with respiratory tract
Infections. M. SC.
Thesis, Pediatr., Tanta University.

Shronts, Eva P. (1993):
Basic concept of Immunology and its application to clinical nutrition.
Immunology ant Clinical Nutrition, 8, 4 : 177 – 83.

Siegel, J.D.(1985):
Neonatal septicemia.
Perinatol., 9: 20.

- Silvers, K.M.; Jibson, A.T.; Russell J.M. and Powers H.J.(1998):
Antioxidant activity, packed cell transfusions and outcome in
premature infants.
Arch. Dis. Child. Fetal neonatal Ed, 78 : 214 – 19.
- Simon, C.; Schorder, H.; Beyer, C. and Zerbst, T.(1991):
Neonatal sepsis in an intensive care unit and results of
treatment.
Infection, 19 (3): 146.
- Sinclair, A. J. ; Barnett, A. H. and Lumec, J.(1990):
Free radicals and antioxidant systems in health and disease.
Br. J. Hosp. Med.; 43 : 334 – 44.
- Singer, L.A.; Plotz, B.; Elester, G. and Amer, A.(1975):
Estimation of C. reactive protein
Clin. Path., 29 : 611.
- Sklavnu, T.; Gatzola, K. and Hatzioannidis, K.(1991):
Efficacy of aztreonam in the treatment of neonatal sepsis.
Rev. Infect. Dis., 13 (7) : 291.
- Sokol and Hoffenberg (1996):
Antioxidants in pediatric gastro-intestinal diseases.
Pediatric Clinic of North Am. April, vol. 43 No.2: 471 - 84.
- Southorn, P.A., and Powis, G. (1988):
Free radical in medicine.
Mayo Clinic Proc, 63: 381-9.
- Spiegelblatt, L.; Saintonge, J. and Chicoine, R.(1995):
Changing pattern of neonatal streptococcal septicemia
Pediatr. Infect. Dis., 4 : 56

Transfusion, 35: 68 – 73

Strauss, R.G.(1988):

Granulopoiesis and neutrophil function in the neonates. In:
Stoekman, JA and Pochedly, C. (eds) :
Developmental and Neonatal Hematology (Ed 4).
Raven press, New York, : 88.

Strauss, R.G.(1991):

Transfusion therapy in neonates
Biol. Neonates, 63 : 42.

Stuart, J. and Whicher, J. T.(1988):

Tests for detecting and monitoring the acute phase response.
Arch. Dis. Child., 63: 115.

Suadicani, P. (1992):

Serum selenium concentration and the risk. of ischemic heart
disease .
Atherosclerosis, 96 : 33-44.

Suffredini A.f. (1994):

Current prospects for the treatment of clinical sepsis.
Crit. Care Med., 22: 12 – 8.

Suri, M.; Thirupuram, S and Sharma, V.K. (1991):

Diagnostic and prognostic utility of C-reactive protein, alpha –1-
antitrypsin and alpha –2 macroglobulin in neonatal sepsis.
Indian Pediatr., 28 (10) : 1159 – 94.

Susan, S.; Alicia, A. and Kennard, I.(1991):

Factors associated with umbilical catheter - related sepsis in
neonates.
A J D C, 145 : 675.

- Taeusch (1991):**
Schaffer and Avery's Diseases of newborn 6th (ed) vol. 3 p.1054
W.B. Saunders Company. Philadelphia, London, Toronto,
Montreal, Sydney and Tokyo.
- Tauber, A.L. ; Simson, E. and Wright, J. (1983):**
Chronic granulomatous disease.
J. Medicine. 62: 286 - 309.
- Taylor (1986):**
Lipoxygenase pathways
Br. Med. Bull, 39 : 219 – 222.
- Thomas, Hansen and Anthony Corbet (1991):**
Hyaline Membrane disease. In:
Schaffer and Avery's Diseases of the newborn sixth Edition W.B.
Saunders Company. Philadelphia, London, Toronto,
Montreal, Sydney and Tokyo : 501.
- Tollner, U. (1982):**
Early diagnosis of septicemia in the newborn : clinical studies
and sepsis score.
Eur. J. Pediatr., 138 : 331.
- Tracey, K.J. and Cerami, A. (1993):**
Tumour necrosis factor: An up dated review of its biology.
Crit. Care Med. 21: 415 - 22.
- Trenam, Ch. W.; Blanke, D.R. and Morris, Ch.J. (1992):**
Skin inflammation: Reactive oxygen species and role of iron.
J. Invest. Dermatol.; 99 : 675 - 82.
- Tullus, B.; Lundstrom, G. and Burman, G.(1991):**
Antibiotic susceptibility of 629 bacterial blood and CSF isolates
from Swedish infants and therapeutic implication.

Acta. Pediatr. Scand., 80:205.

Uavy, R; Fanaroff, A.A. and Korones S, (1991):
Necrotizing enterocolitis in very low birthweight infants.
Biodemographic and clinical correlates.
J. Pediatr., 119: 630-638.

Udall JN (1990):
✓ Gastrointestinal host defence and necrotizing enterocolitis.
J. Pediatr., 117 : 533.

Vain, N.; Mazlumian, J.R. and Swarner, O.W. (1980):
Role of exchange transfusion in the treatment of severe
Septicemia.
Pediatr., 66 : 693

Vanpee M, Blennow M, and Linne T, (1992):
Renal function in very low birth weight infants.
J. Pediatr., 121 : 784 – 8.

Victor Y.H. (1994):
After care of high risk infants and long term outcome In:
Schaffer & Avery's Diseases of the Neonatology
6th ed vol. 1 W.B. Saunders Company. Philadelphia, London,
Toronto, Montreal, Sydney and Tokyo : 293.

Victor, Y.H. (1994):
Prematurity and lowbirth weight In:
Robinson, M.D., Robertson, M.D. and Phelan, P.D.(Eds.):
Practical Pediatrics, 3rd (Ed.) : 237.

Visser, V.E. and Hall, R.T. (1980):

Lumbar puncture in the evaluation of suspected neonatal sepsis.
J. Pediatr., 96 : 1663.

Volpe, J.J.(1991):
Cognitive deficits in premature infants.
N. Engl. J. Med., 325: 276.

Wahlgren, N.G. and Lindqvist, C. (1987):
Haem derivatives in Cerebrospinal Fluid after intracranial hemorrhage.
Eur. J. Neural., 26 : 216.

Walchiers, D.; Van-Loock, F.; Yeanfils, G. and Vandepitte, J. (1995):
Epidemiological study on neonatal septicemia and meningitis in Belgium.
Acta Clin. Belg., 50(6):326.

Walling (1982):
The nature of primary oxidants mediated by metal ions.
Mastonhs and Morrison M. (eds) .
Oxford, Progoman Press, 85 - 97.

Ward, R. J. and Peter T. J. (1995):
Free radical In :
Marshall, W. J. and Bangert, S. K. (eds) .Clinical, Metabolic and Clinical aspects, Churchill Livingstone 765 - 77.

Ward, R.M.(1994):
Pharmacologic enhancement of fetal lung maturation
Pediatric Clin. North Amer, 21:523

Watchko, J.F.(1992):
Kernicterus in preterm new born.
Pediatrics 90: 707

Weaver, L. T. (1998):

Free radical in clinical biochemistry, metabolic and clinical aspects (Eds): William and Marshall Stephen. 77-106.

Weiss, S.J. (1989):

Tissue destruction by neutrophil.
N Engl. J. Med., 320: 365 – 76.

Whitelaw, A.(1990):

Treatment of sepsis with IgG in very low birth weight infants.
Arch. Dis. Child.; 65:347

Winklhofer, P.M.; Eilemunter, H. and Fruhwirth, M (1997):

Plasma vitamin C concentration in patients with cystic fibrosis
Am. J. Clin. Nutr., 65 (6): 1858 – 66.

Yamakage, A.(1980):

Occupational scleroderma-like disorder occurring in men engaged
in the polymerization of epoxy resins.
Dermatology; 161:33-44

Yoder, P.C. and Polin, R.A.(1986):

Immunotherapy of neonatal septicemia.
Ped. Clin. Nor. Am., 33 (3) : 481.

Yoder, P.R.; Gibbs, R.S. and Balanco, J.D.(1983):

A prospective, controlled study of maternal and perinatal
outcome after intra – amniotic infection at term.
Am. J. Obstet. Gyn., 145 : 695.