

Philips, SM.; Lin, JJ.; Galal, N.; Linette, GP.; Walker, DJ. and Perrin, PJ. (1990):

The regulation of resistance to *Schistosoma mansoni* by auto-anti-idiotypic immunity. III. An analysis of effects on epitopic recognition, idiotypic expression, anti-idiotypic reactivity at the clonal level.

J. Immunol., 145(7): 2272.

Pinkus, GS.; Etheridge, CL. and O'connor, EM. (1986):

Are keratin proteins a better tumor marker than epithelial membrane antigen? A comparative immunohistochemical study of various paraffin-embedded neoplasms using monoclonal and polyclonal antibodies.

Am. Clin. Pathol., 85: 269.

Pinkus, GS. and Kurtin, PJ. (1985):

Epithelial membrane antigen-adiagnostic discrimination in surgical pathology.

Hum. Pathol., 16: 929.

Prakash, S. and Wyler, DJ. (1992):

Fibroblast stimulation in schistosomiasis.

J. Immunol., 148: 3583.

Purtilo, DT. and Yunis, EJ. (1971):

Alpha-Fetoprotein: Its immunofluorescent localization in human fetal liver and hepatoma.

Lab. Invest., 25: 291.

Racadot, J. ; Olivier, L. ; Porcile, E. ; De Grye, C. and Klotz, HP. (1964):

Adenome hypophysaire de type "mixte" avec symptôme de l'acromégalie. II. Étude au microscope optique et au microscope électronique.

Ann. Endocrinol. (paris.), 25 : 503.

Ramaekers, FCS.; Puts, JJG.; Moesker, O.; Kant, A.; Huysmans, A.; Haag, D.; JAP, PHK.; Herman, CJ. and Vooijs, GP. (1983):

Antibodies to intermediate filament proteins in the immunohistochemical identification of human tumors: An overview.

Histochem. J., 15: 691.

Ramzi, SC.; Vinary, K. and Tucker, C. (1999):

Pathologic Basis of Disease.

W B Saunders Company.

Raziuddin, S.; Shetty, S.; Ibrahim, A. and Patil, K. (1990):

Activated CD4-Positive T-lymphocytes and impaired cell-mediated immunity in patients with carcinoma of urinary bladder with schistosomiasis.

Cancer, 65(4): 931.

Redman, CA.; Kennington, S.; Spathopoulou, T. and Kusel, JR. (1997):

Interconversion of sphingomyelin and ceramide in adult *Schistosoma mansoni*.

Mol. Biochem. Parasitol., 90(1): 145.

Reid, GD.; Sturroch, RF.; Harrison, RA. and Tarara, RP. (1995):

Schistosoma haematobium in baboon (*papio anubis*): assessment of protection levels against either a single mass challenge or repeated trickle challenges after vaccination with irradiated schistosomula.

J. Helminthol., 69(2): 139.

Roncalli, M.; Borzio, M.; De Biagi, G.; Ferrari, AR.; Macchi, R.; Tombesi, VM. and Servida, E. (1986):

Liver cell dysplasia in cirrhosis. A serologic and immunohistochemical study.

Cancer, 57: 1515.

Ruoshlati, E. and Terry, WD. (1976):

Alpha-Fetoprotein and serum albumin show sequence homology.

Nature, 260: 804.

Saad, EA.; Corutinho, SG.; Barbosa, FJ. and Silva, J. (1967):

Studies on schistosomiasis *mansoni*: correlation between clinical findings, liver pathology and portal circulation hemodynamics, In "Bilharziasis" . Edited by Mostofi FK P: 167. Springer Verlag Berlin, NewYork.

Sadun, EH.; Von Lishtenberg, F.; Cheever, AW.; Erickson, DG. and Hickman, RL. (1970):

Experimental infection with *Schistosoma haematobium* in chimpanzees. Parasitologic, clinical, serologic and pathological observations.

Am. J. Trop. Med. Hyg., 19: 429.

Sadun, EH.; Williams, JS.; Witherspoon, C. and Martin, LK. (1969):

The relative role of egg and adult worms in the development of liver damage in mice infected with *Schistosoma mansoni*.

Ann. Ny. Acad. Sci., 660: 841.

Sauer, HI.; Salman, H.; Ambrosius, H. and Schffner, H. (1990):

Effect of *Schistosoma mansoni* infection on the immune reactions against heterologus antigens in mice.

Allerg. Immunol. Lepiz., 36(1): 25.

Schlegel, R.; Banks, SS.; Mcleod, JA. and Pinkus, GS. (1980):

Immunoperoxidase localization of keratin in human neoplasms.

Am. J. Pathol., 101: 41.

- Schmidt, D.; Harms, D. and Lang, W. (1985):**
Primary malignant hepatic tumors in childhood.
Virchows Arch (pathol. Anat.), 407: 387.
- Sells, NM.; Becker, FF. and Leffer, HL. (1974):**
Hepatocyte proliferation and alpha 1-fetoprotein in neonatal, and partially hepatectomized rats.
Cancer Res., 34: 865.
- Sheppard, HW.; Sell, S.; Trefts, P. and Bahu, R. (1977):**
Effects of alpha-fetoprotein on murine immune response.I. Studies on mice.
J. Immunol., 119: 91.
- Shimkin, MB.; Mustacchi, PO.; Cram, EB. and Wright, WH. (1955):**
Lack of carcinogenicity of lyophilized *Shistosoma* in mice.
J. Natl. Cancer Inst., 16: 471.
- Sieinski, W.; Dorsett, B. and Ioachim, HL. (1981):**
Identification of prekeratin by immunofluorescence staining in the differential diagnosis of tumors.
Hum. Pathol., 12: 452.
- Simpson, AJ.; Hagan, P.; Hackett, F.; Omer, I.; Ali, P. and Smithers, SR. (1990):**
Epitopes expressed on very low Mr *Schistosoma mansoni* adult tegumental antigens conform to a general pattern of live-cycle cross-reactivity.
Parasitology, 100(1): 73.
- Smith, CJP. and Kellehear, PC. (1980):**
Alpha-Fetoprotein molecular heterogeneity. Physiologic correlations with normal growth, carcinogenesis, and tumor growth.
Biochim. Biophys. Acta., 605: 1.

Smithers, SR. (1972):

Recent advances in the immunology of schistosomiasis.
Academic Press London and New York.

Smithers, SR. and Tery, RK. (1967):

Resistance to experimental infection with *Schistosoma mansoni* in rhesus monkeys induced by the transfer of adult worms.

Trans. Roy. Soc. Trop. Med. Hyg., 61: 515.

Sobh, M.; Moustafa, F.; Ramzy, R.; Saad, M.; Deelder, A. and Ghoneim, M. (1991):

Schistosoma mansoni nephropathology in syrian golden hamster: effect of dose and duration of infection.

Nephron, 59(1): 121.

Souhel, OH.; Harushige, NT.; Toshi, HH.; Kozo, A.; Kazuhiro, N.; Masahiko, I.; Naoki, H.; Yoshiyuki, K.; Shuji, U. and Takao, T. (1998):

Expression of alpha-fetoprotein and albmin genes in human hepatocellular carcinomas: limitations in the application of the genes for targeting human hepatocellular carcinoma in gene therapy.

Hepatology, 599.

Standen, OD. (1949):

Experimental schistosomiasis. II. Maintenance of *Schistosoma mansoni* in laoratory, with some notes on experimental infection with *s. haematobium*.

Ann. Trop. Med. Parasitol., 43 : 268.

Steffens, J.; Friedman, W. and Nagle, K. (1984):

Immunohistochemical demonstration of tumor associated antigen with monoclonal antibodies.

Urol. Res., 13: 55.

Stenberger, L.; Hardy, P. and Meyer, H. (1970):

The unlabeled antibody enzyme method of immunohistochemistry.

J. Histochem. Cytochem., 18: 315.

Stenger, R.J.; Warren, K.S. and Johnson, E.A. (1967):

An ultrastructural study of hepatic granulomas and schistosome egg shells in murine hepato splenic *Schistosomiasis mansoni*.

Exp. Mol. Path., 7: 116.

Sun, T.T. and Green, H. (1978):

Keratin filaments of cultured human epidermal cells: formation of intermolecular disulfide bonds during terminal differentiation.

J. Biol. Chem., 253: 2053.

Sun, T.T.; Shih, C. and Green, H. (1979):

Keratin cytokeratins in epithelial cells of internal organs.

Proc. Natl. Acad. Sci. USA., 76: 2813.

Symmers, WSTC. (1904):

Note on a new form of liver cirrhosis due to the presence of the ova of bilharzial *haematobium*.

J. Path. Bact., 9: 237.

Taylor, M.G.; Nelson, G.S.; Smith, M. and Andrews, B.J. (1973):

Studies on heterologic immunity in schistosomiasis 1. observations on the development of acquired homologous and heterologous immunity to *Schistosoma mansoni* in baboons.

Bull. Wrld. Hlth. Organ., Geneva.

Teixeira, R.; Ferreira, MD. ; Coelho, PMZ.; Filho, B.; Azevedo Junior, GM. and Lambartucci, JR. (1996):

Pyogenic liver abscesses and acute schistosomiasis mansoni: report on 3 cases and experimental study.

Trans. Roy. Soc. Trop. Med. Hyg., 90: 280.

Thung, SN.; Gerber, MA.; Sarno, E. and Popper, H. (1979):

Distribution of five antigens in hepatocellular carcinoma.

Lab. Invest., 41: 101.

Tuczek, HV.; Fritz, P.; Wagner, T.; Braun, U.; Grau, A. and Wenger, G. (1981):

Synthesis of alph-fetoprotein (AFP) and cell proliferation in regeneration liver of NMRI mice after partial hepatectomy. An immunohistochemical and autoradiographic study with 3H-Thymidine.

Virchows Arch (cell Pathol.), 38: 229.

Tungeskar, M.; Aladnani, M. and Gultesk, F. (1987):

Schistosomiasis associated bladder lesions, expression of cytoperlin.

B. J. Urol., 60: 423.

Vadas, MA.; David, JR. ; Butterworth, AE.; Pisani, NT. and Siongok, TA. (1979):

A new method for the purification of human eosinophils and neutrophils and a comparison of the ability of these cells to damage schistosomes of *Schistosoma mansoni*.

J. Immunol., 122: 1228.

Van Muizen, GNP.; Ruiter, DJ.; Ponc, M.; Huiskens-Vander Mer, C. and Warnaar, SO. (1984):

Mono-Clonal antibodies with different specificities against cytokeratins: an immunohistochemical study of normal tissues and tumors.

Am. J. Pathol., 114: 9.

Vasconcelos, EG.; Nascimento, PS.; Meirelles, MN.; Verjovski, AS. and Ferreira, St. (1993):

Characterization and localization of an ATP-diphosphohydrolase on the external surface of the tegument of *Schistosoma mansoni*.

Mol. Biochem. Parasitol., 58(2): 205.

Velupillai, P. et al. (1997):

B-1 cell (CD5+B220+) out growth in murine schistosomiasis is genetically restricted and is largely due to activation by polylactosamine sugars.

J. Immunol., 158: 338.

Warren, KS. (1962):

The influence of treatment on the development and course of murine hepatosplenic schistosomiasis.

Trans. Roy. Soc. Trop. Med. Hyg., 56: 510.

Warren, KS. (1963):

The pathogenesis of clay pipe-stem cirrhosis in mice with chronic schistosomiasis *mansoni*, with a note on the longevity of schistosomes.

Am. J. Path., 49: 477.

Warren, KS. and Klein, L. (1969):

Chronic murine hepatosplenic *schistosomiasis mansoni*; relative irreversibility after treatment.

Trans. Roy. Soc. Trop. Med. Hyg., 63: 333.

Warren, KS. (1972):

The immunopathogenesis of schistosomiasis. A multi-disciplinary approach.

Trans. Roy. Soc. Trop. Med. Hyg., 66: 417.

Warren, KS. (1973):

The pathology of schistosome infections.

Helminth. Abstract, 42: 591.

- Warren, KS. (1976):**
Schistosomiasis. A multiplicity of immunopathology.
J. Invest. Dermatol., 67: 464.
- Warren, KS. and Peters, PA. (1967):**
Comparison of penetration and maturation of *Schistosoma mansoni* in hamster, mouse guineapig, rabbit and rat.
Am. J. Trop. Med. Hyg., 16: 778.
- Wats, AE. and Socks, K. (1977):**
Squamous metaplasia and invasive epidermoid carcinoma of bladder.
Urology, 9: 317.
- Webbe, G. and James, G. (1971):**
The importance and maintenance of schistosomes of human and veterinary importance.
The 9th Symposium of British Society Parasitol.
- Werner, J.; Johannes, B.; Andre, D.; Marc, W. and Eric, VM. (1997):**
Adult *Schistosoma mansoni* worms positively modulate soluble egg antigen-induced inflammatory hepatic granuloma formation in vivo.
Am. J. Pathol., 150(6): 2033.
- Wiener, DP.; Koss, LG.; Sablay, B. and Freed, SZ. (1979):**
The prevalence and significance of Brun's nests, cystitis cystica and squamous metaplasia in normal bladder.
J. Urol., 122: 317.
- Wolfson, RL.; Maddison, SE. and Kagan, IC. (1972):**
Migration inhibition of peripheral leukocytes in human schistosomiasis.
J. Immunol., 109: 123.

Xiao, SH.; Shen, BG.; Horner, J. and Catto, A. (1996):

Tegument changes of *Schistosoma japonicum* and *Schistosoma mansoni* in mice treated with artemether.

Chung. Kuo. Yao. Li. Hsueh. Pao., 17(6): 535.

Zhang, C.; Yang, Y. and Xiao, S. (1995):

Pathological changes in the livers of rabbits infected with schistosome cercariae and treated with artemether or praziquantel in the early stage of infection.

Chung. Kuo. Chi. Sheng. Chung. Hsueh. Yu. Chi. Sheng. Chung. Ping. Tsa. Chih., 13(1): 51.

Zohair, MM. (1976):

The immunology of Schistosomiasis. In compiled review on schistosomiasis (bilharziasis).

Published NIDOG Cairo.