

References

REFERENCES

- Acharya G, AL-Sammaral MT, Patel N, Al-Habib A, Kiserud T. (2001):* A randomized controlled trial comparing effect of oral misoprostol and intravenous syntocinon on intra-operative blood loss during cesarean section. *Acta Obstet Gynecol Scand*; 80(3): 245-50.
- ACOG Practice Vulletin. (2001):* Clinical management guidelines for obstetrician and gynecologists: Medical Management of abortion. *Obstet Gynecol*, 97(4): 1-13.
- Adekanmi OA, Purmessur S, Edwards G. (2001):* Intrauterine misoprostol for the treatment of severe recurrent atonic secondary postpartum hemorrhage. *Br J Obstet Gynecol*; 108:541-2.
- Akhan SE, Iyibozkurt AC, Turfanda A. (2001):* Unscarred uterine rupture after induction of labor with misoprostol: a case report. *Clin Exp Obstet Gynecol*; 28(2):118-20.
- Akiyama H, Kohzu H, Matsuoka M (1981):* An approach to detection of placental separation and expulsion with new clinical signs: A study based on hemodynamic method and ultrasonography. *Am J Obstet Gynecol*; 140:505-9.
- Al-Hussaini TK. (2001):* Uterine rupture in second trimester abortion in a grand multiparous woman. A complication of misoprostol and oxytocin. *Eur J Obstet Gynecol Reprod Biol*; 96(2):218-9.
- Amant F, Spitz B, Timmerman D, Corremans A, Van Assche FA (1999):* Misoprostol compared with methylergometrine for the prevention of postpartum hemorrhage: a double-blind randomized trial. *Br J Obstet Gynecol*, 106 (10): 1066-70.
- Aronsson A, Bygdeman B, Gemzell-Danielsson K. (2004):* Effects of misoprostol on uterine contractility following different routes of administration. *Human Rep*; 19 (1): 81-4.

- Aubeny E, Baulieu EE. (1991):** Active contragestivie de L'association au RU 468 d'une prostaglandin active par voie oral CR Acad Sci (III); 312:539.
- Babinszki A, Kerenyi T, Torok O, Grazi V, Lapinski RH,, Berkowitz RL. (1999):** Perinatal outcome in grand and great-grand multiparity: effects of parity on obstetric risk factors. Am J Obstet Gynecol; 181:669-72.
- Bamigboye A, Derek A, Hofmyer G, Mitchell R. (1998):** Randomized comparison of rectal misoprostol with syntometrine for management of third stage of labour. Acta Obstet Gynecol Scand; 77: 178-81.
- Baskett TF (1999):** Complications of the third stage of labor. In essential management of obstetrical emergencies. 3rd ed. Bristol, UK: Clinical Press; 196-201.
- Bauer TA, Brom D, Chai LK. (1997):** Vaginal misoprostol for term labor induction. Ann pharmacother; 31:1391-3.
- Beazley JM (1986):** Natural labor and its active management. Dewhurst's text book of obstetrics and gynecology for postgraduates 4th edition, Whitfield C.R.; P. 324-46.
- Begley CM (1990):** A comparison of "active" and "physiological" management of the third stage of labor. Midwifery; 6(1):3-17.
- Beischer NA,, Mackay EV (1996):** Obstetrics and the newborn. London Bailliere Tindal; 169-77.
- Benedetto MT, De cicco F, Rossiello F, Nicosia AL lupi G, Acqua S. (1990):** Oxytocin receptor in human foetal membranes at term and during delivery Obstet Gynecol; 70: 35-9.
- Bennett B. (1997):** Uterine rupture during induction of labor at term with intravaginal misoprostol. Obstet Gynecol; 89(3):832-3.
-

- Berghahan L, Christensen D, Droste S. (2001)** Uterine rupture during second trimester abortion associated with misoprostol. *Obstet Gynecol* 98 (2): 976-7.
- Birlain M (2001):** Comparison of low doses of misoprostol with the traditional use of oxytocin for effective cervical ripening and labor induction *Obstet Gynecol*; 97(4): 567-9.
- Bjornerem A, Acharya G, Oian P, Maltau JM. (2002):** Postpartum hemorrhage prophylaxis and treatment in Norway. *Tidsskr Nor Laegeforen*; 122(26): 2536-7.
- Boulvain M, Stan C, Irion O; (2001):** Membrane sweeping for induction of labor; *Cochrane Data Base Syst. Rev.*; (2): CD000451.
- Brandt M. L. (1985):** Mechanism and management of the third stage of labor. *Am J obstet Gynecol*; 25:622-6.
- Brindley BA, Sokol RJ. (1998):** Induction and augmentation of labor. Basis and methods for current practice. *Obstet Gynecol*; 43:730-4.
- Bullough CH, Msuku RS, Karonde L (1989):** Early suckling and postpartum hemorrhage: controlled trial in deliveries by traditional birth attendants. *Lancet*; 2(8662): 522-5.
- Calder AA; (1997):** Review of prostaglandin use in labor induction. *Br J Obstet Gynaecol*; 104(15):2-7.
- Calder AA; (1999):** Induction and augmentation of labor. in Edmonds DK. *Dewhurst's Textbook of Obstetrics and Gynaecology for Postgraduates*; 6th ed; blackwell science; London and Cambridge; 252-9.
- Caliskan E, Dilbaz B, Meydanil MM, Ozturk N, Narin MA and Haberal A. (2003):** Oral misoprostol for the third stage of labor: a randomized controlled trial. *Obstet Gynecol* 101 (2): 921-8.
- Carroli G, Bergel E (2002):** Umbilical vein injection for management of retained placenta (cochrane Review). In: *The Cochrane Library*, Issue 2. Oxford, UK:
-

- Challis JRG, Vaughan M. (1997):** Steroid synthesis and PG metabolizing activity present in different cell population from human foetal membranes and deciduas. *Am Obstet Gynecol*; 157:74.
- Chestnut DH, Eden SD, Gall SA, (1985):** Peripartum hysterectomy: A review of cesarean and postpartum hysterectomy. *Obstet Gynecol*; 65:365-8.
- Clark S, Koonings PP, Phelan JP. (1985):** Placenta previa/accreta and prior cesarean section. *Obstet Gynecol*; 66:89-91.
- Clark SL, Yeh S-Y, Phelan J (1984):** Emergency hysterectomy for the control of obstetric hemorrhage. *Obstet Gynecol*; 64:376-7.
- Clarke J (1975):** Medical induction of labor in the active management of labor *Obstet Gynecol*; 2:49-79.
- Coleman RA, Smith WL,, Narumiya S, (1999):** International union of pharmacology classification of prostanoid receptors. *Pharmacol Rev*; 46:205-29.
- Collins PW, Casey ME, Varelal. (1993):** Historical review of misoprostol. *Contraception*; 55:330-2.
- Collins PW. (1990):** Misoprostol discovery, development and clinical application *Contraception*; 53:321-7.
- Combs C.A., Murphy E.L., Laros R.K., Jr (1991):** Factors associated with postpartum hemorrhage with vaginal birth. *Obstet Gynecol*; 77:69-76.
- Cook CM, Spurrett B, Murray H. (1999):** A randomized clinical trial comparing oral misoprostol with synthetic oxytocin or syntometrine in the third stage of labor. *Aust N Z J Obstet Gynaecol*; 39(4):414-9.
- Corta G, Vigilaute M, Di-Cesare S, Mascaretti G, Caserta D. Mosconini M. (1997):** Cervical maturation and induction of labor with PGE2 gel. *Minerva Gynecol*; 427-32.
-
-

- Costa SH. (1998):** Commercial availability of misoprostol and induced abortion in Brazil *Int J Gyn Obstet*; 63 (1): 131-9.
- Crane JM, Young DC, Butt KD, Bennett KA, Hutchens D. (2001):** Excessive uterine activity accompanying induced labor. *Obstet Gynecol*; 97(6): 926-31.
- Cunningham F, MacDonald PC, Gant NF (1989):** Abnormalities of the third stage of labor. *Williams obstetrics*, 18th edition, pub. N.Y: Appleton-century-crofts; P. 415-6.
- Cunningham FG, Gant NF, Leveno KJ. (2001):** Conduct of normal labor and delivery. *Williams Obstetrics*. 21st ed. New York NY: McGraw-Hill; 320-5.
- Cunningham, Mac Donald, Gant, Leveno, Gilstrap, Hankins, Clark (2001):** Obstetrical hemorrhage *Williams Obstetrics*, 20th. Edition, published by appletion & Lange, P. 745-82.
- Daisley H Jr. (2000):** Maternal mortality following the use of misoprostol. *Med Sci Law*; 40(1): 78-82.
- Danielsson RG, Marions L, Radriguez A, Spur BW, Wong PY, Bygdeman M (1999):** Comparison between oral and vaginal administration of misoprostol on uterine contractility *Obstet Gynecol*; 93(2):275-80.
- Davies S (1999):** Amniotic fluid embolism and isolated disseminated intravascular coagulation. *Can J Anaesth*; 56:456-60.
- Diab KU, Ramy A.R, Yehia M.A. (1999):** the use of rectal misoprostol as active pharmacological management of the third stage of labor. *J Obstet Gynecol*; 25(5):327-32.
- Donald I (1979):** Postpartum hemorrhage. *Practical Obstetric Problems*. 5th edition, Lloyd-Luke; P. 748.
- Drife J O. (1996):** Choice and instrumental delivery. *Br J Obstet Gynaecol*; 103:608-11.

- Fernando Arias. (1993):** Postpartum complications. In practical guide to high-risk pregnancy and delivery, ed. By Fernando Arias; 2nd ed., P. 433-51.
- Forman A.G, Andersson K. E, Ulmsten U. (1983):** Effect of nifedipine on oxytocin and prostaglandin F2 α induced activity in postpartum uteri. Clin Obstet Gynecol; 26(1)56-77.
- Forna AM, Villar J, Hofmeyr GJ (2002):** Prostaglandins for prevention of postpartum hemorrhage (Cochrane Review) in the Cochrane Library, Issue 2 oxford, UK.
- Fuchs AR, Fuchs F, Hussein P, Soloff MS, Fernstrom MJ (1982):** Oxytocin receptor and human parturition. A dual role for oxytocin in the initiation of labor. Science; (215):1396-400.
- Fuchs AR, Hussein P, Fuchs F. (1981):** Oxytocin and the initiation of human parturition and stimulation of Prostaglandin production in the human decidua by oxytocin. Am J Obstet Gynecol; 141:694-6.
- Fuchs AR, Periyasamy S. Alexandrova M, Soloff MS (1983):** Correlation between oxytocin receptor concentration and responsiveness to oxytocin in pregnant rat myometrium. Effect of ovarian steroids. Endocrinology; 113:742-7.
- Gainer H, Alstein M, Whitnall MA, Wray S. (1988):** The biosynthesis and secretion of oxytocin and vasopressin In Krobil E, Neill J (eds). The physiology of reproduction, Vol. II, New York, Rave; 2265.
- Genest DR, Di Salvo D, Rosenblatt MJ, Holmes LB. (1999):** Terminal Transverse limb defects with tetherin and amphocele in a 17 weeks fetus following first trimester misoprostol exposure. Clin. Dysmorphol 8(1): 53-8.
- Gerstenfeld TS, Wing DA. (2001):** Rectal misoprostol versus intravenous oxytocin for the prevention of postpartum hemorrhage after vaginal delivery. Am J Obstet Gynecol; 185(4): 878-82.

- Gilbert L, Porter W, Brown VA (1987):* postpartum hemorrhage: A continuing problem. *Br J Obstet Gynecol*; 94:67-71.
- Gilbert WM, Danielsen B. (1999):* Amniotic fluid embolism: Decreased mortality in a population-based study. *Obstet Gynecol* 93:973-7.
- Gilstrap L.C; Hauth JC; Hankins GDV., Patterson AR. (1987):* Effect of type of anesthesia on blood loss at cesarean section *Obstet Gynecol*; 69:328.
- Gonzalez C, Marques-Dias M, Kim C, Sugayama S, Da-Paz J Juson S, Holmes L. (1998):* Congenital abnormalities in Brazilian children associated with misoprostol misuse in first trimester of pregnancy. *Lancet*; 351:1624-7.
- Goto M (1984):* A survey of placental separation by real time B-mode scanning *Nippon Sanka Fujinka Gakkai Zasshi*; 36(8): 1171-6.
- Green J R, Soho S.L. (1989):* Factors associated with rectal injury in spontaneous deliveries. *Obstet Gynecol*; 73:732-6.
- Greenhil S, Friedmdn RK (1979):* Conservative management of third stage of labor. *Brit. J Obstet Gynecol*; 8:117-24.
- Gries J B, Bieniarz J, Scommegna A. (1981):* Comparison of maternal and foetal effects of vacuum extraction with forceps or cesarean deliveries. *Obstet Gynecol*; 92:732-5.
- Gruikshank DP, Pitkin RM, Donnelly E, Reynolds WA. (1981):* Urinary magnesium, calcium and phosphate excretion during magnesium sulfate infusion. *Obstet Gynecol*;; 103:375-8.
- Gulmezoglu Am (2000):* Prostaglandins for prevention of postpartum hemorrhage *Cochrane Database Syst Rev*; (2):CD000494.
- Gulmezoglu Am (2001):* WHO multicentre randomized trial of misoprostol in the management of the third stage of labor. *Lancet*,358:689-95.

- Gülmezoglu AM, Forna F, Villar J, Hofmeyr GJ (2002):** Prostaglandins for prevention of postpartum haemorrhage (Cochrance Review). In: The Cochrane Library, Issue2. Oxford, UK.
- Gulmezoglu Am, Villar J, Ngoc NT, Piaggio G, carroli G, Adetoro L, et al., (2001):** WHO multicentre randomised trial of misoprostol in the management of the third stage of labor. *Lancet*, 358:689-95.
- Gupta R, Ramji S (2002):** Effect of delayed cord clamping on iron stores in infants born to anaemic mothers a randomized controlled trial. *Indian pediatri*; 39(2): 130-5.
- Guyton A. C.; MD. (1991):** The heart as a pump In Text book of medical physiology, 8th edition by Guyton A.C., published by W.B. Saunders Co., P. 106, U.S.A.
- Hall M. H, Halliwell R, Carr-Hill R. (1985):** Concomitant and repeated happening of complications of the third stage of labor. *Br J Obstet Gynecol*; 92:732-41.
- Harris R. E. (1970):** An evaluation of the median episiotomy. *Am J Obstet Gynecol*; 96:1-8.
- Hayashi R. H.; Castillo M.S., Noah M.L (1981):** Management of severe postpartum hemorrhage due to uterine atony using an analouge of prostaglandin F2. *Obstet Gynecol.*; 58: 426-9.
- Healy DL, Quinn AM, Pepperell RJ. (1982):** Rotational delivery of the foetus. Kielland's forceps and two other methods. *Brit J Obstet Gynecol*; 89:501-6.
- Henriksen T.B, Bekk M, Hedegaard (1992):** Episiotomy and perineal lesions in spontaneous vaginal deliveries. *Br J Obstet Gynecol*; 99:950-3,
- Hibbard BM (1988):** Mechanism and management of normal labor. *Clinical obstetrics & Gynecol* 1st edition, pub. Lippincot, J.B, Washington; P. 449.
-
-

- Hofmeyr GJ, Gulmezolgu AM; (2003):* Vaginal Misoprostol for cervical ripening and induction of labor; cochrane database Syst. Rev.; (1): CD00941.
- Hofmeyr GJ, Nikodem VC, de Jager M, Drakely A. (2001):* Side-effects of oral misoprostol in the third stage of labor-a randomized placebo-controlled trial. S Afr Med J; 91(5): 432-5.
- Hofmeyr GJ, Nikodem VC, de Jager M, Gelbert Br (1998):* A randomized placebo controlled trial of oral misoprostol in third stage of labor. Br J Obstet Gynecol; 105: 971-5.
- Inman WHW. (1991):* Report on current PEM studies: drugs for peptic ulcer or reflux. Presser Event Monitor News; 7:32.
- Iwamura S, Kambegawa A (1987):* Determination of methylergometrine and Dihydro ergonovine in biological fluids. J pharm. Dyn; 4: 275-81.
- Jackson KW. Jr, Allbert JR, Schemmer GK. (2001):* A randomized controlled trial comparing oxytocin administration before and after placental delivery in the prevention of postpartum hemorrhage. Am J Obstet Gynecol 185(4): 873-7.
- Jing Song; (2000):* Use of Misoprostol in Obstetrics and Gynecology; Review Article-Obst & Gyn Survery; 55(8):503-10.
- Jones SA, Brooks AN, Challis JRG; (1989):* Steroids Modulate corticotropin-releasing hormone production in human foetal membranes and placenta; J Clin Endocrinol Metab; 68:825-33.
- Joy SD, Sanchez-Ramos L, Kaunitz Am (2003):* Misoprostol use during the third stage of labor. Int J Gynaecol Obstet 82(2): 143-52.
- Keirse MJNC. (1993):* Eicosanoids in human pregnancy and parturition. In Mitchell M; Eicosanoids in reproduction. CRC press Boca Raton, FL, USA, P. 199.

- Koonin LM, Mackay AP, Berg CJ, Atrash HK, Smith JC (1997):** Pregnancy-related mortality surveillance-USA1987-1990. *MMWR*;46:17.
- Koren G, Pastuszak A, Ito S. (1998):** Drugs in pregnancy. *N Engl J med*; 338-63.
- Kundodyiwa TW, Majoko F, Rusakaniko S. (2001):** Misoprostol versus oxytocin in the third stage of labor. *Int J Gynaecol Obstet*; 75(3): 235-41.
- Lam H, Tang Os, Lee CP, Ho PC (2004):** A pilot-randomized comparison of sublingual misoprostol with syntometrine on the blood loss in third stage of labor. *Acta Obstet Gynecol Scand*. 83(7): 647-50.
- Laurence DR, Bennett PN (1987):** "Oxytocinase and its pharmacodynamics" *Clinical pharmacology in Obstetrics* 6th edition, pub. Bristol-London-Boston; p. 1176.
- Leake RD. (1990):** Oxytocin in the initiation of labor. In Carsten ME, Miller JD (eds). *Uterine function. Molecular and cellular aspects*. Plenum, New York; 361.
- Letourneur B, Parant O, Tofani V, Berrebi A. (2002):** Uterine rupture on unscarred uterus following labor induction for 2nd trimester termination of pregnancy with oral misoprostol: conservative management. *J Gynecol Obstet Biol Reprod (Paris)* 31(4): 371-3.
- Levine R. J, Hauth J C, Current L. B. (1997):** Trial of calcium to prevent preeclampsia. *N Engl J Med*; 337:69-76.
- Liu HS, Chu Ty, Chang YR, Yu MH, Chen WH. (1999):** Intracervical misoprostol as an effective method of labor induction. *Int. J Gynecol Obstet*, 64:49-53.
- Lokugamage AU, Sullivan KR, Niculescu I, Tigere P, Onyangunga F, El Refaey H. et al., (2002):** A randomized study comparing rectally administered misoprostol versus Syntometrine combined with an oxytocin infusion for the cessation of primary post partum hemorrhage. *Acta Obstet Gynecol Scand*; 80 (9): 825-9.
-
-

- Lumbiganon P, Hofmeyr J, Gulmezoglu Am, Pmol A, Villar J. (1999):** Misoprostol dose related shivering and pyrexia in the third stage of labor. WHO collaborative trial of misoprostol in the management of the third stage of labor. *Br J Obstet Gynecol*; 10(4):304-8.
- Lumbiganon P, Villar J, Piaggio G, Gulmezoglu AM, Adetoro L, Carroli G. (2002):** Side effects of oral misoprostol during the first 24 hours after administration in the third stage of labor. *Br J Obstet Gynecol*; 109(11):1222-6.
- Mantyla R, Kanto J. (1987):** Clinical pharmacokinetics of methylergometrine (methylergonovine). *Int J Clin Pharmacol Ther Toxicol*; 19(9):386-91.
- Mategrano VA, Gabay MP. (2001):** Misoprostol in the prevention of postpartum hemorrhage. *Ann Pharmacother*, 35(12):1648-52.
- Maxwell AW (1985):** A comparison of buccal and intravenous oxytocin. *J Obstet Gynecol*; 71:37-44.
- McColgin SW, Bennett WA, Roach H, Cowan BD, Martin JN, Morrison JC; (1993):** Paturitional factors associated with membrane stripping; *Am J Obstet Gynecol*; 169:71.
- McDonald S, Prendivill W, Blair E. (1993):** Randomized controlled trial of oxytocin alone versus oxytocin and ergometrine in active management of third stage of labor. *BMJ*; 307:1167-71.
- McDonald S, Prendiville WJ, Elbourne D (2002):** Prophylactic syntometrine versus oxytocin for delivery of the placenta (cochrane Review). In the cochrane Library, Issue 2.Oxford, UK:
- Mercer JS, Skovgard RL (2002):** Neonatal transitional physiology: a new paradigm. *J Perinat Neonatal Nurs*; 15 (4): 56-75.
- Michaela R. (2000):** Pharmacology of prostaglandin in drugs facts and comparisons. St. Louis. A Wolters Kluwer Company; 1140-1.
-

- Mitche LL BF, Rogers K, Wong S; (1997):** The dynamics of prostaglandin metabolism in human foetal membranes and decidua around time of parturition. *J. Clin. Endocrinol Metab*; 77:759-62.
- Moran M, Mozes MF, Maddux MS. (1990):** Prevention of acute graft rejection by prostaglandin E1 analogue misoprostol in renal-transplant recipients treated with cyclosporine and prednisone. *N Engl J Med*; 322:1183-88.
- Mousa HA, Alfirevic Z. (2003):** Treatment for primary postpartum hemorrhage *Cochrane Database Syst. Rev*; (1):CD003249.
- Myerscough P.R. (1982):** Uterine action in labor in: Munro kerr's operative obstetrics. 10th edition. By myerscough, published by bailliere Tindall, P. 36-46. London.
- Nakla S, Skinner K, Mitchell BF, Challis JRG; (1986):** Changes in PG transfer across human foetal membranes obtained after spontaneous labor; *Am J Obstet Gynecol*; 179:1210-17.
- Ng PS. Chan AS, Sin WK, Tang LC, Cheung KS, Yuen PM. (2001):** A multicentre randomized controlled trial of oral misoprostol and I.M. syntometrine in the management of the third stage of labor. *Hum Reprod*, 16(1):31-35.
- Ngai SW, Yik Ming Chan, Sze wing Lam, Terence Tlao (2000):** Labor characteristics and uterine activity: misoprostol compared with oxytocin in women at term with pre-labor rupture of the membranes. *Br J Obstet Gynecol*; 102-109.
- Nordstrom; Fogelstam K. Friedman G. Larsson A, Rydhstroem H. (1997):** Routine Oxytocin in the third stage of labor *Br J Obstet Gynecol*; 104:781-6.
- Norman J, thong K, Baird D. (1991):** Uterine contractility and induction of abortion in early pregnancy by misoprostol and mifepristone. *Lancet*, 338:1233-6.
-

- Pierre F, Mesnard L, Body G (1992):** For a systematic policy of i.v. oxytocin induced placental deliveries in a unit where a fairly active management of third stage of labor is yet applied: results of a controlled trial. *Eur J Reprod Biol* Jan 31; 43(2): 131-5.
- Plaut MM, Schwartz ML, Lubarsky SL. (1999):** Uterine Rupture associated with the use of misoprostol in the gravid patient with a previous cesarean section. *Am J Obstet Gynecol*; 180:6(1): 799-806.
- Prendiville W., Elbourne D. (1987):** Care During the third stage of labor. In cholmers I, Enkin M, Keirse MJNC, editors. *Effective care in pregnancy and childbirth*. Oxford: Univdrstiy Pres; 1145-69.
- Prendiville WJ, Elbourne D, McDonald S (2002):** Active versus expectant management in the third stage of labor (Cochrane Review). In: *The Cochrane Library, Issue 2*. Oxford, UK.
- Ramwell PW, Foegh M, Loeb R, Leovey EMK; (1980):** Synthesis and Metabolism of prostaglandins, prostacyclins and thromboxane: The arachidonic acid cascade; seminars *Perinatol*; 4:3.
- Reddy VV, Carey J C (1989):** Effect of umbilical vein oxytocin on puerperal blood loss and length of the third stage of labor. *Am J Obstet Gynecol*; 16:206-11.
- Reichler A, Romen Y, Divon MY; (1995):** Induction of labor; current opinion in *Obstet Gynecol*; 7:432-6
- Robert A, Nezamis J E., Phillips J P. (1967):** Disoccovery of misoprostol. *Am J Dig Dis*; 12:1073-79.
- Roman WM, Lukash LA, Challis JRG, Mitchell BF; (1986):** Substrate utilization for estrogen synthesis by human foetal membranes and deciduas; *Am J Obstet Gynecol*; 155:1170-3.
-

- Sanchez-Ramos L, Chen AH, Kaunitz Am, Caudier Fl, Delke I; (1997):** Labor induction with intravaginal misoprostol in term premature rupture of membranes: A randomized study; *Obstet Gynecol*; 89:909-12.
- Satin AJ, Leveno KJ, Sherman ML. (1994):** High-dose oxytocin 20-versus 40 minutes dosage interval. *Obstet Gynecol*; 83:234-6.
- Sciscione A, Nguyen L, Manley J, Shlossman P, Colorgen G. (1998):** Uterine Rupture during preinduction cervical ripening with misoprostol in a patient with a previous cesarean delivery. *Aust. NZJ Obstet Gynecol.*; 38:96-7.
- Secher NJ, Arnso P, Wallin L (1978):** Haemodynamic effects of oxytocin (syntocinon) and methylergometrine (methergine) on the systemic and pulmonary circulation of pregnant anaesthetized women. *Acta Obstet Gynecol Scand*; 57:97-101.
- Seitchek J (1987):** The management of functional dystocia in the first stage of labor. *Clin. Obstet Gynecol.*; 30:42-5.
- Sellers Sm.; Hodgson H.T.; Mitchell M.D.; Andersson A.B., Turnbull A.C. (1982):** Raised prostaglandin level in the third stage of labor. *Am J Obstet Gynecol*; 114:202-9.
- Shah-Hosseini R. Evrard Jr (1989):** Puerperal uterine inversion *obstet Gynecol*; 73:567-70.
- Shetty A, Mackie L, Danielian P, Rice P, Templeton A. (2002):** A comparison of two dosage regimens of oral misoprostol for labor induction at term. *Acta. Obstet Gynecol Scand*; 81(4):337-42.
- Shojai R, Piechon L, D'Ercole C, Boubil L, Ponties JE. (2001):** Rectal administration of misoprostol for delivery induced hemorrhage. Preliminary study. *J Gynecol Obstet Biol. Reprod (Paris)* 30(6):572-5
-

- Sleep J (1993):* Physiology and management of the third stage of labor In: Bennett VR, Brown LK, eds. *Myles' Textbook for Midwives*. 12th ed. London, UK: Churchill Livingstone; 216-29.
- Spencery PM, Filegner JR, Hibbard BM (1966):* Management of third stage of labor. *Br Med J*; 2: 622-6.
- Speroff L, Glass RH, Kase NG; (1999):* The endocrinology pregnancy. In: speroff L, Glass RH, Kase NG (eds): *Clinical Gynecologic endocrinology and infertility* 6th (ed); Williams & Wilkins. Baltimore, Mary Land, USA; P. 251.
- Srisomboon J, Plyamongkol W, Aiewsakul P. (1997):* Comparison of intracervical and intravaginal misoprostol for cervical ripening and labor induction in patients with unfavorable cervix. *J Med Assoc Thai*; 80(3):189-94.
- Studd J (1985):* Management of labor 1st edition, Blackwell Scientific Publication; P 288.
- Surbek DV, Fehr PM, Hosli I, Holzgreve W. (1999):* Oral misoprostol for third stage of labor: a randomized placebo-controlled trial. *Obstet Gynecol*; 94(2):255-8.
- Tang OS, Ho Pc. (2001):* Pilot study on the use of sublingual misoprostol for medical abortin. *Contraception* 64(5):315-7.
- Tang OS, Lee SW, Ho PC. (2002):* A prospective randomized study on the measured blood loss in medical termination of early pregnancy be three different misoprostol regimens after pretreatment with mifepristone. *Hum Reprod*.17(11):2865-8.
- Tapozada M, Baird BY, Grimes DA, Van Look P. (1995):* Termination of pregnancy after 14 weeks, *Modern Methods of inducing abortion*. Oxford, U.K.; Blackwell Science;70-95.
- Tappozada M. El-Bossati M., El-Rahman HA. (1995):* Control of intractable postpartum hemorrhage by 15-methyl prostaglandin F₂ *Obstet Gynecol.*; 58:327-30.

- Templeton A. (1998):** Misoprostol for all. *Br J Obstet Gynecol*; 105:937-9.
- Turman T. (1996):** Report from symposium on the prevention and management of anaemia in pregnancy and postpartum hege. World Health Organization. Zurich 1996;1-13.
- Van Dongen PW, de Groot AN (1995):** History of ergot alkaloids from ergotism to ergometrine. *European journal of Obstet Gynecol Reproductive Biology*; 60(2): 109-16.
- Van-Coever Den, De-Groot HA, Howl, R C et al., (1982):** Management of the third stage of labor in the midwife obstetric units in cape town. *S Afr Med. J*; 62(14):478-81.
- Vartner M., (1991):** Postpartum hemorrhage. *Critical care clinics*; 4(4): 883-97.
- Villar J, Gulmezoglu AM, Hofmeyr GH, Forna F. (2002):** Systematic review of randomized controlled trials of misoprostol to prevent postpartum hemorrhage. *Obstet Gynecol* 100(6): 1301-12.
- Vimal N, Mittal S, Kumar S. (2003):** Sublingual misoprostol for preabortion cervical ripening in first trimester pregnancy termination. *Contraception* 67(4):295-7.
- Vimala N, Mittal S, Kumar S, Dadhwal V, Mehta S. (2004):** Sublingual misoprostol versus methylergometrine for active management of the third stage of labor. *Int J Gynaecol. Obstet*; 87(1): 1-5.
- Wagaarachchi PT, Ashok PW, Smith NC, Templeton A (2002):** Medical management of early foetal demise using sublingual misoprostol *BJOG* 109(4): 462-5.
- Walder J (1997):** Misoprostol: preventing postpartum hemorrhage. *Mod Midwife* 7 (9):23-7.
- Walker H. (1974):** Median episiotomy and the occurrence of post partum hemorrhage. *Br J Obstet Gynecol*; 19:316-9.

- Walley RL, Wilson JB, Crane JM, Matthews K, Sawyer E, Hutchens D. A (2000):** A double-blind placebo controlled randomized trial of misoprostol and oxytocin in the management of the third stage of labor. *Br J Obstet Gynecol*; 107(9): 111-5.
- Weir Mr, Klassen DK, Hall PS. (1991):** Minimization of Indomethacin-induced reduction in renal function by misoprostol. *J Clin Pharmacol*; 31:729-35.
- Wilkinson C, Enkin MW (2002):** Manual removal of placenta at caesarean section (Cochrane Review). In the Cochrane Library, Issue 2. oxford UK.
- Wilson RF, Mammem E,, Walt AJ (1971):** Eight years of experience with massive blood transfusions, *Trauma Obstet Gynecol*; 11:275-8.
- Wing D, Lovett K, Paul R. (1998):** Disruption of prior uterine incision following misoprostol for labor induction in women with previous cesarean delivery. *Obstet Gynecol*; 91:828-30.
- Wing DA, Ham D, Paul RH. (1999):** A comparison of orally administered misoprostol with vaginally administered misoprostol for cervical ripening and labor induction. *Am J Obstet Gynecol*; 180(5):1155-60.
- Wing DA, Omphroy Go, Paul RH (1997);** A comparison of intermittent vaginal administration of misoprostol with continuous dinoprostone for cervical ripening and labor induction; *Am J Gynecol Obstet*, 1177(3): 62-8.
- Winkler CL, Gray SE, Hauth JC. (1991)** Mid-second-trimester labor induction: Concentrated oxytocin compared with prostaglandin E₂ suppositories. *Obstet Gynecol*; 77:297-300.
- Yuen PM, Chan NS, Uim, SF et al., (1995):** A randomised double blind comparison of syntometrine and syntocinon in the management of the third stage of labor. *Br J of Obstet Gynecol*; 102(5):377-80.
-

- Zelop CM, Harlow BL, Frigelletto FD, Safon LE, Saltzman DH. (1993):** Emergency peripartum hysterectomy. *Am J Obstet Gynecol*; 168:1443-7.
- Zhao Y, Li X, Peng Y. (1998):** Clinical study on reduction of postpartum bleeding in cesarean section by misoprostol. *Zhonghua Fu Chan Ke Za Zhi*. 33(7): 403-5.
- Zieman M, Fong SK, Benowitz NL, Banks D, Darney PD, (1997):** Absorption Kinetic of misoprostol with oral or vaginal administration. *Obstet Gynecol*; 90(1):88-92.
- Zinner MJ, Rypins EB, Martin LR. (1989):** Misoprostol versus antacid titration for preventing stress ulcers in postoperative surgical ICU patients. *Ann Surg*; 210:590-5.