

R E F E R E N C E S

R E F E R E N C E S

- **Abuelo, J.G. (1986):**
Proteinuria: Diagnostic principles and procedures.
Annals of Internal Medicine, 98: 186-191.
- **Anderson, N.G.; Anerson, N.L. and Tollaksen, S.L. (1979):**
Proteins of human urine. Concentration and analysis
by two dimensional electrophoresis.
Clin. Chem., 25, 199.
- **Antoine, B.; Patte, D. and Gourdin, M.F. (1970):**
Hypokalemic nephropathy.
Am. J. Clin. Pathol., 54: 53.
- **Baker, F.J. and Silverton, R.E. (1982):**
Introduction to Medical Laboratory Technology.
5th ed., p. 246.
Butterworth & Co., London (Publishers).
- **Barnett, H.L. (1978):**
Classification of the nephrotic syndrome.
Am. J. Med., 4: 760.
- **Barratt, T.M. (1983):**
Renal disorders. In: Clayton, B.E.; Round, J.M., eds.
Chemical Pathology and the Sick Child, 120-49.
Oxford: Blackwell Scientific Publications.
- **Batsford, S.R. and Hardwicke, J. (1975):**
The spectrum of urinary basement membrane antigen
excretion in normal and pathological urine.
Clin. Neph., 3, 60.

- **Behrman, R.E. and Vaughan, V.C. (1987):**
Nelson Textbook of Pediatrics. Proteinuria, p. 1308.
Membranous and membranoproliferative glomerulonephritis, p. 1327-1328.
Lowe's Syndrome, p. 1661.
W.B. Saunders Company, Philadelphia, London, Tokyo.
- **Bergstein, J.M. (1982):**
Proteinuria.
Pediatric Clinic of North America, Vol. 29, No. 1.
- **Bonsnes, R.W. and Taussky, H.H. (1945):**
J. Biol. Chem., 158-581.
- **Bourdeau, J.E.; Carone, F.A. and Ganste, C.E. (1972):**
Plasma protein turn over.
J. Cell. Biol., 54: 382.
- **Brenner, B.M. and Dworkin, L.D. (1985):**
Glomerular permselectivity. In Proteinuria, by Avram, M.M., p. 3.
Plenum Medical Book Company, New York and London.
- **Brensilver, J.; Albright, R. and Cortell, S. (1981):**
Proteinuria in congestive heart failure.
Eighth International Congress of Nephrology, Athens, Greece, 291.
- **Brodehl, J. and Krohn, H.P. (1979):**
Alternate day versus intermittent prednisone in frequently relapsing nephrotic syndrome.
Lancet, 1: 401.

- **Buchanec, J.; Kilment, J. and Javorks, K. (1983):**
X-ray changes in kidneys of childrens with
orthostatic proteinuria.
Int. Urol. Nephrol., p. 3-9.
- **Butler, E.A. and Flynn, F.V. (1958):**
The proteinuria of renal tubular disorders.
Lancet, 2: 978-980.
- **Cameron, J.S. (1978):**
The problem of focal segmental glomerulosclerosis.
Argoress in Glomerulonephritis, ed. Kincaid-Smith,
P.; d'Apice, A.F. and Athins, R.C.
New York, John Wiley and Sons, p. 209.
- **Castenfors, J.; Mossfeldt, F. and Piscator, M. (1967):**
Effect of prolonged heavy exercise on kidney
function and urinary protein excretion.
Acta Physiol. Scand., 70: 194-206.
- **Chang, R.L.S.; Aeki, L.F. and Troy, J.L. (1975):**
The plasma proteins.
Biophys. J., 15: 887.
- **Deen, W.M.; Myers, B.D. and Brenner, B.M. (1982):**
The glomerular barrier to macromolecules:
Theoretical and experimental considerations.
In: Brenner, B.M. (ed.): Contemporary Issues
in Nephrology: Nephrotic Syndrome.
New York, Churchill Livingstone, pp. 1-29.
- **Ellis, D. and Buffone, G. (1982):**
Protein clearances and selectivity determination
in childhood nephrosis.
Clin. Chem., 27, 1987.

- **Elseses, U.S.; Griffiths, P.D. and Hocking Taylon, C.M. (1988):**
White RHR children in Hosp. Birmingham, England.
Simplified quantification of urinary protein
excretion in children.
Clin. Nephrol., 30: 225.
- **Farguhar, M.G. (1975):**
Chemistry and physiology of the human plasma proteins.
Kidney Int., 8; 197.
- **Galaske, R.G. (1983):**
Mechanism of proteinuria: filtration and resorp-
tion of plasma proteins.
Fortschr. Med., p. 657-9.
- **Gill, P.G. (1977):**
The progression of acute proliferative post-
streptococcal glomerulonephritis to severe
epithelial crescent formation.
Clin. Neph., 8: 449.
- **Ginsberg, J.M.; Chang, B.S. and Matarese, R.A. (1983):**
Use of single voided urine samples to estimate
quantitative proteinuria.
The New England Journal of Med., Vol. 309, No. 25,
Dec. 22, p. 1543.
- **Gitlin, D.; Janeway, C.A. and Farr, L.E. (1952):**
Studies on the metabolism of proteins in nephrotic
syndrome.
J. Clin. Invest., 35: 44.
- **Habib, R. and Kleinknecht, C. (1971):**
Glomerular disorders.
Path. Ann. New York, Appleton, p. 417.

- Haurowitz, F. (1983):
The Chemistry and Function of Proteins, edition 2.
Academic Press Inc., New York, NY.
- Heymann, W.; Makker, S. and Post, R. (1974):
Proteinuria in congenital nephrotic syndrome.
Ped., 50: 814.
- Hirtick, D.E. and Kassirer, J.P. (1982):
The nephrotic syndrome.
Year Book Medical Publisher, I.W.C., Chicago, London.
- Houser, M. (1984):
Assessment of proteinuria using random urine samples.
J. Pediatr., 104 (6), p. 845, 8.
- International Study of Kidney Disease in Children (ISKDC) (1980):
A controlled therapeutic trial of cyclophosphamide
plus prednisone versus prednisone alone in children
with focal segmental glomerulonephritis.
Pediatric Research, 14, 1006.
- Jeery, M.B. (1982):
Proteinuria in children: Management and Prognosis.
Pediat. Clin. North Amer., Vol. 29, No. 1.
- Jones, J.H.; Peters, D.K. and Morgan, D.B. (1967):
Observations on calcium metabolism in the nephrotic
syndrome.
Q. J. Med., 36: 301.

- Kampmann, J.; Siersbaek-Nielsen, K. and Kristensen, M. (1974):
Rapid evaluation of creatinine clearance.
Acta. Med. Scand., 196: 517-20.
- Karlsson, F.A. and Hellisine, K. (1979):
Urinary protein excretion in early infancy.
J. Pediatr., 89.
- King, S.E. and Gronbeck, C. (1952):
Benign and pathological albuminuria.
Ann. Intern. Med., 36: 765-85.
- Kluthe, R.; Hagemann, V. and Klein, N. (1976):
The turnover of 2 macroglobulin in the nephrotic
syndrome.
Vox. Sang., 12: 308.
- Koopman, M.G.; Krediet, R.T.; Koomen, G.C.; Stracke, J.
and Arisz, L. (1989):
Circadian rhythm of proteinuria; consequences of
the urinary protein : creatinine ratios.
Nephrol. Dial. Transplant., 4 (1) p. 9-14.
- Krishna, K.S.; Pandey, A.P.; Kirubakaran, M.G.;
Kanagasabapa, A.S. (1987):
Urinary protein/creatinine ratio as an indication
of allograft function following live related
donor renal transplantation.
Clinic. Chim. Acta., Feb. 27, 163 (1): p. 51-61.

- Kristal, B.; Shasha, S.M.; Labin, L and Cohen, A. (1988):
Estimation of quantitative proteinuria by using
the protein-creatinine ratio in random urine
samples.
Am. J. Nephrol., 8 (3): 198-203.
- Lawler, W.; Williams, G. and Tarpey, P. (1978):
IgA localization in glomerular disease.
Eng. J. Med., 299: 200.
- Leber, P.D. and Marsh, D.J. (1970):
Plasma protein in normal urine.
Am. J. Physiol., 219: 358.
- Maher, J.F. (1985):
Tubular proteinuria: Clinical Implications.
In: Proteinuria, by Avram, M.M., p. 57.
Plenum Medical Book Company, New York and London.
- McCracken, B.H. (1984):
Estimations of quantitative proteinuria.
N. Engl. J. of Med., May 31.
- McEnery, P. and Strife, C. (1982):
Nephrotic syndrome in childhood. In: Symposium
on Pediatric Nephrology from Ped. Clinics of
North America, 875-895.

- **Militenyi, M. (1979):**
Urinary protein excretion in healthy children.
Clin. Neph., Vol. 12, No. 5, pp. 216-221).
- **Mogensen, C.E. and Vittinghus, E. (1975):**
Urinary albumin excretion during exercise in
juvenile diabetics.
Scand. J. Clin. Lab. Invest., 35: 295-300.
- **Murphy, W.M.; Moretta, F.L. and Jukkola, A.F. (1979):**
Epithelial foot process effacement in patients with
proteinuria.
Am. J. Pathol., p. 529.
- **Newman, W.J.; Tisher, C.G.; McCay, R.C. and Robinson,
R.R. (1976):**
Focal glomerulosclerosis: Contrasting clinical
patterns in children and adults.
Medicine, 55, 67.
- **Oken, D.E. and Flamenbaum, W.J. (1971):**
Osmotic pressure of plasma and serum albumin.
J. Clin. Invest., 50: 1498.
- **Paul, T.M. and Frederic, S. (1982):**
Nephrotic syndrome associated with other disorders.
Ped. Clin. of North Am., Vol. 89, No. 4.
- **Pece, A.J. and Furst, M.R. (1985):**
Proteinuria: an integrated review.
New York: Marcel Dekker Inc.
- **Penn, G.M. (1981):**
Clinical use and interpretation of Agarose-Gel
Electrophoretic Patterns.
Amer. Soc. Clin. Path., Chicago III.

- **Pesce, A.J.; Gaisutis, M. and Pollak, V.E. (1970):**
Quantitative determination of protein in urine.
J. Lab. Clin. Med., 75: 586.
- **Reuben, D.B.; Wachtel, T.J. and Brown, P. (1982):**
Transient proteinuria in emergent medical admissions.
N. Engl. J. Med., 306: 1031-3.
- **Righard, L. (1980):**
Urinary screening in children 3-7 years old.
Lancet, 1: 1369.
- **Robinson, R.R. (1985):**
Clinical significance of isolated proteinuria.
In: Proteinuria, by Avram, M.M., p. 67.
Plenum Medical Book Company, New York & London.
- **Ryan, G.B. and Karnovesky, M.J. (1975):**
Clinical aspects of the plasma proteins.
Kidney Int., 219.
- **Sargent, P. and Johnson, R.E. (1956):**
The effects of diet on renal function in healthy men.
Am. J. Clin. Nutr., 4: 466-81.
- **Shaw, A.B.; Risdon, P. and Levis-Jackson, J.D. (1983):**
Protein creatinine index and albustix in assessment
of proteinuria.
Br. Med. J., Vol. 287, Oct., 1.
- **Siwach, S.R.; Kalra, O.P.; Sharma, R.; Singh, V. and
Chapra, J.S. (1990):**
Estimation of 24 hour protein excretion from single
urine specimen.
Indian J. Med. Res., Apr., 92, p. 105-8.

- **Stewart, D.W.; Gordon, J.A. and Schoolwerth, A.C. (1984):**
Evaluation of proteinuria.
Am. Fam. Physician, 29 (4), p. 218-25.
- **Tamm, I. and Horsfall, F.L., Jr. (1952):**
A mucoprotein derived from human urine which reacts
with influenza mumps and newcastle disease viruses.
J. Exp. Med., 95: 71-97.
- **Tiggler, R.G.; Hulme, B. and Wijdeveld, P.G. (1979):**
Effect of indomethalin on glomerular permeability
in the nephrotic syndrome.
Kidney Int., 16: 312.
- **Varley, F.R., A.G. and Bell, F.R. (1980):**
Practical clinical biochemistry, determination of
protein in urine, p. 406.
William Heineman Medical Books LTD., London.
- **Vehaskari, V.M. and Rapola, J. (1982):**
Isolated proteinuria. Analysis of a school-age
population.
J. of Pediatr., Vol. 101, No. 5, pp. 661-668.
- **Viberti, G.C.; Jarrett, R.J. and McCartney, M. (1978):**
Increased glomerular permeability to albumin
induced by exercise in diabetic subjects.
Diabetologia, 14: 293-300.
- **White, R.H.R. (1973):**
The familial nephrotic syndrome.
European Survey.
Clin. Nephrol., 1: 215.

- Wilkin, T.. Mohamed, A.; Gatling, W. and Rowe, D. (1983):
Protein creatinine index and albustix in assessment
of proteinuria.
Br. Med. J., Vol. 287, Dec., 17.
- Williams, D.G.; Peters, D.K.; Fallows, J. et al. (1974):
Studies of serum complement in the hypocomplementemic
nephritis.
Clin. Exp. Immunol., 18: 391.
- Woodroffe, A.J.; Gommly, A.A. and Mackenzie, P.E. (1980):
Immunologic studies in IgA nephropathy.
Kidney Int., 18: 366.
- Yamada, T.; Ishihara, M. and Ichikawa, K. (1980):
Proteinuria and renal function during antihyper-
tensive treatment for essential hypertension.
J. Am. Geriatr. Soc., 28: 114-7.
- Yoshikatsu, M.D. (1983):
Nephrotic syndrome with diffuse mesangial sclerosis
in identical twins.
The Journal of Pediatrics, pp. 586.
- Yu, H.; Yanagisawa, Y.; Forbes, M.A. and Cooper, E.H. (1983):
Alpha-1-microglobulin, an indicator protein for
renal tubular function.
J. Clin. Path., 36, 253.
- Zilleruelo, C.; Abilbol, G.; Freundlich, M. and Strauss,
J. (1990):
Quantitation of proteinuria with urinary protein/
creatinine ratios and random testing with dipsticks
in nephrotic children.
J. Pediatr., Feb., 116 (2), p. 243-7.