

References

Abdul - Karim, R. and Assali, N.S.:

Renal function in human pregnancy and the effect of oxytocin on renal hemodynamics and water and electrolyte excretion. J. Lab. Clin. Med. 57:522-531. 1961.

Adams, J.A.; Hey, D.J. and Hall R.T.:

Incidence of hyperbilirubinemia in breast vs formula-fed infants. Clin. Pediatre. 24:69-73, 1985.

Amanullah, A.:

Neonatal jaundice. Am. J. Dis. Child. 130:1274. 1976.

Anderson. A.; Gawson, M. and Turnbull, A.C.:

Neonatal jaundice and maternal oxytocin infusion. Br. Med. J., 11:476. 1973.

Austin, R.F.; Desforbes, J.F.:

Hereditary elliptocytosis: An unusual presentation of hemolysis in the newborn associated with transient morphologic abnormalities. Pediatrics. 44:196. 1969.

Bakken, A.F.; Thaler, M.M. and Schmid, R.:

Metabolic regulation of heme catabolism and bilirubin production. In: Hormonal control of hepatic heme-oxygenase activity. J. Clin. Invest. 51:530. 1972.

Barrett, P.V.:

Bilirubinemia and fasting. N. Engl. J. Med. 283:823. 1970.

Bartoletti A.L.; Stevenson, D.K.; Ostrander, C.R., Johnson, J.D.:

Pulmonary excretion of carbon monoxide in the human infant as an index of bilirubin production. J. Pediatr. 91:952. 1979.

Beazley, J.M. and Alderman, B.:

Neonatal hyperbilirubinemia following the use of oxytocin in labour. Br. J. Obst. Gyn. 82:265-271, 1975.,

Behrman, R.E.; Kliegman, R.M.:

Jaundice and hyperbilirubinemia in the newborn, In: Behrman R.E.; Vaughn, V.C. III (eds):Nelson Textbook of Pediatrics (12th ed). Philadelphia W.B.Saunders Co.:3778-381. 1983.

*Berk, P.D.; Blaschke, T.K.; Scharschmidt, B.F.; Waggoner J.G.;
Berlin, N.I. :*

A new approach to quantitation of the various sources of
bilirubin in man. J. Lab. & Clin. Med. 87:767. 1976.

Bhargava, M.M.; Arias, I.M.:

Ligandin. Trends in Biochemical Science 6:131. 1981.

Blum, R.M. and Foster, R.E.:

The water permeability of erythrocytes. Biochemica et
Biophysica. Acta. 203:410-423. 1970.

Boylan P.:

Oxytocin and neonatal jaundice Br. Med. J.; 2 (5986)
564-65., 1976.

Bracci R; Buonocore G; Garosi G; Bruchi S; Berni S. :

Epidemiologic study of neonatal jaundice. A survey of
contributing factors. Acta. Pediatr. Scand. Suppl; 360;
P87-92, 1989. (Abstract)

Brodersen, R.:

Competitive binding of bilirubin and drugs to human serum albumin studied by enzymatic oxidation. J. Clin. Invest. 54:1353. 1974.

Brown, S.B.; King, R.F.:

The mechanism of heme catabolism. Biochemical Journal 170:297. 1978.

Buchan, B.C.:

Pathogenesis of neonatal hyperbilirubinemia after induction of labour with oxytocin. Br. Med. J. 11:1255-1257. 1979.

Calder, A.A.; Moar, V.A.; Ounsted, M.K. and Turnbull, A.C.:

Increased bilirubin levels in neonates after induction of labour by intravenous prostaglandin E₂ or oxytocin. Lancet 2:1339-1342, 1974.

Campbell, N.; Harvey, D. and Norman, A.P.:

Increased frequency of neonatal jaundice in a maternity hospital. Br. Med. J. 2:548-552, 1975.

Cashore, W.J.; Oh, W.:

Unbound bilirubin and kernicterus in low birth weight infants. *Pediatrics* 69:485, 1982.

Chalmers, I.; Campbell, M. and Turnbull A.C.:

Use of oxytocin and incidence of neonatal jaundice. *Br. Med. J.* 2:116-118, 1975.

Chalmers, I.; Campbell, M. and turnbull, A.C.:

Oxytocin and neonatal jaundice. *Br. Med. J.* 1:647-648, 1976.

Chard, T.; Hudson, C.N.; Edwards, C.R.W and Boyad, N.R.H.:

Release of oxytocin and vasopressin by the human fetus during labour. *Nature "London"* 234:252, 1971.

Quoted from: Dawood M.Y., Wang, C.F.; Gupta, R. and Fucks,

F:Fetal contribution to oxytocin in human labour. *Obst. Gyn.* 52:205-209, 1978.

Chew, W.C and Swann, I.L.:

Influence of simultaneous low amniotomy and oxytocin

infusion and other maternal factors on neonatal jaundice.
Br. Med. J. 1:72-73, 1977.

Clark, D.A and Landsaw, S.A.:

Bupivacaine alters red blood cell properties:A possible explanation of neonatal jaundice associated with maternal anaesthesia. *Pediatr. Res.* 19:341-343, 1985.

Cole, P.V.; Hawkins, L.H and Roberts, D.:

Smoking during pregnancy and its effects on the fetus. *J. Obst. Gyn. Br. Comm.* 79:782-787, 1972. Quoted from Diwan V.K.; Vaughan T.L.Yang C.Y.: Maternal smoking in relation to the incidence of early neonatal jaundice. *Gynecol. Obstet. Invest;* 27(1); P22-5, 1989.

Coombs R.R.A.; Mourant, A.E. and Race, R.R.:

A new test for the detection of weak and incomplete Rh-agglutinins. *Brit. J. Exp. Path.* 26:225, 1945.

Cottrell, B.H and Anderson, G.C.:

Rectal or axillary temperature measurement:Effect on plasma bilirubin and intestinal transit of meconium. *J. Pediatr. Gastroenterol. Nutr.* 3:734-739, 1984. Quoted form: Maisels,

M.J. and Gifford K.:Normal serum bilirubin levels in the newborn and the effect of breast feeding. *Pediatrics* 78:837, 1986.

Dacie, J.V and Lewis, S.M. (eds).:

Practical Hematology:Blood groups and laboratory aspects of blood transfusion. Chap. 19, (5 th ed). Churchill livingstone. New York. 1975.

Dahlenburg, G.W.; Burnell, R.H and Braybrook, R.:

The relation between cord serum sodium levels in newborn infants and maternal intravenous therapy during labour. *Br. J. Obst. Gyn.* 87:519-522, 1980.

Davidson, D.C. Ford, J.A and McIntosh, W.:

Neonatal jaundice and maternal oxytocin infusion. *Br. Med. J.* 4:106,1973.

Davies, D.P.; Gommersall, R.; Robertson, R.; Gray. O.P and Tyrnbull. A.C.:

Neonatal jaundice and maternal oxytocin infusion. *Br. Med. J.* 3:476-477, 1973.

Dawood, F.S.K and Dawood, M.Y.:

Oxytocin content of human pitutary gland. Am. J. Obst. Gyn.
148 (4):420-422, 1984.

Dawood, M.Y.; Wange, C.F.; Gupta, R. and Fucks, F.:

Fetal contribution to oxytocin in human labour Obst. Gyn.
52:205-209. 1978.

Dawson, J.B.:

Instruments using measurement of light intensity. Varly's,
praetical clinical biochemistry, 6th ed. Heineman Medical
books, london chapter 7 P:129-175., 1978.

De Hemptienne, D.; Schuddinck, L.; Thiery. M and Martens, G.:

Neonatal hyperbilirubinemia following the use of
prostaglandin E₂ in labour. International Research
Communication System. 37:12, 1973. Quoted from:Lang A.P.;
Neonatal jaundice after labour induced or augmented by
oxytocin. Lancet 1:991-994, 1982.

Dhumeaux D. and Berthelot, P.:

Chronic hyperbilirubinemia associated with hepatic uptake
and storage impairment. Gastroenterology; 69:988., 1975.

Diwan V.K.; Vaughan T.L., Yang C.Y.:

Maternal smoking in relation to the incidence of early neonatal jaundice. Gynecol. Obstet. Invest; 27(1); P 22-5, 1989. (Abstract).

Drabkin, D.L. and Austin, J.H.:

Spectrophotometric studies. J. of biological chem. 98:719., 1932 (Q) from Dacie, J.U. and Lewis, S.M. 1984.

Drew J.H. and Kitchen W.H.:

The Effect of Maternally Administered Drugs on Bilirubin Concentrations in the Newborn Infant. The Journal of Pediatrics:657-661, 1976.

D'souza, S.W. Black, P. Macfarlane. T and Richard, B.:

The effect of oxytocin in induced labour on the neonatal jaundice. Br. J. Obst. Gyn. 86:133-138, 1979.

Du Vigneaud, V. Ressler, C. and Tripelt, S.:

The pharmacology and clinical application of various oxytocic drugs. J. Biol. Chem. 205:949, 1953. Quoted from:Munsick, E.A. The pharmacology and clinical application of various oxytocic drugs. Am. J. Obst. Gyn. 93(3):442, 1965.

Eggers, T.R., Fliegner, J.R.:

Water intoxication and syntocinon intoxication. Aust. Nz J. Obstet. Gynaecol. 19:59, 1979. Qouted from Feeney, J.G.:Water intoxication and oxytocin, Br. Med. J. 285:243, 1982.

Embrey, M.:

The effect of prostaglandins on the human pregnant uterus. J. Obst. Gyn. Brit. Commonw. 76:783, 1969.

Embey, M.P. :

Oxytocics and the uterus. In:Scientific foundations of obstetrics and Gynacology. (Ed). Philpp., E., Barnes, J. & Newton, M. Second Edition. P. 798-810. London; William Heinemann. 1977.

Eriksen, L.:

Congenital erythropoietic porphyria, the effect of light shielding. Acta. Pediat. Scand. 62:385. 1973.

Fanaroff, A.A and Martin, R.J. (eds):

Jaundice and liver disease. In:Diseases of The fetus and Neonatal-Perinatal Medicine (4th ed), Chap, 14, The C.V. Mosby Company. 1987.

Feeney, J.G.:

Water intoxication and oxytocin. Br. Med. J. 285:243, 1982.

Fessas, P.; Doxiadis, S. and Valaes. T.:

Neonatal jaundice in glucose-6-phosphate dehydrogenase-deficient infants. Br. Med. J. 2:1359, 1962.

Folliot, A.; Ploussard, J.P.; Housset, E. and Christo Porov, B.:

Breast milk jaundice: In vitro inhibition of rat-liver bilirubin diphosphate glucuronyl transterase activity and z protein-bromosulphaphthlen binding by human breast milk. Pediatr. Res. 10:594, 1976.

Francis, J.C.; Turnbull, A.C. and Thomas, F.F.:

Automatic oxytocin infusion equipment for induction of labour. J. Obst. Gyn. Br. Commonw 77:594-602, 1970.

Friedman, E.A. and Sachtleben, M.R.:

Oxytocin and neonatal jaundice. Lancet 11:600, 1974.

Friedman, E.A. and Sachtleben, M.R.:

Neonatal jaundice in association with oxytocin stimulation

of labour and operative delivery. Br. Med. J. 1:198-199, 1976.

Friedman, L.; Lewis, P.J.; Clifton, P. and Bulplit, C.I.:
Factors influencing the incidence of neonatal jaundice. Br. Med. J. 1:1235-1237, 1978.

Gale R; Seidman D.S.; Dollberg S; Stevenson D.K.:
Epidemiology of neonatal jaundice in the jerusalem population J. Pediatr. Gastroenterol. Nutr; 10(1); P 82:6, 1990. (Abstract).

Gartner, L.M.; Lee, K.S.:
Unconjugated hyperbilirubinemia. In Fanaroff A, Martin R 12th edition:Behrman's Neonatal-Perinatal Medicine, St Louis, Cv Mosby, 1983.

Gartner, L.M.; Lee, K.S.; Vaisman, S. et al.:
Development of bilirubin transport and metabolism in the newborn Rhesus monkey. J. Pediatr. 90:513, 1977.

Gartner, L.M., and Whittington, P.F.:

Disorders of bilirubin metabolism. In: Nathan D.S., Oski, F.A. (eds). Hematology of Infancy and Children Chap. 4 (3rd ed). W.B. Saunders. Co. 1987.

Gemelli M; Deleo S; Tuccio G :

Neonatal hyperbilirubinemia. Lancet 20; 335 (8682); P 179, 1990.

Ghosh, A. and Hudson, F.P.:

Oxytocic agents and neonatal hyperbilirubinemia. Lancet 2:823, 1972.

Gollan, J.L.; Schmid, R.:

Bilirubin update: Formation, transport and metabolism. In: Popperlt, Schaffner, F. (eds). Progress in liver Diseases Vol. III, Grune and Stratton, New York. 261, 1982.

Gould. S.R.; Mountrose. V.; Brown, D.J. Whitehouse, W.L. and Barnardo, D.E.:

Influence of previous oral contraceptives and maternal oxytocin infusion on neonatal jaundice. Br. Med. J. 3:228-230, 1974.

Gourley, G.R. and Arend, R.A.:

B-Glucuronidase and Hyperbilirubinemia in Breast-fed and formula-fed Babies *lancet* 1:644, 1986.

Gray H.G. Mitchell R.:

Neonatal hyperbilirubinemia and oxytocin. *Lancet* 2:1144, 1974.

Gross, S.J.:

Vitamin E and neonatal bilirubinemia. *Pediatrics* 64:321-323, 1979.

Hamad S.A; El-Domiaty B.A, Nayel S.A.:

Neonatal Hyperbilirubinemia in Oxytocin Augmented Labour, *Asia-Oceania J. Obstet. Gynaecol.* Vol. 11, No. 1:69-73, 1985.

Hancock, D.J.; Setzer, E.S. and Beydown, S.N.:

Physiologic and biochemical effects of ritodrine therapy on the mother and perinate. *Am. J. Perinatol.* 2:1-6, 1985.
Quoted from Linn, S.; Schoenbaum, S.C.; Monson, R.R.; Rosner, B.; Stubblefield, P.G. and Ryan, K.J.: *Epidemiology*

of neonatal hyperbilirubinemia. Pediatrics 75(4):770-774, 1985.

Hardy, J.B. and Mellits. E.D.:

Does maternal smoking during pregnancy have a long-term effect on the child? Lancet 2:1332-1336. 1973.

Hopper, A.O.; Cohen, R.S.; Ostrander, C.R.; Brackman, F.S.;

Uekand, K. and Stevenson. D.K.:

Maternal adrenergic tocolysis and neonatal bilirubin production. Am. J. Dis. Child. 137:58-60,1983.

Hubbell, J.P.; Drorbaugh. J.E. and Rudolph. A.J.:

Early versus late feeding of infants of N. Engl. J. Med. 265:835. 1961.

Huszar, G.:

Cellular aspect of labour. In premature labour. Mead. Johnson Symposium on Perinatal and Developmental Medicine 15:16-25. 1980.

Ingemar, L.; Finnstrom. O.; Hedenskog, S.; Ryden. G and Tyllskar, J.:

Spontaneous labour and elective induction. Bilirubin levels in the neonatal period. Acta. Obstet. Gynecol. Scand.

59:103-106, 1980.

James, L.S.:

Onset of breathing and resuscitation. *Pediatr. Clin. North Am.* 13:621, 1966.

Jansen, P.L.M.; Chowdlaury, J.R. and Fischberg, E.G.:

Enzymatic conversion of bilirubin monoglucuronide to diglucuronide by rat liver plasma membranes. *J. Biol. Chem.* 252:2710-2716. 1977.

Jendrassik, L. and Coll, E.:

Measurement of Bilirubin by Colorimetric Method. *Biochem. Z* 297:81, 1938.

Johnson, J.D.; Aldrich, M.; Angelus, P.; Stevenson, D.K. Smith,

D.W.; Herschel M.J.; Papagaroufalis, C. and Valaes, T.:

Oxytocin and neonatal hyperbilirubinemia. Studies of bilirubin production. *Am. J. Dis. Child.* 138:1047-1050, 1984.

Karim, S.:

Appearance of prostaglandins F₂ in human blood during labour. *Br. Med. J.* 4:618, 1968.

Kawade, M.; Onishi, S.:

The prenatal and postnatal development of UDP-glucuronyl transferase activity towards bilirubin and the effect of premature birth on this activity in the human liver.

Biochemical Journal 196:257, 1981.

Kenepp, N.B.; Shelley, W.C.; Kumar, S. Gutsche. B.B.; Gabbe, S.

and Delivoria-Papodopoulos. M.:

Effects on newborn of hydration with glucose in patients undergoing caesarean section with regional anaesthesia.

Lancet 1:645. 1980.

Kivlahan C, James E.J.P:

The natural history of neonatal jaundice. Pediatrics 74:364,

1984.

Kontras, S.B.:

Polycythemia and hyperviscosity syndromes in infants and children. Ped. Clin. North Am. 19:919, 1972.

Kozuki, K.; Cashore, W.J.; Widness, J; Ohw.:

Increase in bilirubin-albumin binding affinity with correction of neonatal acidosis. Acta. Pediat. Scand.

68:213. 1979.

Kuhr M. Paneth N.:

Feeding practices and early neonatal jaundice. J.Pediatr. Gastroenterol. Nutr. 1:485-488, 1982. Qouted from: Maisels, M.J. and Gifford, K.: Normal serum bilirubin levels in the newborn and the effect of breast feeding. Pediatrics 78:837, 1986.

Lange, A.P.; Secher, N.J.; Westergard, J.G. and Skovgard, I.B.:

Neonatal jaundice after labour induced or stimulated by prostaglandin E₂ or oxytocin. Lancet 1:991-994, 1982.

Leibson C; Brown M; Thibodeau S; Stevenson D; Verman H; Cohen R; Clemons G; Callen W; Moore L.G.:

Neonatal hyperbilirubinemia at high altitude. Am. J. Dis. Child; 143(8); 983-7, 1989.

Levine, P.:

Blood group antigens and antibodies as applied to hemolytic disease of the newborn. Orthordiagnosics. Raritan, New Jersey. 1968. Quoted from: Valentine W.N.; Haemolytic anemia and inborn errors of metabolism. Blood 54:549, 1979.

Levitt, M.; Schacter B.S.; Zipursky, A. et al.:

The non-erythropoietic component of early bilirubin. J. Clin. Invest. 27:1281, 1968.

Lewis, P.J. and Friedman, L.A.:

Prophylaxis of neonatal jaundice with neonatal antipyrine treatment. Lancet 1:300-302, 1979.

Li, A.M.C.; Yeung, C.Y.; Chang, W.K. and Soo, H.N.:

Epidemiological aspects of neonatal jaundice in Chinese infants. Trop. Geogr. Med. 31:537-546, 1979.

Liggins, G.C. and Howie, R.N.:

A controlled trial of antepartum glucocorticoid treatment for prevention of the respiratory distress syndrome in premature infants. Pediatrics 50:515, 1972.

Linn, S.; Schoenbaum, S.C.; Monson, R.R.; Rosner, B.;

Stubblefield, P.G. and Ryan, K.J.:

Epidemiology of neonatal hyperbilirubinemia. Pediatrics 75 (4):770-774, 1985.

Liston, W.A. and Campbell, A.J.:

Dangers of oxytocin induced labour to fetuses. Year Book of Obst. Gyn. 138-139, 1975.

Lucas, A.; Adrian, T.E.; Aynsley-Green. A and Bloom, S.R.:

Iatrogenic hyperbilirubinism at birth. Lancet 1:144, 1980.

Lux S.E.:

Disorders of red cell membrane. In Nathan D.G. Oski F.A. (eds). Haematology of infancy and childhood 3rd ed. Philadelphia, W.B. Saunders company, 1987.

Luzzatto. I.:

Genetic heterogeneity and pathophysiology of G6PD deficiency. Brit. J. Hematology 28:151. 1974.

MacGillivray, M.H.; Crawford, J.D. and Robey, J.S.:

Congenital hypothyroidism and prolonged neonatal hyperbilirubinemia. Pediatrics 40:283. 1967.

Maisels, M.J.:

Bilirubin: On understanding and influencing its metabolism in

the newborn infant. *Ped. Clin. North. Am.* 19:447. 1972.

Maisels, M.J.:

Neonatal Jaundice. In: Avery, G.B. (ed.): *Neonatal Pathophysiology and Management of the Newborn*. 3rd ed., P. 484, Philadelphia. Lippincott J.B. Co., 1987.

Maisels, M.J. and Gifford, K.:

Normal serum bilirubin levels in the newborn and the effect of breast feeding. *Pediatrics* 78:837, 1986.

Maisels, M.J.; Gifford K.; Antie, C.E. and Leib, G.R.:

Jaundice in the healthy newborn infant; A new approach to an old problem. *Pediatrics* 81:505, 1988.

Maisels, M.J.; Leib. G.; Gifford. K. and Antie, C.:

The yellow baby syndrome (or why well babies are jaundiced). *Pediatr. Res* 19:240 A, 1985.

Mast. H.; Quakermack. K.; Lenfers. M. and Hagen. C.:

Der Einfluss des Gburtsverlaufes anf den Ikterus neonatorum.
Geburtshille and Frauenheilk 31:443, 1971. Quoted
from:Hibbard B.M.: Inductrion of labour. Obstetric Textbook.
Butterworth. London. Boston, Durban Singapore. Sydney.
Torouto, Wellington, 439-448, 1988.

Mc Connell, J.B.; Glasgow, J.F.T. and Mc Nair, R.:

Influence of previous oral contraception and maternal
oxytocin infusion on neonatal jaundice. Br. Med. J.
3:605-607. 1973.

Mee, W.M.; Friddian, R.V. and Rothwell-Jackson, R.L.:

Oxytocin and neonatal jaundice Br. Med. J. 1:293, 1978.

Meisel, P.; Jahring, D.; Beyersdor, E. and Jahring. K.:

Bilirubin binding and acid-base equilibrium in newborn
infants with low birth weight. Acta Pediat. Scand. 77:496.
1988.

Miller, C.A. and Reed. H.R.:

Oxytocics and neonatal hyperbilirubinemia. Pediatr. 21:362,
1958. Quoted from Ghosh. A and Hudson. F.P.: Oxytocic agent

and neonatal hyperbilirubinemia. Lancet. 2:823, 1972.

Mollison, P.L.:

Blood transfusion in clinical medicine (6th ed). Chap, 16
Oxford Blackwell. Scientific Publication. 666-722. 1979.

Moore, L. G.; Newherry, M.A.; Freeby, G.M.; Crnic, L.S.:

Increased Incidence of Neonatal Hyperbilirubinemia at 3,100
m. in Colorado. Am. J. Dis Child. 138:157. 1984.

*Morgan, D.B.; Kirwan, N.A.; Hancock, K.W.; Robinson, D.; Howe,
J.G. and Ahmed. S.:*

Water intoxication and oxytocin infusion Br. J. Obst. Gyn.
84:6-12. 1977.

Munroe, M.; Shah. C.P.; Badgley, R. and Bain. H.W.:

Birth weight, length head circumference and bilirubin level
in indian newborns in the Sioux lookout zone. Northwestern
Ontario. Cand. Med. Assoc. J. 131:453-456, 1984

Murphy, B.E.P.:

Does the human fetal adrenal play a role in parturition? Am. J. Obst, Gyn. 115:521-525, 1971.

Nathenson, G.; Cohen, M.I.; Litt, I.F. and McNamara. H.:

The effect of maternal heroin addiction on neonatal jaundice. J. Pediatr. 81:899-903, 1972.

Nelson, T.; Jacobsen, J.; Wennberg, R.P.:

Effect of pH on interaction of bilirubin with albumin and tissue culture cells. Pediatr. Research 8:963. 1974.

Nemeth, I. and Szeleczki, B.D.:

Hyperbilirubinemia and urinary D-glucose and excretion in premature infants following antepartum dexamethasone treatment. J. Perinat. Med. 9:35-39, 1981. (Abstract)

Newman T.B.; Maisels M.J.:

Does hyperbilirubinemia damage the brain of healthy full-term infants? Pediatrics 83(6); P 1062-64; 1989.

Nobel, A.D.:

Oxytocin and neonatal jaundice Br. Med. J. 1:239, 1978.

Norman, A.P.; Harvey, D.R.; Kovar, I. and Campbell.:

Oxytocin Agents and Neonatal Hyperbilirubinemia. Lancet, 2, 972, 1972.

Nymand, G.:

Maternal smoking and neonatal hyperbilirubinemia. Lancet 2:173. 1974.

Ode11, G.B.:

Neonatal jaundice, In Popper H. and Schaffner, F. (eds) Progress in liver disease. New York, Grune and Stratton. Inc. 457-475, 1976.

Odiever, M. and Luzeaur. R.:

Lipolytic activity in milk from mother of unjaundiced infants. Acta Pediatr. Scand. 67:49054, 1978.

O'Driscoll, K. and Meagher, D.:

Oxytocin in labour in: Active Management of Labour; Edited by O'Driscoll, K. and Meagher, D.; Second edition; condon, Rio de Janiro Bailliere Tindall Company; P.42-46, 1986.

Oski F.:

Oxytocin and neonatal hyperbilirubinemia. Am J. Dis child, 129; 1139-1140, 1975.

Oski, F.A.:

Neonatal jaundice. General consideration. In: Schaffer's Disease of the Newborn. 5th ed. Avery and Taeasch. W.B. Saunders Co., Philadelphia, P. 622-624, 1984.

Owens, D.; Sherlock, S.:

Diagnosis of Gilbert's syndrome; role of reduced caloric intake test. Br. Med. J. 3:559, 1973.

Perlman, M. and Frank J.W.:

Bilirubin beyond the blood brain barrier. Pediatrics 81:304, 1988.

Philipp, K.; Pateisky, N. and Endler, M.:

Effects of smoking on enteroplacental blood flow. Gyn. Obst. Invest. 17:179-182, 1984.

Pimstone, N.R.; Engel, P.; Tenhunen, R.; Seitz, P.T.; Marver, H.S.; Schmid, R.:

Inducible heme-oxygenase in the kidney: A model for the homeostatic control of hemoglobin catabolism. J. Clin. Invest. 50:2042, 1971.

Platto, Nathan D.G.:

Sickle cell disease In Nathan D.G., Oski F.A. (ed): Haematology of infancy and childhood 3rd ed. Philadelphia, W.B. Saunders Company, 1987.

Pocock, S.J. and Turner, T.L.:

Neonatal hyperbilirubinemia, oxytocin and sex of infant. Lancet 11:43-44, 1982.

Poland, R.L. and Odel, G.R.:

Physiologic jaundice. The enterohepatic circulation of bilirubin. N. Eng. J. Med. 284:1-6, 1971.

Pritchard, J.A.; MacDonald, P.C. and Gant, N.F.:

Oxytocic Agents in: Williams obstetrics; 18th edition, Appleton-century-Crofts, New York, P 345-347. 1989.

Rall, T. and Schleifer, L.:

Oxytocin, prostaglandins, ergot alkaloids and other agents.

Quoted from: Gillman, A.F.; Goodman, L.S. and Gilman, A.

(eds.): The Pharmacological Basis of Therapeutics. ed. 6. New York, Macmillan, P. 935-950, 1980.

Reyes, H.; Levi, A.J.; Gaatmaitan, Z. et al.:

Studies on Y and Z, two hepatic cytoplasmic organic anion-binding proteins. Effects of drugs, chemicals, hormones. J. Clin. Invest. 50:2242, 1971.

Ritter, D.A.; Kenny, J.D.:

Bilirubin binding in premature infants from birth to 3 months. Arch. Dis. Child 61:352. 1986.

Ritter D.A. Kenny J.D.; Norton H.J.:

A prospective study of free bilirubin and other risk factors in the development of kernicterus in premature infants. Pediatrics 69:260, 1982.

Schwartz, R.H. and Jones, R.W.A.:

Transplacental hyponatremia due to oxytocin. Br. Med. J.
1:152-153, 1978.

Sigstrom, L.:

Role of active sodium and potassium transport in
hyponatremic states in infancy and childhood. Acta Paediat.
Scand. 70:353-359, 1981.

Sims, D.G. and Neligan, G.A.:

Factors affecting the increasing incidence of severe
nonhemolytic neonatal jaundice. Br. J. Obst. Gyn.
82 : 863-867. 1975.

Singhi, S.; Choo-Kang, E. and Hall, J. ST. E.:

Hazards of maternal infusion with 5% dextrose. Lancet
11:335-336, 1982.

Singhi, S.; Choo-Kang, E. and Hall, J. ST. E.:

Intrapartum infusion of aqueous glucose solution, transpla-
cental hyponatremia and risk of neonatal jaundice.
Br. J. Obst. Gyn. 91:1014-1018, 1984.

Singhi, S.; Choo-Kang. E.; Hall, J. ST. E. and Kolghatgi, S.:

Iatrogenic neonatal and maternal hyponatremia following oxytocin and aqueous glucose infusion during labour. Br. J. Obst. Gyn. 92:356-363, 1985.

Singhi. S. and Singh, M.:

Pathogenesis of oxytocin induced neonatal hyperbilirubinaemia. Arch. Dis. Child. 54:400-402. 1979.

Singhi, S. and Singh, M.:

Erythrocytic osmotic fragility in neonates born following oxytocin infusion during labour. Indian J. Med. Res. 72:843-845, 1980. Quoted from: Johnson, J.D.; Aldrich, M.; Angelus. P.; Oxytocin and neonatal hyperbilirubinemia. Studies of bilirubin production. Am. J. Dis. Child. 138:1047-1050, 1984.

Spencer, S.A.; Mann, N.P.; Smith, M.L.; Woolfson, A.M.H. and Benson, S.:

The effect of intravenous therapy during labour on maternal and cord serum sodium levels. Br. J. Obst. Gyn. 88:480-483, 1981.

Stroup, M. and MacIlroy, M.:

Evaluation of the albumin antiglobulin technic in antibody detection. *Transfusion* 5:184, 1965.

Tarnow-Mordi, W.O.; Shaw, J.C.L.; Liu, D.; Gardner, D.A. and Flynn, F.V.:

Iatrogenic hyponatremia of newborn due to maternal fluid overload, a prospective study. *Br. Med. J.* 283:639-642, 1981.

Taylor, P.M.; Bright, N.H. and Birchard, E.L.:

Effects of early versus delayed clamping of the umbilical cord on the clinical condition of the newborn infants. *Am. J. Obstet. Gynec.* 84:893, 1963.

Theobald, G.:

Induction of labour *J. Reprod. Med.* 6:1-34, 1971.

Theobald G.W.:

The induction of labour. *Obstetric therapeutics*. Ed. Hawkins. D.F. London P. 431, 1974.

Thiery M, de Hemptinne D, Schuiddinck L, et al:

Neonatal jaundice after induction of labour. Lancet, 1:161, 1975.

Tricomi, V.:

Induction of labour A contemporary view, clinical obstetrics and Gynaecology vol. 16, 4, 226-242, 1973.

Trolle, D.:

Phenobarbitone and neonatal icterus. Lancet 1:251, 1968.

Tuppy H.:

The influence of enzymes on neurohypophysial hormone and similar peptides. In: Handbook of experimental pharmacology-edited by Berde. New york:Springer verlag, 23:13., 1968.

Turkel S.B., Guttenberg M.E, Moynes D.R., et al.:

Lack of identifiable risk factors for kernicterus, Pediatrics 66:502, 1980.

hyperbilirubinemia. J. Pediatr; 117(2 pt 1); P 288-91, 1990.

Weiland, A. and Langer, H.:

Geburtsch. Frauenheilk 25:320, 1965. Quoted from:Calder, A.A.; Moar, V.A.; Qunsted. M.K. and Turnbull, A.C.:Increased

bilirubin levels in neonates after induction of labour by intravenous prostaglandins E₂. or oxytocin. Lancet 2:1339, 1974.

Weisiger, R.; Gollan, J.; Ockner, R.:

Receptor for albumin on the liver cell surface may mediate uptake of fatty acids and other albumin bound substances. Science 211:1048, 1981.

Wennberg, R.D.:

Bilirubin physiology, Part 1, Gastroenterology, Chap. 18, In:Roberton N.R.C. (ed.) Textbook of Neonatology, Churchill Livingstone, Edinburgh. 383-390, 1986.

Wennberg, R.D.; Schwartz, R. and Sweet, A.Y.:

Early versus delayed feeding of low birth weight infants:Effects on physiologic jaundice. J. Pediatr.

68:860-866, 1966.

Wintrobe, M.M.; Richard, G.; Boggs, D.R.; Bithe11, T.C.;

Foerster, J.: Athens, J.W. and Lukens, J.N. (eds.):

Isoimmune hemolytic anemia. In: *Clinical Hematology* (8th ed).

Chap. 37, LEA & Febiger, 904-920, 1981.

Wong, Y.K. and Wood, B.S.P.:

A multifactorial survey of neonatal jaundice. *Br. Med. J.*
4:403, 1971.

Wood, B.; Culley, P.; Roginski, C. Powell, J.; Waterhouse, J.:

Factors affecting neonatal jaundice. *Arch. Dis. Child.*
54:11, 1979.

Wu, P.Y.K.; Teilman, P. and Goble, M.:

Early versus late feeding of low birth weight neonates:
Effect on serum bilirubin, blood sugar and responses to
glucagon and epinephrine tolerance tests. *Pediatrics*
39:733-739, 1967.

*Wysowski, D.K.; Flynt, J.W.; Goldried, M.; Altman, R. and Davis,
A.T.:*

Epidemic neonatal hyperbilirubinemia and use of a phenolic disinfectant detergent. *Pediatrics* 61:165, 1978.

Yao, A.C. and Lind, J.:

Placental transfusion. *Am. J. Dis. Child.* 127:128, 1974.

Zeilder, R.B and Kim, H.D.:

Effect of low electrolyte media on salt loss and hemolysis of mammalian red blood cells. *J. Cell. Physiol.* 100:251-265, 1979.

Zinkham, W.H.:

An-in vitro abnormality of glutathione metabolism in erythrocyte from normal newborns. mechanisms and clinical significance. *Pediatrics* 23:18, 1959.

Zueler, W.W. and Brown, A.K.:

Oxytocic agents and neonatal hyperbilirubinemia. *Am. J. Dis. Child* 101:87, 1961.