

# Contents

<b>Abstract</b>	<b>VIII</b>
<b>Chapter I : Introduction</b>	
Previous work	2
Aim of work	5
Location of the studied wells	5
Methodology	5
<b>Chapter II : Lithostratigraphy</b>	
Masajid Formation	8
Burg El Arab Formation	9
Bahariya Formation	11
<b>Chapter III : Systematic palynology</b>	
Introduction	15
Systematic description	
Microspores	16
Gymnosperm pollen	40
Angiosperm pollen	56
Incertae sedis	72
Miscellaneous	76
Microplankton	78
<b>Chapter IV : Palynostratigraphy</b>	
Introduction	100
Present work	100
Palynozone I (Middle-Late Jurassic)	101
Palynozone II (late Albian)	103
Palynozone III (late Albian/early Cenomanian)	106
<b>Chapter V : Palynofacies analysis, organic thermal maturation and source rock evaluation</b>	
Introduction	114
Present work	114
Methodology	115
Palynofacies and kerogen types	115
Organic thermal maturation	118
Source rock potentiality	121
<b>Chapter VI : Palaeoenvironmental interpretation</b>	
The Masajid Formation	127
The Burg El Arab Formation	130
The Bahariya Formation	133
Cretaceous provincialism	133
<b>Chapter VII : Summary of conclusions</b>	<b>135</b>
<b>References</b>	<b>137</b>

## List of figures

Figure 1.1. Generalized sketch map of Egypt showing the location of the Sharib-1X and Ghoroud-1X wells .....	6
Figure 1.2. The laboratory equipments used during the different steps of this study .....	7
Figure 2.1. Lithologic column and samples positions, Sharib-1X well .....	12
Figure 2.2. Lithologic column and samples positions, Ghoroud-1X well .....	13
Figure 2.3. Lithologic correlation between the Sharib-1X and Ghoroud-1X wells showing the occurrence of the studied rock units .....	14
Figure 3.1. Schematic drawings illustrating the basic morphologic features of a trilete spore .....	16
Figure 3.2. Schematic drawings showing the basic types of surface sculpture of spores .....	25
Figure 3.3. Sporoderm stratification in perinate spores as defined by Traverse (1988) .....	38
Figure 3.4. Schematic drawings illustrating the basic morphologic features of a monolete spore ...	39
Figure 3.5. Orientation of <i>Equisetosporites</i> pollen grain .....	50
Figure 3.6. Phylogenetic relationships of the different <i>Afropollis</i> species .....	64
Figure 3.7. Measurements of a tricolpate pollen grain .....	66
Figure 3.8. Positions of apocolpia and mesocolpia in an angiosperm pollen as defined by Erdtman (1952) .....	68
Figure 3.9. Basic anatomy of a thecate dinoflagellate cyst .....	79
Figure 3.10. The relation between processes length, as a percentage of the shortest diameter of the central body, and the major cyst kinds .....	79
Figure 3.11. The four important archeopyle types .....	80
Figure 3.12. The basic morphologic terminology .....	81
Figure 3.13. Schematic drawings illustrating most of the encountered dinoflagellate cyst genera in the present study .....	83
Figure 3.14. Some special distal ornamentation of a process in dinoflagellate cysts .....	85
Figure 4.1. Observed ranges of some selected palynomorphs in Africa .....	104
Figure 4.2. Correlation of selected late Albian to middle Cenomanian miospore palynozonations from Egypt and other northern Gondwana regions .....	107
Figure 4.3. Semi-quantitative distribution of the recorded palynomorphs in the studied interval of the Sharib-1X well .....	108
Figure 4.4. Stratigraphic distribution of the recorded palynomorphs in the studied interval of the Sharib-1X well .....	111
Figure 5.1. Percentages of (POM) groups within the defined palynofacies of the Sharib-1X well ..	116
Figure 5.2. Percentages of (POM) groups within the defined palynofacies of the Ghoroud-1X well	117
Figure 5.3. Oil and gas generation as a function of increasing sediment burial .....	118
Figure 5.4. Pearson's (1984) colour chart compared with organic thermal maturity,	

TAI and vitrinite reflectance .....	119
Figure 5.5. Palynofacies, kerogen type, spore/pollen colour, TAI, vitrinite reflectance (Ro %), thermal maturation and source rock potentiality of the Sharib-1X well .....	122
Figure 5.6. Palynofacies, kerogen type, spore/pollen colour, TAI, vitrinite reflectance (Ro %), thermal maturation and source rock potentiality of the Ghoroud-1X well .....	123
Figure 6.1. A) Ternary AOM-Phytoclast-Palynomorph kerogen plot. B) Ternary Microplankton-Spore-Pollen palynomorph plot .....	127
Figure 6.2. A ternary AOM-Phytoclast-Palynomorph kerogen plot of Palynofacies 3, Masajid Formation, Sharib-1X well .....	128
Figure 6.3. A ternary Microplankton-Spore-Pollen palynomorph plot of the Masajid Formation, Sharib-1X well .....	128
Figure 6.4. Vertical trends in palynomorphs and marine/non-marine ratio in the Sharib-1X well	129
Figure 6.5. A ternary AOM-Phytoclast-Palynomorph kerogen plot of the Ghoroud-1X well .....	130
Figure 6.6. A ternary Microplankton-Spore-Pollen palynomorph plot of the Kharita Member and the Bahariya Formation, Sharib-1X well .....	131
Figure 6.7. A ternary AOM-Phytoclast-Palynomorph kerogen plot of Palynofacies 2, Sharib-1X well .....	132
Figure 6.8. A ternary AOM-Phytoclast-Palynomorph kerogen plot of Palynofacies 1, Sharib-1X well .....	132
Figure 6.9. A schematic palaeoenvironmental model for the Masajid, Burg El Arab and Bahariya formations .....	133
Figure 6.10. Distribution of the most distinctive species in the Albian-Cenomanian Elaterates Province of Herengreen <i>et al.</i> (1996) .....	134