

REFERENCES

**** R E F E R E N C E S ****

- ALANKO, H.I. and AIRAKSINEN, P.J. (1983) :*
Effects of topical timolol on corneal endothelial cell morphology in-vivo.
Am.J.Ophthamol., 96 : 615.
- ANDERSON, J.A.; DAVIS, W.L. and WEI, C.P. (1980) :*
Site of ocular hydrolysis of a produg, dipivefrin, and a comparison of its ocular metabolism with that of the parent compound, epinephrine.
Invest.Ophthalmol.Vis.Sci., 19(7) : 817-23.
- ARMALY, M.F. and JEPSON, W.C. (1982) :*
Accommodation and the dynamics of the steady state I.O.P.
Ophthalmol., 1 : 480.
- ARTHUR, B.W.; HAY, G.J.; WASAN, S.M. and WILLIS, W.G. (1983):*
Ultrastructural effects of topical timolol on the rabbit cornea.
Arch.Ophthalmol., 101 : 1607-1610.
- ASSEFF, C.F.; WEISMAN, R.L.; PODOS, S.M. and BECKER, B. (1973) :*
Ocular penetration of pilocarpine in patients.
Am.J.Ophthal., 75 : 212.
- BALACCO-GABRIELI, C.; REIBALDI, A. and SALAMANNA, S. (1980):*
Preliminary study of acidification of PH of the aqueous humor in rats treated with timolol.
Boll.Soc.Ital.Biol.Sper., 56 : 2071-2075.

BALLIN, N.; ECKER, B. and GOLDMAN, M. (1966) :

Systemic effects of epinephrine applied topically to the eye.
Invest.Ophthalmol., 5 : 125.

BARTELS, S.P.; ROTH, O.; JUMBLATT, M.M. and NEUFELD, A.H.
(1980) :

Pharmacological effects of timolol in the rabbit eye.
Invest.Ophthal.Vis.Sci., 19 : 1189.

BECKER, B.; MONTGOMERY, S.W.; KASS, M.A. and SHIN, D.H.
(1977) :

Increased ocular and systemic responsiveness to epinephrine in primary open-angle glaucoma.
Arch.Ophthalmol., 95 : 789.

BECKER, B.; PETTIT, T.H. and GAY, A.J. (1961) :

Topical epinephrine therapy of glaucoma.
Arch.Ophthalmol., 66 : 219.

BECKER, B. and LEY, A.P. (1958) :

Epinephrine and acetazamide in the therapy of the glaucomas.
Am.J.Ophthalmol., 45 : 639.

BERNSTEIN, L.P. and HENKIND, P. (1981) :

Additional information on adverse reactions to timolol.
Am.J.Ophthalmol., 92 : 295.

BIGAN, F. and WITMER, R. (1981) :

Effect of high I.O.P. on the corneal endothelial density.
Am.Acad.Ophthalmol., November 1-6.

BIGGER, J.F. (1980) :

Dipifevryn and glaucoma.
Prespect.Ophthalmol., 4 : 85.

BILL, A. and PHILIPS, C.I. (1971) :

Uveoscleral drainage of aqueous humour in human eyes.
Exp.Eye Res., 12 : 275.

*BIRSS, S.A.; LONGWELL, A.; HECKBERT, S. and KELLER, N.
(1978) :*

ocular hypotensive efficacy of topical epinephrine in normotensive and hypertensive rabbits : Continuous drug delivery Vs. eye drops.
Ann.Ophthalmol., 10 : 1045.

BOGER, W.P. (1979) :

Timolol : Short term "escape" and long term "drift".
Ann.Ophthal. 11 : 1239.

*BOGER, W.P.; PULIAFITO, C.A.; STEINERT, R.F. and PAVAN-
LANGSTON, D. (1978) :*

Long term experience with timolol ophthalmic solution in patients with open angle glaucoma.
Ophthalmology, 85 : 259-267.

BOGER, W.P.; STEINERT, R.F.; PULIAFITO, C.A. and PAVAN-
LANGSTON, D. (1978) :

Clinical trial comparing timolol ophthalmic solution to
pilocarpine in patients with open angle glaucoma.
Am.J.Ophthal., 86 : 8-18.

BOURNE, W.M. and KAUFMAN, H.E. (1976) :

Specular microscopy of human corneal endothelium in-
vivo.
Am.J.Ophthalmol., 81 : 319-323.

BOURNE, W.M.; NAGATAKI, S. and BRUBAKER, R.F. (1984) :

The permeability of the corneal endothelium to
fluorescein in the normal human eye.
Curr.Eye Res., 3 : 509.

BROMBERG, B.B.; GREGORY, D.S. and SEARS, M.L. (1980) :

Beta-adrenergic receptors in ciliary processes of the
rabbit.
Invest.Ophthal.Vis.Sci., 19 : 203.

BRUBAKER, R.F.; NAGATAKI, S. and BOURNE, W.M. (1982) :

Effects of chronically administered timolol on aqueous
humor flow in patients with glaucoma.
Ophthalmology, 89 : 280.

BUCKLEY, R.J. and SHERRARD, E.S. (1982) :

A new application of the corneal specular microscope.
Arch.Ophthalmol., 100 : 296.

BURNS, R.R.; BOURNE, W.M. and BRUBAKER, R.F. (1981) :

Endothelial function in patients with cornea guttata.
Invest.Ophthalmol.Vis.Sci., 20 : 77.

CAIRNS, J.E. (1976) :

Goniospasis : A method designed to relieve canalicular blockade in primary open angle glaucoma.
Ann.Ophthal., 8 : 1417.

CHI, H.H.; TENG, C.C. and KATZIN, H.M. (1960) :

Healing process in the mechanical denudation of the corneal endothelium.
Am.J.Ophthamol., 49 : 693.

COAKERS, R.L. and BRUBAKER, R.F. (1978) :

The mechanism of timolol in lowering the I.O.P. in normal eye.
Arch.Ophthal., 96 : 2045.

COLES, W.H. (1975) :

Effects of pilocarpine on the vitro rabbit corneal endothelium.
Invest.Ophthalmol., 14 : 246-250.

COUCH, J.M.; GULLEN, P.; CASEY, T.A. and FABRE, J.W. (1987):

Mitotic activity of corneal endothelial cells in organ culture with recombinant epidermal growth factor.
Ophthalmology, 94 : 1-6.

DAVIES, P.D.; KIRKHAM, J.B. and VILLANUEVA, S. (1976) :

Surface ultrastructure of human donot corneal endothelium.
Trans.Ophthalmol.Soc.U.K., 96 : 69-96.

DeCEVALLOS, A.; DOHLMAN, C.H. and REINHART, W.J. (1976) :

Corneal thickness in glaucoma.
Ann.Ophthalmol., 8 : 177.

DRANCE, S.M. (1970) :

Symposium on ocular pharmacology and therapeutics.
Transactions of New Orleans Academy of Ophthalm. C.V.
Mosby Company. St. Louis, P. 106.

DRANCE, S.M. and NASH, P.A. (1971) :

The dose response of human I.O.P. to pilocarpine.
Canad.J.Ophthalmol., 6 : 9.

*EDELHAUSER, H.F.; HYNDIUK, R.A.; ZEEB, A. and SHULTZ, R.O.
(1982) :*

corneal edema and the intraocular use of epinephrine.
Am.J.Ophthalmol., 93 : 327-333.

EHLERS, N. (1970) :

On corneal and intraocular pressure II : A clinical
study on the thickness of the corneal stroma in
glaucomatous eyes.
Acta.Ophthalmol., 48 : 1107.

EL-RIFAI, M. (1988) :

Text Book of Clinical Ophthalmoptics, p.402, Al-Azhar
University press.

*FABRICANT, R.N.; ALPAR, A.J.; CENTIFANTO, Y.M. and KAUFMAN,
H.E. (1981) :*

Epidermal growth factor receptors on corneal
endothelium.
Arch.Ophthalmol., 99 : 305-308.

FERRY, A.P. and ZIMMERMAN, L.E. (1964) :

A complication of topical use of epinephrine.
Am.J.Ophthalmol., 58 : 205.

FINE, B.S. and YANOFF, M. (1979) :

Ocular histology.
A Text and Atlas 182. Medical Department, Harper and Row, Inc., New York.

GARNER, L.L.; JOHNSTONE, W.W.; BALLINTINE, G.J. and CARROLL, M.E. (1959) :

Effect of 2% levo-rotary-epinephrine on the intraocular pressure of the glaucomatous eye.
Arch.Ophthalmol., 62 : 230.

GARZIA, R. (1982) :

An advance in ophthalmic pharmacology. The use of the epinephrine pro-drug dipivalyl epinephrine in the treatment of glaucoma.
J.Am.Optom.Assoc., 53(9) : 727-30.

GASSETT, A.R.; ISHU, Y.; KAUFFMAN, H.E. and MILLER, T. (1974) :

Cytotoxicity of ophthalmic preservatives.
Am.J.Ophthalmol., 78 : 98-105.

GEROSKI, D.H. and EDELHAUSER, H.F. (1984) :

Quantitation of Na/K ATPase pump sites in the rabbit corneal endothelium.
Invest.Ophthalmol.Vis.Sci., 25 : 2056.

GEROSKI, D.H.; MATSUDA, M.; YEE, R.W. and EDELHAUSER, H.F.
(1985) :

Pump function of the human corneal endothelium: Effects
of age and cornea guttata.
Ophthalmology, 92 : 759-763.

GOLDBERG, I.; ASHBURN, F.S.Jr.; KASS, M.A. and BECKER, B.
(1979) :

Efficacy and patient acceptance of pilocarpine gel.
Am.J.Ophthal., 88 : 843.

GOSPODARWICZ, D.; MESCHER, A.L. and BIRDWELL, C.R. (1977) :

Stimulation of corneal endothelial cell proliferation
in-vitro by fibroblast and epidermal growth factors.
Exp.Eye Res., 25 : 75.

GRAVES, B. (1924) :

A bilateral chronic affection of the endothelial face
of the cornea of elderly persons, with an account of
the technical and clinical principles of its slit-lamp
observation.
Br.J.Ophthamol., 8 : 502.
Quoted from: Mayer (1984)

GREEN, K.; HULL, D.S.; VAUGHN, A.; MALIZIA, A. and BOWMAN,
K. (1977) :

Rabbit endothelial response to ophthalmic response to
ophthalmic preservatives.
Arch.Ophthamol., 95 : 2218-21.

GRIERSON, I.; LEE, W.R. and ABRAHAM, S. (1978) :

Effects of pilocarpine on the morphology of the human
outflow apparatus.
Br.J.Ophthal., 62 : 302.

GRIERSON, I.; LEE, W.R.; MOSELEY, H. and ABRAHAM, S. (1978):

The trabecular wall of Schlemm's canal : A study of the effect of pilocarpine by scanning electron microscopy. Br.J.Ophthalmol., 63 : 9.

HANNU, I.; ALKANO, M.D. and AIRAKSINEN, P.J. (1983) :

Effects of topical timolol on corneal endothelial cell morphology in-vivo. Am.J.Ophthalmol.. 96 : 615-621.

HARRIS, L.A.; GALIN, M.A. and LERNER, R. (1970) :

The influence of low dose L-epinephrine on intraocular pressure. Ann.Ophthalmol.. 2 : 253.

HAVENER, W.H. (1983) :

Ocular Pharmacology. 5th ed., C.V.Mosby Co., St.Louis. P.325-340.

HEFNI, W.; ABDEL-KHALEK, L.M.R. and EMAM, A.A. (1982) :

The effect of aging on the morphology of the corneal endothelium. Bull.Ophthalmol.Soc.Egypt, 75 : 35-39.

HEILMANN, K. (1978) :

Special Pharmacology. Cited in : Glaucoma, conception of a disease, pathogenesis, diagnosis, therapy. Ed. by Heilmann, K. & Richardson, K.T.. George Thieme Publishers, Stuttgart. P.267-283.

HIGGINS, R.G. and BRUBAKER, R.F. (1980) :

Acute effect of epinephrine on aqueous humor formation in the timolol-treated normal eye as measured by fluorophotometry. Invest.Ophthalmol.Vis.Sci.. 19 : 420.

HOGAN, M.J.; ALVARADO, J.A. and WEDDELL, J.E. (1971) :

Histology of the Human Eye, An Atlas and Text Book,
Philadelphia, W.B. Saunders Co., p.75

HOSKINS, H.D.Jr. and KASS, M.A. (1989) :

Becker-Shaffer's diagnosis and therapy of the glaucomas:
Transection of the C.V. Mosby Co., St. Louis, Missouri,
P.373.

HULL, D.S.; CHEMOTTI, M.T.; EDELHAUSER, H.F.; VAN HORN, D.L.
and HYNDIUK, R.A. (1975) :

Am.J.Ophthalmol., 75(1) : 245-250.

JAY and McDONALD (1978) :

The effect of intraocular mitosis on cultured bovine
cornea.
Brit.J.Ophthalmol., 62 : 815-820.

JOHNSON, D.H.; EPSTEIN, D.H.; ALLEN, R.C.; SMITH, J.R.;
CAMPBELL, D.; ROSENQUIST, R. and VANBUSKEIK, M. (1984):

A one-year multicenter clinical trial of pilocarpine
gel.
Am.J.Ophthalmol., 97 : 723-729.

KASS, M.A.; MANDELL, A.L.; GOLDBERG, I.; PAINE, M. and
BECKER, B. (1979) :

Dipivefrin and epinephrine, Treatment of elevated
intraocular pressure : A comparative study.
Arch.Ophthalmol., 97 : 1865.

KATZ, I.M.; HUBBARD, W.A.; GESTON, A.J. and GOULD, A.L.
(1976) :

I.O.P. decrease in normal volunteers following timolol
ophthalmic solution.
Invest.Ophthalmol., 15 : 489-498.

KAUFMAN, H.E.; KATZ, J.L. and BOURNE, W.M. (1966) :

Pathology of the corneal endothelium.
Investigative Ophthalmology & Visual Science, 16 : 265.

KAYE, G.I.; SIBLEY, R.C. and HOEFLE, F.B. (1973) :

Recent studies on the nature and function of the
corneal endothelial barrier.
Exp.Eye Res., 15 : 585-613.

KOLKER, A.E. and HETHERINGTON, J.Jr. (1983) :

Becker-Shaffer's diagnosis and therapy of the glaucoma.
5th ed., C.V.Mosby Co., St.Louis, Toronto, p.21-54, 59-
107, 317-400.

KOREY, M.; GIESER, D.; KASS, M.A.; WALTMAN, S.R.; GORDON, M.
and BECKER, B. (1982) :

Central corneal endothelial cell density and central
corneal thickness in ocular hypertension and primary
open-angle glaucoma.
Am.J.Ophthamol., 94 : 610-616.

KROHN, O.L. (1978) :

Flux of topical pilocarpine to the human aqueous.
Trans.Am.Ophthal.Soc., 76 : 502.

LAING, R.A. (1980a) :

Specular microscopy of the cornea.
In: Zodunaisky, J.A. and Davson, h.h. (Eds.): Recent
progress in eye research. Vol.3, pp.157-217, NewYork,
Academic press Inc.

LAING, R.A. (1980b) :

Specular microscopy of the cornea. Zodunaisky, J.A. and Davson, h.h. (Eds.): Curr. Eye Res., 3 : 157-218.

LAING, R.A.; NEUBAUER, L. and OAK, S.S. (1984) :

Evidence for mitosis in the adult corneal endothelium. Ophthalmology, 91 : 1129.

LAING, R.A.; SANDSTROM, M.M. and LEIBOWITZ, H.M. (1979) :

Clinical specular microscopy I. Optical principles II. Qualitative evaluation of corneal endothelial photomicrographs. Archives of Ophthalmology, 97 : 1714-1720.

LAULE, A.; CABLE, M.K.; HOFFMAN, C.E. and HANNA, C. (1978) :

endothelial cell population changes of human cornea during life. Arch.Ophthalmol., 96 : 2031-2035.

LAWRENCE, W.; HIRST, M.D.; CHERYL AUER, M.S.; HELEN ABBEY,

Sc.D.; JULIE, B.A. and KUES, H. (1984) :

Quantitative analysis of wide-field endothelial specular photomicrographs. Am.J.Ophthalmol., 97 : 488-495.

LAZARE, R. and HORLINGTON, M. (1975) :

Pilocarpine levels in the eyes of rabbits following topical applications. Exp.Eye Res., 21 : 281.

LEE, P.A.; BRUBAKER, R.F. and NAGATAKI, S. (1983) :

Acute effects of thymoxamine on aqueous humor formation in the epinephrine treated normal eye as measured by fluorophotometry.
Invest.Ophthalmol.Vis.Sci., 24 : 165.

LEIBOWITZ, H.M.LAING, R.A.; SANDSTROM, M.M. and BERROSPI, A.R. (1976) :

Changes in the corneal endothelium as a function of age.
Exp.Eye Res., 22 : 587-594.

LIU, H.K.; CHIOU, C.Y. and GARG, L. (1980) :

Ocular hypotensive effect of timolol in cat eyes.
Arch.Ophthalmol., 98 : 1467.

LIU, G.S.; TROPE, G.E. and BASU, P.K. (1989) :

Ultrastructural effects of topical betopic, betayan, and timoptic on the rabbit corneal endothelium.
J.Ocul.Pharmacol., 5(4) : 329-342.

LOHMAN, L.E.; RAO, G.N. and AQUAVELLA, J.V. (1981) :

Optics and clinical applications of wide-field specular microscopy.
Am.J.Ophthamol., 92 : 43.

LOHMAN, L.E.; RAO, G.N. and AQUAVELLA, J.V. (1982) :

In-vivo microscopic observations of human corneal epithelial abnormalities.
Am.J.Ophthamol., 93 : 210.

LOWENSTEIN, O. (1956) :

The Argyl Robertson syndrome.
Am.J.Ophthalmol., 42 : 105.

MANDELL, A.L.; STERTZ, F. and KIFABCHI, A.E. (1978) :

Dipivalyl epinephrine : A new pro-drug in the treatment of glaucoma.
Ophthalmology, 85 : 268.

MARTOLA, E.L. and BAUM, J. (1968) :

Central and peripheral corneal thickness.
Arch.Ophthalmol., 79 : 28-30.

MAURICE, D.M. (1968) :

Cellular membrane activity in the corneal endothelium of the intact eye.
Exp.Eye.Research, 24 : 1094.

MATSUDA, M.; ISHII, Y.; AWATA, T. and FUKUDA, M. (1989) :

The effect of long-term topical administration of commercial beta-blockers on the rat corneal endothelium.
Int.Ophthalmol., Jan., 13(1-2) , p.67-73.

MAYER, D.J. (1984) :

Clinical wide-field specular microscopy.
Chapter 1, page 9-10, Bailliere Tindall London.

McMAHON, C.D.; SHAFFER, R.N.; HOSKINS, H.D.Jr. and

HETHERINGTON J.Jr. (1979) :

Adverse effects experienced by patients taking timolol.
Am.J.Ophthalmol., 88 : 736.

MILLIS, K.B. (1983) :

Blind randomised non-crossover long-term trial comparing timolol 0.25%, timolol 0.5% in the treatment of simple chronic glaucoma.
Br.J.Ophthalmol., 67 : 219.

MINDELL, T.S. (1982) :

Who, what, where, when and how.
Ann.Ophthalmol., 14 : 10.

*MINDEL, T.S.; YABLONSKI, M.E.; TAVITIAN, H.O.; PODOS, S.M.
and ORELLANA, J. (1981) :*

Dipivefrin and echothiophate : Efficacy of combined use
in humans.
Arch.Ophthalmol., 99 : 1583.

MISHI, O. and HANASAKI, K. (1988) :

Automated morphometry of corneal endothelial cell : Use
of video camera and video tape recorder.
Br.J.Ophthalmol., 72(1) : 68-73.

MISHIMA, S. (1982) :

Clinical investigations on the corneal endothelium.
Am.J.Ophthalmol., 93 : 1-29.

MOSES, R.A. and GRODZKI, W.J.Jr. (1977) :

Choroid tension and facility of aqueous outflow.
Invest.Ophthalmol.Vision.Sci., 16 : 1082.

NESHER, R.; KASS, M.A. and GANS, L.A. (1990) :

Corneal endothelial changes in ocular hypertensive
individuals after long-term unilateral treatment with
timolol.
Am.J.Ophthalmol., Vol.110 : 3, 309-310.

NEUFELD, A.H. (1979) :

Experimental studies on the mechanism of action of
timolol.
Surv.Ophthalmol., 2 : 363.

NEUFELD, A.H.; JAMPAL, L.M. and SEARS, M.L. (1972) :

Cyclic AMP in the aqueous humor : The effects of
adrenergic agents.
Exp.Eye Res., 14 : 242.

NEUFELD, A.H. and SEARS, M.L. (1972) :

Adenosine 3',5'-monophosphate analogue increases the
outflow facility of the primate eye.
Invest.Ophthalmol., 14 : 688.

OBSTBAUM, S.A.; GALIN, M.A. and KATZ, I.M. (1978) :

Timolol : Effect on IOP in chronic open-angle glaucoma.
Ann.Ophthalmol., 10 : 1347-51.

O'CONNOR DAVIES, P.H. (1981) :

The action and uses of Ophthalmic drugs.
2nd ed., Butterworth, London, Boston, Sydney, Wellington,
Durban, Toronto, p.189.

OH, J.O. (1963) :

Changes with age in the corneal endothelium of normal
rabbits.
Acta.Ophthalmol., 41 : 568.

OLSEN, R.J. (1980) :

Specular microscopy study in cats.
Arch.Ophthalmol., 98 : 1283-1284.

OLSEN, R.J. and KAUFMAN, H.E. (1978) :

Intraocular pressure and corneal thickness after
penetrating keratoplasty.
Am.J.Ophthalmol., 86 : 97.

OLSEN, T.; NIELSEN, C.B. and EHLERS, N. (1980) :

The optical measurement of corneal thickness : The measuring conditions and sources of error.
Acta.Ophthalmol., 58 : 976-984.

PHILIPS, C.I.; BARTHOLOMEW, R.S.; KAZI, G.; SCHMITT, C.J.
and VOGEL, R. (1981) :

Penetration of timolol eye drops into human aqueous humour.
Br.J.Ophthalmol., 65 : 593-595.

PODOS, S.M; BECKER, B.; ASSEFF, C. and HARTSTEIN, J. (1972):

Pilocarpine therapy with soft contact lenses.
Am.J.Ophthalmol., 73 : 306

PODOS, G.J. and KRACHMER, J.H. (1980) :

Comparison of endothelial cell density in diabetics and a control population.
Am.J.Ophthalmol., 90 : 172.

PRAGER, T. (1981) :

Comparative material on commercially available fluoro-photometer.
Ophthalmology, 84 : 88.

RADIUS, R.L.; DIAMOND, G.R.; POLLAK, P. and LANGHAM, M.E.
(1978) :

Timolol : A new drug for management of chronic simple laucoma.
Arch.Ophthalmol, 96 : 1003-8.

RAMER, R.M. and GASSET, A.R. (1974) :

Ocular penetration of pilocarpine. The effect of hydrophobic contact lenses on ocular penetration of pilocarpine.
Ann.Ophthal., 6 : 1325.

RAO, G.N.; LOHMAN, L.E. and AQUAVELLA, J.V. (1982) :

Cell size-shape relationship in corneal endothelium.
Invest.Ophthalmol., 4 : 22-27.

RENARD, G.; HIRSCH, M.; FAURE, J.P.; USUI, M. and POULIQUEN, Y. (1976) :

Ultrastructural study on the regeneration of the corneal endothelial cells.
Jap.J.Ophthamol., 20 : 243.

RICHARDSON, K.T. and KURTZMAN, C. (1981) :

Evaluation of medical therapy in chronic glaucoma.
Cited in: Symposium on glaucoma. Transaction of the New Orleans Acad. of Ophthal., C.V.Mosby Co., St.Louis, Toronto, London, p.322-324.

RUBEN, M. and WATKINS, R. (1975) :

Pilocarpine dispersion for the soft hydrophilic contact lens.
Br.J.Ophthalmol., 59 : 455.

SAAAMOTO, K.; AKAGI, Y.; KODAMA, Y. and itoi, m. (1984) :

Corneal endothelial changes caused by ophthalmic drugs.
Cornea , 3 : 37-41.

SAMPLES, J.R.; BINDER, P.S. and NAYAK, S. (1989) :

The effect of epinephrine and benzalkonium chloride on cultured corneal epinephrine and trabecular meshwork cells.

Exp.Eye Res., 49 : 1-12.

SLACK, J.W.; EDELHAUSER, H.F. and HELANEK, M.J. (1990) :

Abisulfite-free intraocular epinephrine solution.

Am.J.Ophthalmol., Jul.. 15 ; 110(1), p.77-82.

SASAKI, Y. (1985) :

E.M localization of the Na/K ATPase activity in Fuch's corneal dystrophy.

Invest.Ophthalmol.Vis.Sci.. 26(suppl.) : 53, ARVO abstracts.

SATO, T. (1978) :

Studies on the endothelium of the corneal graft.

Jpn. J.Ophthalmol.. 22 : 114-126.

SETALA, K. (1979) :

corneal endothelial cell density after an attack of acute glaucoma.

Acta.Ophthalmol.. 57 : 1004.

SHENKER, H.E.; YABLONSKI, M.E.; PODOS, S.M. and KINDER, L. (1981) :

Fluorophotometric study of epinephrine and timolol in human subjects.

Arch.Ophthalmol.. 99 : 1212.

SHERRARD, E.S. (1978) :

Characterization of changes observed in the corneal endothelium with the specular microscope.
Investigative Ophthalmology & Visual Science, 17 : 322

SHERRARD, E.S. and BUCKLEY, R.J. (1984) :

Clinical specular microscopy of the corneal endothelium
I: Intraocular lens implication (Eds.) Rosen. E.S.;
Haining, W.M. and Arnott, E.J. The C.V.Mosby Company,
pp.131-149.

SHERRARD, E.S. and BUCKLEY, R.J. (1980) :

Endothelial wrinkling : A complication of clinical
specular microscopy.
VI Congress of European Soc. of Ophthalmology, London,
Academic press.

SHIELDS, M.B. (1982) :

A study guide for glaucoma.
Williams and Wilkins. Baltimore, London. p.7-20, 47-70,
152. 392-432.

SLINGSBY, J.G. and FORSTOT, S.L. (1981) :

Effect of blunt trauma on the corneal endothelium.
Arch.Ophthalmol., 99 : 1041-1043.

SMITH, S.A.; SMITH, S.E. and LAZARE, R. (1978) :

An increased effect of pilocarpine on the pupil by
application of the drug in oil.
Br.J.Ophthalmol., 62 : 314.

SQUIRES, E.L. and WEINAR, V.L. (1980) :

Stimulation of repair of human corneal endothelium in
organ culture by mesodermal growth factor.
Arch.Ophthalmol., 98 : 1462.

STAAZ, W.D.; RADIUS, R.L.; VAN HORN, D.H. and SCHULTZ, R.O.
(1981) :

Effects of timolol on bovine corneal endothelial cultures.
Arch.Ophthalmol., 99 : 660-663.

STOKER, F.W. (1971) :

The endothelium of cornea and its clinical implications
Trans.Am.Ophthal.Soc., 15 : 669-686.

STREMPEL, I. (1982) :

Role of the B-adrenergic receptor blocking agent
(Timolol) in the treatment of open angle glaucoma.
Ophthalmologica [Basel], 184 : 198-203.

STURROCK, G.D.; SHERRARD, E.S. and RICE, N.S.C. (1978) :

Microscopy of the corneal endothelium.
Brit.J.Ophthalmol., 62 : 809-814.

SVEDBERGH, B. (1975) :

Effects of artificial intraocular pressure elevation on
the corneal endothelium in the vervet monkey.
Arch.Ophthalmol., 53 : 839-844.

SVEDBERGH, B. and BILL, A. (1972) :

Scanning electron microscopic studies of the corneal
endothelium in man and monkeys.
Acta.Ophthalmol., 50 : 321.

TAY and McDONMALD (1978) :

The effect of intraocular miotics on cultured bovine
cornea.
Brit.J.Ophthalmol., 62 : 815-820.

TICHO, U.; BLUMENTHAL, M.; ZONIS, S.; GAL, A.; BLANK, I. and
MAZOR, Z.W. (1979) :

Piloplex : A new long-acting pilocarpine polymer salt :
A long-term study.
Br.J.Ophthalmol., 63 : 45.

VAN BUSKIRK, E. (1978) :

Changes in the facility of aqueous outflow induced by
lens depression and I.O.P. in excised human eyes.
Am.J.Ophthalmol., 82 : 736.

VAN BUSKIRK, E.M. (1980) :

Adverse reactions from timolol administration.
Ophthalmology, 87 : 447.

VAN HOOSE, M.C. and LEADERS, F.E. (1974) :

The role of the cornea in the biologic response to
pilocarpine.
Invest.Ophthalmol., 13 : 377.

VANHORN, D.L. and HYDIUK, R.A. (1975) :

Endothelial wound repair in primate cornea.
Exp.Eye Res., 21 : 113.

VANNAS, A.; STELALA, A. and RUUSUVAARA, P. (1977) :

Endothelial cells in capsular glaucoma.
Acta.Ophthalmol., 55 : 951.

VAUGHN, D.; SHAFFER, R. and RIEGELMAN, S. (1961) :

A new stabilized form of epinephrine for the treatment
of open-angle glaucoma.
Arch.Ophthalmol., 66 : 108.

VEIRS, E.R. and McGREW, J.C. (1963) :

Ocular complications from topical epinephrine therapy
of glaucoma.
EE NT Mthly, 42 : 46.

VOGT, A. (1920) :

Die sichtbarkeit des lebenden hornhautendothels. Ein
Beitrag Zur Methodik der spaltlampenmikroskopie.
Graefes Archives Klin. Ophthalmology, 101 : 123.
Quoted from: Mayer (1984).

VON SALLMAN, L. (1961) :

Studies on the corneal endothelium of the rabbit.
Am.J.Ophthamol., 51 : 955.

WAGATAKI, S. and BRUBAKER, R.F. (1981) :

Early effect of epinephrine on aqueous formation in the
normal human eye.
Ophthalmology, 88 : 278.

WALTMAN, S.R. and KAUFMAN, H.E. (1970) :

In-vivo studies of human corneal endothelial
permeability.
Am.J.Ophthalmol., 70 : 45.

WALTMAN, S.R.; YARIAN, D.; HEART, W. and BECKER, B. (1977) :

Corneal endothelial changes with long-term topical
epinephrine therapy.
Arch.Ophthalmol., 95 : 1357-8.

WARING, G.O. (1980) :

The corneal endothelium.
Ophthalmology, 89 : 531-590.

WARING. G.O. (1982) :

Posterior collagenous layer of the cornea. Ultra-
structural classification of abnormal collagenous
tissue posterior to Descemet's membrane in 30 cases.
Arch.Ophthalmol., 100 : 122-134.

WARING, G.; BOURNE, W.M.; EDELHAUSER, H.F. and KENYON, K.
(1982) :

The corneal endothelium : Normal and pathologic
structure and function.
Ophthalmology. 89 : 531.

WARING, G.O.; GEORGIA, A.; FORD, G.E. and ROSENBLANTT, L.S.
(1980) :

Individual corneal endothelial changes with age:
A specular microscopic study.
IV. Congress of the European Soc. of Ophthalmology.
40 : 75.

WEI, C.P.; ANDERSON, J.A. and LEOPOLD, I. (1978) :

Ocular absorption and metabolism of topically applied
epinephrine and a dipivalyl ester of epinephrine.
Invest.Ophthalmol.Vis.Sci., 17 : 315.

YTTEBORG, J. and DOHLMAN, C.H. (1965) :

Corneal oedema and intraocular pressure II. Clinical
results.
Arch.Ophthalmol., 74 : 477.

ZIMMERMAN, T.J.; HARBIN, R.; PETT, M. and KAUFMAN, H.E.
(1977) :

Timolol and facility of outflow.
Invest.Ophthalmol.Vis.Sci., 16 : 623.

ZIMMERMAN, and KAUFMAN, H.E. (1977a) :

Timolol : A beta-adrenergic agent for the treatment of
glaucoma.
Arch.Ophthal., 95 : 601-604.

ZIMMERMAN, and KAUFMAN, H.E. (1977b) :

Timolol : Dose response and duration of action.
Arch.Ophthal., 95 : 605.