

REFERENCES

- Barson, G.; Renn, N. and Bywater, A.F. (1994):** Laboratory evaluation of six species of entomopathogenic fungi for the control of the house fly (*Musca domestica* L.), a pest of intensive animal units. J. Invertebr. Pathol. 64 : 2, 107-113.
- Bedidimadani, N.; Favier, C.; Kodjo, A. and Richard, Y. (1993):** protein profiles of coagulase negative *Staphylococcus* of goat origin- taxonomic application. Revue. De. Medecine Veterinaire 144, 183-189.
- Bent K.J. (1967):** Electrophoresis of proteins of 3 *Penicillium* species on acrylamide gels. J. Gen Microbiol 49:195-200.
- Berny J.F. and Hennebert G.L. (1990):** Variants of *Penicillium expansum*: an analysis of cultural and microscopic character as taxonomic criteria. In: Samson RA, Pitt JI, eds. Modern Concepts in *Penicillium* and *Aspergillus* classification. NATO AIS series A, Life Sciences 185. London: Plenum Press, 49-65.
- Bidochka, M. J., McDonald, M. A., St. Leger, R. J., and Roberts, D. W. (1994):** Differentiation of species and strains of entomopathogenic fungi by random amplification of polymorphic DNA (RAPD). Curr. Genet. 25, 107-113.

REFERENCE

- Bidochkha, M.J. and Khachatourians, G.G. (1994)** : Basic proteases of entomopathogenic fungi differ in their adsorption properties to insect cuticle. *J. Invertebr. Pathol.* 64 : 1, 26-32.
- Bidochkha, M.J.; St – Leger, R.J. and Roberts, D.W. (1997)**: Mechanisms of deuteromycete fungal infections in grasshoppers and locusts: An overview *Memoirs-of-the-Entomological-Society-of-Canada* (Mem . Entomol . Soc. Can.) . 171 : 213-224.
- Bohmfolk, G.T. (1986)**: Practical factors influencing the utilization of baculoviruses as pesticides. In the biology of baculoviruses, (eds. Granados, R.R. and Federici, B.A.). PP. Vol. II, PP. 223. CRC press, Boca Raton, FL.
- Boucias, D.G. and J.C. Pendland. (1988)**: "Nonspecific factors involved in the attachment of entomopathogenic deuteromycetes to host insect cuticle." *Appl. Environ. Microbiol.* 54(7): 1795-1805.
- Boucias, D.G.; Hung, S.; Mazet, I. and Azbell, J. (1994)**: Effect of the fungal pathogen , *Beauveria bassiana* , on the lysozyme activity in *Spodoptera exigua* larvae. *J. Insect physiol.* 40(5) : 385-391.
- Burke, J.J.; Hatfield, J.L.; Klein, R.R. and Mullet, J.E. (1985)**: Accumulation of heat shock proteins in field-grown cotton. *Plant Physiol.* 78, 394-398.

REFERENCE

- Carruthers, R.I. (1997):** Diet critical to mass rearing deneficials. Agric. Res.; 45:2.
- Chang, L.O., Srb A.M. and Steward F.C. (1962):** Electrophoretic patterns of the soluble proteins of *Neurospora*. Nature 193:756-759.
- Charnley, A. K. (1984):** Physiological aspects of destructive pathogenesis in insects by fungi a speculative review in.' Invertebrate – microbial interactions' (J. M. Anderson, A. D. M. Ráyner, and D. W. H Walton, eds.) 229-270. Camberidge University Press, Camberidge.
- Chiuo, W.C. and Hou, R.F. (1993):** Infection of the Asian corn borer, *Ostrinia furnacalis Guenee* (Lep., Pyralidae), with entomopathogens under screen house condition *S.littoralis*. J. Appl. Entomol. 115(3): 246-253.
- Christensen, C.M. (1957):** Deterioration of stored grains by fungi. Biotanical Review, 23:108-134.
- Cilek, J.E., and G. L. Greene. (1994):** Stable fly (Diptera: Muscidae) insecticide resistance in Kansas cattle feedlots. J. Econ. Entomol. 87:275-279.

REFERENCE

- Clarkson, J.M. and Charnley, A.K. (1996):** New insights into the mechanisms of fungal pathogenesis in insects, Trends in Microbiology 4: 197-204.
- Cobb B.D. and Clarkson J.M. (1993):** Detection of molecular variation in the insect pathogenic fungus *Metarhizium* using RAPD-PCR. FEMS Microbiol Lett 15, 114(3):355.
- Cravanzola, F.; Piatti, P.; Bridge, P.D. and Ozino, O.I. (1997):** Detection of genetic polymorphism by RAPD - PCR in strains of the entomopathogenic fungus *Beauveria brongniartii* isolated from the European cockchafer (*Melolontha spp.*). Lett. App. Microbiol. 25 : 4, 289-294.
- David, W. A. L. (1967):** The physiology of the insect integument in relation to the invasion of pathogens. In 'insects and physiology' (J.W.L. Beament and J.E. Treherne, eds.) Oliver and Boyd, Edinburgh: 17-35.
- Davila, M.; Zambrano, K. and Castillo, M.A. (2001) :** Use of RAPD technique for the Identification of DNA fragments likely related to virulence in entomopathogenic fungus. Bioagro. 13 : 3, 93-98.
- Devi, K.U.; Padmavathi, J.; Sharma, H.C. and Seetharama, N. (2001):** Laboratory evaluation of the virulence of *Beauveria bassiana*

REFERENCE

isolates to the Sorghum Shoot borer *chilo partellus* Swinhoe (Lipidoptera: Pyralidae) and their characterization by RAPD-PCR. World. J. Microbiol. Biotechnol. 17 : 2, 131-137.

Doyle, J.J. and J.L. Doyle, (1987): A rapid DNA isolation procedure for small quantities of fresh leaf tissue. Phytochem. Bull. 19:11-15.

Elsworth, J.F. and J.F. Grove (1977): "Cyclodepsipeptides from *Beauveria bassiana* Bals. Part 1. Beauverolides H and I." J. Chem. Soc [Perkin 1] 3: 270-3.

Fargues, J., Robert, P.H., and Vey, A. (1976): Role du tegument et de la defence cellulaire des Coleopteres hotes dans la spcificite des souches entomopathogenes de *Metarhizium anisopliae* (Fungi Imperfecti) C. R. Acad. Sci. Paris Ser. D 282: 2223-2226.

Fargues, J. and Maniania, N.K. (1992): Variation in the susceptibility of *Spodoptera littoralis* (Lep: Noctuidae) to the entomopathogenic hyphomycete *Nomuraea rileyi*. Entomophaga. 37(4) : 545-554.

Ferron, P. (1975): Les champignons entomopathogenes: Evolution des recherches au cours des dix dernieres annees. Bul. Sci. Biol. Organ. Int. Lutte boil., WPRS Bul. 1975/3.

Ferron, P. (1978): Biological control of insect pests by entomogenous fungi. Annu. Rev. Entomol. (23): 409-442.

REFERENCE

- Finney, D.J. (1964):** Probit analysis: a statistical treatment of the sigmoid response curve, 2nd ed. Cambridge University press, London.
- FRANSEN, J.J and Butt, T.M. (1990) :** Natural enemies of whitefly : fungi. In : GERLING,D.(ed.): whiteflies : their bionomics, pest status and management. Intercept Ltd., Andover, UK, 187-210.
- Freire, L.L.C; Costa, A.B.I; Goes, LB and Oliveira, N.T. (2001):** DNA polymorphism and total protein in mutants of *Metarhizium var. anisopliae* (Metsch.) Sorokin strain E₉. Braz. J. Microbiol 32(2).
- Fungaro, M.H.P.; Vieira, M.L.C.; Pizzirani, Kleiner, A.A; Azevedo, J.L; de. and De-Azevedo, J.L. (1996):** Diversity among soil and insect isolates of *Metarhizium anisopliae var. anisopliae* detected by RAPD. Lett. App. Microbiol. 22 : 6, 289-392.
- Gillespie, J.P. and Khachatourians, G.G. (1992):** Characterization of the *Melanoplus sanguinipes* hemolymph after infection with *Beauveria bassiana* or wounding . COMP.BIOCHEM .PHYSIOL. B., 103B(2): 455-463.
- Glare, T.R. and Inwood, A.J. (1998) :** Morphological and genetic characterization of *Beauveria spp.* from New Zealand. Mycol .Research. 102 : 2, 250-256.

REFERENCE

Gottlieb. (1971): Gel electrophoresis: new approach to the study of evolution. *Bio-Science* 21:939-944.

Graystone, J.L.; Charnley, A.K.; Albajes, R(ed.). and Carnero, A. (1997):

Some fungi grow profusely in the insect before death and disrupt vital processes. Others have limited growth. Integrated control in protected crops. *Bulletin.OILB.SROP.*; 20 : 4, 263-267.

Gupta, S.; Leathers, T.D.; El-Sayed, G.N. and Ignoffo,C.M. (1992) : Insect

cuticle - degrading enzyme from the entomogenous fungus *Beauveria bassiana*. *Exp. Mycol.*, 16(2): 132-137.

Hamill, R.L. and H.R. Sullivan (1969): "Determination of pyrrolnitrin and derivatives by gas-liquid chromatography." *Appl. Microbiol* 18(3): 310-2.

Hawksworth, D.L.; Kirk, P.M.; Sutton, B.C. and Pegler, D.N. (1995):

Ainsworth and Bisby's Dictionary of the fungi, 8th edn. International Mycological Institute, CAB International, Wallingford. 616 PP.

Hegedus, D.D.; Bidochka, M.J.; Miranpuri, G.S. and

Khachatourians, G.G. (1992) : A comparison of the virulence, stability and cell wall surface characteristics of three spore types produced by the entomopathogenic

REFERENCE

fungus *Beauveria bassiana*. Appl. Microbiol. Biotechnol., 36(6): 785-789.

Hegedus, D.D. and Khachatourians, G.G. (1995): The impact of biotechnology on hyphomycetous fungal insect biocontrol agents. Bio-technol. Adv. 13, 455-490.

Hegedus, D.D. and Khachatourians, G.G. (1996): Identification and differentiation of the entomopathogenic fungus *Beauveria bassiana* using polymerase chain reaction and single -strand conformation polymorphism analysis. J. Invertebr. Pathol. 67 : 3, 289-299.

Herrera, F.; Carballo, M. and Shannon, P. (1999): Efficacy of native entomopathogenic fungal strain against *Bemisia tabaci* in the laboratory. Manejo.Integrado.de.Plagas. 54, 37-43.

Hugh, G. G. and Michael, J. G. (1995): Purification and cloning of DNA fragments fractionated on Agarose Gels. Molecul. Biotechnol. 3 : 2, 135-138.

Hung, S. and Boucias, D.G. (1992): Influence of *Beauveria bassiana* on the cellular defense response of the beet armyworm, *Spodoptera exigua*. J. Inverteb. Pathol., 60(2): 152-158.

REFERENCE

- Jeffs, L.B.; Feng, MingGuang, Falkowsky, J.E.; Khachatourians, G.G. and Feng, M.G. (1997):** Infection of the migratory grasshopper (Orthoptera: Acrididae) by ingestion of the entomopathogenic fungus *Beauveria bassiana*. J. Economic Entomol. 90: 2, 383-390.
- Jeyaratnam, J. (1990):** Acute pesticide poisoning: A major global health problem. World Health Stat. G., 43:139-144.
- Kassa, A.; Zimmermann, G.; Stephan, D. and Vidal, S. (2002) :** Susceptibility of sitophilus Zeamais (Motsch.) (Coleoptera: Curculionidae) and prostephanus truncatus (Horn) (Coleoptera: Bostrichidae) to entomopathogenic fungi from Ethiopia. Biocontrol .Science . Technol. 12 : 6, 727-736.
- Kerwin, J. L. and Washino, R. K. (1986):** Oosporogenesis by *Lagenidium giganteum* Induction and maturation are regulated by calcium and calmodulin. Can. J. Microbiol. 32(8): 663-672.
- Khachatourians, G.G.; Valencia, E.P.; Miranpuri, G.S.; Koul, O.(ed.). and Dhaliwal, G.S. (2002):** *Beauveria bassiana* and other entomopathogenic fungi in the management of insect pests. Microbial . biopesticides. 239-275.

REFERENCE

Kirschbaum, J.B. (1985): Potential implication of genetic engineering and other biotechnologies to insect control Ann. Rev. Entomol. 30, 51-70.

Laemmli, U.K. (1970): Cleavage of structural proteins during the assembly of the head of Bacteriophage T4. Nature. 227, 680-685.

Leal, S.C.M.; Bertioli, D.J.; Butt, T.M. and Peberdy, J.F. (1994): Characterization of isolation of the entomopathogenic fungus *Metarhizium anisopliae* by RAPD-PCR. Mycol. Research. 98 : 9, 1077-1081.

Lin, Huafeng, Huang, B.O.; Li-ZengZhi, Hu-Cui, Lin-HF, Huang, B.; Li-ZZ. and Hu-C. (1999) : Analysis of the relationship between DNA amplified fingerprints and source of *Beauveria bassiana* strains. Mycosystema. 18 : 1, 73-78.

Louela A. Castrillo and Wayne M. Brooks (1998): Differentiation of *Beauveria bassiana* isolates from the Darkling Beetle, *Alphitobius diaperinus*, using isozyme and RAPD analysis. J.Invertebr. Pathol., 60:185-187. J.Invertebr. Pathol., 72, 190-196.

Luthe, D.S. (1992): Electrophoretic analysis of seed proteins in the Dicotyledoneae. Plant Mol. Biol. Repr. 10, 254-262.

REFERENCE

- Maniania, N.K. and Fargues, T. (1984) :** Specificity of entomopathogenic hyphomycetes to *lepidopterous noctuid* larvae.. *Entomophaga*., 29(4): 451-464.
- Maniania, N.K. and Fargues, J. (1992):** Susceptibility of *Mamestra brassicae* (L.) and *Spodoptera* (Boisd.) larvae (Lep., Noctuidae) to the hyphomycetes *Paecilomyces fumosoroseus* (Brown and Smith) and *Nomuraea rileyi* (Samson) at two temperature *S. littoralis*. *J. Appl. Entomol.* 113: 518-524.
- Marcelo F. Berretta, roberto E. Lecuona, RUBEN O. Zandomeni, and Oscar Grau. (1997):** Genotyping isolates of the entomopathogenic Fungus *Beauveria bassiana* by RAPD with fluorescent labels. *J. Invertebr. Pathol.* 71, 145-150.
- Maria, S.G., Luigi De Micheli and Attilio Scienza (1996):** Characterization of *Vitis* germplasm using random amplified polymorphic DNA markers. *Genetic Resources and Crop Evolution* 43: 187-192.
- Marrone, P.G. and Machntosh, S.C. (1993) :** Resistance to *B. thuringiensis* and resistance management, In *B. thuringiensis an Environmental Biopesticide; Theory and practice*; Ebrwistfe, P.E.; Cory, J.S.; bailey, M.J. and Higgs, R. (eds.); John Wiley & Sons, Ltd., Chichester, UK.

REFERENCE

- Marshall, H.F.; Shirer, M.A. and Cherry, J.P. (1984):** Characterization of glandless storage proteins by sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Cereal chem.* 61,166-169.
- McCoy, C. W.; Samson, R. A. and Boucias, D. G. (1988):** Entomogenous fungi in 'CRC Handbook of Natural Pesticides, Part A. Entomogenous Protozoa and fungi.' (C. M. Ignoffo, ed.),
- Miranpuri, G.S. and Khachatourians, G.G. (1994):** The fungal toxins were possibly involved in the early mortality observed with some *B. bassiana* isolates. *J.Appl.Entomol.* 118 : 1 , 103-110.
- Mullis, K.B. and Faloona, F.A. (1987):** Specific synthesis of DNA in vitro via a polymerase-catalyzed chain reaction. *Methods Enzymol.* 155:335-350.
- Nadeau, M.P. and Biosvert, J.L. (1994) :** Larvicidal activity of the entomopathogenic fungus *Tolypocladium cylindrosporum* (Deuteromycotina: Hyphomycetes) on the mosquito *Aedes triseriatus* and the black fly *Simulium vittatum* (Diptera: Simuliidae). *J.American.Mosquito.Control.Association.* 10 : 4, 487-491.
- Negasi, A.; Parker, B.L. and Brownbridge, M. (1998) :** Screening and bioassay of entomopathogenic fungi for the control of silverleaf

REFERENCE

whitefly, *Bemisia argentifolii*. Insect.Science.and.its.Application.
18 : 1, 37-44.

Neuveglise, C.; Brygoo, Y.; Vercambre, B. and Riba, G. (1994):

Comparative analysis of molecular and biological characteristics
of strains of *Beauveria brongniartii* isolated from insect
S.littoralis. Mycol. Re *S.littoralis*. 98 : 322-328.

Neves, P.J. and Alves, S.B. (2000) : Selection of *Beauveria bassiana*

(Bals.) Vuill. and *Metarhizium anisopliae* (Metsch.) Sorok.
Strains for control of *Cornitermes cumulans* (Kollar). Brazilian.
Archives. Biol. Technol. 43 : 4, 373-378.

Newbury, H. J. and B. U. Ford-Lloyd (1993): The use of RAPD for
assessing variation in plants. Plant growth ., 12:43-51.

Obornik, M. and Landa, Z. (1997) : Molecular phylogeny of
entomopathogenic fungus *Paecilomyces fumosoroseus* based on
RAPD analysis. Sbornik - Jihoceska - Univerzita - Zemedelska -
Fakulta-Ceska-Budejovice-Fytotechnicka-Rada. 14 : 2, 57-62.

Obornik, M.; Klic, M. and Zizka, L. (2000) : Genetic variability and
phylogeny inferred from random amplified polymorphic DNA
data reflect life strategy of entomopathogenic fungi . Canadian. J.
Botany. 78 : 9, 1150-1155.

REFERENCE

- Oliveira, N.T.; de.; Messias, C.L.; De. and Olivera, N.T. (1996) :** Variability of strains of *Metarhizium anisopliae* var. *anisopliae* (Metsch.) Sorokin in terms of protein content and spore germination. *Revista. De. Microbiologia.* 27 : 4, 268-272.
- Onofre, S.B.; Miniuk, C.M.; Barros, N.M.; de.; Azevedo, J.L.; de. and Barros, N.M. (2001):** Pathogenicity of four strains of entomopathogenic fungi against the bovine tick *Boophilus microplus*. *American. J. Veterinary . Research.* 62 : 9, 1478-1480.
- Ouellet, T. and Seifert, K.A. (1993):** Genetic characterization of *Fusarium graminearum* strain using RAPD and PCR amplification . *Phytopathol.* 83:1003-1007.
- Ozino, O.I.; Piatti, P.; Cravanzola, F.; Bridge, P(ed).; Couteaudier, Y(ed). and Clarkson, J. (1998) :** RAPD - PCR for the molecular characterization of *Beauveria brongniartii* isolates useful as biocontrol agents of the European cockchafer *Melolontha melolontha*. *Molecul.Variability. Fungal. Pathogens.* 239-251.
- Paccola, L. D. and Araujo, W. L. (1993):** Mycelial development and nuclear differentiation in the deuteromycete *Beauveria bassiana* cultivated in different liquid media. *Microbiologica.,* 16: 73-78.

REFERENCE

- Pfeifer, T.A. and Khachatourians, G.G. (1993):** Isolation of DNA from entomopathogenic fungi grown in liquid cultures. *J. Invertebr. Pathol.* 61:2, 113-116.
- Piatti, P.; Cravanzola, F.; Bridge, P.D. and Ozino, O.I. (1998):** Molecular characterization of *Beauveria brongniartii* isolates obtained from *Melolontha melolontha* in Valle d'Aosta (Italy) by RAPD-PCR. *Lett. Appl. Microbiol.* 26 : 4, 317-324.
- Rashed, E. (1993):** In the biopesticide *Bacillus thuringiensis* and its applications in developing countries Eds. Salama, S.; Morris, O.N. and E. Rashed proceedings of an international workshop organized by NRC-Cairo, Agriculture Canada and IDRC, 4-6 November., 399: (1991).
- Riba, G.; Bouvier-Foucade, J. and Caudal, A. (1986):** Isozymes polymorphism *Metarhizium anisopliae* (Deuteromycotina: Hyphomycetes) entomopathogenous fungi. *Mycopathologia* 96,161-169.
- Saiki, R.K.; Gelfand, D.H. and Stoffel, S. (1988):** Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239:487-491.

REFERENCE

- Sambrook, J., E.F. Fritsch and T. Maniatis (1989):** Molecular cloning. A laboratory manual, Second edition. 6: 3-9.
- Schilling, A.G.; Molle, E.M. and Geige, H.H. (1994):** RAPDs of *Fusarium culmorum* and *F. graminearum*: Application for genotyping and species identification. In Modern Assays for Plant Pathogenic Fungi: Identification, Detection and Quantification, ed. A. Schots, F. M. Dewey, and R.Oliver, PP. 47-56, CAB international, Oxford.
- Sharma, S.; Agarwal, G.P. and Tajak, R.C. (1994) :** Toxins of the entomogenous fungus *Beauveria bassiana* destroy the natural physiological balance in the host. J. Exp. Biol. 32 : 3, 168-171.
- Shimizu, S.; Nishida, Y.; Yoshioka, H. and Matsumoto, T. (1991):** Separation of chromosomal DNA molecules from *Paecilomyces fumosoroseus* by Pulsed-field electrophoresis. J.Invertebr. Pathol. 58:3, 461-463.
- Shimizu, S.; Arai, Y. and Matsumoto, T. (1992):** Electrophoretic Karyotype of *Metarhizium anisopliae*. J.Invertebr. Pathol., 60:185-187.

REFERENCE

- Shimizu, S.; Yoshioka, Y. and Matsumoto, T. (1993):** Electrophoretic characterization of chromosomal DNA from *Paecilomyces farinosus*. Transactions-of-the-Mycological-Society-of-Japan. 34:2, 239-244.
- Snider, R.D. (1973):** Electrophoresis and the taxonomy of phytopathogenic fungi. Bull Torrey Bot Club 100:272-276.
- St.; Leger, R.J.; Butt, T.M.; Goettel, M.S.; Staples, R.C. and Roberts, D.W. (1989):** Production invitro of appressoria by the entomopathogenic fungus *Metarhizium anisopliae*. Exp. Mycol. 13 : 3, 274-288.
- Thom, C. and Raper, K.B. (1945):** Amanual of Aspergillus Willamas and Wilkins Co. Batimore, Mol.
- Urtz, B.E. and Rice, W.C. (1997):** RAPD-PCR characterization of *Beauveria bassiana* isolates from the rice water weevil *lissorhoptrus oryzophilus*. Lett. Appl. Microbiol. 25 : 6, 405-409.
- Valadares-Inglis, M.C. and Peberdy, J.F. (1998):** Variation in the electrophoretic karyotype of Brazilian strains of *Metarhizium anisopliae*. Genet. Molecul. Biol. 21 : 1, 11-14.

REFERENCE

Valderrama, F.; A.M.; Cristancho, A.; M.A. and Chaves, C.B. (2000):

Analysis of genetic variability of the entomopathogenic fungus *Beauveria bassiana* by RAPD markers. Revista. Colombiana.de. Entomologia. 26 : 1-2, 25-29.

Van Frankenhuyzon, K. (1993): The challenge of *B. thuringiensis*. In

B. thuringiensis an environmental biopesticide: Theory and practice; J.S. Cory; M.J. bailey, and S.R. Higgs, (eds.) PP. 1-35; John Wiley Sons, Ltd., Chichester, UK.

Westwood, G; Huang, S and Keyhani, N (2005): Allergens of the

entomopathogenic fungus *Beauveria bassiana* Clinical and Molecular Allergy 3:1.

Whitney P.J.; Vaughan J.G. and Heale J.B. (1968): A disc electrophoretic

study of the proteins of *Verticillium albo-atrum*, *Verticillium dahliae* and *Fusarium oxysporum* with reference to their taxonomy. J. Exp. Bot 19:415-426.

Wilcox, D.R.; Shivakuman, A.Q.; Melin, B.E.; Miller, M.F.; Benson, T.A;

Shopp, C.W.; Casuto, D., Gundling, G.J.; Bolling, T.j.; Spear, B.B.

and Fox, J.L. (1986): Genetic engineering of bioinsecticides. In: Protein engineering: application in science, Medicine and industry, (eds. Inouye, M. and sarma, R.), PP. 395-413.

REFERENCE

Williams, J. G., A. R. Kubelik, K. J. Livak, J. A. Rafalski and S. V. Tingey

(1990): DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids* 18:6531-6535.

Zurek, L. and Keddie, B.A. (2000) : *Beauveria bassiana* (Balsamo)

Vuillemin- -a promising microbial control agent of the *Satin moth* (Lepidoptera: Lymantriidae). *Biocontrol. Science. Technol.* 10 : 5, 641-644.