

REFERENCES

REFERENCE

- Ahmad, V. U.; Aliya, R.; Perveen, S and Shameel, M. (1992):
"Asterol Glycoside from marine Green-Algae *Codium iyengurii*"
phytochemistry 31:1429-1431.
- Ahmed ,V.U.; Aliya .R.;perreens .and shameel . N. (1993). "Streols
from marine Green algae cadium decorticatum." Phyto chemistry,
33 : 1189 – 1192 .
- Abbott, I.A ., (1996) "Ethnobotany of sea weeds : clues to uses of sea
weeds " Hydro biologica, 326 / 327 : 15-20 .
- Abd El-Fattah, A. F. and sary, H. H. (1987): "Glycoproteins from ulva
lactuca". Phytochemistry 26:1447-1448
- Aknin, M.; Nzaou, R. M.; Cisse, E.; Kornprobst, J. M.; Gaydou, E.
M.; Samp, A.and Miralles, J. (1992): "Fatty acid composition of
twelve species of chlorophyceae from the seegalese" Coast,
Phytochemistry .3:2739.
- Aleem, A. A. (1993): (Ed.) "The marine algae of the Alexandria"
Egypt.Ed University of Alexandria, Egypt.135pp +55 plates
- Arieli, A.; sklan, D. and Kissil, G. (1993): "Anote on the nutritive
value of *ulve lactuca* for ruminants" Animal production, 57(2),
329 – 331.
- Attaway, D. H.and Zaborsky, O. R. (1993): "Marine biotechnology" I
Pharm. Bioac. Nat. Prod.

- Awad, N. E. (1998):** "Phytochemical and Biological studies on the green-algae *Enteromorpha intestinalis* link" Egypt. J. Pharm. Sci. 39:303-322.
- Awad, N.E. (2000)** “ Biologically active steroid from the green algae *ulva lactuca*” phyto therepy – Res14 (8): 641 – 643 .
- Baller. I;L Maran la, K.A; Young; y. shimizu.; C. Fairchild; L. cornell; J. Macbeth and S. Huang. (1995).** “Isolation and structure of carbenolide from a cultured free- swimming caribbean dino flagellate, *Amphidinium* Sp S, - 36 – S” journal of organiz chemistry. 60: 1084 1086 .
- Ballesteros .E . , martin D., and uriz . M.J. (1992)** “ Biological activity of extracts from some mediterranean macro phytes” *Botanica marina* . 35 : 481 – 485 .
- Bandara, B. M. R.; Gunatilaka, A. A. L.; Kumar, N.s.and Wimalasiri, W. R. (1988):** "Antimicrobial activity of some marine algae of Sri lanka." J. the national. Science. Council. of .Srilanka. 16(2): 209-221.
- Barbier, P.; Guise, S.; Huitorel, P.; Amade, P.; Pesando, D.; Briand, C.and Peyrot, V. (2001):** “Caulerpenzne from *CaulerPa Taxifolia* has an antiproliferative activity on tumor cell line SK-N-SH and modifies the microtubule net work”. life Sciences. 70:415-429.

- Biosvert, C. (1984):** (ed.) "La cuisine des plantes sauvages". Dargaud Éditeur, Paris, pp. 302.
- Biosvert, C. (1988):** (ed.), "Les Jardins de La mer du bon usage des algues". Terre Vivante, Paris, pp. 149.
- Bird, k. T. (1987):** "Cost analysis of energy from marine biomass. In: Seaweed cultivation for renewable resources". Bird, K. T., and Benson, P. H., (ed), 327-350. Elsevier, Amsterdam.
- Blunden, G. (1991):** "Agricultural uses of seaweeds and seaweed extracts. In: Seaweeds Resources in Europe: Uses and Potential". Guiry, M. D. and Blunden, G. (ed.), John Wiley and Sons, Chichesier 65-81.
- Blunden, G., Binns, W. W., and Perks, F. (1975):** "Commercial collection and utilization of marin". Econ. Bot. 29:140-145.
- Bonotto, S. (1979):** "List of multi-cellular algae of commercial use. In: Marine Algae in Pharmaceutical Science" (Ed by H. A. Hoppe, T. Levringly. Tanaka) pp. 121-137.
- Brieskorn, C.H.; Khuger, H. and Polonills, W. (1961):** "Triterpenes and sterols in leaves of *Salvia trilobe* and *Pyrus malus*". Arch. Pharm, 294,389.
- Burrows, E.M. (1991).** Seaweeds of the British Isles. Volume 2. Chlorophyta. London: British Museum (Natural History).

- Ermakova, S. P.; Burtseva, Y. V.; Sova, V. V.; Kratchun, V. V., and Zvyagintseva, T. N. (2001):** Proteins of brown seaweeds as inhibitions of endo-1,3-beta-D-glucanases of marine invertebrates. *Biochemistry Mose*, 66:188-94.
- Fahmy, I.R.; Khafagy , C.F.; Islam, a.M.; Olana, H. Y and Gizgis, A.N. (1933):** “Preliminary verification of the chemical constitution of fourty-one desert plants from the north – western coastal region”. *Drug Research, Egypt*, 6:39.
- Fieser, L.F. and Fieser, M. (1959):** “Sterols”. Rain hold publishing Co., New Yourk.
- Finical, W. (1975):** Halogenation in the Rhodophyta, Review. *J. Phycol.* 11:245-259.
- Flodin, C., and Whitfield, F. (1999):** 4-Hydroxybenzoic acid:alikely precursor of 2,4,6-Tribromophenol in *Ulva lactuca*, *Phytochemistry* 51:249-255.
- Flurence, J and Gueant, JL. (1999):** “Algae a new source of proteins”. *Biofuture*, No : 191, 32 – 36.
- Galal, A.M; El-Ayouty, Y.M; Hashem. S.A and El-Sabbahy, G., (1999):** “Screening of different algal extracts for their efficiency against mechanical transmission of two phyto viruses”. *Acta. Hydrobiol*, 41 : 2, 155-163.

- Glombitza K.W. (1979)** “ Antibiotics from algal . In Hoppe H.A. Learning T. Tanaka y (eds) marine algae in pharmaceutical science . water de Gruyter. “ Berlin pp 303 – 42 .
- Gomorri, G. (1942):** Calcium; phosphorus and phosphatase. In “Practical clinical Biochemistry” **PP. 431-466**. Edited by Varley, H. A. H.
- Guiry, M.D. and Blunden, G. (1991).** (ed) “sea weed resources in Europe : Uses and potential”. John Wiley, England.
- Guiry, M. D. and Blunden, G. (1991):** (ed.) Seaweed resources in Europe: Uses and Potential. John Wiley, England.
- Hagurag, R. (2000):** Diabetes mellitus and life style for the primary prevention of diabetes mellitus: the role of diet. Br. J. Nutr., 84:191-194.
- Harvey, W. (1988):** Biotechnology 6:488-492.
- Hegazi, M.M.; Perez, R.A.; Almela, L.; candela, M.E. (1998):** “Separation and identification of chlorophylls and carotenoids from *Caulerpa prolifera*, *Jania nubens* and *Padina pavonica* by reversed– phase high – performance liquid chromatography”. J. of- chromatography, A. 829: 1-2, 153-159.
- Hellio, C.; Bremer, G.; Pons, A.M.; Gal, y.L; Bourgougnon, N. (2000):** “Inhibition of the development of microorganisms

(bacteria and fungi) by extracts of marine algae from Brittany, France". *Applied. Microbiol Biotechnol* 54 (4) 543 – 549.

Hoppe, H. A. (1979): Marine Algae and Their products and Constituents in Pharmacy, Berlin. 1:25-119.

Hoppe, H.A. (1979) "Marine algae and their products and constituents in pharmacy In Hoppe H A . Learning T . Tamake y (rds) marine algae in phamacelitical science water de Gnyen . " Bernlin, 1: 25 – 119 .

Hornsey, I.Y. D. and Hide.(1874) . "the production of antimicrobial compounds by British marine algae I : Antibiotic – producing marine algae . "B.r phycol . J. 9 : 353 – 361 .

Hornsey , I.Y.D and Hide. (1976) . " the production of antimicrobial compounds by British marinee algae II: seasonal variation in production of antibiotics" . Br. Phycol. I. 11 : 63 – 67 .

Hurtado, A. Q., Agbayani, R., Sanares, R., and de Castro-Mallare, M. T., (2001): The seasonality and economic feasibility of cultivating *Kappaphycus alvarezii* in Panagatan Cays, Caluya, Antique, Philippines. *Aquaculture*, 199:295-310.

Ivanova, E.; Toshkova, R.; Ivanva, V.; Rouserva, R. and Koarova, M. (1994)a: "Stimulating effect of *Ulva lactuca* Heteropolysaccharide on the Immune Response in mice". *Doki. Bulg. Akad. Nauk.* 47:109-112.

- Ivanova, V.; Rouseva, R.; Kolarova, M.; Serkedjieva, J.; Rechev, R. and Malonova, N. (1994)b:** "Isolation of polysaccharide with antiviral effect from *Ulva lactuca* " *Prep. Biochem.* 24:83-97.
- Ivanova, V.; Gegova, G.; Manolova, N.; Serkedjiva, J. and Belomatove, D. (1991):** "Biologically active substances from marine organisms, 1. Isolation of a substance from marine algae with antiviral effect". *Herba – Hungarica*, 30(3), 47-53.
- Kalaiselvi, M; Manoharan, M; Kathiresan, K., (1993):** "Effect of maleic hydrazide on the total chlorophyll content sea weeds". *J. of-Environmental –Biology*, 14(1), 17-20.
- Kamat, S. Y.; wahidulla, S.; D.sollza,L.; Naik, C. G.; Ambiy, V.; Bhakuni, D. S.; Geol, A. K.; and Garg, H. S. (1992):** "Bio activity of marine organisms, VI Antiviral evaluation of marine algal extracts from the Indian coast" *Bot. Marina*, 35 :161 – 164.
- Kim, J. H.; Hudson, J. B.; Huang, A. M.; Bannistes, K.; Jin, H.; Choi, T. J.; Towers, G. H. N.; Hong, Y. K.; and De wreede, R. E. (1997):** "Biological activities of seaweed extracts from British Columbia, Canada, and Korea. Antiviral activity." *Can. J. Bot. Rev. Can. Bot.* 75(10), 1656-1660.
- Kuo, I. M.; Hwang, A. and yeh, D. B. (1997):** "Purification, substrate specificity, and products of a Ca^{2+} stimulating lipoxygenase from sea

the same as active gradient". Biotechnology, Abstract No 2752818.

Neushul, P. (1987): Energy from marine biomass: The historical records. In: Seaweed cultivation for renewable resources. (ed.), 1-37. Bird, K. T. and Benson, P. H. Elsevier, Amsterdam.

Newton, L. (1951): (ed.) Seaweed Utilisation. Sampson Low, London, pp. 188.

Nisiwaza, K.; Noda, H.; Kikuchi, R. and Watanabe, T. (1979): The main seaweed foods in Japan. Proc. Intl. Seaweed Symp. 12.5-29.

Ohno, M. and Critchley, A. T. (1993): (ed.) Seaweed cultivation and marine ranching. Japan Intr. Cooperative Agency, Nagai, Yokosuka.

Okai, Y.; Higashi, O. K.; Ishizaka, S.; and Yamashita, U. (1997): "Enhancing effect of polysaccharide from Edible Brown alga, *Hijikia Fusiforme* (Hijiki), on Release of tumour necrosis factor alpha from macro phages of Endotoxin non responder C₃H/He J. Mice" Nutr. Cancer. 27(1), 71-79.

Olessen, P. E.; Maretzki, A. and Almodovor, L. A. (1963): An investigation of antimicrobial substances from marine algae, Bot. Marina, 6:226-232.

Paech, K. and Tracey, M. V. (1956): Modern methods of the plant analysis. Springer- Verlag, Berlin., 1 (4): 643.