

REFERENCES

- Agodi, A; Jones, C; Threlfall, EJ; D'Angelo, M and Marranzano M (1990)** : Molecular characteristics of trimethoprim resistance in *Shigella sonnei* in Sicily. *Epidemiol. Infect.* 105 : 29-40.
- Agodi, A; Marranzano, M and Jones C (2000)** : Characterization of trimethoprim resistance genes in K-aerogense isolate in Sicily. 1985-1988. *Eur. J. Epidemiol.* 11 : 33-38.
- Agodi, A; Marranzano, M and Jones C (1995)** : Characterization of trimethoprim resistance genes in *Salmonella* isolated in Sicily. 1985-1988. *Eur. J. Epidemiol.* 11 : 33-38.
- Ahmed F.M (2004)**Antimicrobial studies of some essential oils on some bacteria; Master Degree; Botany Department; Faculty of Science; Botany Department .
- Alves, TM; Silva, AF; Brandao, M; Grandi, T; Smania, E; Junior, A and Zani, C (2000)** : Biological screening of Brazilian medicinal plants. *Mem. Inst. Oswaldo; Cruz. Rio de Janeiro.* 95 : 367-373.
- Arakawa, T and Osawa, K (2000)** : Pharmacological study and application to food of mint flavor-antibacterial and antiallergic principles. *Aroma Research.* 1 (1) : 20-23.
- Araki, W.B and Tomai (1999)**: Quantitative relationships in human serum and non-vascular fluids; T.B.Jr. *Gamma Globulins Science*; 142; 1080.

- Asscher, A; Kokkini, S and Lanaras, T (1995) :** Antifungal activities of *Origanum Vulgare* subsp. *Hirtum*; *Mentha spicata* *Lavandula angustifolia* and *salvia fruticosa* essential oils against human pathogenic Fung. *J. Agric. Food Chem.* 46 : 1739-1745.
- Asscher, D.B and Finkel, G (1998):** *Candida pyelonephritis* in Kass B.H.; editor: *Progress in Pyelonephritis*; Philadlephia; F.A. Davis Co.; pp. 179-184.
- Asscher, N; Legakis, N.J; Papavassilion, J and Sifalakis, J (1999):** Sterilization of urological instruments infected with *Pseudomonase aeroginosa*; Dept. Microbiol. Fac. Med.; Univ. Athens. *Actampicrobiol Hellen* (285-291).
- Baeza, D.A and Berruti, J.D (1998):** Method for detection significant bacteriuria a large group of patients. *J. Clin. Pathology*; 117: 298.
- Bailey, R.J and Kirby, W.M (1996):** Simple tests for significant bnacteriuiia. *Arch. Intern.*; 112: 742.
- Baron, E.J and Finegold, S.M (1990) :**General microbiology ; Baily and Scotts *Diagnostic Microbiology*. 8th. Ed. Mosby Co. St. Louis. Balitimore.
- Barr, S.T (1997):** Hereditary interstitial nephritis. In: *Biology of pyelonephritis*; edited by B.I. Nas and Boston; Bittle; Browen and Marry; pp. 427-432.
- Barry, L.B and Thrnsberry, Y (1987) :** Inhibition of growth and aflatoxin production by cinnamon and Clove oils. *Cinnamic aldhyle and eugenol*. *J. Food Sci.* 42: 1107-1109.

- Bauer, A.W; Kirby, W.M; Sherris, J.C and Turk, M (1986)** : Antibiotic susceptibility testing by a standerized single disk method Am. J. of Cl. Pathology; 45: 493-496.
- Beeson, J; Filloy, L and Galindo, E (1996)** : Resistance of Shigella dysenteria type I to ampicillin and other antimicrobial agents; strains isolated during a dysentery outbreak in a hospital in Mexico. J. infect Dis. 133: 572.
- Ben-Hamida, N; Abdelkefi, M.M; Aissa, R.B and Chaabouni, M.M (2001)** : Antibacterial screening of Origanum majorana L. oil from Tunisia. Journal of Essential Oil Research. 13 (4) : 295-297.
- Beuchaut, J (1994)** : Effects of variety and ontogenic stage on the essential oil composition of biological activity of fennel (Foeniculam Vulgare Mill) J. Essent. Oil Res. 6: 57-62.
- Blasser, M.J; Huq, M.I; Glass, R.I; Zimicki, S and Brikness, KA (1982)**: Salmonellosis in rural and urban clinics in Bangladesh. Epidemiologic and clinical characteristics. Amer. J. Epidemiology; 116 : 266-275.
- Blasser, G.K.; Glass, A.R. and Edwarders, J.E (1991)**: Pathology of acute and healed experimental pyelonephritic kidneys. Brit. J. S Urg.; 41: 38.
- Borrows, L.R (1993)**: Studies of urinary tract infection in diabetes. Bacteriuria in patients with diabetes mellitus and in control subjects; II. Significant bateriuria in relationship to long term diabetic management. Acta Med.; 179: 173; 183.

- Boston, HC (1987)** : Resistance of fluoroquinolones. Rev. Inf. Dis; 10: 57.
- Boston, J and Sandys, G.H (1992)**: Diagnosis urinary infection in general practice by means of dipinoculum. Transport medium. Brit. Med. J.; 2: 1286.
- Bouanchaud, D.H and Chabbert, Y.A (1981)** : Practical effectiveness of agents curing of R plasmids NY. Acad. Sci. 182 : 305-311.
- Brenner, R.M and DeNavasquez, S.J (1997)**: Experimental pyelonephritis in the mouse produced by *Escherichia coli*; *Pseudomonas aeruginosa* and *Proteus mirabilis*. J. Path. Bact.; 87: 79-89.
- Brogan, O.M (1994)**: The quantitative significance of bacteria visualized in the constained urinary sediment. New. ENgl. J. Med.; 265: 589-570.
- Brostoff, R.K and Male, E.J (1985)** : Nosocomial urinary tract infection: A microbiological study. Ann Acad. Med.; Singapore 22: 6: 873-7.
- Brown, M.R; El-Beih, F.M and El-Satoury, E.H (1996)** : Total lipid and pattem of certain antibiotic resistant and sensitive strains of *E.coli* and *pseudomonas aeuroginosa* .Egypt.J.Microbiol.; 9:37-50.
- Brumfitt, K; Sivropoulou, A; Lanaras, T and Arsenakis, M.B (1984)** Antifungal activities of *origanum vulgare* Subsp. *hirtum* ; *Mentha spicata* ;*salvia fruticosa* essential oils against human pathogenic fungi.J.Agric .Food.Chem.46:1739-1745.

- Buffenmyer, M; Benesuva, D and Rossmann, P (1986):** Pathogenecity of various strains of *Escherichia coli* for the intract rabbit kidney and the effect of repeated passage of renal tissue progress in pyelonephritis Philadelphia; S.A.; Davis Co.; pp. 155-196.
- Burrows, M.O (1963):** Role of ureter in pathogenesis of ascending pyelonephritis. JAWA; 168: 1983.
- Burton, Y.U and Borisova, G.N (1978) :** Antibiotic 7 (11): 975 (CF. Egorov; N; 1985; P. 377).
- Campbell, P; Yoshida, Y and Terakwi, Y (1985) :** Antibiotic resistance and R plasmid in salmonella isolated from human in Japan (1966-1972). In microbiol Drug Resistance Mitshhashi S and Hashimoto H (eds). University Park press; Tokyo; London. Pp. 237-252.
- Campbell, V.K and Kozreva, L.F (2005) :** Experience with the new chemotherapeutic nalidixic (Negram) against Gram-negative pathogens praxis;Epidemiol 53: 15-19.
- Cavalito, A.C (2000) :** Synthesis of amino-Carboxyvaleryl glycine and amino-Carboxuvaleramide .Biochem.J.58:266.
- Chara, J.C and Balter, J (1996) :** The action of essential oils on phytopathogenic fungi. Plant Dis. Dep. Rep; 43; 1143-1147.
- Chen, W.L; Dommelhofcomplex; Eindkovenklin and Geriat, B.L (1999):** The indwelling catheter in chronically ill patients; No. I; 78-89.

- Chodirker, C.R; Hewitt, W.L and Lenguze, L.B (1996):** An out break of infection caused by gentamicin resistant staphylococcus aureus ;Pyelonephritis; *Nedecine*; 39: 3.
- Collee, J.G and Marr, W (1989) :** Culture containers and culture media. In practical Medical Microbiology 13th ed. Vol. (2). Collee; JG : Duguid JP : Fraser AG; and Marmion; B. P. (eds); Churchill Livingstone. New York. Pp. 100-120.
- Collee, N.D; Duguid, S; Lt, A.M; Verrier, N.B (1984):** The bacteriology of U.T.I. in Kuwait J. of the Medical Association vol. 7. No. 3; page 175; September.
- Cosentino, S; Tuberoso, C; Pisano, B; Satta, M; Mascia, V; Arzedi, E and Palmas, F (1999) :** In-vitro antimicrobial activity and chemical composition of Sardinian thymus essential oils. *Letters in Applied bacteria*. 29 (2) : 130.
- Courvalin, P (1999) :** Transfer of antibiotic resistance genes between Gram-positive and Gram-negative bacteria. *Antimicrob. Agent and Chemother*. 38 : 1447-1451.
- Dabbah, C; Koutsoumanis, K and Nychas, G.J (1987) :** Inhibition of Salmonella enteritidis and staphylococcus aureus in nutrient broth by mint essential oil special issue. Improved traditional foods for the next century food research international. 33 (3 – 4) : 273-280.
- Dapkevicius, A; Van Beek, T.A; Lelyveld, G; Van Veldhuizen, A; de Grot Linssen, J and Venskutonis, R (2005) :** Isolation and

structure elucidation of radical scavengers from thymus Vulgaris leaves. J Nat Prod.; 65 (6) : 892-6.

Da-Silva, J.P; Kaber, R; Robins-Brown, V; Prado, P; And Levine, M.M (1989) : Patterns of adherence of diarrheagenic Escherichia Coli to Hep2 Cells. *Pediatr. Infect. Dis.* 6: 829-831.

Datta, N (1969) : Plasmid classification in compatibility grouping. In plasmids of Medical; Environmental and Commercial importance; KN; Timmis and A. Puhler (Editors) P; 3-12. Elsevier/North Holland Biomedical press.

Davis, N.D and Maston, L.T (1994): Evaluation of Dip inceculum transport medium in diagnosis of U.T.I. *Medical Jou. Of Cairo University* Vol. 44; No. 3.

Deans, AN and Ritchie, K (1987) : Antibacterial; antidermatophytic and antitoxigenic of onion (*Allium Cepa* L) oil *Microbiological research* 150 (2) 167-172.

Deans, M (1997) : Antimicrobial activity of some essential oils from Lamiaceae and compositae. *International Journal of food Microbiology.* 67 (3) : 187-195.

Donneberg, M and Kapper, J (1992) : Minire view : Entero pathogenic E. Coli. *Infection. Immunity* 60 : 93-53.

Douria, D and Finkel, C (1983) : Antioxidant activity of chemical components from sage (*Salvia Offivinalis* L) and thyme (thymus vulgaris L) measured by the oil stability index method. *J Agric food chem* : 50 (7): 1845-51.

- Dubos, T.L; Eried, F.A and Goven B.A (1998):** Dept. Surg.; Univ. North Carolina; Champel Hill; N.C. 27514-J. Surg. Res. 1813 (277-284): Evidence for Obst. Ractive Etiology in Intravneous Bacterial Pyeloneph.
- Duguid, D.M and Staney, T.A (1994):** Bacteriologic localization patterns in bacterial. Prostatitis and Invest. Urol.; 5: 412.
- Dyer, R and Cotchin, B (1999):** Studies on coryne bacterium; venale II. The experimental pathogenicity for uric. J. Comp. Path.; 56: 205-214.
- Edmondson, R.A and Cooke, M.C (1999):** A pathological study of 5 cases of pyelonephritis in the newborn. Arch. Dis. Children; 31: 303.
- Edward, M.F and Little, J.D (1997):** Ureteral obstruction in infancy a study of 74 cases. J.A.M.A.; 92: 544.
- El-Gayyar, M; Draughon, F.A; Golden, D.A and Mount, J.K (2003) :** Antimicrobial activity of essential oils from plants against selected pathogenic and saprophytic microorganisms. Journal of Food Protection. 64 (7) : 1019-1024.
- El-Mishad, J (1991) :** Microorganisms isolated from urinary tract infections and their beta lactamase production and evaluation of clinical efficacy of sulperazone. Hinyokika Kivo 34: 8: 1503-14.
- Erioesson, N; Juven, B.J; Shaaya, E; Menasherov, M; Nitzan, R; Weiss Lowicz, H and Ravid, U (1998) :** Inhibitory effect of oregano and thyme essential oils on moulds and food borne bacteria letters in Applied microbiology; 11 (1): 33-37.

- Farag, M.C and Juven, M.A (2003) :** Spasmolytic and antimicrobial activity of essential oil from thymus *Leptophyllus* and *T. Webbii* anus. *Fitoterapia*; 60: 5; 469 – 473.
- Faser, A and Simango, C.S (1982) :** Cope of urinary pathogens isolated in the public health bacteriology laboratory; Harare : Antibiotic sensitivity patterns of isolates and incidence of Haemolytic Bacteria. *Cent. Afri. J. Med.*; 42: 8244-9.
- Freedman, R.H; Ranade, N.P; Schrempf, H and Vining, C (1988) :** Cloramphenicol resistance in streptomycetes : Cloning and characterization of achloramphenicol hydrolase gene from strep. *Venezuelae J. General Microbiol*; 136: 293.
- Freedman, C (1980) :** Screening of plants used in Argentine folk medicine for antimicrobial activity .*Journal of Ethnopharmacology* 39(2):119-128.
- Famer, A.S (1995):** Some clinical aspects of chronic pyelonephritis. In: Quinn; B.L. and Kass; B.H. *Biology of Pyelonephritis* (Henry Foral hospital international symposium). Boston; 955.
- Fry, L (1999) :** The patient with urinary tract infections. In *Manual of Nephrology; Diagnosis and Treatment* (Ed 5 Chrieder); Little Brown and Co; Boston; P9.
- Gohar, G; Kumar, A; Singh, A.K; Bhattachargo A.K; Singh, K and Sharma, V.D (1997) :** Antimicrobial activity of essential oils of some *Ocimum* species and clove oil. *Fitoterapia VII* (6): 429-432.

- Gora, D.j and Hill, V (2000) :** Clinical correlates of Kimmelstiel-Wilson Lesion; New Eng.J.Med.;154:896.
- Gorrill, B and Love, W.C (1994) :** Acalculous nontyphoidal salmonellal; cholecystitis requiring surgical intervention despite ciprofloxacin therapy: report of three cases. Clin. Infect. Dis. 24: 707-709.
- Gorrill, R.H; Denvasgue, S.J and Hossain, M.A (1987) :** Pivecillinam resistant shigellad dysentaria type 1 infection in Bangladesh. The lancet. 335: 1461-1462.
- Gosling, R.R and Little, P.J (1995):** Renal infection and renal scaring edited by Kincaidsmith and Fairel K.F.; Melbcurne; Mecceeds Publishing Co.; pp. 81.
- Guze, H.C (1981) :** Overview of mechanisms of bacterial resistance. DIAGN. Microbiol. Inf. Dis.; 12: 1095.
- Gymans, J (1986):** Cited from excer[ra medioca; urology Yol. 9. 2.; Dept. Surg. Sect. Urol. Univ. Michigan; Ann. Arbor. Mich. Univ. Mich Medcant. J.; 39/3: 103-112.
- Hall, B.H; Monzon, O.T; Ory, B.M and Brennan, J.C (1991):** The microflora of the urinary tract; IH Gunn; B.L. and Kass; E.H. (eds). Biology of Pyelonephriytis (hennery ford Hospital International Symposium Bston; Little; Brawen; p. 391).
- Harvery, C.V; Johns, U; Abrahao, F.N; Diogenes, A and Benedito, P (1989) :** Antibacterial activity of Ocimum gratissimum L. essential Oil. Mem inst Oswald Cruz. Rio de Janeiro; Vol. 94 (5) : 675-678.

- Harvey, D; Agarwal, K.C; Verma, A.D and Dubey, M.L (1997) :**
Incidence of shigellosis and multidrug resistant shigella : a 10
years study. J.of Trop.Med and Hygiene; 90: 25-29.
- Hassanein, F.M and Eldoksch, H.A (1997) :** Antimicrobial action of
carvone and some plant extracts on certain phytopathogenic
bacteria and pathogenicity of *Agrobacterium tumifaciens*.
Alexandria J of Agriculture Research. 42 (1) : 127-136.
- Hayashi, J.C and Henry, P.A (1995) :** The in vitro antibacterial activity
of essential oils and oil combinations. J. Am. Pharm. Assoc; 47;
294-296.
- Hostettman, K (1991) :** Assays for bioactivity methods in plant
biochemistry; vol 6. Academic press; San Diego; 360 pp.
- Hsieh, J and Liu, G.D (1979) :** Isolation ;Purification and antibiotic
activity of o-methoxy cinnamaldehyde from cinnamon. Pharma
Cognosy; phyto chemistry; Medicinal plants. 2nd Edition.
Lavoisier publishing Intercept. Paris Pg. 545-546.
- Huang, S; Contronic, G and Khan, W (1999) :** Resistance of shigella to
smpecillin and other antibiotics. Its clinical and epidemiological
application. JAMA; 221: 45-47.
- Huch, L.J; Griggs, M.C; Hall and Jin Y.F (1999) :** Ciprofloxacin
resistance in clinical isolates of salmonella typimurium obtained
from two patients Antimicrob. Agents Chemother. 37: 662-666.
- Hughes, D and Hayler (1998):** Dipslide and aid to quantitative urine
culture in general practice. Brit. Med. J.; 3: 343.

- Imai, H; Osawa, A; Yasuda, H; Hamashima, H; Arai, T and Sastsu, M (2000)** : Inhibition by the essential oils of peppermint and spearmint on the growth of pathogenic bacteria. *Microbios*; 106 (supplement 1) : 31-39.
- Javor, N and Ivemark, B.I (2000)**: Renal dysplasia and pyelonephritis in infants and children. Part I *Arch. Path.*; 66: 255-263.
- Jawetz, D; Nachman, S and Tomasz, A (1980)** : Identification of an outer member protein of E. Coli with a role in the Corrdination of deoxyribonucleic acid replication and cell elongation. *J. Bact*; 136 : 1143.
- Jones, A.L (1999)**: The pathogenesis of experimental entrococal pyelonephritis in mice. *J. I of Dis.*; 108: 181-188.
- Kambal, A.M (1996)** : Antimicrobial susceptibility and serogroups of Salmonella isolates from Riyadh; Saudi Arabia. *Int J Antimicob Agents*. 7: 265-269.
- Kass, M and Erkmen, O (1997)** : Antimicrobial activity of the essential oils of turkish plant spices. *European food Research and Technology* 212 (6): 658-660.
- Kass, D.D (1998)**: Relation of urinary tract infection to diabetic children. *Nrit. J. Urol.*; 37: 73.
- Kass, L.R (1997)**: Experimental pyelonephritis. VI. Observation on susceptibility of the rabbit kidney to infection by a virulent strain of staph. Aureus; *Yale. J. Biol. Med.*; 32: 272-279.

- Kassis, W (1997) :** Antimicrobial resistance of salmonella isolates in aspanish hospitals. Lancet; 2; 1059; cited from Br. J. Urol.; 36: 360-363.
- Kessner, J and Noble, W.C (1987) :** Transfer of gentamicin resistance between strains of staphylococcus aurens on skin J Gen. Microbiol 107: 391-393.
- Kessner, P; Doleres, D.M; Rodriguez, C.M; Cercenado, E; Pelaez, T and Bouze, E (1989) :** Antimicrobial resistance of salmonella isolates in aspanish hospital. Antimicrob Agents chemother 37: 1200-1202.
- Khuzaie, R.F; Rhashan, L.J and Al-Douri, N.A (1999) :** In vitro antimicrobial activities of the crude aqueous extracts of some local medicinal plants growing in Jordan J. Appli Sci. 1: 46-52.
- Kippax, B.E (1987) :** Resistance of shigella; salmonella and other selected enteric pathogens to antimicrobial agents Rev. infect. Dis 8 (2): S172-S181.
- Kleeman, L; Hawitt, E.A; EL-Shabrawy, A.D and De-Pooter, H.L (2000) :** Chemical composition and rhizome of Aplinia speciosa grown in Egypt. Flavor fragrance Journal 10 (2) 63-67.
- Koneman, E.W; Allen, S.D; Dowell, V.R and Cho, Y (1983) :** Colour Atlas and Text Book of Diagnostic Microbiology. 2nd ed. J. P. Lippincott Company; N. Y.; London. Pp. 57-125.
- Kumara, P.M (1995) :** Activity of gentamicin against pseudomonas and hospital Staphylococcus. Brit.Med.J.; 1:203.

- Lapidas, C.F (1983)** : Spasmolytic and antimicrobial activity of the essential oil from *thymus leptophyllus* and *T. webbianus*. Microbiology Addison-Wesley publishing Co. USA Pp. 575-612.
- Lau, S and Adetubi (2001)** : Pyelonephritis its relation to vascular lesions and to arterial hypertension. Medicine (Ball) 18: 211.
- Lawson, V.C and Wood, J.C (2003)** : Influence of different Media and bloods on results of diffusion antibiotic susceptibility tests. Antimicrob agents chemother. 1: 116-122.
- Lee, L.L; Puher, N.D; Maloney, E.K; Bean, N.H and Tauxe, R.V (1994)** : Increase in antimicrobial resistant Salmonella infection in the United States; 1989-1990. J Infect Dis. 170 : 128-134.
- Lepper, T.I; Rotimi, V.O and Adeugam, A.A (1992)** : A study of the aetiological agents of childhood diarrhea in Lagos Nigeria. J Med Microbiol. 40: 10-14.
- Ling, J; Zhou, G.M; Woo, T.H and French, G.L (1991)** : Antimicrobial susceptibilities and b-Lactamase production of Hong Kong isolates of gastroenteric Salmonellae and Salmonella typhi. J Antimicrob Chemother. 28 : 877-885.
- Longstaff, A.M (1987)** : Studies on beta – Lactomases producing bacteria belonging to genus Pseudo monas Ph. D. Thesis. Faculty of Science; Zagazig University.
- Lowry, S.J (1975)** : Diarrheal disease : Current concepts and future challenges. Enterodherent E. Coli : Aheterogenous group of E.

- Coli implicated as diarrheal pathogens. Trans. R. Soc. Trop. Med. Hyg. 87: 549-553.
- Malowany, L; Smith, D.J and Klavias, J.V (1989):** Renal papillary necrosis induced by heterologous serum. Arch. Pathol.; 78: 108.
- Marino, M; Bersani, C and Comi, G (2000) :** Antimicrobial activity of some essential oils from Lamiaceae and Compositae. International journal of food Microbiology 67 (3) : 187-195.
- Maslin, M.M and Pascule, J.V (2000) :** Thiopropanal S-oxide : alachrymatory factor in onion. J. Agric food. Chem. 19: 269-272.
- Mata, L.J; Urrutia, J.I and Garcia, B (1969) :** Shigella infection in breast feed Guatemalan Indian neonates. Amer. J. Dis. Child; 117 : 142.
- McColla, D.R; Olive, P; and Fan, M.L (1976) :** Gentamicin resistant Staphylococcus aureus associated with the use of tropical gentamicin .Canad J. Microbiol 21 : 1484.
- Mcgeachie, J and Kennedy, J.E (1989) :** De diego Sierra D; Ruiz Poveda; Gacia Rojo A. Bacteriological and Resistance in Ambulatory urinary infections. Aten Primaria 18 (6): 315-7.
- Mears, I.S (1988) :** Isolation; purification and antibiotic activity of O-methoxy cinnamaldhyde from cinnamon. Appl. Environ. Microbiol 36: 577-583.
- Mikhail, I.A; Fox, E; Haberberger, R.L; Ahmed, M.H and Abbatte, E.A (1990) :** Epidemiology of bacterial pathogens associated

- with infection diarrhea in Djibouti. J. Clin. Microbiol. 28 (5) : 956-961.
- Miles, G.C; Misra, G; Moreddu, M and Arcangelletti, M.C (1999) :** Susceptibility of frequent urinary pathogens to Fosfomycin; Trometamol and eight other antibiotics. Results of an Italian multicentre study. Infection 20 suppl; As : 291-5.
- Miura, K; Kikuzaki, H and Nakatani, N (2002) :** Antioxidant activity of chemical components from sage (*Salvia officinalis* L.) and thyme (*Thymus vulgaris* L.) measured by the oil stability index method. J Agric Food Chem; 50 (7) : 1845-51.
- Mocabe, M.M; Myint, W; Myint, M.M; Bwin, M and Aye, T (1985) :** Antibacterial properties of essential oil from six medicinal plants. Myanmar. Health science Research Journal 8 (2): 62-65.
- Monica, C (1985) :** Medical laboratory manual for tropical countries. ELBS. Ed. English Language Book Society; tropical health Technology. Butterworths. Vol. II. Enteric Gram-negative rods. Chap. 42 : 250-255
- Nakamura, C.V; Tania, U; Nakamura, E.B; Abrahao, F.N; Diogenes, A; Garcia and Benedito, P (1999) :** Antibacterial activity of *Ocimum gratissimum* L. essential oil. Mem Inst Oswald Cruz; Rio de Janeiro; Vol. 94 (5) : 675-678.
- Nakaya, P; Yoshida, Y and Terawki, Y (1975) :** Antibiotic resistance and R plasmid in *Salmonella* isolated from human in Japan (1966-1972). In Microbiol Drug Resistance Mitshhashi; S and

- Hashimoto H; (eds). University park press; Tokyo; London. Pp. 237-252.
- Nataro, J.P and Levine, M.M (1994) :** Bacterial diarrheas; p. 697-752; in A. Z. Kapikian (ed); Viral infection of gastrointestinal tract; 2nd ed. Marcel Dekker; Inc.; New York; NY.
- Neomura (1997):** Mechanisms of phagocytosis in human polymorphonuclear leucocytes. Immunology; 10: 137.
- Oseasohn; N.S.R.; Chalmers; J.A.; Mac-Donald; J.S. and Dewardener; H.E. (1999):** Pyelonephritis. Lancet; 1: 351.
- Ozacan, M and Erkmen, O (2003) :** Antimicrobial activity of the essential oils of Turkish plant spices. European food Research and technology. 212 (6) : 658-660.
- Park, N; Juven, B.J; Shaaya, E; Menasherov, M; Nitzan, R; Weiss, H and Ravid, U (1984) :** Inhibitory effect of oregano and thyme essential oils on moulds and food borne bacteria letters in Applied Microbiology; 11 (1) : 33-37.
- Park, R.W; Blake, P.A; Murlin, A.C; Carter, G.P; Pollard, R.A; Merson, M.H; Allen, S.D and Brenner, D.J (1980) :** Increase in antibiotic resistance among isolated of salmonella in the USA; 1967-1975. J. infect. Dis. 142 (4): 485-491.
- Part, Yen; Hui, Chang and Shyh-Ming, Tsao (1988) :** Inhibitory effects of aqueous garlic extract. Garlic oil and four diallyl sulphid against four enteric pathogens. Journal of Drug and food Analysis. 10 (2): 120-126.

- Paster, N; Juven, B.J; Shaaya, E; Menasherov, M; Nitzan, R; Weisslowicz, H and Ravid, U (1990) :** Inhibitory effect of organo and thyme essential oils on moulds and food borne bacteria. *Letters in Applied Microbiology* 11 (1) : 33-37.
- Perkoff, N (1997) :** Analysis of the essential oil of wild organo from Martinique (*Cdues aromaticus* Bent); evaluation of its bacterios and fungisttic properties. *Journal of Essential oil Research* 7 (2): 165-173.
- Porder, S; Subramanyam, V.R and Rath, C.C (1996) :** Effect of essential oil on the availability and morphology of *E. Coli* (Sp-11). *Microbios* 84 (340): 195-199.
- Rasschou, L.B and Kalmanson, G.M (1998):** Pyelonephritis. I. Observations on the course of chronic non-obstructed enterococcal infection in the kidney of the rats; Yale; *J. Biol. Med.*; 33: 372-385.
- Reinhetz, W.E; Recves, D.S and Turnbull, A.R (1998):** *Postgrad. Med. J.*; 45 (Suppl. Mov.) 56 cited from Garrod and O'Grady and O'Grady 1972; *Ameibicti and Chemotherapy*; p. 49.
- Ress, T; Minney, Y; Niskikira, K and Matsuoka, S (2000):** Bacteria isolated from U.T.I. and their drug sensitivity; statistical studies from 1966 to 1970 (Japanese); *Cent. Clin. Lab.; Univ. Hosp.; Yokohama Jap. CLin. Urol.*; 27/2 (155-162).

- Robbins, J.A; Miller, B.R and Hinman, F (1999):** Perpetuation of infection in unobstructed urinary tract by vesicoureteral reflux. *J. Urol.*; 90: 88.
- Robert, C.R; Valkenburg, H.A; Haverkorn, M.J; Goslings, W.H; (1991) :** A prospective study of urinary tract infection in a Dutch general practice. *Lancet* 2 : 674-7.
- Roitt, P.G; O'shea, B; Flamagan, K.P and Falkner, F.R (1989) :** Urinary tract infection in General infection in General practice; Direct sensitivity testing as a potential Diagnostic Method. *Ir. J. Med. Sci.*; 159: 4: 98100.
- Rolleston, S; Mncianti, F; Cioni, P.L; Flamini, G; Gorelli, I and Pers, C.P (1995) :** Selection of multiple resistance in salmonella enteritidis drug treatment with ciprofloxacin. *Scand. j. Infect. Dis.* 28: 529-531.
- Ross, S; Contronic, G and Khan, W (1972) :** Resistance of Shigella to ampicillin and other antibiotics. Its clinical and epidemiological applications. *JAMA.* 221 : 45-47.
- Ruther, S; Smith, B.K and Mall O.P (1998) :** In vitro antimicrobial efficacy of Thuja orientalis leaf extract *Fitoterapia* LXI (5); 423-455.
- Ryder, R.W; Blake, P.A; Murlin, A.C; Carter, G.P; Pollard, R.A; Merson, M.H; Allen, S.D and Brenner, D.J (1980) :** Increase in antibiotic resistance among isolated of Salmonella in the USA; 1967-1975. *J. Infect. Dis.* 142 (4) : 485-491.

- Sagdic, J (2003) :** American medicinal botany. Vol 1;2 and 3 Cummings and Hilliard Boston U.S.A in Carlson and Douglas (1999) Screening methods for determining antibiotic activity of higher plants J. Bact. 55: 235-240.
- Salamah, A and Zaid, A (2000) :** Antibacterial activity of *Commiphora quadricincta* from Saudi Arabia. J. King Saud Univ. 12 : 1-10.
- SamPsono, Hunponu-Wusu, D and Coker, A.O (1988) :** Bacterial and viral agents associated with diarrheal diseases in Lagos; Nigeria. In : Emejuiwe; D; Ogunbi O; Sanni; SO and editors Global impacts of applied bacteriology (6th international conference). London : Academic press; 427-434.
- Sauces, M.A (1997):** Urinary sepsis in bilharziasis; proceeding to 1st international symposium on bilharziasis part I; pp. 569-573.
- Schneierson, K.G (1981) :** Urinary pathogens and Bacterial sensitivity in Hospitalised urological patients based upon clinical aspects. Scand J Urol Nephrol Suppl; 104: 47-57.
- Shaprio, K.P; Koutsoumanis, K and Nychas, G.J (1994) :** A new Cophalos porin is incorporated into E. Coli Cells via the ton-B-dependent iron transport system. Antimicrobial Agents chemotherapy; 497-504.
- Shih, G.T; Javor and Lau, B.M (1999) :** *Allium sativum* (Garlic) inhibits Lipid synthesis by *Candida albicans*. Antimicrob. Agents chemother 30: 499-501.

- Sinha, J. and Gultai, Y.M (1990) :** Antimicrobial activity of essential oils from some species of poplar (Populus). Fitoterapia: Role of Biogeochemicals. Zhurnal Meditsiny i Biokhimicheskoy Meditsiny; 118-121.
- Sivropoulou, A; Kokkini, S; Lanaras, T and Arsenakis, M (1995) :** Antimicrobial activity of mint essential oils. Journal of Agriculture and Food Chemistry 43; 2384-2388.
- Smith, D.N and Dubos, N.H (1994):** Epidemiologic studies of Gram negative bacilli in the hospital and community. Amer. J. Epidemiol.; 85: 45.
- Smith-Palmer, A; Stewart, J and Fyfe, L (1998) :** Antimicrobial properties of plant essential oils and essences against five important food borne pathogens. Letters in Applied Microbiology; 26 (2) : 118-122.
- Sonnen, H; Minney, N; Mizumoto, H; Mitsui, K and Nonomura, H (1980) :** Bacteria Isolated from urinary tract infections and their susceptibility to New quinolones Hinyokika Kyo 38 (11) : 1215-23.
- Sonnen, S.B and With, N.W (1990):** The cause of death in diabetes is a report of 307 autopsied cases. New. S.J. 231: 865.
- Sonnenwirth, A.C (1980) :** Gradwohl's Clinical Laboratory Methods and Diagnosis Vol. 2; 8th ed. Sonnenwirth; AC and Jarett L. (eds). The CV Mosby Co. London; Toronto.
- Spika, J.S; Waterman, S.H; SooHoo, G.W; St. Louis, M.E; Pager, R.E; James, S.M; Bissett, M.L and Others (1987) :**

Cloramphenicol-resistant *Salmonella* newport traced through
Hamburger to dairy farms. *New Eng. J. Med.* 316 (10) : 565-570.

Stephani, S and Agodi, A (2000) : Molecular epidemiology of antibiotic
resistance. *Int. J. Antimicrob. Agents.* 13 : 143-153.

Strinager, P.J (1997): A comparison of the urinary white cell concentration
with the white cell excretion rate. *Br. J. Urol.*; 36: 360-363.

Sue, G.D; Ciaramonte, J.S and CaroFalo M.M (2001) : Effects of
Vapors of aromatic chemicals on fungi. *J. Amer. Pharm.
Assoc.*; 50 : 665-668.

Talbet, J.P and Levine, M.M (1988) : Bacterial diarrheas; p.697-752. In
AZ. Kapikian (ed); *Viral infection of gastromtestinal tract*; 2nd
ed. Marcel Dekker. Inc.; New York; NY.

Tassios, A.A; Essa, H; Abo-Elezz, F.A and Safun, M (2000): Evaluation
of different tests used for detection of urinary tract infection.
Med. J. of Cairo University XXX VIII; No. 4.

Tenover, F and McGowan, J (1996) : Reasons for the emergence of
antibiotic resistance. *Am. J. Med. Sci.* 311 : 9-16.

**Terakada, N; Azechi, H; Koyama, N; Sato, G and Mitsunashi, S
(1975)**: Demonstration of R factors with chloramphenicol
resistance in *Salmonella* strains isolated from domestic animals.
In *Microbial Drug Resistance*. Mitsunashi; S and Hashimoto H
(eds) University Park Press. Tokyo; London. PP. 253-260.

- Toama, M.A; El-Kersh, T.A and Ahmady, A.M (1983) :** Antibiotic-resistance pattern and resistance curing of E.coli locally isolated. Bulletin of Faculty of Pharmacy; Cairo University; Vol. XXII No. 1 : 129-146.
- Tong, M.J; Martin, M.J; Cunningham, J.I and Gunning, J.I (1970) :** Clinical bacteriological evaluation of antibiotics treatment in Shigellosis. JAMA. 214 : 1841-1844.
- Valkenburg, R.R (1993):** The significance of pyuria in the diagnosis of urinary tract infection. Br. M. J.; 2: 280.
- Vejlasaari, E.H (2000):** Significant and insignificant bacteriuria in the diagnosis of U.T.I. Arch Int. Med.; 100: 709.
- Vejlsgaard, R.H (1999):** Susceptibility of the kidney to experimental infection. In: O'Grady; F. and Brumfitt; W.; editors: urinary tract infection; London; Oxford; University Press; p. 24-36.
- Verman, R.R and Little, P.J (1996):** The significance of proteinuria in the diagnosis of urinary tract infection: The J. of Med. Lab. No.; 1: 336.
- Waffaa, Y.J; Pal, T; Rotimi, V.O and Chugh, T.D (1998) :** Serogroups and antimicrobial susceptibility of clinical isolates of Salmonella species from a teaching hospital in Kuwait. J Diarrheal Dis Res. 16 (3) : 180-186.
- Warren, M.T; Dorman, H.J; Deans, S.G and Ruberto, G (2000) :** Antimicrobial and antioxidant properties of some essential oils. Flav. Fragr J.13:235-244.

- Watanable, T (1971) :** The origin of R-factors. Annu. NY; Acad. Sci. 182 : 126-140.
- Weber, B and Rocha, H (1999):** Bacteriuria in apparently healthy young adults.; Dept. of Med. Hosp. Proi Edgard Santos; Univ of Bahia. Salvador. J. Uro. (Baltinore) 102/3: 342-344.
- Weiss, S.B; Parker, D and Kass, B.H (2000):** Bacteriuria in diabetes mellitus. Med.; 276: 1118.
- Well, B and Stood, R.J (1989) :** The genus shigella. In Bergey's Manual of systematic Bacteriology. Vol. I: 8th ed. Krieg; N. R: Hole; JG and others (eds) Williams and Wilkins. Hong Kong. Pp. 423-427.
- WHO (1980) :** World Health organization .Enteric infection due to Campylobacter; Yersinia; Salmonella and Shigella. Report of the scientific working group Bullein of Who. S8 : 519.
- Wilson, F; Match, J.A; Miller, B.R and Minman, F (1984):** A careful study. Relation of chronic pyelonephritis with diabetes. J. Urol.; 90: 96.
- Wilson, J.W (1994):** Non-obstructive escherchia coli pyelonephritis; in the rat. Proc. Sec. Exp. Biol. Med.; 104: 116-120.
- Wilson, S.C; Sorhati, I; Caspo, G and Balazs (1984):** The relationship between diabetes mellitus and infection of the urinary tract amer. J. Med. Sci.; 240: 48-141.