

## ACKNOWLEDGEMENT

The author would like to express her deep gratitude and sincere appreciation to Prof. Dr. *Hassan Kasam Bekhiet* Professor of Pesticides, Plant Protection Research Institute, for suggesting the topic of the study, direct supervision, guidance, and help during the presentation of this study and his beneficial advices.

The author is also indebted to Dr. *Talaat Ibrahim El Sayed* Assit. Prof. of Microbiology in Botany Dept, Faculty of Science Benha University, for his guidance, advising supporting, assistance and reviewing the manuscript in the final form.

The author is also indebted to Dr. *Mahmoud Moustafa Amer* Assit. Prof. of Microbiology in Botany Dept, Faculty of Science Benha University kind advice and continuous encouragement during the preparation of the manuscript.

The author is also indebted to Dr. *Salah Ali Moustafa* Assit. Prof. of Microbial Molecular Biology, Agricultural Genetic Engineering Research Institute, AGERI, Giza, Ministry of Agricultural, Giza, Egypt, for his guidance helpful suggestions, valuable assistance, and help in so many ways.

Thankful is extended to Dr. *Manal Abel-Kader El-Sawi*. Researcher in the central lab. of Residue Analysis of Pesticide in Ministry of A.R.C, for her helps in detection of aflatoxin using HPLC