

R E F E R E N C E S

Adams, A.R.D. (1952):

Cutaneous Amoebiasis. In: Handbook of Tropical Dermatology and Medical Mycology. Simons, R.D.G.P.(ed). vol.2.Elsevier, Amsterdam. P.402

Adolph, P.E., Kagan, I.G., and Mcquay, R.M.(1962):

Diagnosis and treatment of Acanthocheilonema Perstans filariasis. Am.J.Trop. Med. Hyg. 11:76.

Allain, D.S., and Kagan, I.G.(1961):

The use of formalinized red cells in the serology of hydatid disease.J. Parasitol.47.61.

Andreev, V.C., Angelov,N., and Zlatkov, N.E.(1969):

Skin manifestations in Toxoplasmosis. Arch.Dermatol. 100:196.

Apted, F.I.C.(1962):

Sleeping sickness in Tanganyika-Past, present and future. Trans.Roy.soc. Trop. Med. Hyg. 56:15

Arthur, R.P., and shelley, W.B.(1958):

Larva currens. Arch. Dermatol. 78:186.

Beal, C.B., Viens, P. and Grant, R. (1970):

Technique for sampling duodenal contents. Am.J. Trop. Med. Hyg. 19:349.

Beaver, P.C. (1949):

Methods of Pinworm diagnosis. Am. J. Trop. Med. Hyg.
29:577.

Beaver, P. C., Lepoz villegas, A., and Cuello, C. (1978):

Cutaneous amoebiasis of the eyelid with extension
into the orbit. Am. J. Trop. Med. Hyg. 27:1133.

Berberian, D. A. (1944):

Cutaneous leishmaniasis (oriental sore). 11.
Incubation period. Arch. Dermatol. 50:231.

Berline, C. (1941):

Leishmaniasis recidiva cutis; Leishmanid. Arch.
Dermatol. 41:874.

Berline, C. (1953):

Leshmanid: Report of a case. Brit. J. Dermatol.
65:265.

Biagi, F. (1969):

Cutaneous Amoebiasis. In : Essays on Tropical Derat-
ology. Simons, R.D.G.P., and Marshall, J. (eds).
Vol. I. Excerpta Medica. Amsterdam P. 206.

Biagi, F. F., and Martucelli, A. (1963):

Cutaneous amoebiasis in Mexico. Dermatol. Trop.
2:129.

Botero, D. and Castano, A. (1973):

Comparative study of pyrantel pamoate, oephenium hydroxynaphthoate, and tetrachlorethylene in the treatment of *Necator americanus* infection. Am. J. Trop. Med. Hyg. 22:45.

Bryceson, A. (1976):

Tropical dermatology: Cutaneous leishmaniasis. Brit. J. Dermatol. 94: 223.

Bunnag, T. (1984):

Gnathostomiasis. In: Hunter's Tropical Medicine Strickland, G. T. (ed). sixth ed. W.B.Saunders co. philadelphia. P.980.

Cahill, K. M. (1967):

Bancroftian Filarial nodule. Am. J. Trop. Med. Hyg. 16:636.

Chandrosoma, P. T., and Mendis, K.V. (1978):

Filarial infection of the breast. Am.J. Trop.Med. Hyg. 27:770.

Chaun, H. (1967):

The treatment of chronic strongyloidiasis with thiabendazole. Trans. Roy. Soc. Trop. Med. Hyg. 61:812

Convit, J. and Kerdel- Vegas, F. (1965):

Disseminated cutaneous leishmaniasis. Arch. Dermatol.
91:439.

Convit, J, Reyes, O. and Kerdel- Vegas, F. (1957):

Disseminated anergic cutaneous leishmaniasis. Arch.
Dermatol. 76:213.

Daengsvang, S. (1971):

Infectivity of *Gnathostoma spinigerum* larvae in
primates. J. parasitol. 57:476.

Daengsvang, S. (1980):

A Monograph on the Genus *Gnathostomiasis* in Thailand.
Tokyo, Southeast Asian Medical information center.
Quoted from Mckoy, K., and Moschellas, S.L. (1985):

Davey, D.G. (1958):

Human and animal trypanosomiasis in Africa. Am.J.
Trop. Med. Hyg. 7:549.

Dixon, H. B. F., and Hargreaves, W. H. (1944):

Cysticercosis (*Taenia Solium*). A further ten years
clinical study, covering 284 cases. Q. J. Med.13:107.

Domonkos, A.N.(1971):

Diseases due to animal parasites. In: Andrew's Diseases
of the Skin. 6th ed. W.B.Saunders Co. Philadelphia
P.474.

Dostrovsky, A. (1958):

Steroids as supplementary treatment in late leishmaniasis. A preliminary report. Brit. J. Dermatol. 70:288.

Dostrovsky, A., Sagher, F., and Zukermen, A.(1952):

Isophasic reaction following experimental superinfection of leishmania tropica. Arch. Dermatol.66:665.

Ebiling, F.J. (1979):

Introduction to Cutaneous Parasitology. In: Textbook of Dermatology. Rook,A., Wilkinson, D. S., and Ebiling, F.J.(eds). 3rd ed.Vol.I. Blackwell Scientific Publication. Oxford. P.869.

EL-Mofty, A.M. and Nada, M.M.(1967):

Cutaneous Schistosomiasis. In: Skin Diseases, 1st ed. The scientific Book Centre.Cairo P.260.

EL-Zawahry, M. (1963):

Bilharziasis Cutis. In: Skin Disease in Arabian Countries, Vol. I, Cairo, P. 245.

EL-Zawahry, M. (1964):

Schistosomal Granuloma of the skin. In: Skin Disease in Arabian countries, Vol. II, Cairo, P. 271.

EL-Zawahry, M. (1965):

Schistosomal granuloma of the skin. Brit. J.
Dermatol. 77:344.

EL-Zawarhy, M. (1969):

Schistosomal Granuloma of the skin. In:Essays on
Tropical Dermatology, Simons, R.D.G.P., and Marshall,
J.(eds). Vol. I.Excerpta Medica Amsterdam, P. 211.

EL-Zawahry, M., and EL-Komy, M (1973):

Amoebiasis Cutis. Int. J. Dermatol. 12:305.

Farah, F.S. (1979):

Protozoan and helminthic infections. In:Dermatology
in General Medicine. Fitzpatrick, T.B.; Eisen, E.Z.,
Wolf; K., Freedberg, I.M and Austein, K.F.(eds) 2nd
ed. McGraw-Hill book Co. New York. P. 1638.

Farah, F.S., and Malak, J.A. (1971):

Cutaneous leishmaniasis. Arch. Dermatol. 103:467.

Fasal, P. (1952):

American Leishmaniasis or leishmaniasis mucocutinea &
South American Trypanosomiasis (Chagas' Disease).
In:Hand book of Tropical Dermatology and Medical
Mycology. Simons, R.D.G.P. (ed). Vol. I. Elsevier.
Amsterdam. P.375 & P.394.

Faust, E.C. and Russell, P.F. (1964):

General introduction and orientation & Helminthic infections. In: Graig and Faust's Clinical Parasitology 7 th ed. Lea and Febiger, Philadelphia P.7 & P. 313.

Feldman, H.A. (1968):

Toxoplasmosis. N.Engl. J.Med. 2:475.

Findlay, G.H., and Whiting, D.A.(1971):

Disseminated and Zosteriform Cutaneous Schistosomiasis. Brit. J. Dermatol. 85:98.

Fistein, B., and Sutton, R. N. (1963):

Chaga's disease in the West Indies. Lancet. 1:330.

Fortado, A. (1959):

Clinical result in the treatment of American leishmaniasis with oral and intravenous amphotericin. Antibiot. Annu. P. 631. Quoted from Mckoy, K., and Moschella, S.L. (1985).

Galliard, H. (1953):

Sparganosis. In: Handbook of Tropical Dermatology and Medical Mycology. Simons, R.D.G.P. (ed). Vol. 2.Elsevier. Amsterdam. P.936.

Gelfand, M. (1966):

The early Clinical features of Rhodesian trypanosomiasis with special reference to the "Chancre" (local reaction). Trans. Roy. Soc. Trop. Med. Hyg. 60:376.

Girgla, H.S., Marsden, R. A., and Singh, G.M.(1977):
Post-Kala-azar dermal leishmaniasis. Brit. J. Dermatol. 97:307.

Goodwin, L. G. (1970):

The pathdogy of African trypanosomiasis. Trans. Roy. Soc. Trop. Med. Hyg. 64:797.

Gordon, R.M. (1955):

Symposium on Loiasis. Brief review of recent advances in our knowledge of loiasis and some of the still outstanding problems. Trans. Roy. Soc. Trop. Med. Hyg. 49:98.

Greenspan, G. and Stevens, L. (1961):

Infection with cysticercus cellulosa. N. Engl. J. Med. 264:751.

Griffiths, W.A.D., and Dutz, W. (1975):

Repeated tissue sampling with a dental broach, a trial in cutaneous leishmaniasis. Brit.J.Dermatol.93:43

Grove, D. I., and Davis, R.S. (1978):

Serological diagnosis of Bancroftian and Malayan filariasis. Am. J. Trop. Med. Hyg. 27:508.

Harman, R.R.M.(1977):

Tropical skin diseases in temperate climates. In: Recent Advances in Dermatdeg. Rook, A.(ed). No.4. Churchill Livingstone, Edinburgh, London, P.91.

Harman, R.R.M. (1979):

Parasitic Worms and Protozoa. In: Textbook of Dermatology. Rook, A., Wilkinson, D.S., and Ebling, F.J.(eds). 3rd ed. Vol. I. blackwell Scientific Publication. Oxford. P.873.

Hart, M., Livingood, C.S., and Goltz, R.W. (1969):

Late cutaneous leishmoniasis. Arch. Dermatol. 99:455.

Hawking, F. (1955):

Periodicity of microfilariae of Loa loa. Trans. Roy. Soc. Trop. Med. Hyg. 49:132.

Heiner, D.C., and Kevy, S.V. (1956):

Visceral larva migrans. N. Engl. J. Med. 254:629.

Hodgson, C., and Barrett, D.F. (1964):

Chronic dracunculosis. Brit. J. Dermatol. 76:211.

Kanani, S.R., and Rees, P.H. (1970):

The diagnosis of strongyloidiasis with special reference to the value of the filarial complement fixation test as a screening test. Trans. Roy. Soc. Trop. Med. Hyg. 64:246.

Katzenellenbogen, I. (1944):

Vaccination against oriental sore: report of results of five hundred and fifty-five inoculation. Arch. Dermatol. 50:239.

Katzenellenbogen, I. (1952):

Cutaneous Leishmaniasis. In: Handbook of Tropical Dermatology and Medical Mycology. Simons, R.D.G.P. (ed). Vol. I. Elsevier. Amsterdam. P. 336.

Katzenellenbogen, I. (1954):

Studies on dracunculosis among Yemenite immigrants to Israel, Dermatologica. 108:129.

Katz, A.M., and Pan, C.T. (1958):

Echinococcus disease in the United States. Am. J. Med. 25:759.

Kilpatrick, M.E., Trabalsi, B., and Farid, Z. (1981):

Levamisole compared to mependazole in the treatment of *Ancylostoma duodenale* in Egypt. Trans. Roy. Soc. Trop. Med. Hyg. 4:578.

Komiya, Y. and Kondo, S. (1952):

Trop. Dis Bull. 49:1058 Quoted from Moschella, S.L. (1975).

Kozenvniko, P.V. (1963):

Two nosological forms of cutaneous leishmaniasis. Am. J. Trop. Med. Hyg. 12:719.

Krick, J.A., and Remington, J.S. (1978):

Toxoplasmosis in adult, an overview. N. Engl. J. Med. 298:550.

Kurban, A.K., and Chaglassian, H.T. (1969):

Cutaneous Leishmaniasis. In: Essays on Tropical Dermatology. Simons, R.D.G.P., and Marshall, J. (eds). Vol. I. Excerpta Medica. Amsterdam. P. 187.

Kurban, A.K., Malak, J.A., and Farah, F.S. (1966):

Histopathology of cutaneous leishmaniasis Arch.
Dermatol. 93:396.

Lagerholm, B., Gip, L., and Lodin, A. (1968):

The histopathological and cytological diagnosis of
cutaneous leishmaniasis in two cases of leishmania
Tropica. Acta. Derm. Venereol. (stockh). 48:600

Lever, W.F., and Lever, S.G. (1983):

Diseases caused by protozoa. In: Histopathology of
the skin. 6th ed. J.B.Lippincott Co. Philadelphia.
P.356.

Majumder, T. D. (1969):

Post-Kala- azar dermal leishmaniasis. In: Essays on
Tropical Dermatology. Simons, R.D.G.P., and Marshall,
J. (eds). Vol. I. Excerpta Medica- Amsterdam. P.195.

Manson- Bahr, P.(1953):

Filarial Bancrofti. In: Handbook of Tropical Dermat-
ology and Mycology. Simons, R.D.G.P. (ed). Vol. 2.
Elsevier. Amsterdam. P.976.

Marsden, P.D. (1979):

Current concepts in Parasitology: Leishmaniasis. N.Engl.
J. Med. 300:350.

Marsden, P.D. (1984):

American Trypanosomiasis. In: Hunter's Tropical Medicine. Strickland, G.T. (ed). sixth ed. W.B. Saunders Co. Philadelphia P.698.

Maynard, J.E., and Kagan, I.G. (1964):

Intradermal test in the detection of Trichinosis. Further observation on Two outbreaks due to bear meat in Alaska. N. Engl.J.Med. 270:1

Mckoy, K.C. and Moschella, S.L. (1985):

Parasites, Arthropods, Hazardous Animals, and Tropical Dermatology. In: Dermatology. Moschella, S.L. and Huriely, H.J. (eds). 2nd ed. Vol. 2. W.B. Saunders Co. Philadelphia. P. 1731.

Montenegro, J. (1926):

Cutaneous reaction in Leishmaniasis. Arch. Dermatol. 13:187.

Moschella, S.L. (1975):

Parasitology and Tropical dermatology. In: Dermatology Moschella, S.L., Pillsbury, D.M., and Huriely. H.J. (eds). Vol.2. W.B.Saunders co. Philadelphia P.1487.

Most, H.(1965):

Trichinellosis in the United States: changing

epidemiology during past 25 years. J.A.M.A.
193:871.

Most, H. (1978):

Trichinosis- Preventable yet still with us. N.Engl.
J. Med. 298:1178.

Muller, R. (1984):

Dracunculosis. In: Hunter's Tropical Medicine Strickland,
G.T. (ed). sixth ed. W.B. Saunders co. Philadelphia.
p. 637.

Mulligan, H.W., and Potts, W.H. (1971):

The African Trypanosomiasis. Wiley- Interscience. New York.

'Quoted from McKoy, K., and Moschella, S.L. (1985):

Munro, D.D., Duvivier, A., and Jopling, W.H. (1972):

Post-Kala- azar dermal leishmaniasis. Brit. J. Dermatol.
87:374.

Nitidandhaprabhas, P., Sirimachan, S., and Charnvises,
K. (1978):

A case of penile Gnathostomiasis in Thailand. Am. J.
Trop. Med. Hyg. 27:1282.

Orihel, T.C., and Moore, P.J. (1975):

Loa loa: Experimental infection in two species of

African primates. Am. J. Trop. Med. Hyg. 24:606.

Osuntokun, O., and Olurin, O. (1975):

Filarial worm (Loa loa) in anterior chamber. Brit. J. ophthalmol. 59:166.

Padonu, K.O. (1973):

A controlled Treatment of dracunculosis in Nigeria. Am. J. Trop. Med. Hyg. 22:42.

Paz, E., Weinrauch, L., Livshin, R., El. on, J. and Greenblatt, C.L. (1982):

Rifampicin treatment of cutaneous leishmaniasis. Int. J. Dermatol. 12:112.

Pearson, R.D., and Guerrant, R.L. (1983):

Praziquantel: A major advance in antihelminthic therapy. Ann. Inter. Med. 99:195.

Pearson, R.D., Irons, R.P., Sr., and Iron, R.P.Jr. (1981):

Chronic Pelvic peritonitis due to the pinworm Enterobius vermicularis. J.A.M.A. 245:1340.

Perea, D., Western, K.A., and Schultz, M.G. (1970):

Niclosamid treatment of cestodiasis. Am. J. Trop. Med. Hyg. 19:610.

Pettit, J.H. (1962):

Chronic (lupoid) Leishmaniasis. Brit. J. Dermatol.
74:127.

Pierini, L.E. (1969):

South American Leishmaniasis. In: Essays on Tropical
Dermatology. Simons, R.D.G.P., and Marshall, J.
(eds). Vol. I. Excerpta Medica Amsterdam. P. 175.

Piers, F. (1953):

Loaiasis & Dracunculosis. In: Handbook of Tropical
Dermatology and Mycology. Simons, R.D.G.P. (ed).
Vol. 2. Elsevier. Amsterdam. P. 963. & P. 967.

Pillsbury, D.M., Shelly, W.B., and Kligman, A.T.(1966):

Dermatologic parasitology. In: Dermatology W.B.
Saunders Co., philadelphia. P 1028.

Price, S.M., and Silvers, D.N. (1977):

New world leishmaniasis. Arch. Dermatol. 113:1415.

Ramose e Silva, J., and Netto, M.P.(1973):

Leishmanids. Int. J. Dermatol. 12:104.

Ranque, J., and Quilici, M. (1970):

Recent advances immuno-diagnosis of leishmaniasis.
J. Parasitol. 56:277.

- Rau, R.C., Dubin, H.V., and Taylor, W.B. (1976):
Leishmania Tropica infection in traveller. Arch.
Dermatol. 112:197.
- Reddy, C.R.R.M., Reddy, M.M., and Sivaprasad, M.D.(1969):
Niridazole (Ambilhar) in the treatment of dracu-
nculosis. Am.J. Trop. Med. Hyg. 18:516.
- Rein, M.F., and chapel, T. A. (1975):
Trichomoniasis, Candidiasis and the minor venereal
disease. clin.obstet. Gynecol. 18:73.
- Remeington, J.S., Jacobs, L., and Kaufman, H.E. (1960):
Toxoplasmosis in adult. N. Engl. J. Med. 262:180.
- Rook, A., and Straughton, R. (1972):
The cutaneous manifestations of Toxocariasis.
Dermatologica. 144:129.
- Rydzewski, A.K., Chisolm, E.S., and Kagan, I.G. (1975):
Comparison of serologic tests for human cysticer-
cosis. J. Parasitol. 61:154.
- Sasa, M. (1976):
Human Filariasis. Baltimore, Univeristy Park Press.
Quoted from McKoy, K., and Moschella, S.L. (1985)

Schantz, P.M., and Glickman, L.T. (1978):

Toxocaral visceral larva migrans. N. Engl J. Med.
293:436.

Shaw, J.J., and Lainson, R. (1977):

A simply prepared amastigote leishmanial antigen
for use in indirect fluorescent antibody test for
leishmaniasis J. Parasitol. 63:384.

Simons, R.D.G.P. (1953):

Oxyuriasis. In: Handbook of Tropical Dermatology and
Medical Mycology. Vol. 2. Elsevier. Amsterdam. P.1007

Simpson, M.H., Mullins, J.F., and Ston, O.F. (1968):

Disseminated anergic cutaneous leishmaniasis Arch.
Dermatol, 97:301.

Snow, J.H., Satulsky, E.M., and Kean, B.H. (1948):

American cutaneous leishmaniasis. Arch. Dermatol.
Syphilol. 57:90.

Sondergaard, J. (1972):

Filariasis caused by Acanthocheilonema perstans.
Arch. Dermatol. 106:547.

Sparks, A.K., Neafie, R.C., and Connor, D.H. (1976):

Coenurosis. In: Pathology of Tropical and Extraordinary diseases. Binford, C.H., and Connor, D.H. (eds).
Armed Forces Institute of Pathology. Washington.
D.C. P.543. Quoted from Mckoy, K.C., and Mouschella
S.L. (1985).

Stamm, W.P., Ashley, M.D., and Bell, K. (1976):

Value of amoebic serologic tests in an area of low
endemicity. Trans. Roy. Soc. Trop. Med. Hyg. 70:49.

Stone, O.J. (1964):

Thiabendazole. A probable cure of trichinosis J.A.M.A.
187:556.

Ston, J.J. (1972):

Cutaneous strongyloidiasis: Larvae currens. Arch.
Dermatol. 106:734.

Strauss, W.G., and Manwaring, J.H. (1964):

Sparganosis. A new case in the United States and
review of the literature. Derm. Trop. 3:73.

Swartzwelder, J.C., Beavers, P.C., and Hood, M. (1964):

Sparganosis in Southern United States. Am. J. Trop.
Med. Hyg. 13:43.

Symmers, W.S.C., (1950):

Pathology of oxyuriasis, with Special reference to granuloma due to presence of oxyuris Vermicularis (Enterobios Vermicularis) and its ova in tissue. Arch. pathol. 50:475.

Templeton, A.C. (1968):

Human Coenurus infection. A report of 14 cases from Uganda. Trans. Roy. Soc. Trop. Med. Hyg. 62:251.

Torres, V.M. (1976):

Dermatologic manifestations of Schistosomiasis mansoni. Arch. Dermatol. 112:1539.

Tschen, E.H., Tschen, E.A., and Smith, E.B. (1981):

Cutaneous cysticercosis treated with metrifonate. Arch. Dermatol. 117:507.

Turner, L.H. (1959):

Studies on filariasis in Malaya. The Clinical features of filariasis due to Wuchereria malayi. Trans. Roy. Soc. Trop. Med. Hyg. 53:154.

Urcuyo, F.G., and Zaias, N. (1982):

Oral Ketoconazole in the Treatment of leishmaniasis. Int. J. Dermatol. 21:414.

Walton, B.C., Chinel, L.V., and Eguia y Eguia, O. (1973):

Onset of espundia after many years of occult
infection with *Leishmania braziliensis*. Am. J. Trop.
Med. Hyg. 22:696.

Webster, L.T. (1975):

Suppression of delayed hypersensitivity in schistosome
-infected Patients by niridazole. N.Engl.J. Med.
292:1144.

Wilcocks, C., and Manson- Bahr, P.E.C. (1978):

Disease caused by protozoa. In: Manson's Tropical
Diseases. 17th ed. Balliere Tindoll, London P.39.

Wiseman, R.A., and Woodruff, A.W. (1968):

Toxocariasis in Britain as revealed by skin sensit-
ivity test. Brit. Med. J. 1:677.

Wood, M.G., Srolovitz, H., and Schetman. (1976):

Schistosomiasis. Paraplegia and ectopic skin lesions
as admission symptoms. Arch. Dermatol. 112:690.

Woody, N.C. and Woody, H.B. (1961):

American Trypanosomiasis. J. Pediatr. 58:568.

Yesudian, P., and Thanbiah, A.S. (1974):

Amophnetericin B therapy in dermal leishmanoid.

Arch. Dermatol. 109:720.

Zuckerman, A., and Sagher, F. (1963):

Experimental cutaneous leishmaniasis. The development of multiple cutaneous lesions (leishmanid) following the prophylactic inoculation of living *Leishmania tropica* into a single site. J. Invest. Derm. 40:193.