




REFERENCES

- Abd El- Hady, S. (1996):* Clinical application of urine antigen detection in group B streptococcal neonatal sepsis. Thesis, submitted for partial fulfillment of M.Sc. degree in pediatrics. Benha Faculty of Medicine .
- Abd El-Maksoud, Y.H. (1997):* Some metabolic changes in neonatal sepsis. Thesis, submitted for partial fulfillment of M.D. degree in pediatrics. Benha Faculty of Medicine.
- Ahr, A.; Scharl A.; Goheing, U.J.; Neuhaw, W. and Koufman, M. (1997):* Blind prophylactic administration of antibiotics to mothers with premature rupture of fetal membranes. Zentrable. Gnaecol., 119(3): p111-116.
- Al-Mohamady, A.M. (2000):* A study of some cytokins in neonatal septicemia. Thesis, submitted for partial fulfillment of M.D. degree in pediatrics. Cairo Faculty of Medicine.
- Amara E.E. (1995):* A study of calcium homeostasis in septicemic neonates. Thesis, submitted for partial fulfillment of M.D. degree in pediatrics. Benha Faculty of Medicine.
- Ambrosino, D.M.; Delaney, N.R. and Shamberger, R.C., (1990):* Human polysaccharide specific B cells are responsive to IL-6 J. Immunol . 144 : 1221 –1226 .
- American College of obstetrics and Gynecology Committee Opinion (1996):* Prevention of early onset Group B strep. disease in newborns. Technical Bulletin No. 173 Washington, D.C. American Academy of obstetric and Gynecology.

- Andus , H.; Groll, Anja Meiser, Martina Weise (1987): IL-6 as early mediator in neonatal sepsis . Pediatric . Infect .Dis .J. 11, (6) , 496 – 498.
- Asano, S.; Okano, A. and Ozawa, K.(1990): In vivo effects of recombinant human IL-6 in neonates. Stimulated production of platelets.Blood75, 1602-1605.
- Ashitani, J. Mukae, H. Hirastuka, T. et al. (2002): Elevated levels of alpha – defensins in plasma and BAL fluid of patients with active pulmonary tuberculosis. Chest Feb; 12 (2) : 519 – 26.
- Ashitani, M.D.; Hiroshi, Mukne, M.D.; Haruko Taniguchi, M.D.; Kazuhiko et al., (1998): Elevated pleural Fluid levels of Defensins in Patients with Empyema. Chest. March; 113. 788 – 94.
- Aub-Farrag ,M.S.; Rewash, M.A.; Mahfouz, S.A. and El-Sheikh, M.M. (1996): Interleukin-6 in neonatal sepsis. The Gaz . Egypt. Pediatr. Assoc., 35:36.
- Baker, C.J. and Edwaras, M.S. (1995): Group B streptococcal infection. In : Text Book infectious disease of the fetus and newborn Infant 4th ed. Remington JS and Klein JO. WB Saunders, Philadelphia 981-1054.
- Bals, R.; Weiner, D. and Wilson, J. (2000): The innate immune system in cystic fibrosis lung disease. J. Clin. Invest., 103 : 303 – 307.

- Bals, R.; Weiner, D.; Moscioni, A.; Meegalla, R. and Wilson, J. (1999):** Augmentation of innate host defense by expression of a cathelicidin antimicrobial peptide. *Infect Immune* - 6089.
- Barkovich, A.J. and Lindan, C.E. (1994):** Congenital cytomegalo – virus infection of the brain imaging analysis and embryogenic consideration. *American Journal of Neuroradiology* 15: 703-715.
- Beagley, K.W.; Kiyono, H. and Kishimoto, T. (1989):** Interleukins and IgA synthesis. *J. Exp .Md* . 169: 2133-2148 .
- Becker, Mn.; Diamond, G.; Vergheze, M. W. and Randell, S.H. (2000):** CD14-dependent LPS- induced β - defensin-2 expression in human tracheobronchial epithelium. *J. Biol. Chem*, 275: 29731-29736.
- Behrman, R.; Kliegman, R. and Arvin, A. (1996) :** Infections of the neonatal infants. In : *Nelson textbook of Pediatrics* (15th ed.) W.B. Saunders Company, Philadelphia, P. 514 – 40.
- Behrman, R.; Kliegman, R. and Jenson, H. (2000):** Infections of the neonatal infants. In : *Nelson Textbook of Pediatrics* (16th ed.) W.B. Saunders Company, Philadelphia, P. 538-47.
- Bennet, R.M. (1976):** A solid-phase radioimmunoassay for the measurement of lactoferrin in human plasma variation with age, sex and disease. *J. LAB, clin mED*; 88: 156-66.

- Bennet, R.; Eviksson, M. and Zelterston, R. (1987):** Bacterial etiology of neonatal sepsis in relation to prior antibiotic treatment. *Acta. Pediatr.* 76:673.
- Bensch, K.; Raida, M.; Magert, H.J.; Schulz-Knappe, P. and Foreman, W.G (1995):** hBD-1 : a novel β -defensin from human plasma. *FEBS Lett*, 368:331-335.
- Berger, C.; Uehlinger, D.J.D.; Blau, N. and Fanciness (1995):** Comparison of protein and white blood cell count with differential in at risk for septicemia. *Eur. J. of pediatrics* 154:138-144.
- Berner, R.; Niemeyer, C.M.; Leititis, J.U.; Funke, A.; Schwab, C.; Rau, U.; Richter, K.; Tawfeek, M.S.; Clad, A. and Brandis, M. (1998):** Plasma levels and gene expression of granulocyte colony – stimulating factor, tumor necrosis factor – alpha, interleukin (IL) –1 beta, IL-6, IL-8 and soluble intercellular adhesion molecule –1 in neonatal early onset sepsis. *Pediatric Research*, Vol. 44, 469 – 477 by International Pediatric Research Foundation.
- Blanco, A.; Solis, G.; Arranz, E.; Coto, G.D.; Ramos, A.; Telleria, J. (1996):** Serum levels of CD14 in neonatal sepsis by Gram positive and Gram – negative bacteria. *Acta. Pediatr.*, 85 (6) : 728 – 32.
- Blom, N.A. and Vreede, W.B. (1993):** Infected cephalic hematoma associated with osteomyelitis, sepsis and meningitis. *Pediatr. Infect . Dis . J.* 12: 1015-20.

- Bortollussi, Burbridge, S. and Durnford, P. (1991):** Neonatal *Listeria monocytogenes* infection and gamma interferon. *pediatr. Res.* 29: P 234-239.
- Boyle, R.J.; Chandler, B.D. and Stonestreet, B.C. (1990):** Early identification of sepsis in infants with RDS. *Pediatr.* 62: 744-50
- Bragden, K.A.; Ackermann, M.R.; McCray, P.B.; JR and Huttner, K.M. (1999):** Differences in the concentrations of small anionic antimicrobial peptides in bronchoalveolar lavage fluid and in respiratory epithelia of patients with and without cystic fibrosis. *Inect Immune*, 67: 4256-4259.
- Brown, T.J.; Lie, J.; Shoyab, M. and Rowe, J.M. (1990):** Regulation of IL-6 expression by oncostatin. *M.J. Immunol.* 147 : 2175 –21800.
- Buck, C.; Gallati, M.; Pohlandt, F. and Barman, P. (1994):** Increase levels of tumor necrosis factor alpha (TNF- α) and interleukin -1 beta (IL-1 β) in tracheal aspirates of newborns with pneumonia infection. *Pediatrics*. 22: 238-241.
- Cairo, M.S.; Plunked, J.M.; Nguyen, A. and Van de Ven, C. (1991):** Sequential administration of IL-6 and G-CSF in newborn rats : Modulation of newborn granulopoiesis and thrombopoiesis. *Pediatr. Res*; 30 : 554 – 559 .
- Caltermole, H.E.J.; Rivers, R.P.A. (1987):** Neonatal candidal septicemia diagnosis on buffy smear. *Archives of Disease in Childhood*, 62:302-304.

- Campbell, M.J. and Machin, D. (1998):** Medical statistics A common sense . apparel . 2nd ed . John Wildy & sons New York, chisherter; Brisbon; Toronto & Singapore.
- Castell, GV.; Geiger, T.; Gross V. et al (1988):** Plasma clearance organ distribution and target cells of IL-6 / hepatocyte stimulating factor in the rat. Eur J, Biochem , 177 : 375-361 .
- Catherine, M.H.; Costarring, A.T.; Sullivan, J.S. and Dulkerian, S. (1994):** Cytokine elevations in critically ill infants with sepsis and nercotizing enterocolitis . J. Pediatr., 124 : 105-111 .
- Cebon, J.; Layton, J.E.; Maher, D. et al. (1994):** Endogenous haematopoieticgrowth factors in neutropenia and infection. Br. J. Haematol., 86 : 265 – 274.
- Centers for disease control (1996):** Prevention for neonatal Group B streptococcal disease: A pulic health perspective. MMWR 45(RR7) .
- Chan, D.K.; Ho, L.Y. (1997):** Usefulness of C-reactive protein in the diagnosis of neonatal sepsis. Singapore, Med. J., 38 (6) : 252 –5.
- Chen, I.; Mory , Y. and Revel, M. (1988):** Growth inhibition of human breast carcinoma and leukemia/ lymphoma cell lines by recombinant interferon B2.Proc. Natt . Acad Sci . U.S.A 85 , 8037-8041 .

- Chertov, O.; Michiel, D.F.; Xu, L.; Wang, J.M.; Tani, K.; Murphy, W.J.; Longo, D.L.; Taub, D.D. and Oppenheim, J.J. (1996):** Identification of defensin-1, defensin-2, and CAP37/ azurocidin as T-cell chemoattractant proteins released from interleukin-8- stimulated neutrophils. *J.Bio Chem* 217: 2935-2940.
- Christener, R.D. and Rethstein G. (1995):** Granulocyte transfusion in neonates with bacterial infection, neutropenia and depletion of mature marrow neutrophils. *Pediatr.*, 70 (1) : 1.
- Corrigan, J.J. (1984):** Thrombocytopenia. A laboratory sign of septicemia in infants and children. *J. Pediatr* , 105:219.
- Costa, A.; Guimaraes, H.; Soute, A.; Martin, S.A.; D'orey, C.; Mateus, M. and Santosn, T. (1996):** Sepsis in very low birth weight newborn infants. *Acta. Med. Port* . 9 (10-12): 331-334.
- Damas, P.; Ledaux, D.; DeGroote, D. and Lamy, M. (1992):** Cytokine serum level during sever sepsis in human IL-6 as marker of severity . *Ann. Surg.*; 215 (4) : 356-62.
- Daniel, W.W. (1987):** Biostatistics , A foundation for analysis in the health sciences . 4th ed. John Wiley sons new York; Chichester; Brisbane; Torouto Singapore.
- Dear, P. (1999):** Infection in Newborn. In:Textbook of Neonatology 3rd ed. (eds). By Jannet Rennie and NRC Robertson, Churchil Living Stone Edinburgh London New York Philadelphia Sydney Tornado. P. 1109-202.

- Debets, R.; Rao, S. and Rangnoker, J. (1989): Exchange transfusion in neonatal sepsis. Indian J. Pediatr., 28(1): 39.
- Denuk, M. and Daviad, S. (1993): Diagnostic value of leucocyte indices in neonatal infection. Trop. Geogr. Med., 55 (4) : 360.
- Di -Geronimo, R.J. (1992): Lack of efficacy of urine culture, as part of initial work-up of suspected neonatal sepsis. Pediatr. Infect. Dis. J. 11(9): 764-6.
- Diamond, G.; Zasloff, M.; Eck, H.; Brasseur, M.; Maloy, W.L. and Bevins, C.L. (1991): Tracheal antimicrobial peptide, a cysteine- rich peptide from mammalian tracheal mucosa: peptide isolation and cloning of a cDNA. Proc Natl Acad Sci USA 88 : 3952 - 3956.
- Dinareello, C.A. (1991): Interleukin-1 and interleukin - 1ra . Blood 77:1627-1652.
- Doellner, H.; Arntzen, K.J.; Haereid, P.E.; Aag, S.; Austgulen, R. (1998): Interleukin-6 concentrations in neonates evaluated for sepsis. J. Pediatr., 132 (2) : 295 – 9.
- Dollagramatic, H.D. and Tapratizitim, S. (1993): Comparison of 4 tests used in diagnosis of neonatal septicemia. Arch . Dis . Child., 69:429 .
- Dyke, M.P. and Forsythia, K.D. (1993): Decrease plasma fibronectin concentrations in preterm infants with systemic candidiasis. Journal of pediatrics and Child Health 29:298-301.

- Edwards, M.S.; Rench, M.A.; Hall, M.A. and Baker, C.J. (1993):** Fibronectin levels in premature infants with late – onset sepsis. *Journal of Perinatal* . 13:8-13.
- El Maraghi, S.; Refaat, M.; Eissa, S.; El Rashidi, F. and El Nemr, F. (1994):** Exchange transfusion in the management of neonatal sepsis. *The Egypt. J. Pediatr.* Vol. (11) No. (1-2).
- El-Alfy, M.S. and Awad, H.A. (1993):** The value of IVIG therapy in the management of septic pre-term neonates. *The Egyptian J. Pediatr.* 10(1-2) : 227-36.
- El-Gamal, Y.; Abu Louz S.; Saad EL-Din, K.; El-Shimi and Helmi, W. (1992):** Transplacental entero virus among Egyptian's newborns. *Proccding : The 15th international annual.AinShams.Medical Congress.Cairo Egypt, 21 – 24.*
- El-Gamal, Y.; El-Samahi. M.; El-Shimi, S.; El-Tahau, M. and Magdy, S. (1990):** Alpha-1 antitrypsin in neonatal infection . *Egypt. J. pediatr.* 7(1-2) 43-35.
- El-Maraghi, S.; El-Saifi, A. and Abou Alam, M. (1984):** Diagnosis of neonatal septicemia by gastric aspirate examination. *J. Egypt. Soc.* Vol X, No.1.
- El-Saadany, A.M.; Hassanien, B.M.; Youssef, M.S. and Aronus, A.M. (1992):** Screening tests for neonatal sepsis. *Medical. J. Cairo. Univ.,* 60 (4) : 1137.
- El-Sallab, S.M.; Abdel Aal, A.; El-Tonbary, Y. and El-Kappany, M.A. (1987):** Some aspects of neonatal septicemia of Mansoura Neonatal care unit. Thesis submitted in partial Fulfillment for the Master degree in pediatrics, P. 74 – 94.

- Engle, W.D.; Rosenfeld, C.R. and Mouzinho, A. (1997):**
Circulating neutrophils in septic preterm neonates :
Comparison of two reference ranges. *Pediatrics*, 99 : 10.
- Entman, M.I.; Bossen, R.D. and Anderson, D.C. (1991):**
Inflammation in the course of early myocardial ischemia.
FASEB. J. 5, 2529-2537.
- Ernst, R.K.; Yi, E.C.; Guo, L.; Lim, K.B.; Burns, J.L.; Hackett, M. and Miller, S.I. (1999):** Specific lipopolysaccharide found in cystic fibrosis airway pseudomonas aeruginosa. *Science* 1999, 286: 1561-1565.
- Farid, D.F.A. (1996):** A study of soluble interleukin-2 receptor as predictor of neonatal sepsis. Master thesis in pediatric Ain shams University .
- Farid, M. (1999):** A study of serum transforming growth factor B1 in neonatal sepsis. Master thesis in pediatric Ain shams University .
- Farid, M. (2000):** Procalcitonin as a marker of neonatal sepsis. Master thesis in pediatric Ain shams University
- Feigin, R. and Cherry, J. (1992):** Bacterial infection. Text Book of Pediatric Infectious disease 3rd ed (ed.) Remington JS , Klein JO, Philadelphia, WB, Saunders.
- Fetter, W.P.F.; Bearts, W. and Vonlingen, R.A. (1995):** Surfactant replacement therapy in neonates with respiratory failure due to bacterial sepsis. *Acta. Ped.*, 84 : 14 – 16.

- Figueroa-Damian, R.; Arredondo-Garcia, J.L.; Mancilla - Ramirez, J. (1999):*** Amniotic fluid interleukin-6 and the risk of early - onset sepsis among preterm infants. Arch. Med. Res., 30 (3) : 198 - 202.
- Findlay, R.; Taeusch, H. and Walther, F. (1996):*** Surfactant replacement therapy for meconium aspiration syndrome. Pediatr. 97(1): P 48-52.
- Frohm Nilsson, M.; Sandstedt, B.; Sorensen, O.; Weber, G.; Borregaard, N; & Stahle-Backdahi, M. (1999):*** The human cationic antimicrobial protein (hCAP 18), a peptide antibiotic, is widely expressed in human squamous epithelia and colocalizes with interleukin-6-. Infect Immun., 67: 2561-2566.
- Frohm, M.; Agerberth, B.; Shangri, G.; Stahle-Backdate, M.; Liden, S.; Wigzell, H. and Gudmudsson, G.H. (1997):*** The expression of the gene codin for the antibacterial peptide LL-37is induced in human keratinocytes during inflammatory disorders. J. Biol Chem, 272: 15258-15263.
- Ganz T. and Lehrer, R.I. (1999):*** Antibiotic peptides from higher eukaryotes: biology and applications. MoMed Today, 5292-7.
- Ganz, T.; Selsted, M.E.; Szklarek, D.; Harwig, S.S.; Daher, K.; Bainton, D.F. and Lehrer, R.I. (1985):*** Defensins. Natural peptide antibiotics of human neutrophils . J. Clin Invest 76: 1427-1435.

- Gendrel, D.; Assicot, M. and Raymond, J. (1996):** Procalcitonin as a marker for the early diagnosis of neonatal infections . Journal of pediatrics 128:570-573.
- Gibbs, R. S. and Duffy, P. (1991):** Progress in pathogenesis and management of clinical Intra-amniotic infection. J. Obstet. General . 164(5): 1317-1326.
- Ginver, L.B. and Backer, C.J. (1988):** The prevention and treatment of group B streptococci infection . Adv . Ped . Inf . Dis 3: 65-90.
- Givon, T.; Salvin, S.; Revel, M. et al. (1992):** Antitumor effects of human recombinant IL-6 on acute myeloid leukemia in mice and in cell cultures . Blood 79, 2392-2398.
- Goldmann, D. and Pier, G.I (1993):** Pathogenesis of infection related to intro vascular catheterization . Clin . Microbiol . Rev. P: 176.
- Goldmann, D.; Lecait, J. and Macroane, A. (1978):** Bacterial colonization of neonates admitted to intensive care environment. J . Pediatr . Dis . 449-459.
- Goldmann, M.J.; Anderson, G.M.; Stolzenberg, E.D.; Kari, U.P.; Zasloff, M. & Wilson, J.M. (1997):** Human β -defensin -1 is a salt -sensitive antibiotic in lung that is inactivated in cystic fibrosis. Cell, 88:553-560.
- Goldmann, S.; Ellis, R.; Dhar, A. and Carro, M.S. (1999):** Rational and potential use of cytokines in the prevention and treatment of neonatal sepsis. Clin . Perinatol 25 (3): 699-710.

- Gorter, A.D.; Eijk, P.P.; Van Wetering, S.; Hiemstra, P.S.; Dandert, J. & Van Alphen, L. (1998):** Stimulation of the adherence of *Haemophilus influenza* to human lung epithelial cells by antimicrobial neutrophil defensins. *J. Infect Dis.*, 178: 1067-1074.
- Gotoff, S.P. (1996):** Unique aspects of infection, infections of the neonatal infant : In Nelson textbook of pediatrics 15th ed. (eds), Behrman RE Kliegman RM, and Arvin Am, Philadelphia WB. Saunders: P514-20.
- Gottof, G.L. (2000):** Risk diagnosis of neonatal sepsis. *Pediatrics . Inf. Dis .* 12:205-5.
- Gottof, S.P. (2000):** Infections of the neonatal infant in : Nelson Text Book of pediatrics 16th ed. (eds), Behrman RE, Kliegman RM, and Jenson HB, WB saunders, Philadelphia P 538-551.
- Gottoff, S.P. (1992):** Infectious Diseases of children 9th ed (eds), Krugman S, Katz SL, Gershon SL and Wilfert CM, The cv Mosby company, Saint Luis . P 402.
- Graves, G.G. and Rhodes, P.G. (1984):** Tachycardia as a sign of early -onset neonatal sepsis. *Pediatric Infectious Disease Journal* 3:404-406.
- Gray, A.E. and Steven, A.R. (1998):** Common Neonatal Procedures. In: *Manual of Neonatal care* 4th ed. (eds) Lippcott. Raven, Philadelphia- New York : P 651-675.
- Gregory, J. and Hey, P. (1972):** Neutrophil response to bacterial infection in the first month of life. *Archives of Disease in childhood* 47: 747-753.

- Grio, R.; Porpiglia, M. and Vetro, E. (1994):** Asymptomatic bacteria in pregnancy, maternal and fetal complications. *Panminerava Medecine* 36:193-200.
- Groll, A.H.; Meiser, A.; Weise M. et . al. (1992) :** IL-6 as early mediator in neonatal sepsis. *Pediar . Infect. Dis . J.*, 11:496-8.
- Grossman, R.M.; Krueger, J.; Yourish, D. et al. (1989) :** IL-6 is expressed in high level in psoriasis skin and stimulates proliferation of cultured human kratinocytes. *Proc . Natl. Acad . Sci . USA*, 86: 6367- 6371 .
- Guerina NG (1999):** Bacterial and Fungal infection. In *Manual of neonatal Intensive Care* 5th ed. (eds), cloherty JP and Start AR. Lipinn cott Raven, Philadelphia, 271-300.
- Guillermín, R.; Vargo, T.M. and Rossier, J. (1977):** Beta – endorphin and adrenal corticotrophin and secreted concomitantly by the pituitary gland . *Science*; 197: 1367-70 .
- Guillois, B.; Berthou, C. Awad, H. and Youinou, P. (1989):** The importance of C3d estimation in the diagnosis of generalized bacterial infections in newborn infants *Acta Pediatr.* 78: 368-372.
- Guillois, B.; Donnou, M.D.; Sizun, J.; Bendaoud, B. and Youinou, C. (1994):** Comparative study four tests of bacterial infection in the neonate. Total neutrophil count, CRP, fibrinogen, and C3d. *Biologcy of the neonate* 66:175-181.

- Hachey, W.E. and Wiswell, T.E. (1992):** Limitations in the usefulness of urine latex particle agglutination tests and haematologic measurements in diagnosing neonatal sepsis during the first week of life . Journal of perinatology 12:240-245.
- Hague, R.A.; Rassams and Morgan G 1994:** Early diagnosis of severe combined immunodeficiency syndrome. Archives of disease in childhood 70:260-263.
- Hama, T.; Miyamoto, M.; Tsukui, H. et al. (1989):** IL-6 in clinical medicine. Ann. Hematol. 62, 203 – 210.
- Haque, K.H. (1992):** Infection and immunity in the newborn . Textbook of pediater. P.302-316 .
- Haquew, K.N. ; Remo, C; Bahakin, H. (1995):** Comparison of two types of intra venous immunoglobulins in the treatment of neonatal sepsis. Clinical experimental immunology . 101: 328-232.
- Harder, J.; Gbartels, J.; Christophers, E. & Schroeder, J.M. (1997):** A peptide antibiotic from human skin. Nature 387: 861.
- Harder, J.; Meyer-Hoffert, U.; Teran, L.M.; Schwichtenberg, L.; Bartels, Maune, S. and Schroder, J.M. (2000):** Mucoid pseudomonas aeruginosa, TNF α , and IL-1 β , but not IL-6 induce human β - defensin -2 in respiratory epithelia. Am. J. Respir. Cell Mol. Biol. 22:714 –721.

- Harris, M.C.; Costavino, A.T. and Sullivan, T.S. (1994):* Cytosine elevations in critically ill infants with sepsis and narcotizing enterocolitis. *Journal of pediatrics* 124:105-111.
- Helgott, D.C.; May, L.T. and Sthoeger, Z. (1987):* Bacterial lipopolysaccharide (endotoxin) enhances expression & secretion of B₂ interferon by human fibroblasts. *J. Exp. Med.*, 166:1300.
- Hickey, S.M. and McCracken, G. (1997):* Postnatal bacterial infections. In : *Neonatal perinatal Medicine. Disease of the fetus and infants 6th ed* (eds), Fanaroff AA and Mortin RJ, Mosby Philadelphia: P717-59.
- Hill, H.R. (1997):* Immunomodulation by breast-feeding. *Pediatric infect Dis J.* 16:991-5.
- Hill, R.J.; Warren, M.K. and Levin, J. et al. (1992):* Evidence that IL-6 does not play a role in the stimulation of platelet production after induction of acute thrombocytopenia. *Blood* 80:346-351.
- Hindocha, P.; Campbell, C.A. and Gould, J.D. (1984):* Serial study of C-reactive protein in neonatal septicem. *Arch. Dis. Child*, 59 : 435.
- Hirano, T. and Matsuda, T. (1986):* Complementary DNA for human interleukin that induces B-1 lymphocytes to produce immunoglobulin. *Nature (London)* 324, 73-76.

- Hiratsuka, T, Nakazato, M.; Date, Y.; Ashitani, J.; Minematsu, T.; Chino, N. and Matsukura, S. (1998):** Identification of human β -defensin -2 in respiratory tract and plasma and its increase in bacterial pneumonia. *Biochem Biophys Res Commun*, 249:943-947.
- Hoffmann, D.J. and Harris, M.C. (1996):** Diagnosis of neonatal sepsis. In : *Intensive care of the Fetus and Neonate* . 1st ed. (eds), Spitzer AR, Msby Philadelphia : 940-950.
- Hoover, D.M.; Rajashankar, K.R.; Blumenthal, R.; Puri, A.; Oppenheim, J.J.; chertov, O. and Lubkowski, J. (2000):** The structre of human β -defensin-2 shows evidance of higher order oligomerization . *J Biol Chem.*, 275: 32911-32918.
- Hosfard, D. and Braquet, P. (1990):** The potential role of platelet-activating factor in shock and ischemia . *J. Crit Care* ; 5: 115.
- Houssian, F.A.; Olive, D.;Van Snick , J. et al. (1988):** Synergetic activation of human T-cell by IL-6 and IL-1 *Eur. J. Immunol.* 18:653-656 .
- Ikebuchi, K.; Wong, G.G.; Clark, S. et al. (1987):** IL-6 enhancement of IL-3 dependent proliferation of multipotential hemopoietic progenitors. *Proc. Natl. Acd. Sci. U.S.A.* 84: 9035-9039 .
- Isaacs, D. and Moxon, E.R. (1996):** Etiology of neonatal sepsis . In : *Neonatal Infections*. 2nd ed (eds). Oxford University press, Oxoford, . 5-6.

- Ischimi, Y.; Akatsu, T. and Yamguchi, A. (1990):** IL-6 is produced by osteoblasts and induces bone resumption . J. Immunol. 145:3297-3303.
- Ishibashi, T.; Kimura, H. and Shikawa, Y. (1989):** IL-6 is a potent thrombopoietic factor in vivo in mice. Blood 74:1241-1244.
- Jackson, S.K. (1994):** Endotoxin and septic shock. The antibiotic connection Bayer U.K.
- Jay, D.L. and Waits, K.B. (1997) :** Clinical application of C-reactive protein in pediatrics. Pediatr. Infect J. 16 (8) : 736 – 46.
- Jeffery, H.; Mitchison, R. and Wigglesworth, J.S. (1997):** Early neonatal bacteraemia. Archives of Disease in childhood 52: 683-686.
- Jeffry, S.G. (1991):** Clinicopathological approach to the diagnosis of neonatal sepsis. Clin. Perinatol . 18(2): 361-80.
- Jenson, H.B. and Pollock, B.H. (1998):** The role of intravenous immunoglobulin for the prevention and treatment of neonatal sepsis. Semin. Perinatal., 22 (1) : 50 – 63.
- Jerone. D.O.K. and Micheal, M. (1995):** Bacterial sepsis and meningitis In :Text Book of Infectious Diseases of the Fetus and Newborn infants. 4th ed. (eds), Lacks Remington, Jerom O, Klein, WB saunder's company philadelphia London Tornoto sydney, Tokyo. PP. 836-879.

- Johanson, R.; Spencer, S.; Roif, P.; Janes, P. and Malla, D. (1992):** Effect of post delivery care on neonatal body temperature. *Acta. Pediatr.* 81: 859-863.
- Jourdan, M.; Battaille, B.; Segnin, J. et al. (1990):** Constitutive production of IL-6 and immunology features in cardiac myxomas. *Arthritis rheum.* 33: 398-402.
- Kenom, C.; Overturf, G. and Bessman, S. (1996):** Granulocyte colony stimulating factors a marker for bacterial infection in neonates. *Journal of pediatrics* 128: 765-769.
- Khazbak S.M.A. (2001):** A study of serum endotoxins in neonatal sepsis. Master thesis in pediatric Ain shams University
- Kirnbaur, R.; Kock, A. forster, E. et al. (1991):** Regulation of epidermal cell IL-6 production by UV light and corticosteroid. *J. Invest. Dermatol.* 96:484-489.
- Kite, P.; Millar, M.R.; Gorham, P. and Condone, P. (1988):** Comparison of five used testes in the diagnosis of neonatal bacteraemia. *Archives of Disease in childhood* 36-639-643.
- Klein, J.O. and Marcy, S.M. (1995):** Bacterial sepsis and meningitis. In : *Text Book Infectious Disease of the fetus and Newborn Infants*. 4th. ed. (eds), Remington JJS and Klein JO : WB Saunders, Philadelphia. P 835-890.
- Klein, J.O. and Ramington, J.S. (1995):** Current concept of infection of the fetus and newborn infants. In : *Text Book of Infectious Diseases of the fetus and newborn infant* by 4th ed (eds), Klein JO and Remington JS, Philadelphia WB. Saunders. P 1-19.

- Kliegman, M.B.; Reynold, J.K. and Shreiner, R.L. (1986):** Rapid diagnosis of neonatal bacteria with acridine orange – stained buffy coat smears. *J. Pediatr*; 125:419 .
- Kliegman, R.; Calpp, D. and Berger, M. (1998):** Targeted immunoglobulin therapy for the prevention of neonatal infections . *Rev. Infect . Dis* 12: P 443.
- Kocak, U.; Ezer, U. and Vidinlisan, S. (1997):** Serum fibronectin in neonatal sepsis : is it valuable in early diagnosis and outcome prediction ? *Acta. Pediatr. Jpn. (Australia)* 39 *4) : 428 – 32.
- Kogan, B.L.; Ganz, T. and Lehrer, R.I. (1994):** Defensins, a family of antimicrobial and cytotoxic peptides. *Toxicology*, 87:131-49.
- Koj, A. (1985):** The acute-phase response to injury and infection (A.H. Gordon and A.Kop, Vol. 9, 10:P.139, Amstradam. New York.
- Kopp, E. and Medzhitov, R. (1999):** The Toll-receptor family and control of innate immunity. *Curr Opin Immunol.*, 11:13-18.
- Krause, P.J.; Maderaso, E.G. and Contrine, J. (1991):** Modulation of neonatal neutrophil function by pentoxifylline. *Pediatric* 29: 123-127.
- Kushner, I. (1982):** The phenomenon of acute phase response. *Ann. NY Acad . Sci.*; 389: 39-48 .

- Kuster, H.; Weiss, M.; Willeinter, A.E.; Detlefsen, S.; Jeremias, I.; Zbojan, J.; Geiger, R.; Lipowsky, G.; Simbruner, G. (1998):** Interleukin-1 receptor antagonist and interleukin-6 for early diagnosis of neonatal sepsis 2 days before clinical manifestation. *Lancet (England)* 352 (9136) : 1271-7.
- Lacy, J.B. and Ohlsson, A. (1995):** Administration of intravenous immunoglobulin for prophylaxis or treatment of infections in preterm infants. *Arch. Dis . Child .* 72: 151-155.
- Lagrutta, E.F.; Drusin, I.M. and Mackles, A.W. et. Al. (1993):** Neonatal infection . *An J.Dis . child*, 137:838-41
- Lau, Y.L. and Hey, E. (1991):** Sensitivity and Specificity of daily tracheal aspirate cultures in preceding organisms causing bacteraemia in ventillated infants . *Pediatr. Infect. Dis. J.* 10: P 290.
- Lauterbach, R.; Pawlik, D.; Kowalczyk, D.; Ksyc'inski, W.; Helwich, E. and Zembala, M. (1999):** Effect of the immunomodulating agent, pentoxifylline, in the treatment of sepsis in prematurely delivered infants : a placebo – controlled, double – blind trial (see comments). *Crit. Care Med.*, 27 (4) : 807 – 14.
- Lehrer, R.; Ganz, T. and Selsted, M. (1991):** Defesins, endogenous antibiotic peptides of animal cells. *Cell*, 64:229-203 .
- Lehrer, R.I.; Szklarek, D.; Ganz T. and Selsted, M.E. (1985):** Correlation of binding of rabbit granulocyte peptides to candida albicans with candidacidal activity. *Infect Immun* 1985; 49:207-11.

- Lehrnbecher, T.; Schord, T.; Krous, D. and Troos, Martius, J. (1995):** Interleukin-6 and soluble interleukin -6 receptor in cord blood in the diagnosis of early onset sepsis in neonates. *Acta. Pediatr.*, 84:806-8 .
- Lemaitre, B.; Reichhart, J.M. & Hoffmann, J.A. (1997):** Drosophila host defense, differential induction of antimicrobial peptide genes after infection by various classes of microorganisms . *Proc Natl Acad Sci USA*, 94: 14614-14619 .
- Lencer, W.; Cheung, G.; Strohmeier, G.; Currie, M.; Ouellette, A.; Selsted, M.; & Madara, J. (1997):** Induction of epithelial chloride secretory by channel forming cryptodins 2 and 3. *Proc Natl Acad Sci USA*, 94: 8585-8589.
- Linzmeier, R.; Ho, C.H.; Hoang, B.V. and Ganz, T. (1999):** A450 – kb contig of defense genes on human chromosome 8P23. *Gene*. 211.
- Lipursky, A. and Jaber, H. (1978):** The Hematology of bacterial infection in newborn infants. *Clin. Haematol.*, 7:175.
- Lucas, M.J. and Cunningham, F.G. (1993):** Urinary tract infection in pregnancy. *Clin. Obstet. Gynecol.* 36: 855-867.
- Ludtke, S.J.; He, K.; Heller, W.T.; Harroun, T.A.; Yang, L. and Huang, H.W. (1996):** Membrane pores induced by magainin. *Biochemistry*, 35: 13723-13728 .
- Lysenko, E.S.; Gould, J.; Bals, R.; Wilson, J.M. and Weiser, J.N. (2000):** Bacterial phosphorylcholine decreases susceptibility to the antimicrobial peptide LL-37/h CAP18 expressed in the upper respiratory tract. *Infect Immun*, 68: 1664 – 1671.

- Maisels, M.T. and Kring, E. (1992):** Risk of sepsis in newborn with severe hyperbilirubinaemia. *Pediatr.* 90: 741.
- Malech, H.L. and Gallin, J.I. (1987):** Neutrophils in human diseases. *N Engl Med*; 317: 687-49.
- Manro, B.L.; Weinberg, A.G.; Rosenfeld, C.R. and Browre, R. (1979):** The neonatal blood count in health and disease. *J. Pediatr.*, 95:89.
- Manrocet, B.L.; Weinberg, A.G.; Brown, R. and Rosenfield, .CR. (1979):** The neonatal blood count in health and disease. Reference values for neutrophilic cells. *Journal of pediatrics* 95: 89-98.
- Martens, J.; Samson, W.P.F. and Fetter, A. (1999) :** TNF- α , IL-6 and IL-1B plasma level in neonatal sepsis. *Pediatr .Res .* 33(4) : 380-383.
- Maruo, N.; Morita, I. And Matota, S. (1992):** IL-6 -increases endothelial permeability in vitro . *J. Indocrinology* 134:710-714.
- Mason, W.H.; Andrew, R.L. and Wright, H.T. (1989):** Omphalitis in the newborn infant. *Pediatr. Infect . Dis J.* 8: 521-525.
- Mathur, N.B.; Kalil, A. and Sarker, R. (1991):** Mortality in neonatal sepsis with involvement of mother in management . *Indian . Pediatr.* 28: 1259-63.
- Matsuda, Y.; Maruyama, H. and Kuraya, K. (1995):** Relationship between granulocyte elevates levels and perinatal infections. *Gynecologic and obestetric In vestigation* 39: 162-166.

- McCue, J. (1989):** Changes in therapeutic protein caused by preparation technique. *Ann. Med.* 3: P 271-272.
- McIntosh, N.(1992):** The Newborn. In : Chanphell AGM and McIntosh, N and Arneil's Textbook of pediatrics, 4th ed. WB Saunders. Company, Philadelphia, London, Toronto, P.109.
- Medzhitov, R. and Janeway, C.J. Jr (2000):**Advances in immunology: innate immunity. *N Engl. J. Med.* 343:338-344.
- Miki, S.; Yamamoto, M.Sonoda, T.et al. (1989):** IL-6 functions as in vitro autocrine growth factor in renal cell carcinoma. *FEBS Lett.* 250:607-891.
- Miles, S.A.; Stevens, R.H.; Longan, D.M. et al. (1990) :** Kaposi Sarcoma derived cells produce and respond to IL-6 *Proc . Natt . Acd. Sci. U.S.A.* 87: 4068-4072.
- Miller, L.C.; Isa, S.; Loprestd G. et. al. (1990):** Neonatal interleukin- 1B , interlukin -6 and tumour necrosis factor: Cord blood levels and Cellular production . *J. Pediatr.* 22:281-285
- Mock, B.A.; Kozak, C. and Clark, S.C. (1989):** The murine IL-6 gene maps to the proximal region of chromosomes. *J. Immunol.* 142: 1372-1376.
- Molyneux, R.; Schleiffenbaum, B. and Fehr, J. (1991) :** IL-1 and TNF stimulate human vascular endothelial cells to promote transendothelial neutrophil passage . *J.Clin. Invest* , 83:444.

- Morena, Z.A.; Naqui, S.H. and Muzaffar, T. (1994):** Neonatal septicemia in Pakistan : Presentation and Pathogens. Acta. Pediatr. Scand; 80 : 596 – 601.
- Morinaga, Y.; Suzuki, H. and Takasuki, F. (1989) :** Contribution of IL-6 to the anti-proliferative effect of IL-1 and TNF- α on the tumor cell lines . J. Immunol. 143:4538-4542.
- Mustafa, M.M.; Lebel, M.H. and Ramilo, O. (1989):** Correlation of IL-1B and TNF concentrations in CSF and outcome from bacterial meningitis. J. Pediatr; 115 : 208 –13 .
- Naeye, R.L. and Blanc, W.A. (1970):** Relation of poverty and race to antenatal infection. N Eug J. Med , 283-555.
- Naitoh, Y.; Tominaga, T.; Imura, H. et al. (1998):** IL-6 stimulates the secretion of adrenocorticotrophic hormone in conscious, freely moving rats. Bioch. Biophys . Res Commun. 155:1459-1463.
- Nijsten, N.w.N.; DeGroot, R.; Ten Duis, H.J. et al. (1987):** Serum levels of IL-6 and acute phase responses. Lancet 2: 921 .
- Nishimoto, N.; Yoshizaki, K.; Tagoh, M. et al. (1989):** Elevation of serum IL-6 prior to acute phase proteins on the inflammatory by surgical conditions. Clin. Immunopath. 50:399-401.
- Nuijens, J.N.; Abbink, J.J.; Wachtfogel, Y.T. et al. (1992):** Plasma elevates antitrypsin and lactoferrin in sepsis : evidence for neutrophils as mediators in fatal sepsis. J LAB CLN MED; 119: 159-68.

- O'Garra, A. (1989):* Interleukins and the immune system, *Lancet* 2 : 943 – 946.
- O'Neil, D.A.; Porter, E.M.; Elewaut, D.; Anderson, G.M.; Eckmann, L.; Ganz, T. and Kagnoff, M.F. (1999):* Expression and regulation of the human β – defensins hBD-1 and hBD-2 in intestinal epithelium. *J Immunol*, 163:6718-6724.
- Offer levy, sic Eichenwald, Tomas Ganz et al.(1999):* Impaired Innate immune in the newborn pediatrics 104; 1327-32.
- Ohlsson, A. (1989):* Treatment of premature rupture of membranes. A meta-analysis . *A.M.J. Obstet. Gynecol.*, 160:P 890-906.
- Okusawa, S.; Gelfand, J.A. and Ikejima, T. (1988):* IL-1 induces a shock – like state in rabbits synergism with TNF and the effect of cyclooxygenase inhibitors. *J.Clin . Invest.*, 81:1162-72.
- Opal, S.M.; Fisher, G.J.; and Dhainaut, J.F. (1997):* Confirmatory IL- Ira trial in severe sepsis *Crit. Care Med.* 25(7): 1115-24.
- Panyutich, A.V.; Panyutich, E.A.; Krapivin, V.A. et al. (1993):* Plasma defensin concentrations are elevated in patients with septciemia or bacterial meningitis. *J. lab Clin. Med.* 122: 202-07.
- Panyutich, A.V.; Votinok, N.N.; Lehrer, R.I.; Ganz T. et al. (1991):* An enzyme immunoassay for human defenses . *J Immunol Methods*; 141:149-55.

- Pena, C.; Pujol, M.; Ardanuy, C. et al. (1998):*** Epidemiology and successful control of a large outbreak due to *Klebsiella pneumoniae* producing extended spectrum beta – lactamases. *Antimicrob. Agents Chemother.*, 42 : 53 – 8.
- Pereira, H.A.; Martin, L.E.; Spitznagel, J.K.(1989):*** Quantitation of a cationic antimicrobial granule protein of human polymorphonuclear leukocytes by ELISA. *J Immunol Methods*; 117: 115-20
- Peschel, A.; Otto, M.; Jack , R.; Kalbacher, H.; Jung, G. and Gotz, F. (1999):*** Inactivation of the *dlt* operon in *Staphylococcus aureus* confers sensitivity to defensins, protegrins, and other antimicrobial peptides . *J Biol Chem*, 274: 8405 –8410.
- Peterec, S.M.; Brennan, S.A. and Rinder, H.M. (1996):*** Reticulated platelet value in normal and thrombocytopenic neonates. *J. Pediatr.* 129 – 274.
- Pincemail, J.; Deby-Dupont, G.; Deby, C. et al. (1991):*** Fast double antibody radioimmunoassay of human granulocyte myeloperoxidase and its application to plasma. *J Immunol Methods*; 137:181-91.
- Pizzo, P.A. (1992):*** Modulation of host defenses by cytokines evolving adjuncts in prevention and treatment of serious infections in immunocompromised host. *Clin. Infect. Dis.*, 15:508-524.

- Ralph, D.; Feigin, M.D.; George, H.; McCracken, J.R.M.D. and Jerome, M.D. (1992):** Diagnosis and management of meningitis. *J. Pediatr. Infect. Dis.*, 11 (9) : 758 – 813.
- Roberton, N.R. (1992):** Neonatal infection. *Ped. Infect. Dis. J.*, 13 : 65
- Rodriguez, A.F.; Kaplan, S.L. and Mason, E.O. (1990):** Cerebrospinal fluid values in the very low birth weight infants. *J. Pediatr.* 116:971.
- Rodweel, R.; Lesil, A. and Tudeinope, D. (1988):** Early diagnosis of neonatal sepsis heamatological scoring system. *J. Pediatr.* 112:761-767.
- Rodwell, R.L.; Tudehope, D.I. and Gray, P.H. (1993):** Hematologic scoring system in early diagnosis of sepsis in netropenic newborns. *Pediatr. Infect. Dis. J.*, 12 : 372-376.
- Roine, I.; Fonces, L.M. and Cofre, J. (1992):** C-reactive protein, tumor necrosis factor α and interleukin – 1 beta of the CSF cell count. *Pediatr Infect. Dis. J.* 11: P1057.
- Rooney, J.C.; Hill, D.J. and Danks, D.M. (1971):** Jaundice associated with bacterial infection in newborn . *American Journal of Diseases of children* 122:39-41.
- Rouse, D.J. (1994):** Strategies for the prevention of early –onset neonatal group B streptococcal sepsis: A decision analysis. *Obstet. Gynecol* 83:483.

- Rusconi, F.; Parizzi, F.; Garlaschi, et al. (1991)** : IL-6 activity in infants and children with bacterial meningitis. *Pediatr. Infect. Dis. J.*; 10:117-21 .
- Saez-Liorens, X. and McCracken, G.H. (1993)**: Sepsis syndrome and septic shock in pediatrics: current concepts of terminology, pathophysiology, and management. *J. Pediatr.* 123:497-508.
- Samuel, P.G. (1992)**: Sepsis in the newborn. In : Krugman S, Kat SL, Gershon SI and Wilfert CM. (eds). *Textbook of infectious Disease of children*, 9th ed. The CV & Mosby Company, Saint Louis, P.402.
- Schieve, L.A.; Handler, A. and Hershow, R. (1994)**: Urinary tract infection during pregnancy and its association with maternal morbidity and perinatal outcome. *American Journal of public Health* 34: 405-410.
- Schuchat, A.; Cowworker, B.; Oxtoby, M. and Cochi, S. (1990)**: Population based risk factors for neonatal group B streptococcal disease (results of cohort study in Metropolitan Atlanta) . *J. Inf . Dis* 162: 672.
- Selsted, M.E.; Harwig, S.S.; Ganz, T.; Schilling, J.W. and Lehrer, R.I. (1985)**: Primary structures of three human neutrophil defensins. *J. Clin Invest* 1985, 76: 1436-1439.
- Shackelford, P.G. (1997)**: Infection of the fetus and neonates. In : *Principle and practice of pediatric infection diseases* (ed. Long, SS; Pickering, LK and Prober, CC) 1st edition, Section N; Page 596, Chirchill Livingstone, New York, Edinburgh, London, Madrid, Melbourne San Francisco, and Tokyo.

- Shafer, W.M.; Qu, X.; Waring, A.J. and Lehrer, R.i (1998):** Modulation of *Neisseria gonorrhoeae* susceptibility to vertebrate antibacterial peptides due to a member of the resistance/ modulation / division efflux pump family. *Proc Natl Acad Sci USA*, 95: 1829-1833.
- Shaheen, A.M. (1998):** IL-6 as an early marker for diagnosis of neonatal septicemia Thesis submitted for partial full-filament of M.S. degree in pediatrics. Benha faculty of Medicine
- Shamseldin El-Shafie, S.; Smith, W. and Donnelly, G. (1995):** An outbreak of gentamicin – resistant *Klebsiella pneumoniae* in a neonatal ward. *Eur. J. Public. Health*, 3 : 129 – 31.
- Shatla, H.; El-Khayat, I.I. and Ibrahim, M.A. (1989):** The predictive value of gram stained tracheal aspirate smears in diagnosing neonatal bacteremia. *Ain shams Medical J.*, 49 : 31.
- Sheagren, J.N. (1991):** Corticosteroids for treatment of septic shock. *Infect. Clin. North . Am .* 5: 875-82.
- Shimizu, S.; Hirano, Y.; Yoshiok, B. et al. (1988) :** IL-6 dependent growth of alert's lymphoma-derived T-cell line *Blood* 72: 1826-1828.
- Shiomi, K.; Nakazato, M.; Thi, T. (1993):** Establishment of radioimmunoassay for human neutrophil peptides and their increases in plasma and neutrophil in infection. *Biochem Biopsy Res. Commum* 1993: 195: 1336-14.
- Siegel, J.D. (1985):** Neonatal septicemia. *Perinatol.* 9:20.

- Silveria, R.C.; Procianoy, R.S. (Find other articles with these Authors) (1999):** Evaluation of interleukin-6, tumour necrosis factor alpha and interleukin-1 beta for early diagnosis of neonatal sepsis. *Acta. Pediatr.*, 88 (6) : 647 – 50.
- Singer, L.A.; Plotz, B.; Elester, G. and Amer, A. (1975):** Estimation of C-reactive protein. *Clin. Path.*, 25 : 611.
- Somen, M.; Green, B. and Daling, J. (1993):** Risk factors for early neonatal sepsis . *American Journal of Epidemiology* 121: 712-19.
- Soong, L.; Ganz, T.; Ellison, A. and Caughey, G. (1997):** Purification and characterization of defensins from cystic fibrosis sputum. *Inflamm Res.*, 46: 98-102.
- Spangelo, B.L.; Isakson, P. and MacLead, R.M. (1989):** IL-6 stimulates anterior pituitary hormone release in vitro *Endocrinology* 125:575-577.
- Springer, T.A. (1990):** Adhesion receptors of the immune system. *Nature*; 64:425-434
- Squir E.; Favara, B. and Todd, J. (1979):** Diagnosis of Neonatal Bacterial Infection. Hematologic and Pathologic Finding in Fatal and Non Fatal cases. *Pediatrics* 64 : 60 – 64.
- St Gem, J.W.; Bewll, L.M. and Baumgart, S. (1990):** Distinguishing sepsis from blood cultre contamination in young infants with blood cultures growing coagulase – negative staphylococci 86: 157-162.

- St Geme, J.W. and Polin, R.A. (1996):** Prevention and treatment of neonatal sepsis . In : Text Book of Intensive care of the Fetus and Neonate 1st ed. (eds). Spitzer AR. Mosby, Philadelphia. 927-939.
- Stataua, J. and Wicher, J.T. (1988):** Tests for detecting and monitoring the acute phase response. Archives of Disease in Childhood 14: 804-805.
- Stoll, B.J. and Wiesman, L.E. (1997):** Infection in perinatology . Clin. Perinatol. 24:1.
- Stoll, B.J.; Gordon, T. and Kornos, S.B. (1996):** Early onset sepsis in VLBW neonates. A report from National Institute of Child Health and Human Development . Neonat. Res . J ped. 129:72.
- Sullivan, J.S.; Kilpatrick L. and Costarino A.T. (1992):** Correlation of Plasma cytokine elevations with mortality rate in children with sepsis. J pediar., 120:510-5 .
- Taga, T.; Kamanishi, K. and Hardy, R.R. (1987) :** Receptors for B-cell stimulatory factor (BSF-2) : Quantiations, specificity, distribution and regulation of the expression. J.Exp. Med., 166:967.
- Takai, Y.; Wong , G.G. and Clark, S.C. (1988):** B cell stimulatory factor- 2 is involved in the differentiation of cytotoxic T lymphocytes. J. Immunol. 140,508-512.
- Tang, Y.Q.; Yaun, J.; Osapay, G.; Osapay, C.; Tran, D.; Miller, C.; Quелlette, A. and Selsted, M. (1999):** A cyclic antimicrobial peptide produce in primate leukocytes by the ligation of two truncated a defensins. Scince, 286: 498-502.

- Territo, M.C.; Ganz, T.; Selsted, M.E. & Lehrer, R. (1989):** Monocyte- chemotactic activity of defensins from human neutrophils. *J Clin Invest*, 84:2017-2020.
- Thomas, N.J.; Carcillo, J.A.; Doughty, L.A.; Sasser, H. and Heine, R.P. (2002):** Plasma Concentrations of defensins and Lactoferrin in children with severe sepsis. *Pediatr Infect Dis. J.* 21 (1) : 34 – 8.
- Thompson, P.I.; Greenough, A. and Gamsa, H.K. (1992):** Congenital bacterial sepsis in preterm infant. *J. Med Microbial* . 36(2): P117-120.
- Thurlbeck, S.M. and McIntoch, N. (1987):** Preterm blood counts vary with sampling site. *Archives of Disease in childhood* 62:72-75.
- Tollner, U.; Pohlandt and Rike, S.R. (1982):** Septicemia in the newborn due to gram-negative bacilli: Risk factors, clinical symptoms and hematological changes. *Europ J. Pediatr.*; 123:243-254.
- Toolner, U. (1982):** Early diagnosis of septicemia in newborn. Clinical studies and sepsis score. *Eur. J. Pediatric*. 138/(10): 331-7.
- Tracey, K.J.; Beutler, B. and Lowry, S.F. (1986) :** Shock and tissue injury induced by recumbent human cachectin-*Science*, 234 : 470-4.
- Trugo, W.E.; Gibson, R.L. and Henderson, W.R. (1990):** Tumour necrosis factor induced neonatal pulmonary hypertension. *Pediatr. Res.*, 27:466.

- Valore, E.V.; Park, C.H.; Quayle, A.J.; Wiles, K.R.; McCray, P.B. Jr and Ganz, T. (1998):* Human β -defensin -1 : an antimicrobial peptide of urogenital tissues . J Clin invest., 101:1633-1642.
- Van Damme, J.; Fibbe, W.F. and Fiksenbury, L.H.(1989):* Simultaneous production of IL-6 , interferon - B and colony stimulating activity by fibroblasts after viral and bacterial infection. Eur. Immunol. 19, 163-168.
- Van Deventer, S.J.H.; Buller, H.R. and Catem, J.W. (1990):* Experimental endotoxemia in human : analysis of cytosine release and coagulation, fibrinolytic and complement pathways. Blood; 76:2520.
- Van Wetering, S.; Mannesse - Lazeroms, S.P.; Van Sterkenburg, M.A.; Daha, M.R.; Dijkman, J.H. and Hiemstra, P.S. (1997):* Effect of defensins on interleukin -8 synthesis in airway epithelial cells . Am J Physiol Lung Cell Mol physiol., 272: L 888 - L896.
- Van Wetering, S.; Mannesse- Lazeroms, S.P.; Dijkman, J.H. and Hiemstra, P.S. (1997):* Effect of neutrophil serine proteinases and defensins on lung epithelial cells : Modulation of cytotoxicity and IL-8 production . Leukoc Biol., 62: 217-226.
- Van Wetering, S.; Rahman, I.; Hiemstra, P.S. and MacNee, W. (1998):* Role of intra-cellular glutathione in neutrophil defensin- induced IL-8 synthesis and cytotoxicity in airway epithelial cells. Eur Respir J., 12: 420s .

- Van Wetering, S.; Sterk, P.J.; Rabe, K.F. and Hiemstra, P.S. (1999):* Defensins: players or bystanders in infection, injury, and repair in the lung. *Allergy Clin Immunol.*, 104: 1131-1138.
- Van Wetering, S.; Van der Linden, A.C.; Van Sterkenburg, M.A.; de Boer, W.I.; Kuipers, A.L.; Schalkwijk, J. and Hiemstra, P.S. (2000):* Regulation of SLP and elfin release from bronchial epithelial cells by neutrophil defensins. *Am J Physiol Lung Cell Mol Physiol.*, 278:L51-L58.
- Vesikari, T.; Isolauri, Tuppurainen, N.; Renlund, M.; Koivisto, M. and Kunnas, M. (1985):* Neonatal Septicemia in Finland 1981 – 1985. *Acta. Pediatr. Scand.*, 78 : 44.
- Vink, A.; Coulie, P.G. and Stevens, M. (1990):* Mouse plasmacytoma growth in vivo Enhancement by IL-6 and inhibition by antibodies direct against IL-6 or its receptor. *J. Exp. Med.* 172,997-1000.
- Voora, Strinivasan, G. and Lilien, L.D. (1982):* Fever in full – term newborns in the 1st 4 days of life. *Pediatrics* 69:40-44.
- Watkinson, M. and Mupanemunda, R.H. (1999):* Infection – General Key Topics in neonatology Bios Scientific Pul. Oxford, Washington, D.C.
- Webber, S.; Lindell, D. and Wilkinson, A.R. (1990):* Neonatal pneumonia. *Arch . Dis . Child .* 65: 207-211.

- ❖❖ **Weisbart, P.H.; Gasson, J.C. and Golde, D.W. (1989):** Colon – stimulating factors and host defence. *Ann Inter Med.*, 110:297-303.
- Weisman, L.E. and Lorenzetti, P.M. (1989):** High intravenous doses of human immunoglobulin suppress neonatal GBS immunity in rat. *J. Pediatr.* 115:445-450.
- Welling, M.; Pieter, S.; Maria, T.H.; peter, H.; Ernest, K.; Pauwels, and Wim Calame. (1998):** Antibacterial activity of human neutrophil Defensins in experimental infections in mice. *J. Clin. Invest.* 102 : 1583 – 1590.
- Whitfield, J.M. (1993):** Neonatal sepsis. In : Holbrook PR (eds.) . *Textbook of pediatric critical care 3rd ed.* WB Saunders Company , Philadelphia, London, Toronto, P.98.
- Wilkins, I.; Mezrow, G.; Lynchi Boltone, E.J. and Berkowitz R.L. (1989):** Amnionitis and life threatening respiratory distress after percutaneous umbilical blood sampling . *Am . J. Obstet. Gynecol .* 160 (2) : 424- 428.
- William, W.; Groothin, J.R.; Hayward, A.R. and Levin, M.J. (1997):** Infection control current pediatric diagnosis and treatment (13th ed), P. 59.
- Wilmott, R.; Fiedler, M. and Strak, J. (1998):** Host defense mechanisms. In disorders of the Respiratory Tract in Children. Edited by Chernick V, Boat TF. Philadelphia : Saunders; 238-264.

- Wilson, C.L.; Ouellette, A.J.; Satchell, D.P.; Ayabe, T.; Lopez-Boado, Y.S.; Stratman, J.L.; Hultgren, S.J.; Matrisian, L.M. and Parks, W.C. (1999): Regulation of intestinal defense activation by the metalloproteinase matrilysin in innate host defense. *Science*, 286: 113-117.
- Wood, J.R. (1993): Drug abuse in pregnancy. *Clin. Obstet. Gynecol.* 36(2) : 221-231.
- Wright, S.D. and Jong. M.T.C. (1986): Adhesion Promoting reception on human macrophages recognize E-coli by binding to polysaccharides. *J.Exp.Med*; 164:187-88.
- Yachie, A.; Takano, N.; Ohta K. et al. (1992): Defective production of IL-6 in very small premature infants in response to bacterial pathogens. *Infect immune*, 60:749-35.
- Yancey, M.O.; Duff, P.; Kubitis, P.; Cark, P. and Fontana, B.M. (1996): Risk factors for neonatal sepsis. *Obstet. General* 87 (2):
- Yang, D.; Chen, Q.; Chertov, O. and Oppenheim, J.J. (2000): Human neutrophil defenses selectively chemoattract naïve T and immature dendritic cells. *J Leukoc Biol.*, 68:9-14.
- Yang, D.; Chertov, O.; Bykovskaia, S.; Chen, Q.; Buffo, M.; Shogan, J.; Anderson, M.; Schroder, J.; Wang, J.; Howard, O. and Oppenheim, J. (1999): β - defensins: linking innate and adaptive immunity through dendrite and T cell CCR6. *Science*, 286:525-528.

Yang, K.; Bohansack, J. and Hill, H. (1993): Fibronectin in host defense: Implication in the diagnosis and prophylaxis therapy of infectious diseases. *Pediatr. Infect. Dis. J.*, 12: P 234- 239.

Zipursky, A.; Palko, J.; Milner, R. and Akenzua, G.I. (1967): The hematology of bacterial infection in premature infants. *Pediatrics* 57: 839-53.