

REFERENCES

Abrams, J.S. and Reines, H.D. (1979) :

Increasing incidence of right-sided lesions in colorectal cancer. Am, J. surg.; 137 : 522.

Abrams, J.S. (1980) :

Clinical staging of rectal cancer. Am. J. surg.; 139:539.

Abdi, E.A.; Hanson, J.; Harbora, D.E. and Young, D.G. (1989):

Adjuvant chemoimmuno and immunotherapy in Duke's stage B2 and C colorectal carcinoma : A 7-year follow-up analysis. J. surg. Oncol.; 40 : 205.

Adachi, W.; Sugeno, A. and Lida, F. (1991) :

Immune competent cells of regional lymph nodes in colorectal cancer patients. Flowcytometric analysis of lymphocyte subpopulation. J. Surg. Oncol.; 46: 110.

Adloff, M.; Arnaud, P.; Schleogel, M. and Bermaschi, R. (1986):

Colorectal cancer in patients under 40 years of age. Dis. Colon Rectum; 29: 322.

Adloff, M.; Arnaud, J.P.; Bermaschi, R. and Schloegel, M. (1989):

Synchronous carcinoma of the colon and rectum : Prognostic and therapeutic implications. Am. J. Surg.; 157: 299.

Ahfeld, F. (1880):

Zur Cavistik der congenitalen Neoplasmen. Arch. Gynakol; 16: 135. In: Br. J. Surg; 66: 621.

Akwari, O.E. and Kelly, K.A. (1980) :

Anterior resection for adenocarcinoma of the distal large bowel. Am. J. Surg.; 139 : 88

Al-Fallouji, M.A.R. and McBrien, M.P. (eds.) (1986) :

Tumour markers and cancer screening in surgery.; page 152, chap 1.44. In: Postgraduate Surgery, The Candidate's Guide; printed in Great Britain by Alden press, Oxford.

Almersjo, O. (1972):

Evaluation of hepatic dearterialization in primary and secondary Cancer of the liver. Am. J. Surg.; 124: 5.

Albridge, MC.; Phillips, R.K.S. and Hittinger R.; Fry, JS. (1986):

Influence of tumour site on presentation, management, and subsequent outcome in large bowel cancer. Br. J. surg.; 73 : 663.

Allee, P.E.; Tepper, J.E.; Gunderson, L.L. and Munzenrider, J.E. (1990):

Postoperative radiation therapy for incompletely resected colorectal carcinoma. Int. J. Radiat. Oncol. Biol. Phys.; 17 : 1171.

American Joint Committee for Cancer Staging and End Results (1977):

Manual for staging of cancer (Chap 2). Chicago, 25 : 80

Ambros, RA.; Pawel, BR.; Meshcheryakov, I. and Kotrotsios, J. (1990):

Nuclear morphometry as a prognostic indicator in colorectal carcinoma resected for cure. Anal-Quant cytol-Histol; 12 : 172.

Anonymous (1990):

Rectal cancer should be treated by experts. Lancet; I: 1476.

Armitage, N.C.; Davidson, A. and Wood, CB. (1984):

A study of the reliability of CEA blood levels in following the course of colorectal cance. Clin. Oncol.; 10: 141.

Arnaud, JP.; Schloegel, M.; Ollier, JC. and Adloff, M. (1991):

Colorectal carcinoma in patients over 40 years of age. Dis Colon Rectum ; 34 : 896.

Astler, V.B. and Collier, F.A. (1954) :

The prognostic significance of direct extension of carcinoma of the colon and rectum. Ann. Surg.; 139 :846.

AU, F.C.; Stein, B. and Tang, Ch.K. (1986) :

Carcinoembryomic antigen levels in colonic lesions. Am. J. Surg.; 151 : 61.

Autrup, H. and Williams, G.M. (eds.) (1985):

Experimental colon carcinogenesis. Boca Raton Press Inc. 1983.

Axtell, L.M.; Cutler, SJ. and Myers, MH (1972):

End Result in Cancer Report no. 4, HEW publication (NIH), 73-272: 1.

Baretton, G.; Gile, J. and Lohrs, U. (1991):

Flow-cytometric analysis of the DNA-content in paraffin embedded tissue from colorectal carcinomas and its prognostic significance Virchows. Arch. B. Cell. Pathol.; 60: 122.

Barillari, P.; de-Angelis, R.; Valabrega, S. and Indinnimeo, M. (1989) :

Relationship of symptom, duration and survival in patients with colorectal carcinoma. Eur. J. Surg. Oncol., 15 : 441.

Beckman, E.N., Gathright, J.B, and Ray, J.E. (1984):

A potentially brighter prognosis for colon carcinoma in the third and forth decades. Cancer; 54: 1478.

Bear, H.D.; Macintyre, J.; Burns, HJ. and Jarrett, F. (1984):

Colon and rectal carcinoma in the west of scotland. Symptoms, histologic characteristics and outcome. Am. J. Surg.; 147 : 441.

**Beart, R.W.; Moerter, C.G.; Wieand, H.S. and Leigh, J.E.
(1990) :**

Adjuvant therapy for resectable colorectal carcinoma with fluorouracil administered by portal vein infusion. A study of Mayo clinic and North Central Cancer Treatment Group. Arch. Surg.; 125 : 897.

Bedikian, A.Y.; Kantarjian, H. and Nelson, R.s. (1981) :

Colorectal Cancer in young adults.
South Med. J.; 74 : 920.

Begenet, R.H.; Keep, P.A.; Green, A.G. and Cooke, T.G. 1986):

Radioimmunolocalization and select for surgery in recrrtent CRC. Br. J. Surg. 73; 64: 67.

Beynon,J.; Mortensen, N.J., Foy,D.M. and Channer, J.L.(1986):

Pre-operative assessment of local invasion in rectal cancer : digital examination, endoluminal sonography or computed tomography? Br. J. Surg.; 73 :1015.

Belli, L.; Beati, C.A.; FRangi, M. and Aseni, P. (1988) :

Outcome of patients with rectal cancer treated by stapled anterior resection. Br. J. Surg.; 75 : 422.

**Behbehani, A.I.; Al-Nageeb N.; Omar, Y.T. and El-Nas, S.A.
(1990) :**

Serial determinations of CEA in monitoring and management of patients with colorectol carcinoma. Oncology; 47 : 303.

**Biert, J.; Eggink, W.F.; Van Berg, G.P. and Reuvers, C.B.
(1990):**

The lynch syndrome; A high-risk factor for colorectal carcinoma. Neth. J. Surg.; 41: 108.

**Bischof, D.A.; Delaloye, B.; Buchegger, F. and Gilgien, W.
(1989):**

Clinical value of immunoscintigraphy in colorectal carcinoma patients : A prospective study.
J. Nucl. Med; 30 : 1646.

Blair, SD.; Theodorou, NA.; Begent, RH. and dawson, PM. (1990):

Comparisom of anti-foetal colonic microvillus and anti-CEA antibodies in preoperative radioimmunolocalization of colorectal cancer. Br. J. Cancer; 61 : 891.

Block, I.R. and Enquist (1964):

A more radical resection for carcinoma of the rectum in the female, Surg. Gynaecol. Obstet., 150: 85.

Blumberg, N.; Agarwal, M.M. and Chuang, C. (1985):

Relation between recurrence of cancer of the colon and blood transfusion. Br. Med. J.; 290 :1037.

Bohm, B.; Nouchirvani, K.; Hucke, HP. and stock, W. (1990) :

The effect of age and sex on localization, tumour stage and prognosis of colorectal carcinoma. Med. Clin; 85:586.

Bown, S.G.; Barr, H.; Mathewson, K. and Boulos, P.B. (1986):

Endoscopic treatment of inoperable colorectal carcinoma with the Nd YAG Laser. Br. J. Surg.; 73: 949.

Bonnheim, D.C.; Petrelli, N.J.; Herrera, L. and Walsh, D. (1986):

Osseous metastases from colorectal carcinoma. Am. J. Surg.; 151 : 457.

Boxall, T.A. (1962):

The blood supply of the distal segment of the rectum in anterior resection, Br. J. Surg.; 50: 399.

Bromdt, L.J., Katz, H.J. and Wolf, E.L. (1985) :

Stimulation of Colonic Carcinoma by ischemia. Gastroenterology ; 88 : 1137.

Bresalier, RS. and Kim, YS. (1985):

Deer and Colon Cancer. N. Engl. J. Med.; 313: 1413.

Broders, A.C. (1925) :

Grading of carcinoma. Minn Med.; 8 : 726.

Brown, S.C. and Sykes, P.A. (1988) :

Operative mortality rate and surgery for colorectal cancer. Br. J. Surg.; 75 : 645.

Brandao, O.; Sorbrinho, M.A. and Serrao, D. (1985) :

Prognosis in colorectal carcinoma. A reassessment of the pathologist's role. Path. Res. Pract.; 180 : 506.

Brzeniski, W.; Orrom, W.J. and Wiens, E. (1990):

The prospective and retrospective analysis of colonoscopy findings in patients with a history of colorectal carcinoma in first-degree relatives. Can. J. Surg.; 33: 314.

Buechter, K.J.; Boustany, CH. and Caillouette, R. (1988):

Surgical management of the acutely obstructed colon. Am. J. Surg.; 156 : 163.

Bulow, S.; Svendsen, L.B. and Mellemegaard, A. (1990) :

Metachronous colorectal carcinoma.
Br. J. Surg. 77 : 502.

Bulow, S. (1980):

Colorectal cancer in patients less than 40 years of age in Denmark. Dis. Colon Rectum; 23: 327.

Bull, A.W.; Nigro, ND. and Crissman, J.D. (1984):

In vivo stimulation of DNA synthesis and induction of ornithine decarboxylase in rat colon by fatty acid hydroperoxidase, autoxidation products of unsaturated fatty acids. Cancer Res.; 44: 4927.

Bundred, N.J. (1985):

Gastric surgery and the risk of subsequent colorectal cancer. Br. J. Surg.; 72: 618.

Buyse, M.; Zeleniuch-Jacquotte, A. and Chalmers, TC. (1988) :

"Adjuvant therapy of colorectal carcinoma"
JAMMA; 259 : 3571.

Carretier, M.; Coste, G.; Jardel, P. and Reynauld, J. (1983):

Generalized peritonitis induced by non-traumatic colonic disorders. Sixty-two cases. Ann. Chir.; 37 : 417.

Castro, E.B.; Rosen, P.P. and Quan, S.H. (1973):

Carcinoma of large intestine in patients irradiated for carcinoma of cervix and uterus. Cancer; 31: 45.

Cedermark, B.J.; Schultz, S.S.; Baksslu, S. and Evans, J.T. (1977):

The value of liver scan in the follow-up study of patient with adenocarcinoma of the Colon and rectum. Surg. Gynecol. Obstet.; 144: 745.

Chapuis, PH.; Dent, O.F.; Fisher, R. and Newland, R.C. (1985):

A multivariate analysis of clinical and pathological variables in prognosis after resection of large bowel cancer. Br. J. Surg.; 72 : 693.

Chen, M.; Chuan, C. and Chan, P. (1980):

Evolution of colorectal cancer in schistosomiasis. Transitional mucosal changes adjacent to large intestinal carcinoma in colectomy specimens. Cancers, 46: 1661.

Cheever, A.W. (1981):

Schistosomiasis and colon cancer. Lancet; 2: 1369.

Chifflet, A. (1964):

Surgery for cancer of the lower rectum: The perirectal fascia with reference to conservative surgery and technic, Dis. Colon Rectum.; 7: 493.

Chua, L.; Ardalan, B.; Sridhar, K.S. and Richman, S. (1990):

A phase II study of short-term continuous infusion with high-dose 5-FU with Lecucovorin in Colorectal Carcinoma. Proc. Annu. Meet. Am. Soc. Clin. Oncol.; 9: A 401.

Clarke, D.N.; Jones, P.F. and Needham, CD. (1980):

Outcome of in Colorectal Carcinoma; seve-year study of a population. Br. Med. J.; 280: 431.

**Crucitti, F.; Sofo, L.; Doglietto, GB. and Bellantone, R.
(1991):**

Ptohnodyic factors in colorectal cancer : Current status
and new trends. I. Surg. Oncol. Suppl; 2 : 76.

Cohn, I.Jr. and Bronside, G.H. (1984):

Infections. In: Principles of surgery; edited by Mc
Graw-Hill Book Company, 4th edition, Page 173.

**Cochrane, J.P.; Williams, J.T.; Faber, R.G. and Slack, W.W.
(1980) :**

Value of follow-up after curative surgery for carcinomas
of the large bowel. Br. Med. J. 280 : 593.

Co, E.; HSU, TC. and HSU, K.C. (1990) :

Diagnosis of colorectal carcinoma:is colonoscopy
superior to barium enema?
Chung-Hua -1- Hsueh-Tsa-Chih.; 45 : 60.

Connelly, J.H.; Robey-Caferty, SS. and Clearly, KR. (1991) :

Mucinous carcinomas of the colon and rectum. An analysis
of 62 stage B and C lesion. Arch. Pathol Lab. Med.;
115 : 1022.

Crowther, J.S.; Drasar, B.S. and Hill, M.G. (1976):

Faecal steroids and bacteria and large bowel cancer in
Hong Kong by socioeconomic groups. Br. J. Cancer; 34:
191.

Craft, C.F.; Mendelsohn, G. and Cooper, H.S. (1981):

Colonic Pre-Cancer in Crohn's disease. Gastroenterology;
80: 578.

Cummings, B.J. (1983) :

Symposium on the diagnosis and treatment of common
cancers. 3 : colorectal carcinoma. Can. J. Surg.; 26 :
271.

**Daniel, T.M.; Gesrge, H.H.; Tuttle, S. and Thurston, M.C.
(1985):**

Intraoperative Radioimmuno detection of colorectal
carcinoma with hand-held detector. Am. J. Surg.,150,672.

Davis, N.C. and Newland, R.C. (1983) :

Terminology and classification of colorectal adenocarcinoma: the Australian Clinico-pathological staging system. Aust. NZ. J. Surg.; 53 : 211.

Davis, N.C.; Evans, E.B.; Cohen, J.R. and Job, D. (1985) :

Clinicopathological staging of colorectal cancer : has the time arrived? Br. J. Surg.; (Suppl).; 72 : S 47.

Dawson, P.M.; Habib, N.A. and Rees, H.C. (1987) :

Mucosal field change in colorectal cancer.
Am. J. Surg.; 153 : 281.

de Mascarel, A.; Coindre, J.M.; Maree, D. and Hoerni, B. (1981):

The prognostic significance of specific histologic features of colorectal carcinoma. Surg. Gynecol. Obstet.; 153: 511.

Danapoulou, ED.; Danapoulou, IE. (1981):

Eleven years experience of oral urea treatment in liver malignancies. Clin Oncol; 7: 281.

Decker, GAG. and du Plessis, DJ. (1986) :

"The laege Bowel, Anal Canal and Ischiorectal Fossa" In: Lee Mc Gregors' Synopsis of Surgical Anatomy, page 41, chapter 5, edited by Decker and duplessis, 12th edition, Wight Bristol (Engl).

Decosse, J.J. (1989) :

Conservative treatment of distal recal cancer by local excision. Cancer; 63 : 219.

Denise, Sh. (1985):

Genetic aspects of Carcinogenesis.
Br. J. Surg. 72; Suppl : 39.

Deveney, K.E. and Way, L.W. (1984):

Follow-Up of patients with colorectal cancer. Am. J. Surg.; 148 : 717.

Devine, T.J.; Myers, K.A. and Slattery, P.G. (1980):

Severe leg ischemia caused by anterior resection of the rectum. Br. J. Surg.; 67: 52.

Devereux, D.F. and Deckers, P.J. (1985) :

Contributions of pathologic margins and Dukes' Stage to local recurrence in colorectal carcinoma.
Am. J. Surg.; 149 : 323.

Deutsch, A.A.; Zelikovski, A.; Sternberg, A. and Reiss, R. (1983) :

One-stage Subtotal colectomy with anastomosis for obstructing carcinoma of the left colon. Dis. Colon Rectum; 26 : 227.

Domergue, J.; Ismail, M. and Astre, C. (1988):

Colorectal Carcinoma in patients younger than 40 years of age. Cancer; 61: 835.

Dukes, CE. and Bussey, HJR. (1958) :

The Spread of rectal cancer and its effect on prognosis. Br. J. Surg.; 12 : 309.

Durdey, P.; Cooper, J.C.; Switala, S.; King, R.F. and Williams, N.S. (1985) :

The role of peptidases in cancer of the rectum and sigmoid colon. Br. J. Surg.; 72 : 378.

Durivage, H.J.; Rosenberg, A.H.; Rosman, M. and Bobrow, S.N. (1990) :

Methotrexate (MTX), 5-fluorouracil (5 Fu), and mitomycin C (MMC) for the treatment of metastatic Colorectal Carcinoma.
Proc. Ann. Meet. Am. Soc. Clin. Oncol.; 9 : 449.

Dunlop, M.G. (1990):

Inheritance of colorectal susceptibility. Br. J. Surg.; 77: 245.

Dykes, P.W.; Hine, K.R. and Bradwall, A.R. (1980):

Localization of tumour deposits by external scanning after injection of radiolabelled anti-carcinembryonic antigen. Br. Med. J.; 280: 220.

Eisenberg, S.B.; Kraybill, W.G. and Lopez, M.G. (1990) :

Long-term results of surgical resection of locally advanced colorectal carcinoma. Surgery.; 108 : 779.

Eisenberg, H.; Sullivan, P.D. and Foot, F.M. (1967):

Trends in survival of digestive system cancer patients in connecticut, 1935-1962. Gastroenterolgy; 53: 528.

Elliot, M.S. and Louw, J.H. (1979):

10 year survey of large bowel carcinoma at Groote schur Hospital with particular referece to patients under 30 years of age. Br. J. Surg.; 66: 621.

Enker, W.E.; Laffer, U.T. and Block, G.E. (1979):

Enhanced survival of patients with colon and rectal cancer based upon wide anatomic resection. Ann. Surg.; 190: 350.

Enblad, G.; Enblad, P.; Adami, H.O. and Glimelius, B.(1990) :

Relationship between age and survival in cancer of the colon and rectum with special reference to patients less than 40 years of age. Br. J. Surg.; 77 : 611.

Erlichman, C.; Fine, S.; Wong, A. and Elhakim, T. (1988) :

A randomized trial of flurouracil and folinic acid in patients with metastatic colorectal carcinoma. J. Clin. Oncol.; 6 : 469.

Everhart, J.E. (1989) :

A longitudinal survey of self-reported bowel habits in the United States. Dis. Dis. Sci.; 34 : 115.

Ezzo, J.A.; Sullivan, J.F. and Mack, R.E. (1958):

Carcinoma of the colon under the age of 40. Ann. Intern. Med.; 49: 321.

Fenns, Y.S.; HSU, H. and Chen, S.S. (1987) :

One stage operation for obstructing carcinoma of the left colon. Dis. Colon Rectum; 30 : 29.

Feischer (1989) :

Detection and Surveillance of colorectal Carcinoma.
JAMMA; 261 : 580

Finlay, I.G. and McArdle, C.S. (1986) :

Occult hepatic metastases in colorectal carcinoma.
Br. J. Surg.; 73 : 732.

Filipe, Ml. (1969):

Value of histochemical reactions for mucosubstances in the diagnosis of certain pathological conditions of the colon and rectum. Gut.; 8: 577.

Fisher, D.E. (1988) :

Postoperative adjuvant chemotherapy or radiation therapy for rectal cancer. J. Natl. Cancer Inst.; 80 : 21.

Fork, F.T. (1988):

Radiographic findings in overlooked colon carcinomas.
Acta. Radiol.; 29: 331.

Foutch, PG.; Mai, H.; Pardy, K. and Disario, JA. (1991):

Colorectal cancer in asymptomatic, average-risk men.
Dig. Dis Sci.; 36 : 924.

Gastrointestinal Tumour Study Group (1984):

Adjuvant therapy of colon cancer-results of a prospectively randomized trial. New England Journal of Medicine 310 : 737.

Garland, C.F. and Garland, F.C. (1989):

Do Sunlight and Vitamin D reduce the likelihood of colon cancer?. Int. J. Epidemiol.; 9: 227.

Gilbertsen, V.A. (1959):

Adenocarcinoma of the large bowel: 1; 340 cases with 100 percent follow-up-surg; st. Louis; 46: 1027.

Goldthorn, J.F.; Powars, D. and Hays, D.M. (1983) :

Adenocarcinoma of the colon and rectum in the adolescent surgery; 93 : 409.

Goldberg, S.M.; Gordon, P.H. and Nivatongs, S. (eds.) (1980):

Malignant Neoplasms of the rectum. In: Essentials of Anorectal Surgery, 2nd ed., Chap 16, P 182 Philadelphia; Lippincott.

Goldenberg, DM.; Kim EE.; Bennett, SJ. and Nelson, MO. (1983):

Carcinoembryonic antigen radioimmuno-detection in the evaluation of colorectal cancer and in the detection of occult neoplasms. Gastroenterology 84 : 524.

Goligher, J.C.; Garham, N.G. and Dombal, F.T. (1970):

Anastomotic dehiscence after anterior resection of rectum and sigmoid. Br. J. Surg.; 57: 109.

Goligher, J.C. (1975):

Surgery of the Anus, Rectum and Colon. 3rd ed., London: BAilliere Tindall.

Goligher, J.C. (1985) :

Diagnosis of diseases of the anus, rectum and colon. In Surgery of the Anus, Rectum and colon, 5th. ed., P. 659 London : Balliere Tindall.

Goh, HS.; Soong, TW.; Lee, YS. and Goh, CR (1990):

C-myc oncogene expression in colorectal cancer: its use in prognosis and role in colorectal carcinogenesis. Ann. Acad. Med. Sigapor; 19: 139.

Grinnel, R.S. (1953) :

The grading and prognosis of carcinoma of the colon and rectum. Ann. Surg.; 109 : 500.

Greaney, MG. and Irvin, TT. (1977) :

Criteria for the selection of rectal cancer for local treatment. Dis. Colon Rectum; 20 : 463.

Greaves, P.; Filipe, MI. and Branfoot, AC. (1980):

Transitional mucosa and survival in human colorectal cancer. *Cancer*; 46: 764.

Graft, W.; Glimelius, B.; Pahlman, L. and Bergstrom, R. (1991):

Determinant of prognosis in advanced colorectal cancer. *Eu. J. Cancer* (1991); 27:1119.

Grossman, S. (1988) :

Colonoscopic screening of persons with suspected risk factors for colon cancer : I. Family history. *Gastroenterology*. 94 : 395.

Gryska, P.V. and Cohen, A.M. (1987):

Screening asymptomatic patients at high risk for colon cancer with full colonosc. *Dis. Colon Rectum*; 30: 18.

Gottrup, F.; Diederich, P.; Sorensen, K. and Nielsen, S.A. (1985) :

Prophylaxis with whole gut irrigation and antimicrobials in colorectal surgery. *Am. J. Surg.*; 149 : 317.

Guadagni, F.; Roselli, M.; Amato, T.; Cosimelli, M. and Manella, E. (1991) :

Clinical evaluation of serum Tumour-associated glycoprotein-72 as a novel tumour marker for colorectal cancer patients. *J. Surg. Oncol. Suppl.*; 2 : 16.

Habib, N.A.; Peck, M.A.; Sawyer, C.N., Blaxland, J.W. and Luck, R.J. (1983) :

Does fixity affect prognosis in colorectal tumours? *Br. J. Surg.*; 70 : 423.

Hafstrom, L.; Rudenstam, C.M.; Domellof, L. and the Sweden Gastrointestinal Tumour Adjuvant Therapy Group (1985):

A randomized trial of oral 5-FU versus placebo as adjuvant therapy in colorectal cancer Duke's B and C: results after 5 years observation time. *Br. J. Surg.* 72: 138.

Heald, R.J. and Ryall, R.D. (1986):

Recurrence and survival after total mesorectal excision for rectal cancer. Lancet; i: 1479.

Hermanek, P.J. and Gall, F.P. (1981) :

Die aborale sicherheitsabstand beiderr sphinktererhaltenden Rectumresektion. Chirurg; 52: 25.

Heslov, S.F. and Frost, D.B. (1988) :

Extended resection for primary colorectal carcinoma involving adjacent organs or structures. Cancer; 62 : 1637.

Hill, M.J. and Aries, V. (1971) :

Faecal steroid composition and its relationship to cancer of the larg bowel. J. Pathol.; 104 : 129.

Hill, M.J. (1985) :

Cancer of the large bowel : human carcinogenesis. Br. J. Surg. Suppl.; 72 : S 37.

Hine, K.R. and Dykes, P.W. (1984) :

Serum CEA testing in the postoperative surveillance of colorectal carcinoma. Br. J. Cancer ; 49 : 689.

Hocking, G.R.; Rolland, J.M.; Nairn, R.C. and Pihl, E. (1982):

Lymphocyte fluorescence polarization changes after phytohaemagglutinin stimulation in the diagnosis of colorectal carcinoma. J. Natl. Cancer. Inst.; 68 : 579.

Homes, E.C.; Eilber, F.R. and Morton, D.L. (1975):

Immunotherapy of malignancy in humans. Current status. JAMA; 232: 1052.

Howard, M.L. and Greene, F.L. (1990) :

The effect of preoperative endoscopy on recurrence and survival following surgery for colorectal carcinoma. Am. J. Surg.; 56 : 124.

Holm, A.; Bradley, E. and Aldrete, J.S. (1989) :

Hepatic resection of metastasis from colorectal carcinoma. Ann. Surg.; 209 : 428.

Hojo, K.; Koyama, Y. and Moriya, Y. (1982) :

Lymphatic spread and its prognostic value in patients with rectal cancer. Am. J. Surg.; 144 : 350.

Hoekstra, HJ.; Verschueren, RCJ. and Oldhof, J. (1985) :

Palliative and curative electrocoagulation for rectal cancer. Cancer; 55 : 210.

Holdsworth, P.J.; Johnston, D.; Chalmers, A.G.; Dixon, M.F. and Finan, P.J. (1988) :

Endoluminal ultrasound and computed tomography in the staging of rectal cancer. Br. J. Surg.; 75 : 1019.

Hughes, E.S.R. and Cuthbertson, A.M. (1965) :

Subtotal colectomy for obstructing carcinoma of the upper left colon. Dis. Colon Rectum; 8 : 411.

HSU, Y.H. and Guzman, L.G. (1982):

Carcinoma of the colon and rectum in young adults. AM. J. Proctol. Gastroenterol. Colon Rectal Surg.; 33: 7.

Irvin, T.T. and Greaney, M.G. (1977):

The treatment of colonic cancer presenting with intestinal obstruction. Br. J. Surg.; 64: 741.

Irving, A.D. and Scrimgeour, D. (1987) :

Mechanical bowel preparation for colonic resection and anastomosis.
Br. J. Surg.; 74 : 580.

Irvin, T.T. (1988) :

Prognosis of colorectal cancer in the elderly.
Br. J. Surg.; 75 : 419.

Isacson, P. and Attwood, PRA. (1979):

Failure to demonstrate specificity of the morphological and histochemical changes in mucosa adjacent to colonic carcinoma. J. Clin. Path.; 32: 214.

Itzkowitz, SH.; Bloom, EJ.; Kokal, WA. and Kim, YS. (1990):

Sialosyl-n. Anovel mucin antigen associated with prognosis in colorectal cancer patients. Cancer; 66: 1960.

Jacobsen, BK. (1987):

Coffe, Cholesterol and colon cancer: Is there a link? Br. Med. J.; 294: 4.

Jarvinen, H.J. and Turunen, M.J. (1984) :

Colorectal carcinoma below 40 years of age, prognosis and predisposing conditions. Scand. J. Gastroenterol.; 19 : 634.

Jarvinen, H.J.; Ovaska, J. and Mecklin, J.P. (1988) :

Improvement in the treatment and prognosis of colorectal carcinoma. Br. J. Surg.; 75 : 25.

Jensen, J.; Kewenter, J.; Haglind, E. and Lycke, G. (1986) :

Diagnostic accuracy of double contrast enema and rectosigmoidoscopy in connection with faecal occult blood testing for the detection of rectosigmoid neoplasms Br. J. Surg.; 73 : 961.

Karashima, S.; Kataoka, H.; Itoh, H.; Maruyama, R. and Kono, M. (1990) :

Prognostic significance of alpha-1-antitrypsin (alpha 1-AT) in early stage of colorectal carcinomas. Int. J. Cancer; 45 : 244.

Kato, I.; Tominaga, S.; Matsuura, A. and Yoshii, Y. (1990) :

A comparative case-control study of colorectal cancer and adenoma. Jpn. J. Cancer Res.; 81 : 1101.

Karesen, R.; Hertzberg, J. and Johannesen, J. (1980) :

Carcinoembryonic antigen in the diagnosis and follow-up of colorectal carcinoma. Am. J. Proctol.; 3 : 18.

Kern, W.H. and White, W.C. (1958) :

Adenocarcinoma in a 9 months infant. Cancer; 11 : 855.

Kemeny, N. (1980):

Therapy for metastatic colorectal carcinoma with a combination of methyl CCNU, 5 FU, Vincristine and streptozotocin. Cancer; 45: 44.

Ketcham, A.S. (1985):

The management of reecurrent rectal cancer. The Canad. J. Surg. 28: 422.

Kiefhaber, P.; Huber, F. and Kiefhaber, K. (1987):

Palliative and preoperative endoscopic Neodymium-YAG Laser treatment of colorectal carcinoma. Endoscopy, 99 Suppl 1: 43.

Kirwan, W.O.; Turnbull, R.B. and Fazio, V.W. (1978):

Pullthrough operation with delayed anastomosis for rectal cancer. Br. J. Surg.; 65: 695.

Kirwan, W.O.; Drumm, J.; Hogan, J.M. and Keohane, C.K. (1988):

Determining safe margin of resection in low anterior resection for rectal cancer. Br.J. Surg.; 75 : 720.

Klompje, J.; Petrelli, N.J.; Herrera, L. and Mittelman, A. (1987) :

The prognostic value of preoperative alkaline phosphatase for resection of solitary liver metastasis from colorectal carcinoma. Eur. J. Surg. Oncol.; 13:345.

Kodner, I.J. (1988):

Endocavitary irradiation for rectal cancer: Better results when combined with external irradiation. Paper accepted at the ASCRS annual meeting, Anaheim, Calif.

Kodner, I.J. (1989) :

Preoperative irradiation for rectal cancer. Improved local control and long-term survival. Ann. Surg.209:194.

Koss, G.K. and Greenbaum, L. (1986) :

Measuring DNA in human cancer. JAMA.; 255 : 3158.

Kouri, M.; Lassonen, A.; Mecklin, J.P. and Jarvinen, H. (1990):

Diploid predominance in hereditary nonpolyposis colorectal carcinoma evaluated by flowcytometry. *Cancer*; 65:1825.

Korenaga, D.; Ueo H. and Moshida, K. (1991) :

Prognostic factors in Japanese patients with colorectal cancer; the significance of large bowel obstruction. Univariate and multivariate analysis. *J. Surg. Oncol*; 47: 188.

Kratzer, G.L. and Alia, R. (1980) :

Low anterior resection for rectal cancer, technique and results. *Am. J. Surg.*; 139 : 221.

Kumar, D. and Phillips, S.F. (1989) :

Human myenteric plexus confirmation of unfamiliar structures in adults and neonates. *Gastreenterology*; 96 : 1021.

Kune, G.A. and Kune, S. (1987) :

New design to examine colorectal cancer cause and survival *Dig. Surg.*; 4: 156.

Kuroki, T.; Kubota, A.; Miki, Y.; Yamamura, T. and Utsunomiya, J. (1991) :

Lectin staining of neoplastic and normal background colorectal mucosa in nonpolyposis and polyposis patients. *Dis. Colon Rectum*. 34 : 679.

Kyllonen, L.E.J. (1987) :

Obstruction and perforation complicating colorectal carcinoma. *Acta chir scand.*; 153 : 607-14.

Last, R.J. (1981):

Anatomy of colon, Rectum. In: *Last's Anatomy Regional and Applied*, 6th. edition, P. 327, McMin R.M.H. (ed.) Churchill Livingstone Edinburgh, London.

Lau, W.Y.; Chu K.W.; Poon, G.P. and HO. K.K. (1988) :

Prophylactic antibiotics in elective colorectal surgery *Br. J. Surg.*; 75 : 782.

Lawrance, R.J.; Cooper, A.G. and Loizidou, M. (1990) :

Blood transfusion and recurrence of colorectal cancer : the role of platelet derived growth factor. *Br. J. Surg.*; 77 : 1106.

Larson, G.M.; Bond, Sh. J.; Shallcross, C. and Polk, H.C.
(1986) :

Colonoscopy after curative resection of colorectal cancer. Arch. Surg.; 121 : 535.

Leaper, D.J. (1990) :

Tumours of the colon. In: Maingot's Abdominal Operations; Seymour I, Schwartz, and Harold Ellis (edit), 9th. edition, volume II, Ch. 46, P.1040 by Appleton and Lange. Printed in USA.

Lea, J.W.; Covington, K.; McSwain, B. and Scott, H.W.
(1982):

Surgical experience with carcinoma of the colon and rectum. Ann. Surg.; 195: 600.

Lehman, GA.; Buchner, DM. and Lappas, JC. (1983) :

Anatomical extent of fiberoptic Sigmoidscopy. Gastroenterology; 84 : 803.

Lions, D.A.; Beard, C.M. and Difallon, W.M. (1981):

Cholecystectomy and carcinoma of the colon. Lancet; 2:379.

Loser, C.; Folsch, UR.; Paprotny, C. and Greutzfeldt, W.
(1990) :

"Polyamines in human gastrointestinal malignancies" Digestion; 46 Supple 2 : 430.

Lointier, P.; Meggouth, F.; Dechelotte, P.; Pezet, D. and Ferrier, Ch. (1991) :

1.25-Dihydroxy vitamin D3 receptors and human colon adenocarcinoma. Br. J. Surg.; 78 : 435.

Long, R.T. and Edward, R.H. (1989) :

Implantation metastasis as a cause of local recurrence of colorectal carcinoma. Am. J. Surg.; 157 : 194.

Lofberg R.; Tribukait, B.; Ost, A. and Reichard, H. (1987):

Flow-cytometric DNA analysis in long standing ulcerative colitis: a method of prediction of dysplasia and carcinoma development. Gut; 28: 1100.

Lundy, J. (1980) :

Anaesthesia and surgery: a double-edged sword for the cancer patient. J. Surg. Oncol.; 14: 61.

Madden, J.L. and Kandalaft, S.I. (1983) :

Electrocoagulation as a primary curative method in the treatment of carcinoma of the rectum. Surg. Gynecol. Obstet.; 157 : 164-79.

Madden, J.I. and Mc Veigh, G.J. (1954):

Extension of operation in the treatment of carcinoma in the region of the splenic flexure. Surg. Clin. North. Am.; 34: 523.

Martin, E.W.; Joyce, S. and Lucas, J. (1981) :

Colorectal carcinoma in patients less than 40 years of age: pathology and prognosis. Dis. Colon Rectum ;24 : 25.

Malcolm, A.W.; Perencevich, N.P.; Olson, R.M. and Hanley, J.A. (1981):

Analysis of recurrent pattern following curative resection for carcinoma of the colon and rectum. Surg. Gynecol. Obstet.; 152: 131.

Malmberg, M. (1986):

Recurrence and survival after anterior resection of rectum using EEA stapler. Surg. Gynecol.Obstet.;163:231.

Maxfield, R.G. and Hampshire. C. (1984) :

Colonoscopy as a routine preoperative procedure for carcinoma of the colon. Am. J. Surg.; 147 : 477.

Mann, C.V.; Gallagher, P. and Frecker, P.B. (1988):

Rigid sigmoidoscopy : an evaluation of three parameters regarding diagnostic accuracy. Br. J. Surg.; 75: 425.

Marson, B.C. (1970):

Some leads to the etiology of cancer of the large bowel. Proc. R. Soc. Med. 64: 959.

Mavligit, G.M.; Gutterman, J.V.; Burgess, M.A. and Hersh, E.M. (1975):

Prolongation of postoperative disease-free interval and survival in human colorectal cancer by BCG or BCG plus 5-FU. Lancet; 1: 871.

Mcdermott, FT.; Hughes, ESR.; Pihl, E.; Milne, BJ. and Price, AB. (1981) :

Comparative results of surgical management of single carcinomas of the colon and rectum : a series of 1939 patients managed by one surgeon. Br. J. Surg. 68 : 850.

Mcdermott, S.F.; Hughes, E.S.; Pihl, E.; Johnson, W.R. and Price, A.B. (1985) :

Local recurrence after potentially curative resection for rectal cancer in a series of 1008 patients. Br. J. Surg.; 72 : 34.

Mcardle, C.S.; Hole, D.; Hansell, D. and Blumgart, L.H. (1990):

Prospective study of colorectal cancer in the west scotland : 10-year follow-up. Br. J. Surg.; 77 : 280.

Miles, W.E. (1908) :

A Method of performing abdomino-perineal excision for carcinoma of the rectum and of the terminal portion of the pelvic colon. Lancet; 2 : 1812.

Milnes-Walker, R. (1971):

Annual Report of South Western Regional cancer Bureau. Published by South Western Regional Board, U.T.F. House, Bristol. Engl.

Mitmayer, B.; Begin, LR. and Gordon, PH. (1991) :

Nuclear shape as a prognostic discriminant in colorectal carcinoma. Dis. Colon. Rectum.; 34 : 249.

Miller, M.P. and Stanley, T.V. (1988) :

Results of mass screening program for colorectal cancer Arch. Surg.; 123 : 63.

Mlasowsky, B.; Duben, W. and Jung, D. (1985) :

Cryosurgery for palliation of rectal tumours.
J. Exp. Clin. Cancer Res.; 4 : 81.

Mokhtar, N. (Edit) (1991):

Cancer pathology registry of the National cancer
Institute, Cairo University (1985-1989). page 5, AL-
Asdekaa Graphics centre, Cairo, Egypt.

Morson, B.C. and Day, D. (1981) :

In Decosse J.J. (ed) : pathology of adenomas and cancer
of the large bowel, in large bowel cancer.
London : Chirchill Livingstone, P 34.

Moore, PA.; Dilawri, RA. and Ridler, WJ. (1984) :

"Adenocarcinoma of the colon and rectum in patients less
than 40 years of age". Am. J. Surg.; 50 : 10.

Morgan, C.N. (1968):

Treatment of cancer of the rectum, Am. J. Surg. 115: 442.

Morgan, W.P.; Jenkins, N.; Lewis, P. and Auberg, D.A. (1985):

Management of obstructing carcinoma of the left colon by
extended right hemicolectomy Am. J. Surg. 149: 327.

Morgenstern, L. (1978):

Spatial distribution of colonic carcinoma. Arch. Surg.;
113: 1142. Ann. J. Surg.; 147 : 246.

Moskowitz, M.; Zimmermann, H. and Felson, B. (1964):

The meandering mesenteric artery of the colon. Am. J.
Roentgenol. Radium Ther. Nucl. Med. 92: 1088.

Mossa, A.R.; Ree, P.C. and Marks, J.E. (1979):

Factors influencing local recurrence after APR for
cancer of the rectum and rectosigmoid. Br. J. Surg.; 62: 727.

Muller, R.F. and Figley, M.M. (1957):

The arteries of the abdomen, pelvis, and thigh: I. Normal roentgenographic anatomy. II. Collateral circulation in obstructive arterial disease. Am. J. Roentgenol. Radium Ther. Nucl. Med. 77: 296.

Newland, RC.; Chapuis, PH.; Pheils, MT. and Macpherson, JG. (1981) :

The relationship of survival to staging and grading of colorectal carcinoma : a prospective study of 503 Cases. Cancer.; 47 : 1424.

Nesbitt, J.C.; Moise, K.J. and Sawyers, J.L. (1985) :

Colorectal carcinoma in pregnancy. Arch. Surg.; 120 : 636.

Neugut, AI. and Pita, S. (1988) :

Role of sigmoidoscopy in screening of colorectal neoplasms. Gastroenterology.; 95 : 492.

Ngala Kinda, J.F. (1976) :

Cancer of the large bowel in the African : a 15-year survey at kinshasa university hospital, Zaire. Br. J. Surg. 63 : 966.

Nicholls, R.J.; Mason, A. and Morson, BC. (1982) :

The clinical staging of rectal cancer.Br.J.Surg.;69:404.

Nishida, O.; Sano, F. and Sato, N. (1984) :

Clinical study and management of colon and rectal cancer in the young adults. Nippon shokaki Geka Zasshi;17:1758.

Nigro, ND.; Singh, DV. and Campbell, RL. (1975):

Effect of dietary beef fat on intestinal tumour formation by azoxymethane in rats.J.Natl Cancer Inst. i: 95.

Nigro, N.D. and Bull, A.W. (1985) :

Expermental intestinal carcinogenesis. Br. J. Surg.; 72: Supplement. 36.

Nio, Y.; Ohgaki, K.; Tsuchitani, T.; Imai, S. and et al. (1990) :

Orally administered streptococcal preparation, OK-432 augments the antitumour immunity of patients with gastric or colorectal cancer. Biotherapy.; 2 : 213.

Northover, J. (1986) :

Carcinoembryonic antigen and recurrent colorectal cancer. Gut.; 27 : 117.

Oates, G.D. (1988):

Colloquium on surgical management of cancer of the lower rectum: The feasibility of surgery in advanced case J. R. Coll. Edinb.; 33: 280.

Ohman, U. (1982) :

"Colorectal cancer in patients less than 40 years of age". Dis. Colon Rectum.; 25 : 209.

Okuno, M.; Ikehara, T.; Nagayama, M. and Sakamotok. (1987) :

"Colorectal carcinoma in young adults. Am. J. Surg.; 154 : 264.

Oren, J. W.; Folse, R.; Kraudel, K.L. and Lewis, D.B. (1986):

Preoperative liver scan and surgical decision-making in patients with CRC. Am. J. Surg.; 151: 452.

Osborne, DR.; Higgins, AF. and Hobbs, KEF. (1978) :

Cryosurgery in the management of rectal tumours. Br. J. Surg.; 65 : 859.

Ovaska, J.; Jarvinen, H.; Kujari, H. and Perttinen, I. (1990):

Follow-up of patients operated on for colorectal carcinoma. Am. J. Surg.; 159 : 593.

Papillon, J. (1982) : (edit)

Rectal and Anal cancers, Springer-Verlag Berlin Heidelberg New York.

Papillon, J. (1984) :

New prospects in the conservative treatment of rectal cancer. Dis colon Rectum.; 27 : 695.

Pemberton, J.H. (1988) :

Colonic absorption. Colon Rectal surgery.; 1 : 89.

Pheils, M.T.; Chapuis, P.H.; Newland, R.C. and Colquhoun, K. (1983):

Local recurrence of following curative resection for carcinoma of the rectum. Dis. Colon. Rectum.; 26: 98.

Phillips, R.; Hihinger, R.; Saunders, V. and Blesovsky, L. (1983) :

Preoperative urography in large bodwel cancer : a useless investigation?

Phillips, R.K.S.; Hihinger, R.; Blesovsky, L.; Fry, J.S., and Fielding, L.P. (1984) :

Local recurrence following "Curative" surgery for large bowel cancer : II. The rectum and rectisigmoid. Br. J. Surg.; 71 : 17.

Phillips, RKS.; Hittinger, R.; Fry, JS. and Fielding, LP. (1985) :

Malignant large bowel obstruction. Br. J. Surg.; 72:296.

Pitluk, H. (1983):

Carcinoma of the colon and rectum in patients less than 40 years of age. surg. Gynaecol. Obstet.; 157: 335.

Polk, H.C. and Spratt, J.S. (1979):

Recurrent colorectal carcinoma. Detection, treatment, and other considerations. Surgery; 69: 9.

Pollard, S.G.; Macfarlane, R. and Everett, W.G. (1989):

Surgery for recrrent colorectal carcinoma: Dis. Colon Rectum.; 30: 879.

Pye, G.; Jackson, J. and Hardcastle, J. (1990):

Comparison of coloscreen self-test and haemoccult faecal occult blood tests in detection of colorectal cancer in asymptomatic patient. Br. J. Surg.; 77:630.

Raftery, T.L. and Samson, N. (1980) :

Carcinoma of the colon : a clinical correlation between presenting symptoms and survival. Ann. Surg. 46 : 600.

Rapaport, A.H. and Burleson, R.C. (1970):

Survival of patients treated with systemic fluorouracil for hepatic metastases. Surg. Gynecol Obstet.; 130:773.

Ravo, B. (1988) :

Colorectal anastomotic healing and intracolonic bypass procedures. Surg. Clin. North Am.; 86 : 1261.

Recalde, M.; Hyolyoke, E.D. and Elias, EG. (1974):

Carcinoma of the colon, rectum, and anal in young patients. Surg. Gynecol Obstet.; 139: 909.

Reilly, J.M.; Ballantyne, G.H.; Fleming, F.X. and Zucker, K.A. (1990):

Evaluation of the occult blood test in screening for colorectal neoplasms. A prospective study using flexible endoscopy. Am. Surg. 56: 119.

Reid, J.D.; Robins, R.E. and Atkinson, K.G. (1984) :

Pelvic recurrence after anterior resection and EEA stapling anastomosis for potentially curable carcinoma of the rectum. Am. J. Surg.; 147 : 629.

Rhodes, J.B.; Holmes, S.F. and Clark, G.M. (1977) :

Changing distribution of primary cancers in the large bowel. JAMA.; 238 : 1641.

Rinner, S. and Tartter, P.I. (1989):

Multivariate analysis of recurrence after anterior resection for colorectal carcinoma. Am. J. Surg.; 157:573.

Rittgers, RA.; Steele, GJr. and Zamcheck, N. (1978):

Transient CEA elevations following resection of colorectal cancer: a limitation in the use of serial CEA levels as an indicator for second-look surgery. J. Natl. Cancer. Inst.; 61: 315.

Rosato, F.E. and Marks, G. (1981):

Changing site distribution patterns of colorectal cancer at Thomas Jefferson University Hospital. Dis. Colon. Rectum; 24: 93.

Rose, R.H.; Axelrod, D.M.; Aldea, P.A. and Beck, A.R. (1988):

Colorectal carcinoma in the literature. Clin. pediatr. (phila). 27 : 105.

Romano, G.; Derosa, P. and Vallone, G. (1985) :

Intrarectal ultrasound and computed tomography in the pre-and postoperative assessment of patients with rectal carcinoma. Br. J. Surg.; 72 : S 117.

Rocklin, MS.; Senagore, AJ. and Talbott, TM. (1991) :

Role of carcinoembryonic antigen and liver function tests in detection of recurrent colorectal carcinoma. Dis. Colon. Rectum.; 34 : 794.

Rozen, P.; Fieman, Z.; Fine, N.; Chetrit, A. and Zajicek, G. (1991) :

Rectal epithelial proliferation in persons post sporadic colorectal neoplasia. Cancer-Lett.; 59 : 1.

Runkel, N.S.; Schlag, P.; Schwartz, V. and Herfarth, C. (1991):

"Outcome after emergency surgery for cancer of the large intestine". Br. J. Surg. 78 : 183.

Sanfelippo, P.M. and Beahrs, O.H. (1972) :

Factors in the prognosis of adenocarcinoma of the colon and rectum. Arch. Surg.; 104 : 401.

Schein, P.H.S. (1985) :

Tumour markers. In : Cecil Text Book of Medicine, 17th. ed., Ch. 15, P. 125. James. B.; Wyngaarden Lloyd H. Smith Jr. Saunders Company.

Schrock, T.R. (1991) :

Cancer of the large intestine. In: Current Surgical Diagnosis and Treatment. 9th. edition.; Ch. 32. P. 633. Way L.W. Appleton and Lange, USA.

Schiessel, R.; Wunderlich, M. and Herbst, F. (1986):

Local recurrence of colorectal cancer; effect of early detection and aggressive surgery. Br. J. Surg. 73: 342-4.

Serpell, J.W.; McDermott, F.T.; Katrivessis, H. and Hughes, E.S.R. (1989) :

Obstructing carcinomas of the colon. Br. J. Surg. 76:965.

Shepherd, J.M. and Jones, J.S.P. (1971):

Adenocarcinoma of the large bowel. Br. J. Cancer, 25:680.

Sherif, M. and Ibrahim, A. (1987):

The Profile of Cancer in Egypt. In: The Cancer Registry for the Metropolitan Cairo Area 1972-1987. P. 89. Arab World Printing House, Cairo, Egypt.

Shimokawa, K. and Inada, K. (1990) :

Advances of immunohistochemistry and its application to histopathology studies on liver metastasis of colorectal carcinoma : venous basement membrane Laminin. Japan. J. Clin. Pathol. 38 : 115.

Shindo, K. (1976):

Significance of schistosoma japonicum in the development of cancer of the large intestine: Report of a case and review of the literature. Dis. Colon. Rectum, 19: 460.

Shiraki, S.; Mori, H. and Kadomoto (1982):

Adjuvant immunotherapy of carcinoma coli with levamisole. Prevention of immune depression following surgical therapy and radiotherapy. Int. J. Immunopharmacol., 4: 73.

Silverberg, E. (1981):

Cancer statistics 1981. New York: American cancer society.

Skandlakis, J.E.; Gray, S.W. and Rowe, J.Jr. (eds.) (1983):

Morphology of The Colon, Rectum, and Anus, Ch. 12. P.231
In: Anatomical Complications in General Surgery. Mc
Graw-Hill Book Company New York.

Skipper, D.; Cooper, A.J.; MKarston, J.E. and Taylor, I. (1987):

Exfoliated cells in vitro growth in colorectal cancer.
Br. J. Surg.; 74 : 1049.

Slaney, G. (1971):

Results of treatment of carcinoma of the colon and
rectum. In: Irvine W.T. (ed.). Modern Trends in Surgery
3. Butterworths; London: P. 9.

**Slater, G.; Greenstein, A.D.; Gelerrnt, I. and Aufses, A.H.
(1985):**

Distribution of colorectal cancer in patients with and
without-ulcerative colitis. Am. J. Surg.; 149: 780.

Sloan, D.A.; Fleiszer, D.M. and Richards, G.K. (1983):

Increased incidence of experimental colon cancer
associated with long-term metronidazole therapy. Am. J.
Surg.; 154: 66.

Smith, D.E.; Mufb, N.S.; Shetabi, H. and Eiseland, H. (1989):

Postoperative Adjuvant chemotherapy and radiotherapy for
cancer of the large bowel and rectum. Am. J. Surg. 157:508.

Smiddy, F.G. and Goligher, J. C. (1957):

Results of surgery in the treatment of cancer of the
large intestine. Br. Med. J. 1: 793.

Sontag, S.J.; Cheifec, G. and Stanley, M.M.. (1985) :

Barrett's oesophagus and colonic tumours.
The lancet (1985).; P. 946.

Speyer, J.L.; Sugarbaker, P.H. and Collins, J.M. (1982):

Portal levels and hepatic clearance of 5 FU after intraperitoneal administration in man. Cancer Res.; 41:1916.

Spicer, S.S. (1965):

Diamine methods for differentiation of mucosubstances histochemically. J. Histochem. Cytochem.; 23: 211.

Spratt, J.S. and Spjut, H.J. (1967):

Prevalence and prognosis of individual clinical and pathologic variables associated with colorectal carcinoma. Cancer; 20: 1976.

Stadler, J.; Holscher, AH. and Adolf, J. (1991) :

Intraoperative ultrasonographic detection of occult liver medtastases in colorectal cancer. Surg. Endosc.; 5: 36.

Stephenson, B.M.; Shandall, A.A.; Farouk, R. and Griffith, G. (1990) :

Malignant left-sided large bowel obstruction managed by subtotal/total colectomy. Br. J. Surg.; 77 : 1098.

Storer, E.H.; Goldberg, S.M. and Navatvongs, S. (1984):

Carcinoma of the Colon and Rectum. In: Principles of Surgery 4th. edition, chap 28, P. 1197. Schwartz, S.I. Shires, G.T. Spencer, F.C. and Storer, E.H. (Eds.), Mc Graw-Hill Book Company, New York.

Sudeck, P. (1907):

Über die Gefassversorgung des Mastdrames in Hinsicht auf die operative Gangran, Muenchen. Med. Wschr.; 54: 1314.
In: Skandalakis, 1983.

Sugarbaker, P.H.; Zamcheck, N. and Moore, F.D. (1976):

Assesement of serial CEA assay in postoperative detection of colorectal cancer. cancer; 38: 2310.

Summerton, J.; Flrin, T.; Cooke, T. and Talor, I. (1983) :

Bile acid receptors in colorectal cancer. Br. J. Surg.;
70 : 549.

Surtees, P.; Ritchie, J.K. and Phillips, R.K.S. (1990) :

High versus low ligation of the inferior mesenteric
artery in rectal cancer. Br. J. Surg.; 77 : 618.

Swift, RI.; Danpure, HJ.; Osman, S. and Gaer, J. (1991) :

Imaging of metastatic colorectal cancer with tumour-
activated killer lymphocytes. Lancet.; 337 : 1511.

Szymendera, J.J.; Nowacki, M.P. and Szawlowski, A.W. (1982) :

The predictive value of plasma CEA levels : preoperative
prognosis and postoperative monitoring of patients with
colorectal carcinoma. Dis. Colon. Rectum.; 25 : 46.

Taylor, I. (1985) :

Colorectal liver metastases-to treat or not to treat.
Br. J. Sutg.; 72 : 511.

Taylor, I. and Northover, J.M. (1990) :

Adjuvant therapy in colorectal cancer : the need for
mega-trial. Br. J. Surg.; 77 : 841.

**Talbot, I.Ch.; Ritchie, Sh.; Leightion, M.H. and Hughes, A.O.
(1981) :**

Spread of rectal cancer within veins : Histologic
features and clinical significance. Am. J. Surg.;141:15.

**Tate, J.J.T.; Tawlinson, J.; Royle, G.T. and Taylor, I.
(1988) :**

Pre-operative or postoperadditive colonic examination for
synchronous lesions in colorectal cancer. Br. J. Surg.;
75 : 1016.

Tachikawa, T.; Watanabe, N. and Soma, S. (1983) :

Cancer of the colon and rectum in persons under forty
year of age. Shokaki Geka.; 6 :649.

Tautu, C.; Pee, D.; Dunsmore, M.; Prokor, JJ. and Alhadebb, JA. (1991) :

Evaluation of serum sialic acid and carcinoembryonic anyigen for the detection of early-stage colorectal cancer. J. Clin. Path. Lab. Anal.; 5 : 247.

Terabe, K.; Sako, T. and Sugimoto, K. (1984):

Clinical and histopathological study on cancer of the colon and rectum in persons under 40 years of age. J. Jpn. Soc. Clin. Surg.; 45: 1574.

Thompson, P.M.; Hill, J.T. and Packham, D.A. (1979):

Colonic carcinoma at the site of uretrosigmoidostomy : What is the risk?. Br. J. Surg. 66: 65.

Tjandra, J.J.; Pietersz, G.A.; Penfold, C. and Sullivan, J.R. (1989):

Phase I in clinical trial of drug-monoclonal antibody conjuges in patients with advanced colonrectal carcinoma: A preliminary report. Surgery; 106: 533.

Tornqvist, A.; Ekelund, G. and Leandeer, L. (1981) :

Early diagnosis of metachronous colorectal carcinoma. Aust N.Z.J. Surg.; 51 : 442.

Tomlinson, E. and McVie (1983):

New directions in cancer chemotherapy. 2 targeting with microspheres. Pharmacy. International; 11: 281.

Turnbull, R.B.; Kyle, K. and Watson, F.R. (1967):

Cancer of the Colon: The influence of the no-touch isolation technic on survival. Ann. Surg.; 166: 420.

Umpleby, HC.; Bristol, JB.; Rainey, JB. and Williamson, RCN. (1984) :

Survival of 727 patients with single carcinoma of the large bowel. Dis. Colon Rectum.; 27 : 803.

Umpleby, H.C.; Ranson, D.L. and Williamson, R.C.N. (1985) :

Peculiarities of mucinous colorectal carcinoma. Br. J. Surg.; 715.

Umpleby, H.C. and Williamson, R.C.N. (1987) :

Anastomotic recurrence in large bowel cancer.
Br. J. Surg.; 74 : 873.

Umpleby, H.C. (1989) :

"The development and treatment of local recurrence in colorectal cancer". In: Progres in Surgery, volume 3 edited by Taylor I., chap. 5, page 59, chirchill Livingstone, Edinburgh, London, Melbourne and New York.

Van de Velde, C.J.H.; Veenhof, A.H.N. and Sugarbaker, P.H. (1984):

Methodology in the clinical study of hepatic metastases.
In: Van de velde CJH, Sugarbaker P.H. (eds.). Liver Metastases. Amsterdam: Martinus Nijhoff; 154.

Vasilevesky, C.A. and Gordon, P.H. (1988) :

Colonoscopy in the follow-up of patients with colorectal carcinoma. Can. J. Surg.; 31 : 188.

Varma, J.S.; Chan, A.C.; Li, M.K. and Li A.K. (1990) :

Low anteriorv resection of the rectum using a double stapling technique. Br. J. Surg.; 77 : 888.

Waldron, R.P.; Donovan, I.A.; Drumm, J.; Mottram, S.N. and Susan Tedman (1986) :

Emergency presentation and mortality from colorectal cancer in the elderly. Br. J. Surg. 73 : 214.

Walton, W.W.; Hagihara, PF. and Griffen, W.O. (1976):

Colorectal adenocarcinoma in patients less than 40 years old. Dis. colon. Rectum; 19: 529.

Walch, N. (1989) :

Cyclophosphamide (C), Oncovin (O), Methotrexate (M), and 5-fluorouracil (F), COMF, in the treatment of advaced colorectal carcinoma. Second international conference on gastrointestinal cancer, Jerusalem, Israes : 47.

- Watson, S.A.; Durrant, L.G. and Morris, D.L. (1988) :**
Growth-promoting action of gastrin on human colonic and gastric tumour cell cultured in vitro. Br. J. Surg.; 75:342.
- Weddle, J.; Meier, Z.P. and Luu, T.H. (1986):**
A retrospective study of serial CEA determinations in the early detection of recurrent colorectal cancer. Dis. Colon. Rectum.; 24: 618.
- Whittaker, M. and Coligher, J.C. (1976) :**
The prognosis after surgical treatment for Carcinoma of the rectum. Br. J. Surg.; 63 : 384.
- Windle, R.; Bell, P.R.F. and Shaw, D. (1987) :**
Five year results of a randomized trial of adjuvant 5-fluorouracil and levamisole in colorectal cancer. Br. J. Surg.; 74 : 569.
- Wolmark, N.; Wieand, H.S. and Rockette, H.E. (1983) :**
The prognostic significance of tumour location and bowel obstruction in Dukes' B and C colorectal cancer. Ann. Surg.; 198 : 743.
- Williams, C. and Slack (1980) :**
A prospective study of sexual function after major colorectal surgery. Br. J. Surg.; 67: 722.
- Williams, N.S.; Durdey, P. and Johnston, D. (1985) :**
The outcome following sphincter saving resection and abdominoperineal resection for low rectal cancer. Br. J. Surg.; 72 : 595.
- Williams, N.S.; Jass, J.R. and Hardcastle, J.D. (1988) :**
Clinicopathological assessment and staging of colorectal cancer. Br. J. Surg.; 75 : 649.
- Williamson, R.C.N.; Davies, P.W.; Bristol, J.B. and Wells, M. (1982) :**
Intestinal adaptation and experimental carcinogenesis after partial colectomy. Gut.; 23 : 316.

Wiggers, T.; Jeekel, J.; Arends, J.W., Brinkhorst, A.P. and et al. (1988) :

No-Touch isolation technique in colon cancer: a controlled prospective trial. Br. J. Surg.; 75 : 409.

Wood, C.B.; Ratclibbe, J.G.; Zirt, R.W. and Malcolm, A.J. (1980) :

The clinical significance of the pattern of elevated serum carcinoembryonic antigen levels in recurrent colorectal cancer. Br. J. Surg.; 67 : 46.

Wood, C.B.; Gillis, C.R.; Hole, D.; Malcolm, A.J. and Blumgart, L.H. (1981) :

Local tumour invasion as a prognostic factor in colorectal cancer. Br. J. Surg.; 68 : 326.

Wohlleb, J.C.; Hunter, C.F.; Blass, B. and Kadlubar (1990):

Aromatic amine acetyltransferase as a marker for colorectal cancer : enviromental and demographic association. Int. J. Cancer.; 46 : 22.

Wright, H.K. and Higgins, E.F. (1982) :

Natural history of occult right colon cancer. Am. J. Surg.; 143 : 169.

Yamada, S.; Takahashi, T. and Hayashi, A. (1972) :

Carcinoma of the colon and rectum in the young. Stomach intest.; 7 : 881.

Yeatmen, T.J. and Bland, K.I. (1989) :

Sphincter-Saving procedures for distal cancer of rectum. Ann. Surg.; 209 : 1.

Zaky, S.A. and Hashem, M. (1962):

Distribution of bilharsial lesions and complications in various organs: a study of 1220 autopsies of bilharsial cases. Gas. Kasr. El-Ainty Fac. Med. 30: 15.