

REFERENCES

REFERENCES

- Abbott, R.D.; Wilson, P.W.F.; Kannel, W.B., et al. (1988):
High density lipoprotein cholesterol, total cholesterol and myocardial infarction. The Framingham study. Arteriosclerosis 18:207.
- Abenavoli, T.A.; Rubler, S.; Fisher, V.J.; Axelrod, H.I. and Zuckerman, K.P., (1981):
Exercise testing with myocardial scintigraphy in asymptomatic diabetic males. Circulation 63:54.
- Adner, M.M. and Castelli, W.P.; (1980):
Elevated high-density lipoprotein levels in marathon runners. J.A.M.A., 243:534.
- Albers, J.J.; Cheurg, M.C.; and Hazard, W.R. (1978):
High density lipoproteins in myocardial infarction survivors. Metabolism 27:479.
- Armstrong, M.L.; Warner, E.D. and Conner, W.E. (1970):
Regression of coronary atheromatosis in rhesus monkeys. Circ..Res., 27:59.
- Asher, P. (1966):
Fat babies and fat children. Arch. Dis. Child., 41:672.
- Astrup, P.; Kjeldsen, K. and Wanstrub, J. (1967):
Enhancing influence of carbon monoxide on the development of atheromatosis in cholesterol FEO rabbits. J. Atheroscler. Res., 7:343.
- Augustin, J.; Freeze, H.; Tejada, P. and Brown, W.V. (1978):
A comparison of molecular properties of hepatic triglyceride lipase and LPL from human post-heparin plasma. J.Biol. Chem., 253(9):2912-20.

Avogaro, P.; Bon, G.B.; Chzolato, G. et al. (1978):

Variation in apolipoprotein B and A₁ during the course of myocardial infraction.
Eur. J. Clin. Invest., 8:121-129.

Avogaro, P.; Bittolo Bon, G.; Gazzolato, G.; and Quinci, G.B. (1979):

Are apolipoproteins better discriminators than lipids for atherosclerosis?
Lancet 1:901-3.

Ayman, D. (1934):

Heredity in arteriolar (essential) hypertension : A clinical study of blood pressure in 1,524 members of 277 families.
Arch. Intern. Med., 53:792.

Baker, T.R.; Metcalf, P.A.; Holdaway, I.M; and Johnson, R.N. (1985):

Serum fructosamine concentration as a measure of blood glucose control in type I (insulin dependent) diabetes mellitus.
Br. Med. J., 290:352-355.

Basu, S.K.; Ho, Y.K.; Brown, M.S. et al. (1982):

Biochemical and genetic studies of the apoprotein E secreted by mouse macrophages and human monocytes.
J. Biol. Chem., 257(16):9788-95.

Beaty, T.H.; Kwiterovich, J.R.; Po, Koury, M.J. et al. (1986):

Genetic analysis of plasma sitosterol, apoprotein B and lipoproteins in a large Amish pedigree with sitosterolemia.
Am. J. Human Genet., 38:492.

Bengtsson, G. and Olivecrona, T. (1979):

Apolipoprotein CII enhances hydrolysis of monoglycerides by lipoprotein lipase, but the effect is abolished by fatty acids.
FEBS LETT., 16(2):345-8.

Berg, K.; Borresen, A.L. and Dahlen, G. (1976):

Serum HDL and atherosclerotic heart disease
Lancet 1:499.

Berenson, G.S.; Blonde, C.V.; Farris, R.P. et al. (1979-a):

Cardiovascular disease risk factor variables during the first year of life.
Am. J. Dis. Child., 10:1049-57.

Berenson, G.S.; Srinivasan, S.R. and Frerichs, R.R. (1979-b):

Serum high density lipoprotein and its relationship to cardiovascular disease risk factor variables in children. The Bogalusa Heart Study
Lipids 14:91.

Biss, K. Ho K.J.; Mikkelsen, B.; Lewis, L. and Taylor, C.B. (1971):

Some unique biological characteristics of the Masai of East Africa.
N. Engl. J. Med., 284:694.

Bjorkem, I. and Skreda, S. (1989):

Familial diseases with storage of sterols other than cholesterol. In Scriver CR., Beaudet AL, Sly WS., and Valle D. (eds): The Metabolic Basis of Inherited disease, 6th ed.
New York McGraw-Hill, 1:1283.

Blankenhorn, D.H.; Nessim, S.A.; Johnson, R.L. et al. (1987):

Beneficial effect of combined colestipol-niacin therapy on coronary atherosclerosis and coronary venous bypass grafts.
J.A.M.A., 257:3233.

Bliden, L.C.; Dsnick, R.; Carter, J. and Moller, J.H. (1974):

Cardiac involvement in inherited disorders of metabolism.
Progr. Cardiovas. Dis., 10:615.

Bloor, C.M. (1969):

Hereditary aspects of myocardial infarction.
Circulation 39 and 40 (Suppl.IV) :130.

Blum, C.B. (1980):

Dynamics of apolipoprotein E metabolism in humans.
Circ., 62:11-194.

Blumenthal, S. and Jesse, M.J. (1975):

Risk factors for coronary heart disease in children of affected families.
J. Pediat. 57:1187.

Breckenridge, W.C.; Little, J.A.; Steiner, G. et al. (1978):

Hypertriglyceridemia associated with deficiency of apolipoprotein CII.
N. Engl. J. Med., 298 (23):1265-73.

Breckenridge, W.C.; Little, J.A.; Alauryan, P. et al. (1982):

Lipoprotein abnormalities associated with familial deficiency of hepatic lipase.
Atherosclerosis 45:161.

Breslow, J.L. (1989):

Familial disorders of high density lipoprotein metabolism. In Scriver CR, Beaudet AL, Sly WS and Valle D (eds): The Metabolic Basis of Inherited disease, 6th ed New York McGraw-Hill, 1:1251.

Brewer, H.B. (1983):

Type III hyperlipoproteinemia : diagnosis, molecular defect, pathology and treatment.
Ann. Intern. Med., 98:623-40.

Brewer, H.B.; Shulmar, R.S.; Herber, P. et al. (1974):

The complete amino acid sequence of alanine apolipoprotein (apo CIII), an apolipoprotein from human plasma VLDL.
J. Biol. Chem., 249:4975-84.

Brook, C.G.D.; Lloyd, J.K. and Wolf, O.H. (1972):

Relation between age of onset of obesity and size and number of adipose cells.
Br. Med. J., 2:25.

Brown, W.V. and Baginsky, M.L. (1972):

Inhibition of lipoprotein lipase by an apoprotein of human VLDL.
Biochem., Biophys. Res. Commun., 46:375-382.

Brown, M.S.; Kovanen, P.T.; Goldstein, J.L. et al. (1978):

Prenatal diagnosis of homozygous familial hypercholesterolemia. Expression of a genetic receptor disease in vitro.
Lancet 1:526.

Brunzell, J.D. (1989):

Familial lipoprotein lipase deficiency and other causes of the chylomicronemia syndrome. In Scriver CR, Beaudet AL, Sly WS and Valle D (eds): The Metabolic Basis of Inherited Disease 6th ed.
New York McGraw-Hill, 1:1165

Brunzell, J.D.; Albers, J.J.; Chait, A. et al. (1983):

Plasma lipoproteins in familial combined hyperlipidemia and monogenic familial hypertriglyceridemia.
J. lipid Res., 24:147.

Brunzell, J.D.; Porte, D. and Bierman, E. (1975):

Reversible abnormalities in post heparin lipolytic activity during the late phase of release in diabetes mellitus.
Metabolism 24:1123-1137.

Brunzell, J.D.; Chait, A. and Bierman, E.L. (1978):

Pathophysiology of lipoprotein transport.
Metabolism 27 (9):1109-1127.

Brunzell, J.D.; Sniderman, A.D.; Albers, J.J. et al. (1984):

Apoprotein B and A-1 and coronary artery disease in humans.
Arteriosclerosis 4:79.

Buell, J.C. and Elliot, R.S. (1979):

Stress and cardiovascular disease.
Mod. Concepts Cardiovas. Dis., 43:49.

Bunn, H.F.; Gabbay, K.H. and Gallop, P.M. (1978):

The glycosylation of haemoglobin relevance to diabetes mellitus.
Science 200:21.

Carlson, L.A. and Ericsson, M. (1975):

Quantitative and qualitative serum lipoprotein analysis 11. Studies in male survivors of myocardial infarction.
Atherosclerosis 21:335-450.

Carroll, K.K. (1977):

The role of dietary protein in hypercholesterolemia and atherosclerosis. Symposium. Lipid therapy of atherosclerosis, AOCs 1977 Annual Meeting, New York New York, 10, 1977.

Chait, A.; Albers, J.J. and Brunzell, J.D. (1980):

Very low density lipoprotein overproduction in genetic forms of hypertriglyceridemia.
Eur J. Clin. Invest., 10:17.

Charney, E.; Goodman, H.C.; McBride, M.; Lyon, B. and Pratt, R. (1976):

Childhood antecedents of adult obesity. Do chubby infants become obese adults?
N. Engl. J. Med., 295:6.

Cheung, M.C. and Albers, J.J. (1977):

The measurement of apolipoprotein A-1 and A-II levels in men and women by immunoassay.
J. Clin. Invest., 60:43.

Chung, H.; Abano, D.A.; Fless, G.M. and Scanu, A.M. (1979):

Isolation, properties and mechanisms of in vitro action of LCAT from human plasma.
J. Biol. Chem., 254 (5):7456-64.

Cofman, J.W.; Jones, H.B.; Lindgren, F.T.; Lyon, T.P.; Elliott, A.H. and Strisower, B. (1950):

Blood lipids and human atherosclerosis
Circulation 2:161.

Cortner, J.A.; Coates, P.M. and Gallagher, P.R. (1990):

Prevalence and expression of familial combined hyperlipidemia in childhood.
J. Pediatr., 116:514.

Cox, D.W.; Wills, D.E.; Evan, F. et al. (1988):

A deletion of one nucleotide results in functional deficiency of apolipoproteins C-II (apo C-II).
Toronto. J.Med. Genet., 25:649.

Cryer, P.E.; Morey, W.H. and Santiago, J.V. (1976):

Norepinephrine and epinephrine release and adrenergic mediation of smoking associated hemodynamic and metabolic events.
N. Engl. J., Med., 295:573.

Darmady, J.M.; Fosbrooke, A.S. and Lloyd, J.K. (1972):

Prospective study of serum cholesterol levels during first year of life.
Br. Med. J., 2:685-8.

Davis, O. and Klainer, M.J. (1940):

Studies in hypertensive heart disease. The incidence of coronary atherosclerosis in cases of essential hypertension.
AM. HEART J., 19:185.

Dennison, B.A.; Kikucki, D.A.; Srinivasan, S.R. et al. (1990):

Serum total cholesterol screening for the detection of elevated low-density lipoprotein in children and adolescents: The Bogalusa Heart Study.
Pediatrics 85:472

Dolphin, P.J. (1985):

Lipoprotein metabolism and the role of apolipoproteins as metabolic programmers.
Can. J. Biochem. Cell. Biol. 63, 850-869.

Elkeles, R.S. (1974):

The role of heparin in LPL activity.
Horm. Metab. Res., 6:151-4.

El-Maghrabi, v.R.; Waite, M.; Rudel, L.L. and King, V.L. (1979):

Metabolism of lipoprotein acylglycerols by liver parenchymal cells.
Biochim. Biophys. Acta 572(1):52-63.

Fielding, C.J. and Havel, R.J. (1977):

Lipoprotein lipase.
Arch. Pathol., 101:225-9.

Fielding, C.J.; Shere, V.G. and Fielding, P.E. (1972):
A protein cofactor of LCAT.
Biochem. Biophys. Res. Commun., 46:1493-8.

Flegg, H.M.; (1973):

An investigation of the determination of serum
cholesterol by an enzymatic method.
Ann. Clin. Biochem., 10, 79-84.

Fojo, S.S.; Beisregal, U.; Beil, U. et al. (1988):

Donor splice site mutation in the apolipoprotein
(apo) C-II gene (apo C-II hornburg) of a patient with
apo C-II deficiency.
J. Clin. Invest., 82:1489.

Fomon, S.J. (1974):

Infant nutrition. 2nd ed.
Philadelphia: WB Sanders, 174-5.

Fredrickson, D.S. (1972):

A physician's guide to hyperlipidemia Mod Concepts
Cardiovasc. Dis., 41:31.

Fredrickson, D.S. and Gordon, R.S. (1958):

Transport of fatty acids.
Physiol. Rev., 38:585.

Fredrickson, D.S.; Levy, R.I. and Bonnell, M. (1974):

The Dietary Management of Hyperlipoproteinemia.
DHEW Publication (NIH), National Heart and Lung
Institutes of Health, PP. 76.

Friedman, G. and Goldberg, S.J. (1975):

Concurrent and subsequent serum cholesterol of
breast-and formula-fed infants.
Am. J. Clin. Nutr., 28:42-5.

Froelicher, V.F. (1972):

Animal studies of effect of chronic exercise on the
heart and atherosclerosis. A review.
Am. HEART J., 84:496.

- Fruchart, J.C. (1982):
Rev. Fr. des. Laboratories, (103), 7:17.
- Gabr, M.; Hashem, N.; Hashem, M.; Fahmi, A. and Safouh, M. (1960):
Progeria: A pathologic study.
J. Pediatr., 57:70.
- Glomset, J.A. (1980):
HDL in human health and disease.
Adv. Int. Med., 25:91-116.
- Glueck, C.J.; Gartside, P.; Fallat, R.W. et al. (1976):
Longevity syndromes: Familial hypobeta and familial hyperalphalipoproteinemia.
J. Lab. Clin. Med., 88:941.
- Glueck, C.J.; Mellies, M.J.; Srivastava, L. et al. (1977):
Insulin, obesity and triglyceride interrelationships
In sixteen children with familial hypertriglyceridemia
Pediatr. Res., 11:13.
- Glueck, C.J. and Tsang, R.C. (1972):
Pediatric familial type II hyperlipoproteinemia
effects of diet on plasma cholesterol in the first
year of life.
Am. J. Clin. Nutr., 25:224-30.
- Glueck, C.J.; Tsang, W.B.; Fallat, R. et al. (1972):
Plasma and dietary cholesterol in infancy : effects
of early low or moderate dietary cholesterol intake
on subsequent response to increased dietary cholesterol.
Metabolism 21:1181-92.
- Gofman, J.W.; Glazier, F.; Tamplin, A. et al. (1954):
Lipoproteins, coronary artery disease and atherosclerosis.
Physiol. Rev., 34:589-607.

Goldstein, J.L. (1983):

The hyperlipoproteinemias and other disorders of lipid metabolism. In: Harrison's principles of internal medicine. 10th edition, edited by Petersdorf, R.G., Adams, R.D. Braunwald, E., Isselbacher, K.J.; Martin, J.B. and Wilson, J.D.

Goldstein, J.L. and Brown, M.S. (1989):

Familial hypercholesterolemia. In Scriver CR, Beaudet AL, Sly WS, and Valle D (eds): The Metabolic Basis of Inherited Disease, 6th ed. New York McGraw-Hill 1:1215.

Goldstein, J.L.; Schrott, H.G.; Hazzard, W.R. et al. (1973):

Hyperlipidemia in coronary heart disease. 11 Genetic analysis of lipid levels in 176 families and delineation of a new inherited disorder combined hyperlipidemia. J. Clin. Invest., 52:1544.

Gordon, T.; Castelli, W.P.; Dawber, T.R.; Hjortland, M.C. and Kannel, W.B. (1977):

Diabetes, blood lipids and the role of obesity in coronary heart disease risk for women. The Framingham study. Ann. Intern. Med., 37:393.

Granot, E. and Deckelbaum, R.J. (1989):

Hypercholesterolemia in childhood. J. Pediatrics, 115:171.

Green, P.H. and Glickman, R.M. (1981):

Intestinal lipoprotein metabolism. J. Lipid. Res., 22:1953-73.

Hamilton, R.L. (1972):

Synthesis and secretion of plasma lipoproteins. In "Pharmacological control of lipid metabolism". New York, Plenum Press, Ch., 26, P., 7 - 24.

Havel, R.J. (1978):

Origin of HDL. In: HDL and atherosclerosis, Gott, A.M. et al. (eds.). El Sevier, Amsterdam. PP. 21-35.

Havel, R.J. (1982):

Approach to the patient with hyperlipidemia.
Med. Clin. North Am., 66(2):319-33.

Havel, R.J. Kane, J.P. and Kashyap, M.L. (1973):

Interchange of apolipoproteins between chylomicrons and high density lipoprotein during alimentary lipemia in man
J. Clin. Invest., 52:32-38.

Havel, R.J. Goldstein, J.L. and Brown, M.S. (1980):

Lipoproteins and lipid transport. In: Metabolic control and disease, Bondy, P.K. et al. (eds.) Philadelphia, W.B. Saunders Co., 8th edi., PP. 393-494.

Herbert, P.N.; Gotto, A.M. and Fredrickson, D.S. (1978):

Metabolic Basis of Inherited Disease (Stanbury, J.S., Wyngaarden, J.B. and Fredrickson, D.S., eds.) 4th ed., PP 544-488. McGraw-Hill New York.

Hobbs, H.A.; Leitersdorf, E.; Goldstein, J.L. et al. (1988):

Multiple CVM-mutations in familial hypercholesterolemia. Evidence for 13 allele, including for deletions.
J. Clin. Invest., 81:909.

Hodge, J.E. (1955):

The Amadori rearrangement.
Advances in carbohydrate chemistry, 10:169-205.

Hodgson, P.A.; Ellefson, R.D.; Elveback, L.R. et al. (1976):

Comparison of serum cholesterol in children fed high, moderate or low cholesterol milk diets during neonatal period.
Metabolism 25:739-46.

Hollander, W.; Madoff, I. and Paddock, J. (1976):

Aggravation of atherosclerosis by hypertension in subhuman primate models with coarctation of the aorta.
Cir. Res., 38:63.

Hulley, S.B.; Rosenman, R.H.; Bawol, R.D. and Brand, R.J. (1980):

Epidemiology as a guide to clinical decisions. The association between triglyceride and coronary heart disease.

N. Engl. J. Med., 302:1383-9.

Innerarity, T.L. and Mahley, R.W. (1978):

Enhanced binding by cultured human fibroblasts of apo E containing lipoproteins as compared with LDL. Biochemistry 17:1440-7.

Innerarity, T.L.; Weisgraber, K.F.; Arnold, K.S. et al. (1987):

Familial defective apolipoprotein B-100: Low density lipoproteins with abnormal receptor binding. Proc. Natl. Acad. Sci. USA 84:6919.

Jackson, R.L.; Baker, H.N.; Gillian, E.B. et al. (1977):

Primary structure of VLDL apolipoprotein C II of human plasma.

Proc. Natl. Acad. Sci. (U.S.A.) 74:1942-5.

Jahn, C.E.; Osborne, J.G.; Schaefer, E.J. et al. (1981):

In vitro activation of the enzyme activity of hepatic lipase by apo A-II. FEBS Lett., 131:366-8.

Jansen, H.; Van Berkel, T.J. and Hulsmann, W.C. (1978):

Binding of liver lipase to parenchymal and non parenchymal rat liver cells.

Biochem. Biophys. Res. Commun., 85(1):148-52.

Jenkins, C.D.; Rosenman, R.H. and Friedman, M. (1967):

Development of an objective psychological test for the determination of the coronary prone behavior pattern in employed men. J. Chronic. Dis., 20:371.

Johnson, R.N.; Metcalf, P.A. and Baker, J.R. (1983):

Fructosamine a new approach to the estimation of serum glycosyl protein-An index of diabetic control. Clin. Chim. Acta., 127:87-95.

Kane, J.P.; Hardman, D.A. and Paulus, H.E. (1980):

Heterogeneity of apolipoprotein B: Isolation of a new species from human chylomicrons.
Proc. Natl. Acad. Sci. (U.S.A.) 77:2465-9.

Kane, J.P. and Havel, R.F. (1989):

Disorders of the biogenesis and secretion of lipoproteins containing the B apolipoproteins. In Scriver CR, Beaudet AL, Sly WS, and Valle D (eds.) : The Metabolic Basis of Inherited Disease, 6th ed. New York McGraw Hill, 1:1139.

Kannel, W.B. and Dawber, T.R. (1972):

Atherosclerosis as a pediatric problem.
J. Pediatr., 80:544-554.

Karathanasis, S.K.; Ferris, E. and Haddad, I.A. (1987):

DNA inversion within the apolipoprotein A-I (C-III and A-II) encoding gene cluster of certain patients with premature atherosclerosis.
Proc. Natl. Acad. Sci., USA 84:7198.

Keys, A.; Grande, F. and Anderson, J.T. (1974):

Bias and misrepresentation revisited: Prospective on saturated fats.
Am. J. Clin. Nutr., 27:188.

Kilcoyne, M.; Ritcher, R. and Alsup, P. (1974):

Adolescent hypertension I. Detection and prevalence.
Circulation 50:758.

Kinnunen, P.K.J.; Jackson, R.L.; Smith, L.C. et al. (1977):

Activation of lipoprotein lipase by native and synthetic fragments of human plasma apolipoprotein CII.
Proc. Natl. Acad. Sci. U.S.A., 74:4848-51.

Klotz, O. and Manning, M.F. (1971):

Fatty streaks in the intima of arteries
J. Pathol. Bacteriol., 16:211.

Koizummi, J.; Mabuchi, H.; Yoshimura, A. et al. (1985):

Deficiency of serum cholesteryl ester transfer activity in patient with familial hyperalphalipoproteinemia
Atherosclerosis 58:175.

Kostner, G. and Holasek, A. (1972):

Characterization and quantitation of apolipoproteins from human chylomicrons.
Biochemistry 1:1217-23.

Kreisberg, R.A. (1983):

Lipids, lipoproteins, apolipoproteins and atherosclerosis.
Ann. Intern. Med., 99(5):713-5.

Kussi, T.; Kinnunen, P.K.J.; Eihnholm, C. et al. (1979):

A simple purification procedure for rat hepatic lipase.
FEBS Lett, 98(2):384-8.

Kwiterovich, JR. PO. (1986):

Biochemical, clinical, epidemiologic, genetic and pathologic data in the pediatric age group relevant to the cholesterol hypothesis.
Pediatr., 78:349.

Kwiterovich, JR. PO. (1988):

Hyperapo B: A pleiotropic phenotype characterization by dense low-density lipoproteins and associated with coronary artery disease.
Clin. Chem., 34:B71.

Kwiterovich, JR. PO. (1989):

Pediatric implications of heterozygous familial hypercholesterolemia. Screening and dietary treatment.
Arteriosclerosis 9:1-111.

Kwiterovich, JR. PO. (1990):

Diagnosis and Management of familial dyslipoproteinemia in children and adolescents.
Ped. Clin. Nor. America Vol. 37, No 6 P. 1489:1522.

Kwiterovich, P.O.; Levy, R.I. and Frederickson, D.S. (1973):

Neonatal diagnosis of familial type II hyperlipoproteinemia. Lancet 1:118.

Kwiterovich, P.O.; Fredrickson, D.S. and Levy, R.I. (1974):

Familial hypercholesterolemia (one form of familial type II hyperlipoproteinemia): A study of its biochemical, genetic and clinical presentation in childhood, J. Clin. Invest., 53:1237.

Kwiterovich, P.O.; Farah, J.R.; Brown, M.V. et al. (1977):

The biochemical, clinical and familial presentation of type V hyperlipoproteinemia in childhood. Pediatr., 59:513.

Kwiterovich, P.O.; Keiss, G.; Johnson, N. et al. (1982):

Assessment of plasma total cholesterol as a test to detect elevated low density (beta) lipoprotein cholesterol levels (type IIa hyperlipoproteinemia) in young subjects from a population-based sample. Am. J. Epidemiol., 115:192.

Kwiterovich, P.O.; White, S.; Forte, T. et al. (1987):

Hyperapobetalipoproteinemia in a kindred with familial combined hyperlipidemia and familial hypercholesterolemia. Arteriosclerosis 7:211.

Lackner, K.J. and Schmitz, G. (1990):

Familial high density lipoprotein deficiency with xanthomas is caused by a point mutation with apolipoprotein A-1 gene. Internal Atherosclerosis Congress, Eur. Atherosclerosis Society (EAS) Vienna, Austria.

Larosa, J.C.; Levy, R.I.; Herbert, P.M. et al. (1970):

A specific apoprotein activator of lipoprotein lipase. Biochem. Biophys. Res. Commun., 46:375-81.

Lewis, L.A. and Opplot, J.J. (EDS.) (1980):

Handbook of electrophoresis, volume 1, lipoproteinemia, basic principles and concepts, Boca Raton fla. CRC press Inc. Quoted from: Clinical chemistry, theory, analysis and correlation. Edited by Kaplan, L.A. and Pesce. A.J.

Londe, S. and Goldring, D. (1976):

High blood pressure in children: Problems and guidelines for evaluation and treatment. Am. J. Cardiol., 37:650.

Mahley, R.W. and Rall, J.R. Sc. (1989):

Type III hyperlipoproteinemia (Dysbetalipoproteinemia): The role of apolipoprotein E in normal and abnormal lipoprotein metabolism. In Scriver CR. Beaudet AL. Sly WS and valle D (eds.): The metabolic Basis of Inherited Disease, 6th edt. New York, McGraw-Hill 1-1195.

Mahley, R.W.; Hui, D.Y.; Innerarity, T.L. et al. (1981):

Two independent lipoprotein receptors on hepatic membranes of the dog, swine and man: The apo E and apo B receptors. J. Clin. Invest., 68:1197-206.

Manninen, V.; Rissunan, V.T. and Halcoen, P.J. (1970):

Coronary ostium outside the aortic sinus. A factor in the etiology of this ischemic heart disease. Adv. Cardiol., 4:94.

McGill, H.C. (1968):

Introduction to the geographic pathology of atherosclerosis. Lab. Invest., 18:465.

McClean, J.W.; Tomlinson, J.E.; Kuang, W.J. et al. (1987):

cDNA sequence of human apolipoprotein (a) in homologous to plasminogen. Nature (lond.) 330:132.

Miller, A.L. and Smith, L.C. (1973):

Activation of lipoprotein lipase by apolipoprotein glutamic acid. J. Biol. Chem., 248(9):3359-62.

- Montenegro, M.R. and Solberg, L.A. (1968):
Obesity, body weight, body length, and atherosclerosis.
Lab. Invest., 18:594-603.
- Morganroth, J.; Levy, R.I.; McMahan, A.E. et al. (1974):
Pseudohomozygous type II hyperlipoproteinemia.
J. Pediatr., 85:639.
- Myant, N.B. (1982):
Cholesterol transport through the plasma.
Clin. Sci., 62:261-71.
- Naito, H.K. (1984):
Disorders of lipid metabolism In: Clinical chemistry, theory, analysis and correlation by Kaplan, L.A. and Pesce, A.J. published by the C.V. Mosby Co., St. Louis, Toronto, Princeton.
- Nakaya, Y.; Schaefer, E.J. and Brewer, H.B. (1980):
Activation of human post-heparin lipoprotein lipase by apolipoprotein H (B2-glycoprotein 1).
Biochem. Biophys. Res. Commun., 95:1168-1172.
- Neuman, W.P.; Freedman, D.S.; Voors, A.W. et al. (1986):
Serum lipoproteins and systolic blood pressure are related to atherosclerosis in early life: The Bogalusa Heart Study.
N. Engl. J. Med., 314:138.
- Nilsson-Ehle, P.; Carafinkel, A.S. and Schotz, M.C. (1980):
Lipolytic enzymes and plasma lipoprotein metabolism.
Ann. Rev. Biochem., 49:67-93.
- Olivecrona, T.; Bengtsson, G.; Marklund, S. et al. (1977):
Heparin-lipoprotein lipase interactions.
Ped. Proc., 36:60-5.
- Oncley, J.L. (1958):
Plasma lipoproteins. In: Chemistry of lipids as related to atherosclerosis (Symposium). Edited by I.H. Page, Springfield, 111:Thomas, PP. 114-33.

Osion, R.E. and Vester, J.W. (1960):

Nutrition-endocrine interrelationships in the control of fat transport in man.
Phys. Rev., 40:673-733.

Paffenbarger, R.S. and Hale, W.E. (1975):

Work activity and coronary heart mortality.
N. Engl. J. Med., 292:545.

Paffenbarger, R.S.; Wing, A.L. and Hyde, R.T. (1968):

Physical activity as an index of heart attack in college alumni.
Am. J. Epidemiol., 108:161.

Poskitt, E.M.E. and COLE, T.J. (1977):

Do fat babies stay fat?
Br. Med. J., 1:7.

Reckless, J.P.D.; Stock, J. and Holdsworth, G. (1982):

The role of c-apolipoproteins in the metabolism of triglyceride-rich lipoproteins.
Clin. Sci., 62:125-9.

Rhoads, G.G.; Gulbrandsen, C.L. and Kagan, A. (1976):

Serum lipoproteins and coronary heart disease in a population study of Hawaiian-Japanese men.
N. Engl. J. Med., 294:293.

Roberts, W.C. and Crall, F.V. J.R. (1977):

Extramural and intramural coronary arteries in juvenile diabetes mellitus (Abstr).
Circulation 56 (Suppl. 111):111-20.

Sabesin, S.M.; Bertram, P.D. and Freeman, M.R. (1980):

Lipoprotein metabolism in liver disease.
Adv. Int. Med., 25:117.

Schaefer, E.J.; Blum, C.B.; Levy, R.I. et al. (1978):

Metabolism of high density lipoprotein, apolipoproteins in Tangier disease.
N. Engl. J. Med., 299:905.

Schmitz, G.; Assmann, C.; Robenak, H. et al. (1985):

Tangier disease: A disorder of intracellular membrane traffic.

Proc. Natl. Acad. Sci., USA 82:6305.

Schrott, H.G.; Conner, W.E.; Lauer, R.M.; Clarke, W.R. and Wiebe, D.E. (1979):

Increased coronary mortality in relatives of hypercholesterolemic school children.
Circulation 59:320.

Scow, R.O.; Blanchette-Mackie, E.J. and Smith, L.C. (1976):

Role of capillary endothelium in the clearance of chylomicrons. A model for lipid transport for blood by lateral diffusion in cell membranes.
Cir. Rec., 39:149-62.

Shekelle, R.B.; Shyrock, A.M.; Paul, O.; Lepper, M.; Stamler, J.; Liu, S. and Raynor, W.J. (1981):

Diet, serum cholesterol and death from coronary heart disease. The western electric study.
N. Engl. J. Med., 304:65.

Shellbourn, F.; Hanks, J.; Meyers, W. et al. (1980):

Effect of apoproteins on hepatic uptake of triglyceride emulsions in the rat.
J. Clin. Invest., 65:652-8.

Sherrill, B.C.; Innerarity, T.L. and Mahley, R.W. (1981):

Rapid hepatic clearance of the canine lipoproteins containing only the E apoprotein by a high affinity receptor.
J. Biol. Chem., 255(5):1804-7.

Sniderman, A. and Kwiterovich, JR. PO. (1987):

Hyperapobetalipoproteinemia and LDL and HDL2 heterogeneity. Proceedings of the workshop on lipoprotein heterogeneity US Department of Health and Human services, VIH publication No. 87-2646, P 293.

Sniderman, A.; Shapiro, S.; Marpole, D. et al. (1980):

The association of coronary atherosclerosis and hyperapobetalipoproteinemia (increased protein but normal cholesterol content in human plasma low density lipoprotein).
Proc. Natl. Acad. Sci., USA 97:604.

Sniderman, A.; Wolfson, C.; Teng, B. et al. (1982):

Association of hyperapobetalipoproteinemia with endogenous hypertriglyceridemia and atherosclerosis.
Ann. Intern. Med., 97:833.

Sniderman, A.D.; Teng, B.; Genest, J. et al. (1985):

Familial aggregation and early expression of hyperapobetalipoproteinemia.
Am. J. Card., 55:291.

Soria, L.F.; Ludwig, E.A.; Clarke, R.G. et al. (1989):

Association between a specific apolipoprotein B mutation and familial defective apolipoprotein B-100.
Proc. Natl. Acad. Sci., USA 86:587.

Soutar, A.K.; Garner, C.W.; Baker, H.N. et al. (1975):

Effect of human plasma apolipoproteins and phosphatidyl choline acyl donor on the activity of LCAT.
Biochemistry 14:3057-64.

Sprecher, D.S.; Schaefer, E.J.; Kent, K.M. et al. (1984):

Cardiovascular features of homozygous familial hypercholesterolemia: Analysis of 16 patients
Am. J. Cardio., 54:20.

Srinivasan, S.R.; Ellefson, R.D.; Whitaker, C.F. et al. (1978):

Lipoprotein cholesterol in the serum of children as determined independently by two different methods.
Clin. Chem., 21:157-161.

Stafford, J.L. (1968):

Tomorrow's laboratory.
Proc. Roy. Soc. Med., 61:1047.

Stamler, J. (1973):

Epidemiology of coronary heart disease.
Med., Clin. North Am., 57:5.

Stamler, J.; Lindberg, H.A.; Berkson, D.M.; Shaffer, A.;
Miller, W. and Poindexter, A. (1960):

Prevalence and incidence of coronary heart disease in
strata of the labor force of Chicago industrial
corporation
J. Chronic Dis., 11:405.

Stamler, R.; Stamler, J. and Riedlinger, W.F. (1978):

Weight and blood pressure.
J.A.M.A., 240:1607.

Stocks, J.; Holdsworth, G. and Galton, D. (1979):

Hypertriglyceridemia associated with an abnormal
triglyceride-rich lipoprotein carrying excess
apolipoprotein C-111.
Lancet 2:667-71.

Strong, J.P. and McGill, H.C. J.R. (1962):

The natural history of coronary atherosclerosis.
Am. J. Pathol., 40-43.

Strong, J.P. and McGill, JR H.C. (1969):

The pediatric aspects of atherosclerosis.
J. Atheroscler. Res., 9:251.

Surgeon General (1979):

Smoking and health. Department of Health, Education
and Welfare, Publication No. (DHS), 50066, U.S.
Government, Printing Offices Washington, D.C.

Tall, A.R. and Small, D.M. (1978):

Plasma high density lipoproteins.
N. Engl. J. Med., 299:1232.

Taylor, C.B.; Allen, E.S. and Mikkelsen, B. (1976):

Serum cholesterol levels of seventh-day adventists.
Paroi Arterielle/Arterial wall 3:175.

Teng, B.; Thompson, G.R.; Sniderman, A.D. et al. (1983):

Composition and distribution low density lipo-protein fractions in hyperaphobetalipoproteinemia, normolipidemia and familial hypercholesterolemia. Proc. Natl. Acad. Sci., (USA) 80:6662..

Third, J.L.; Montag, J.; Flynn, M. et al. (1984):

Primary land familial hypoalphalipoproteinemia. Metabolism 33:136.

Trinder, P. (1969):

Determination of glucose in blood using glucose oxidase with an alternative oxygen acceptor. Ann. Clin. Biochem., 6-24.

Utermann, G.; Menzel, H.J.; Kraft, H.G. et al. (1987):

LP(a) glycoprotein phenotype: inheritance and relation to LP(a) lipoprotein concentrations in plasma. J. Clin. Invest., 80:458.

Vaith, P.; Assman, G. and Uhlenbruch, G. (1978):

Characterization of oligosaccharide side chain of apolipoprotein CIII from human plasma VLDL. Biochim. Biophys. Acta., 541:234-40.

Vlodaver, Z.; Kahn, H.A. and Neufeld, H.N. (1964):

The coronary arteries in early life in the different ethnic groups. Circulation 29:541.

Wahlefeldt, A.W. (1974):

Triglyceride determination after enzymatic hydrolysis. In: Methods of enzymatic analysis, 2nd edition. Academic press, New York, 1831.

Walker, A.R.P. and Walker, B.F. (1978):

High density lipoprotein cholesterol in African children and adults in a population free of coronary heart disease. Br. Med. J., 2:1336.

Wilhelmsson, C.; Vedin, J.A.; Elmfeldt, D.; Tibblin, G.
and Wilhelmsen, L. (1975):

Smoking and myocardial infarction.
Lancet 1:115.

Wilson, D.E.; Schreiber, P.H.; Brewster, A.C. and Arky,
R.A. (1970):

The enhancement of alimentary lipemia by ethanol in
man.
J. Lab. Clin. Med., 75:264.

Windmueller, H.G.; Herbert, P.N. and Leavy, R.I. (1973):

Biosynthesis of lymph and plasma lipoprotein and
apolipoproteins by isolated perfused rat liver and
intestine.
J. Lipid. Res., 14:215-23.

Ziegler, E.E. and Fomon, S.J. (1980):

Infant feeding and blood levels during childhood. In:
Lauer RM. Shekelle RB. eds. Childhood prevention of
atherosclerosis and hypertension.
New York: Raven. Press., 121-5.