

REFERENCES

REFERENCES

- Abdel- halim, A; Hammad, S; Abdel latif, M.; Abdel- AAL, A.;
Massoud, B.; (1986) . Prediction of sepsis among high- risk newborns.
the Gaz. Egypt. pediatr. Ass.; 34 (1 -4) : 149 .
- Addison, G.M.; (1988). Hypothalamic- pituitary- adrenocortical system . In :
practical clinical biochemistry, 6 th Edition. Gowenlock, A.H (Ed) .
philadelphia, London, New york. W.B. saunders, Co. P : 880 .
- Akenzua, G.I.; Hui, y.t.; Zipursky, A.; (1974). Neutrophil and band
counts in the diagnosis of neonatal infections. pediatr.; 54 (1) : 38 - 42.
- Albritton, W.L.; (1976). Neonatal listeriosis : distribution of serotypes in
relation to age at onset of disease . J. pediatr; 88 : 481 .
- Alexander. J.B; Giacoia. G.P; (1978) . Early onset nonenterococcal group
D streptococcal infection in the newborn infant J. pediatr; 93 : 489 - 490 .
- Alpert, G.; Campos, M.; preblud, S.R.; plotkin, S.A; (1984).
Vancomycin dosage in pediatrics. AJDC.; 138 : 20 - 25 .
- Amato, M.; Howald, H.; von moralt, G; (1988). Quantitative changes of
white blood cells and perinatal diagnosis of infection in high - risk preterm
infants . pediatr. padol .; 23 (2) : 129 - 34 .

- Anagnostakis, D.E; Lardinois, R.; (1971). Urinary catecholamine excretion and plasma NEFA concentration in small - for - date infants. *pediatr.*; 47 : 1000 - 9 .
- Anand, K.J.; Green, A.A.; (1988). Measuring the severity of surgical stress in newborn infants . *J. pediatr . surgery*; 23 (4) : 297 .
- Anast, C.S; (1984). Disorder of the adrenal gland and sodium metabolism. In : Schaffer's diseases of the newborn, 5 th Edition . Avery, M.E. and Taeusch, H.W. (Eds) . W.B. saunders, Co. london, tokyo, sydney, philadelphia . P : 479 .
- Aperia, A.; BroBerger, P.; Herin, o.; Zetterström, R.; (1979). Sodium excretion in relation to sodium intake and aldosterone excretion in newborn; preterm and full-term infants. *Acta. pediatr. scand.*; 68 : 813-817.
- Avery, G.B.; (1987). Bacterial and viral infection of the newborn . In : Avery G.B (Ed.) Neonatology, pathophysiology and management of the newborn 3 rd Edition, JB lippincott. P : 922 - 31 .
- Axelrod, J.; (1966). Methylation reactions in the formation and metabolism of catecholamines and other biogenic amines . *pharmacol . Rev*; 18 : 95 - 113 .
- Backer. C.J and Barret T.F.; (1973) . Transmission of group B streptococci among parturient women and their neonates . *J. pediatr.*; 83 : 919 .
-

- Baden, M.; Baure, C.R.; colle, E; klein, G.; stern, L.; (1973).
Plasma corticosteroids in infants with the respiratory distress syndrome .
pediatr.; 52 (6) : 782 - 788 .
- Baker, C.J.; kasper, D.L.; (1977). Immunological investigation of infants
with septicaemia or meningitis due to GBS. J. infect. dis.; 136 : 998 .
- Baker, C.J.; (1981). Nosocomial septicaemia and meningitis in neonates .
Am. J. Med; 70 : 698 - 701 .
- Baker, C.J.; Rench, M.A.; Edward, M.S.; Hays, B.M.; (1988).
Immunization of pregnant women with polysaccharide vaccine of group B
streptococcus. N. Engl. J. Med.; 319 : 1180 - 5 .
- Balitimore, R.S.; Kasper, D.L.; Baker, C.J.; (1977). Antigenic
specificity of opsonophagocytic antibodies in Rabbit antisera to GBS . J.-
immunol., 118 : 673 .
- Battisti, o.; Mitchinson, R; Davies, P.A; (1981). Changing blood
culture isolates in a referral neonatal intensive care . arch. dis. child.; 56 :
775 - 778 .
- Bauer, C.R.; Stern, L.; Colle, H.; (1974). Prolonged rupture of
membranes associated with a decreased incidence of respiratory distress
syndrome in premature infants. pediatr.; 53 : 7 - 12 .

- Bayard, F.; Ances, I.G.; tapper, A.; (1970). Transplacental passage and fetal secretion of aldosterone. J. clin. invest; 49 : 1389 .
- Baxter, J.D.; and tyrrell, J.B.; (1987). The adrenal cortex . In : endocrinology and metabolism, 2 nd Edition, by Felig, P. (Ed.) . Mc raw- hill .
- Beisel, W.R.; (1977). Magnitude of the host nutritional responses to infection. Am. J. clin. Nutr.; 30 : 1236 - 47.
- Beitin, I.Z.; Bayard, F.; Anceses, G.; kowarski, A; Migeon, J.; (1972). Plasma aldosterone concentration at delivery and during the newborn period. J. clin - invest; 51 : 386 - 394 .
- Beitin, I.; Bayard, F.; Ances, I.; Migeon, C; (1973).The metabolic clearance rate, blood production, interconversion and transplacental passage of cortisol and cortisone in pregnancy near term . pediatri. Research; 7 : 509 - 519 .
- Bell, G.H.; Smith, D.E.; paterson, C.; (1980). The adrenal cortex. In : textbook of physiology. 10 th, Edition, London, the English language Book society and churchil livingstone. P : 476 .
- Bennet. R.; Eriksson M.; Nord C.E and zetterstrom R. (1982) . Suppression of aerobic and anaerobic foecal flora in newborn receiving parenteral gentamycin and ampicillin. Acta . pediatri. scand; 71 : 559 .

- Bennet, R.; Eriksson, M., Zetterstrom. R.; (1987). Neonatal septicaemia Comparison of onset and risk factors during 3 consecutive 5 year periods . Acta. paediatr. scand; 76 : 361 - 2 .
- Bennet, R.; Bergdahl, S.; Erikson, M.; zetterstorm, R.; (1989). The outcome of neonatal septicaemia during fifteen years . Acta. paediatr . scand; 78 : 40 .
- Benuck, L.; David, R.J.; (1983). Sensitivity of published neutrophil indices in indentifying newborn infant with sepsis. J. paediatr.; 103 : 961.
- Bergqvist, G.z Erikson, M.; zetter storm, R.; (1979). Neonatal septicaemia and perinatal risk factors. Acta. paediatr. scand; 68 : 337 - 339 .
- Black, J.; and Williams, D.; (1973). Natural history of adrenal haemorrhage in the newborn. arch. dis. child.; 48 : 183 .
- Blair, J.R.; Coghlan, J.P; Denton, D.A; (1980) . A dose response comparison of the action of angiotensin II and angiotensin III in sheep . J. endocrinol.; 87 : 409 - 417 .
- Blaschko, H.; (1973). Catecholamines biosynthesis. Br. Med. Bull.; 29 : 105.
- Bondy, P.K. and Rosenberg, L.E.; (1980). Control of Renin secretion . In : Metabolic control and disease, 8 th Edition, philadelphia, W.B. saunders, Co.

Boyer K.M; Gadzala. C.A; Burd L.I; Fisher, D.E; Gotoff, S.P;
(1983). Selective intrapartum chemoprophylaxis of neonatal group B streptococcal . J. infect. dis; 148 : 795 - 801 .

Boyer, K.M.; Samuel. P.; Gotoff. S.; (1986). Prevention of early onset neonatal group B streptococcal disease with selective intrapartum chemoprophylaxis . N. Engl. J. Med.; 314 : 1665 - 9 .

Boyle, R.J.; chandler, B.P.; stone, B.S.; (1978). Early identification of sepsis in infants with respiratory distress . pediatr.; 62 : 744 .

Brook, I; (1983). Anaerobic infection in childhood, Boston W.B. saunders, Co. P : 50 .

Cacciari, E.; Cicognani, F.; Bernardi, S.; salardi, F.; (1976). GH,LH,ACTH and FSH behaviour in the first seven days of life . Acta. pediatr. scand.; 65 : 337 .

Cairo, M.S.; Rucker, R.; Bennetts, G.A.; Hicks, D.; Katz, J.; Johnson, S.; (1984) . Improved survival of newborns receiving leucocyte transfusions for sepsis . pediatr.; 74 (5) : 887 - 92 .

Cairo, M.S.; (1989). Neonatal neutrophil host defence. AJDC.; 143 : 40 - 46 .

Campbell, A.; Murphy, B.; (1977). The maternal - fetal cortisol gradient in pregnancy and at delivery . J. clin. endocrin . Meta.; 45 : 435 .

Carr, B.; OHashi, M.; simpson, E.; (1982). Low density lipoprotein binding and de novo synthesis of cholesterol in the neocortex and fetal zones of human adrenal gland. Endocrinol.; 110 : 1994 .

Casimiri,V.; Acker, G.; parvez, S; parvez, H.; Catro, L.; (1991). Characterization of enzymes of catecholamine synthesis and metabolism in human fetal membranes at birth . Am. J. obstet. Gynecol.; 164 : 599 - 3 .

Challis, R.G.; and thorburn, D; (1976). The fetal pituitary - adrenal axis and its functional interaction with the neurohypophysis . In : fetal physiology and medicine. Beard, R. and Nathanielsz, P. (Eds) . W.B. saunders, Co.london, philadelphia, toronto . P : 232 .

Cheesbrough, M.; (1985). Collection and transport and examination of specimens. In : Medical laboratory manual for tropical countries; cheesbrough, M. (Ed.), 5 th Edition. philadelphia, Butterwork pub. P : 176 .

Chevalier, R.L.; (1990) . Special needs of the newborn infant in fluid therapy. ped. clin. Nor. Am.; 37 (2) : 323.

Christensen, R.D.; and Rothstein. G.; (1979). pitfalls in the interpretation of leucocyte counts of newborn infants . Am. J. clin. pathol.; 72 : 608 .

- Christensen, R.D.; Bradly, R; Rothstein, G.; (1981). The leukocyte left shift in clinical and experimental neonatal sepsis. J. pediatr.; 98 : 101 .
- Christensen, R.D.; Rothstein, G.; Harald, B.; Anstall, B.; ByBee, B.; (1982) . Granulocyte transfusion in neonates with bacterial infection, neutropenia, and depletion of mature marrow neutrophils . pediatr.; 70 (1) : 1 - 7 .
- Christensen, R.D.; Hill, H.R.; Anstall, H.B.; (1984). Exchange transfusion as an alternative to granulocyte concentrate administration in neonates with bacterial sepsis and profound neutropenia . J. clin. apheresis; 2 : 177 .
- Christensen, R.D.; Mark, S.; David, C.; Harry, R.; Hill, A; (1991). Effect on neutrophil kinetics and serum opsonic capacity of intravenous administration of immunoglobulin to neonates with clinical signs of early - onset sepsis . J. pediatr.; 118 : 606 - 14 .
- Clapp, D.W.; Kligman, R.M.; Baley, J.E.; shenker, N.; Fanaroff, A.A.; Berger, M.; (1989) . Use of intravenously administered immunoglobulin to prevent nosocomial sepsis in low birth weight infants : Report of a pilot study . J. pediatr.; 115 (6) : 973 - 978 .
- Cooke, B.; Taylor, P.; (1971). Site of dehydroepiandrosterone sulfate biosynthesis the adrenal gland of the previable fetus . J. endocrinol.; 51 : 547 .

- Coovadia, y.m.; Mayosi, B.; Adhikari, M; Solwa, Z; (1989).
Hospital acquired neonatal bacterial meningitis : the impact of cefotaxime
usage on Mortality and of Amikacin usage on incidence . Ann. trop. pediatr.;
9 (4) : 233 - 9 .
- Cornett, L.E.; (1989). The adrenal medulla. In : textbook of physiology, 21
st Edition, by patton, H.D; Fuchs, A.F; hille, B.; scher, A.; steiner, R.
(Eds), philadelphia, sydney, tokyo, W.B. saunders, Co. P : 1229 .
- Corrigan, J.J. (1974). Thrombocytopenia : A laboratory sign of septicemia in
infants and children . J. pediatr.; 85 : 219 - 221 .
- Costa, A.; De Filippis, V.; voglino, M.; Massobrio, M.; (1988).
Adrenocorticotrophic hormone and catecholamines in maternal, umbilical and
neonatal plasma in relation to vaginal delivery . J. endocrinol invest.; 11
(10) : 703 - 9 .
- Coupland, R.E; (1965). The naturol History of the chromaffin cell . london,
longmans, Green. P : 157 .
- Cowett, R.M.; perter. G; Hakonson, D.o.; (1976). Reliability of
bacterial culture of blood obtained from an umbilical arterial catheter . J.
Pediatr.; 88 : 1033 - 1036 .
- Crickard, K.; Ill.C.; Jaffe.R.; (1981). Control of proliferation of human
fetal adrenal cells in vitro . J. clin. endocrinol . Metab. ; 53 : 790 .
-

- Crosson, F.J; Feder, H.M; Bocchini, J.A; Hackell, J.M; Hackell.J.C; (1977). Neonatal sepsis at the Johns hopkins hospital (1969 - 1975) . John. Hopkins. Med. J.; 140 : 37 - 41 .
- Cryer, P.E.; (1980). physiology and pathophysiology of the human sympathoadrenal neuroendocrine system . N. Engl. J. Med.; 303 : 436 .
- Dalvi, R.; Rao, S.; Rangnoker, J.; (1991). Exchange transfusion in neonatal sepsis . Indian - pediatr .; 28 (1) : 39 - 43 .
- Demetre, N.; Anthony, A.; Maria, G.; Constantin, S.; (1969). Urinary excretion of catecholamines by full - term and premature infants . pediatr.; 44 262 - 264 .
- Diczfalusy, E.; (1962). Endocrinology of the fetus. Acta. obstet. Gynecol. scand. (suppl.1); 41 : 45 .
- Dietzman, D.E.; Fischer, G.W.; schoen kencht, D.; (1974). Neonatal Escherichia coli septicaemia J. pediatr.; 85 : 128 .
- Diez, J.A.; Sze, P.Y.; Ginsburg, B.E.; (1976). Postnatal development of mouse plasma and brain corticosterone levels. Endocrinol.; 98 : 1434 - 42.
- Dillon, M.J. Gillin, E.A.; Ryness, J.M.; swift. M.; (1976). plasma renin activity and aldosterone concentration in the human newborn . arch. dis. child.; 51 : 537 - 541 .
-

- Doerr, H.G.; versmold, H.T.; biddingmaier, F.; sippell, W.G.; (1989). Adrenocortical steroid in small- for- gestational- age term infants during the early neonatal period. *pediatr. Res.*; 25 (2) : 115 - 8 .
- Donato, H.; Gebara, E.; de cosen, R.H.; Gioseffio, M.; (1979). Leukocyte alkaline phosphatase activity in the diagnosis of neonatal bacterial infection . *J. pediatr.*; 94 (2) : 242 - 244 .
- Donowitz, L.G; Marsik, F.K.; Fischer. K.A; and weuzel. R.P; (1981). Contaminated Breast Milk : A source of klebsiella bacteremia in a newborn intensive care unit . *Rev. infect. dis.*; 394 : 716 .
- Dubowitz, L.M.; Dubowitz, V.; Goldberg, C.; (1970) . Clinical assessment of gestational age in newborn infant . *J. pediatr.*; 77; 1- 10.
- Dunkle, L.M.; Brotherton, T.J and Feiginm R.D.; (1976). Anaerobic infections in children : a prospective study. *pediatr.*; 57 : 311 - 320 .
- Eberlein, W.R.; (1965). Steroids and sterols in umbilical cord blood. *J. clin. endocrinol.*; 25 : 1101 - 1105 .
- Edelmann, C.; Ogwo. J.; Fine, B. ; Martinez, M.; (1973). The prevalence of bacteremia in full term and premature newborn infants . *J. pediatr.*; 82 : 125 .

- EL Gamal, y.; haroun, A.A.; EL shimi, EL Samahi, M.; Abo Mosalem, M.; (1989). A new look on buffy coat smear examination in high risk pregnancy and neonatal infection. Egypt. J. pediatr. ; 6 (3 - 4) : 301 - 13 .
- Eliot, R.J.; Lam, R.; Leake, R.F.; Hobel, C.G.; (1980). plasma catecholamine concentrations at birth and during the first 48 hours of life . J. pediatr.; 96 (2) : 311 - 315 .
- EL Maraghy, S.; EL saifi, A.; soleman, M.; (1982). Gastric aspirate in early detection of neonatal sepsis. M.sc. thesis, pediatr.; Cairo university .
- Engle, W.D.; Rosenfeld, C.R.; (1984). Neutropenia in high- risk neonates. J. pediatr.; 105 : 982 - 986.
- Eshali, H.; Ringertz, S.; Nystrom, S.; (1989). Septicaemia with coagulase negative staphylococci in a neonatal intensive care unit . Acta. pediatr. scand; 360 (suppl) : 127 - 134 .
- Feigin. R.D; Abramson. SL and Edwards . M.S; (1987). Infection in the fetus and neonate. in : Neonatal- perinatal medicine, 4 th edition- Franaroff. A.A; Martin. RJ (Eds), the C.V.Mosby, Co. st. louis, toronto, washington . P : 763 .
- Ferrieri, P; (1985). GBS infections in the newborn infant : diagnosis and treatment . Antibiot . chemother.; 35 : 211 - 224 .
-

- Finegold, S.M; Baron, E.J; (1986).** Microorganisms Encountered in the Blood . in : Carson, D.C (Ed); Bailey & scott's diagnostic microbiology 14th Edition, cv Mosby, Co. P : 214 .
- Finely, W.E.; McKee, J.I.; (1982).** Serum cortisol levels in severely stressed patients . Lancet.; 1 : 1414 - 1415 .
- Foda, A.I.; El-shazly, A.H.; EL- kannishy, M.H.; EL- Salab, shadia M.; (1982).** Study of fetal plasma cortisol levels in relation to the mode of oncet and method of labour and fetal distress. The Gaz. Egypt. pediatr. Ass.; 30 (1- 2) : 167 .
- Francis, S.J.; Walker, R.F.; Riad- Fahmy, D.; hughes, D.; Murphy, J.F.; (1987).** Assessment of adrenocorticol activity in term newborn infants using salivery cortisol determinations . J. Pediatr.; 111 : 129 - 33 .
- Franks, R.D.; (1967).** Diurnal variation of plasma 17- hydroxycorticosteroids in children. J. clin. endocrinol. Metab.; 27 : 75 .
- Franson, T.R; Sheth, N.K; Rose. H.D; Sohnle, P.G; (1984).** Scanning electrone microscopy of bacteria adherent to intravascular catheters . J. clin. microbiol; 20 : 500 - 5 .
- Freedman, R.M; Ingram, D.L; Gross. L; warshav. J.B; Baltimore, M.; (1981).** A half century of neonatal sepsis at yale (1928 - 1978) . Am. J. Dis. child.; 135 : 140 - 4 .

- Freidman, S.B.; Glasgow, L.A.; Grotta, L.J.; (1970). plasma adrenocorticosteroid levels during experimental infections . (Abst) . The American. pediatr. society; 30 : 69 .
- Freisen, C.A.; (1986) . Characteristic features of neonatal sepsis due to *Haemophilus influenza* . *Rev. infect. dis.* ; 8 : 777 - 80 .
- Fujeda, K.; Faiman, C.; Reyes, F.; winter, J.S.; (1982). The control of steroidogenesis by human fetal adrenal cells in tissue culture . *J. clin. endocrinol . Metab.*; 54 : 89 - 94 .
- Gardner, L.; (1969). Development of the normal fetal and neonatal adrenal. In : *Endocrine and genetic disease of childhood.* by gardner. L (Ed). W.B. saunders, Co. philadelphia . P : 392 .
- George, J.M.; wright, L.; Bell, N.H.; Barter, F.C.; (1970). The syndrome of primary aldosteronism . *AM.J.Med.*; 43 : 343 .
- Gerdes, J.S.; Polin, R.A.; (1987). Sepsis screen in neonates with evaluation of plasma fibronectin. *pediatr. infect. Dis. J.*; 6 : 443 - 6 .
- Gill,G.N.; (1979). AcTH regulation of the adrenal cortex . In : *pharmacology of adrenal cortical hormones.* Gill, G.N. (Ed), pergamon .
- Gill, G.N.; (1987). Biosynthesis, secretion and metabolism of hormones . In : *Endocrinology and metabolism . 2 nd Edition,* Felig, P. (Ed). McGraw - hill

- Gilstrap, L.G.; Leveno, K.J.; Cox, S.M.; Burris, J.S.; Rosenfeld, C.R.; (1988) . Intrapartum treatment of acute chorio-amnionitis : impact on neonatal sepsis . Am. J. obstet. Gynecol.; 159 : 579 - 83 .
- Goldfien, A.; (1991). Adrenal Medulla . In : basic and clinical endocrinology, 3 rd Edition, by Greenspan, F.S. (Ed), Los Altos, California, Appleton & Lange. P: 380 .
- Gomella, T.L.; (1988). Hypotension and shock. in : Neonatology, basic management. edited by Gomella, T.L, Appleton and Lange. Co, U.S.A, Canada. P : 155.
- Gordon, B.; Cutler, J.R.; Lynn.; L.; (1990). Synthesis and secretion of corticosteroid. in : principles and practice of endocrinology and metabolism, by Becker, K.L (Ed) J.B. Lippincott Co., New york, London, sydney, tokyo . P : 579 .
- Gotoff S.P. and Behrman R.E (1970). Neonatal septicaemia. J. pediatr.; 76 (1) : 142 - 153 .
- Gotoff. S.; (1974). Neonatal immunity. J. pediatr.; 85 : 149 .
- Gower, D.B.; Cooke, G.M.; (1983). Regulation of steroid- transforming enzymes by endogenous steroids . J. steroid. Biochem.; 19 : 1527 .
- Granner, D.K.; (1985). Hormones of the adrenal gland. in Harper's Review of biochemistry, los Angeles, california. Lange Med, publication .

Grant, J.K.; (1978). An introductory review of adrenocortical steroid biosynthesis . In : the endocrine function of the human adrenal cortex, by James, V.H.T (Ed), Academic press .

Grael, EL.; Halle. E.; Bollmann- R.; BuchHo 12.P; (1989). Neonatal septicaemia - incidence, etiology and outcome . Acta. pediatr. scand.; 360 (suppl) : 113 - 119 .

Greenberg, R.E. (1975). The physiology and metabolism of catecholamines . In : endocrine and genetic diseases of childhood and Adolescence, by Gardner, L.I. (Ed), 2nd Edition, philadelphia, london, toronto, W.B. Saunders, Co. P : 886 .

Greisen, G.; Ferderiksen, P.S.; Hertel, J.; christensen, J.; (1985). Catecholamine response to chest physiotherapy and endotracheal suctioning in preterm infants. Acta. pediatr. scand.; 74 : 525 - 529 .

Gunnar, M.R.; (1989). Studies of human infant's adrenocortical response to potentially stressful events. New. Dir. child. dev.; 45 : 3 - 18 .

Gutai, J.; George, R.; Koeff.; S.; (1972). Adrenal response to physical stress and the effect of adrenocorticotrophic hormone in newborn infants . J. pediatr.; 81 : 719 .

Guyton, A.C.; (1986). The adrenocortical hormones. In : textbook of medical physiology 7th edition, London, tokyo, New york. W.B. saunders, Co. P : 909 .

- Guyton, A.C.; (1991). The adrenocortical hormones. In : textbook of medical physiology, 8 th Edition, London, toronto, tokyo. W.B. saunders, Co. P : 842 .
- Hahn, H.; (1976). Adaptations mechanismen der nebbenierenrinde. Munich : Ludwig Maximillians university (M.D thesis) P : 166.
- Hall, R.T.; Shigheoka, A.o; hill, H.R; (1983).Serum opsonic activity and peripheral neutrophil count before and after exchange transfusion in infants with early onset group B streptococcal septicaemia . *Pediatr. infect. dis.*; 2 (5) : 356 .
- Hammad, S.; Abdel halim. A.; Abdel latif, M.; Abdel- AAL, A.; Massoud, B.; (1986). Early diagnosis of neonatal septicaemia. *the Gaz. Egypt. pediatr. Ass.*; 34 (1 - 4) : 7 - 15 .
- Haque, K.N.; ziadi, M.H.; Bahakin, H.; (1988). IgM. enriched intravenous immunoglobulin therapy in neonatal sepsis . *Am.J. Dis. child.*; 142 : 1293 - 1296 .
- Harkness, J.L; Hall, L.M; Ilstrup, D.M; washington J.A; (1975). Effects of atmosphere of incubation of routine subcultures on detection of bacteremia in vacuum Blood culture bottles . *J. clin. microbiol.*; 2 : 296 - 299
- Hartman, K; (1978). Glass electrodes for hydrogen and other cations. *Mikrochimica Acta* : 235 - 246.
- Hauth, J.C; Gilstrap, L.C, Hankins, G.D.V; Canner, kiD; (1983). Term maternal and neonatal complication of acute chorioamnionitis . *J. obst. gyn.*; 66 : 59 - 62 .

- Haynes, R.C.; Murad, F.; Fraster, R.; (1985). Adrenocorticotrophic hormone . In : Goodman and Gilman the pharmacologic basis of therapeutics, by Goodman, L.S.; Gilman, A.G (Eds) . 7 th Ed, New york, Macmillan, P : 1459 .
- Helmuth, G.; wolfgang, G.; Hans, T.; Frank, B.; (1987). Plasma aldosterone and 11- deoxycortisol in term neonates : A reevaluation . J. clin. endocrinol. Metab.; 65 : 208 - 211 .
- Hemming V.G; overall. J. C. and Britt. M.E; (1976). Nosocomial infection in a newborn intensive care unit : results of forty- one months of surveillance. N. Engl. J. Med; 294 : 1310.
- Herson, V.C.; Kranse, P.J.; Eisenfeld, L.I and maderozo,E.G.; (1988). Indomethacin. associated sepsis in very low birth weight infants; Am. J. Dis. child.; 142 (5) : 555 - 8 .
- Hill, h.r.; Shigeoka, A.O.; Angustine, N.H.; (1984). Fibronectin enhances the opsonic and protective activity of monoclonal and polyclonal antibody against Group B streptococci . J . Exp. Med.; 159 : 1618 ,
- Hocher, J.R.; Simposn, P.M.; Rabaliais, G.; stewart, L.; (1992). Extracorporeal membrane oxygenation and early onset group B streptococcal sepsis . pediatr.; 89 (1) : 1 - 4 .

- Hoff, R.; (1984). physiology of adrenal hormones. In : clinical chemistry : theory analysis and correlation, kaplan, L.A.; pesce, A.S.; Abbot, L.; and kasper, B. (Eds) , London, tokoyo, New york. C.V. Mosby, Co. P : 816 .
- Holden, K.R.; young, R.B.; piland, J.H.; Hurt, G.; (1972). plasma pressors in the normal and stressed newborn infant . *Pediatr.*; 49 (4) : 495 - 502 .
- Honour, J.W.; Valman. G.B.; Shackleton, H.L.; (1977). Aldodterone and sodium homeostasis in preterm infants. *Acta. pediatr. scand.*; 66 : 103 - 9.
- Horowitz, B.; chang, M.D.; Lippin, A.; (1984). preparation of antihaemophilic factor and fibronectin from human plasma cryoprecipitate . *Transfusion*; 24 : 357 .
- Hyatt, P.J.; (1987). Functional significance of adrenal cortex . In : recent advances in adrenal regulation and function, by Agata, R.; chrosous, G.P. (Eds) . New york, Raven press. P : 35 .
- Ito, T.; Woo, J.; Hening, R.; Horton, R.; (1972). A radioimmunoassay for aldosterone in human peripheral plasma including a comparison of alternate techniques . *J. clin. Endocrinol.*; 34 : 106 .
- Jacobes, R.F.; Darville, T.; Parks, A.J.; (1992). Safety profile and efficacy of cefotaxime for the treatment of hospitalized children . *Clin-infect. dis.*; 14 : 56 - 65 .

- Jaffe, R.; Seron- Ferre, M.; Crikard, K.; Koritnik, D.; (1981).
Regulation and function of the primate fetal adrenal gland and gonad. recent
progress in hormone research; 37 : 41 - 69 .
- Jenkins, H.R.; Feuton, T.R.; Dillon, M.; Milla, P.J.; (1990).
Development of colonic sodium transport in early childhood and its
regulation by aldosterone. Gut ; 31 : 194 - 197 .
- John. J.; Buchino . E; Iewin. K; (1979). Systemic group D streptococcal
infection in newborn infants . AJDC; 133 (3) : 270 - 273 .
- Jonathan F. Donald. A. Golman, NaNcy. E, David. G (1990).
Association of intravenous lipid emulsion and coagulase - negative
staphylococcal bactremia in neonatal intensive care units . N. Engl. J. Med.;
323 (5) : 301 - 8 .
- Jones, M.T.; (1979). Control of adrenocortical hormones. In : the adrenal
gland. by James, V.H.T. (Ed), Raven press .
- Kafetzis, D.A.; Brater, D.C.; Kapiki, A.N.; (1982). Treatment of
severe neonatal infection with cefotaxime : efficacy and pharmacokinetics . J
pediatr.; 100 : 483 .
- Keller, W.; Wood, M.; and Dallaman, M.F.; (1984). Corticosteroid
inhibition of AcTH . Endocrinol. Rev; 5 : 1 .
-

- Kelsey, M.C; Lipscomb. A.P; Mowles, J.M; (1982). Limulus Amoebocyte Lysate Endotoxin test : An aid to the diagnosis in the septic neonates. J. infect; 4 : 69 .
- Kenny, F.M.; Megeon, C.J.; preeyasombat, C.; (1966). Cortisol production rate in normal infants , children and adult . pediatri; 37 (1) : 34 - 42 .
- Khalifa, A.M. Allam, M.; Nowar, F.; Sabry, S.I; (1989). Quantitative and qualitative changes in neonatal bacterial infection. The Gaz. Egypt. pediatri. Assoc.; 37 : 1.
- Kite, P.; Millar, M.R; Gohram, P.; congdon, P.; (1988). Comparison of five tests used in diagnosis of neonatal bacteremia . arch. dis. child; 63 : 639 - 643 .
- Kite. P; langdale. V.; Miller . M.R; Mackay. P; (1989). Direct isolation of coagulase negative staphylococci from neonatal Blood samples . J. Hosp. infect; 14 : 135 - 40 .
- Kite. P; todd.N; MacKay . P; Miller. M.R; (1990). plasma neutrophil elastase alpha - 1 - proteinase inhibitor level in premature neonates with Coagulase- negative staphylococcal bactermia . J. pediatri; 117 (4) : 630 - 1 .
- Kleiman, M.B.; Reynold, J.K.; shreiner, R.L.; (1984). Rapid diagnosis of neonatal bacteremia with acridine orange- stained Buffy coat smears. J. pediatri; 105 : 419 - 21 .

- Klein, R.; Fortunato, J.; Papadatos, C.; (1954). Free blood corticoids in the newborn infant . J. clin. invest. ; 33 : 35 .
- Klein, J.O.; Marcy, S.M.; (1976). Bacterial infection . in : Remington, R.S.; Klein; J.O. (Eds); infections disease of the foetus and Newborn. 2 nd Edition. W.B. Saunders, Co. P : 747 .
- Klein, J.O; Dashafesky, B.; Norken, C.R.; Mayer. A.; (1983). Selection of antimicrobial agents for treatment of neonatal sepsis . Rev. infect. dis; 5 : 55 .
- Klein, J.O.; (1984). Recent advances in management of bacterial meningitis in neonates . infection; 12 (suppl) : 44 - 51 .
- Kotchen, T.A.; strickland, A.L.; Rice, T.; Walters, D.R.; (1972). A study of the renin - angiotensin system in newborn infants . J. pediatr.; 80 : 938 .
- Kovatch, A.L; wald, E.R.; (1985) . Evaluation of febrile neonates. semin. perinatol; 9 : 12 - 3 .
- Kowarski, A; Katz, H.; Migeon, J.C.; (1974). plasma aldosterone concentration in normal subjects from infancy to adulthood. J. clin. endocrinol. Metab.; 38 : 489.
- Kraiem, Z.; Sack, J.; Brish, M.; (1985). Serum cortisol levels : the first 10 days in full- term and preterm infants. Isr. J. Med. sci.; 21 : 170 - 2.

- Kraus, A.N.; Albert, R.F.; Kannan, M.M.;(1970). Contamination of umbilical catheters in the newborn infant . J. pediatr; 77 : 965 - 9 .
- Kreiger, D.T; (1973). The effect of ocular enucleation and altered lighting regimens at various ages on the circadian periodicity of plasma corticosteroid levels in the rats . Endocrinol.; 93 : 1077 - 1091 .
- Kusnetsova, L.V.; (1987). Adrenal glucocorticoid function and its regulation in human prenatal ontogeny and in the 1 st week of life . Biull. Eksp. biol. Med., 104 (7) : 11 - 13 .
- Kusuma. B; Richard. L; James. A; Edwin. L; (1979). Group - D streptococcal septicaemia in the neonate. AJDC; 133 (5) . 493 - 496 .
- Kyllonen, K.S.; Clapp, W.; Robert, M.; Nancy, S.; Fanaroff, A.; (1989) .Dosage of intravenously administered immune globulin and dosing interval required to maintain target levels of immunoglobulin G in low birth weight infants . J. pediatr.; 115 (6) : 1013 - 16 .
- Lagercrantz, H.; Bistoletti, P.; (1973) . Catecholamine release in the newborn infant at birth. pediatr. Res.; 11 : 889 - 93 .
- Lancefield, R.C.; McCarty, M.; Everly, W.; (1975). Multiple mouse protective antibodies Directed against Group B streptococci. J. Exp. Med.; 142 : 165 .

- Landsberg, L.; and young, J.B.; (1985).** Catecholamines and the adrenal medulla. In : williams textbook of endocrinology, 15 th ed., wilson, J.D, Foster, D.W. (Eds), philadelphia, W.B. saunders, Co. P : 932 .
- Lannering, B.; Larsson, L.E; Rojas, J.; stahlman. M; (1983).** Early oncet group B streptococcal disease . seven years of experiance and clinical scoring system. Acta. pediatri. scand; 72 : 597 - 602 .
- Lao, T.T.; and panesar, N.S.; (1989).** The effect of labour on prolactin and cortisol concentration in the mother and the fetus . Eur. J. obstet. Gynecol. Reprod. biol; 30 (3) : 233 .
- Lau, y.L; Hey. E (1991).** Sensitivity and specificity of daily tracheal aspirate culure in predicting organisms causing bactermia in ventilated neonates . pediatri. infect . dis . J; 10 (4) : 290 - 4 .
- Lawrence, C.C.; seron- Ferre, M.; Jaffer, R.; Siteri, P.; (1978).** Steroid production by definitive and fetal zones of the human fetal adrenal gland . J. clin . endocrinol. Metab; 47 : 603 .
- Le Douarin, N.M.; Smith, J.; Le lievre, C.S.; (1981).** From the neural crest to the ganglia of the peripheral nervous system . Annu. Rev. physiol; 43 : 653 .
- Lemos, L.; (1976).** Aspects hematologiques et cliniques des septicemies neonatales . Rev. Med. suisse. Romande.; 96 : 589 .

- Leslie, G.R.; Gallery, D.M.; arnold, J.D.; Nicholson, E.; (1991).
Hyaline membrane disease and early neonatal aldosterone metabolism in
infants of less than 33 weeks gestation. *Acta. pediatr. scand.*; 80 : 628 - 633.
- Liu, C.H.; Lehan. C.; Speer, M.E.; Smith, E.O.; Fernbach, D.J.;
Rudolph, A.J.; (1985). Early detection of bacteremia in an outpatient
clinic. *pediatr*; 75 (5) : 827 - 813 .
- Maberry,M.C.; and Gilstrap,L.C.; (1991). Intrapartum Antibiotic therapy
for suspected intraamniotic infection: impact on the fetus and neonate . *Clin.*
obstet- Gynecol.; 34 (2) : 345 - 51 .
- Macleod, J; (1987). Septic shock. in : Davidson's principles and practice of
medicine. 15 th edition. Macleod, J.; Edward. C.; Douchier, I. (Editors).
churchil livingstone. London. P : 48.
- Manroe, B.L.; Rosenfeld, C.R.; weinberg, A.G.; Brown, R.;
(1977). The differential leukocyte count in the assessment and outcome of
early - oncet neonatal group B streptococcal disease . *J. pediatr.*; 91 : 632 .
- Manroe, B.L; weinberg, A.G; Rosenfeld, C.R.; Browre, R.; (1979).
The neonatal blood count in health and disease. *J. pediatr*; 95 : 89 .
- Mantagos, S.; Koulouris, A.; Vagenakis, A.; (1991). A simple stress
test for the evaluation of hypothalamic pituitary- adrenal axis during the first
6 months of life . *J. clin. endocrinol. Metab.*; 72 : 214 - 216 .

- Margolis, F; Roffi, J.; Jost, A.; (1966).** Norepinephrine methylation in fetal rat adrenals. *Science*; 154 : 275 .
- Marino, J.; Martinez, M.J.; hawkins, F.; Gonzalez, A.; (1990).** Encysted adrenal haemorrhage. Prenatal diagnosis. *Acta. pediatri. scand.*; 79 : 230 - 231 .
- Marodi, L.; Kalmar, A.; Szabo, I.; (1989).** Opsonic activity in serum from septic infants treated with intravenous Immunoglobulin . *arch. dis. child.*; 64 530 - 34 .
- Mason, J.; Hemsell, P.; korte, K; (1983).** Steroidogenesis in dispersed cells of the human fetal adrenal . *J. clin. endocrinol. metab*; 56 : 1057 .
- Maxwell,G.M.; Crompton,S.; Davies,A.; (1985).** Urinary catecholamine levels in the newborn infant . *Eur. J. pediatri.*; 143 (3) : 171 - 4.
- McCaa, R.E.; Langford, H.G.; Montalvo, J.M.; (1981).** Regulation of aldosterone biosynthesis in sodium deficiency. *Hypertension* ; 3 (suppl 1) : 74 - 80 .
- McCracken, G.; (1976).** Neonatal sepsis (Editorial comment). *J. pediatri*; 89 : 203 .
- McCracken, G.H.; (1981).** perinatal bacterial diseases. in : textbook of pediatric infectious disease, 1 st Edition. Freij, R.D. and chervy, J.D (Eds) . philadelphia, W.B. saunders, Co. P : 747 .

McCracken, G.H. Jt; Freij, B.J. ; (1987). Perinatal bacterial disease . In : Feigin, R.D; cherry, J.D. (Eds); textbook of pediatric infectious disease, 2nd Edition, W.B. saunders, Co. P : 940 - 66 .

Meignier, M.; Renaud, P.; Robert. R; Roze. J.C.; Rigal. E; Mouzard. A; (1989). Cephalohaematoma infection in neonatal septicaemia. *pediatr*, 44 (1) : 27 - 9 .

Melby, J.C.; and spink, W.W.; (1958). Comparative studies on adrenal cortical function and cortisol metabolism in healthy adults and in patient with shock due to infection . *J. clin. invest.*; 37 : 1791 - 1798.

Melby, J.C.; (1984). primary aldosteronism. *kidney int.*; 26 : 769 .

Migeon, C.J.; (1972). Adrenal androgens in man. *AM.J. Med.*; 53 : 606 .

Miller, W.L.; (1988). Molecular biology of steroid hormone. *Endocrinol. Rev.*; 9 : 295 .

Miller; M.R.; todd.N; Machay. P.; (1990). Neonatal infection with coagulase negative staphylococci. *arch. dis. child*; 65 (10) : 1015 .

Mirkin, B.; (1972). Ontogenesis of the adrenergic nervous system : functional and pharmacologic implication . *Fed. proc*; 31 : 65 .

Mitchell S. cairo, Ralph Rucker, Geni A. Bennetts and David Hicks
(1984). Improved survival of newborn recieving leukocyte transfusion
for sepsis . *pediatr.*; 74 (5) : 887 - 92 .

Murphy, B.P; (1974). Evidence of cortisol deficiency at birth in infants with
the respiratory distres syndrome. *J. clin. Endocrinol. Metab.*; 38 : 158.

Murphy, J.F.; Joyce, B.G.; - hughes, I.A.; (1983). plasma 17 -
hydroxyprogesterone concentrations in ill newborn infants . *arch. dis. child.*;
58 : 532 - 34 .

Murphy, S.J.; Ryalls, J.D.; Hughes, A.A; (1986). Response to AcTH
in the newborn . *Arch. dis. child.*; 61 : 57 - 60 .

Murphy, J.F.; Hughes, D.; Robinson, J.A.; Riad - Fahmy, D.;
(1987). Blood spot glucocorticoid concentrations in ill preterm infants .
arch. dis-child.; 62 : 1014 - 18 .

Narbona, L.E.; Maldonado, L.J.; Nieto, G.M.; Garcia, D.C.;
(1990). plasma concentration of atrial natriuretic peptide, vasopressin and
aldosterone in the umblical cord blood : its relation to perinatal asphyxia .*An.*
Esp. pediatr.; 32 (1) : 49 - 52 .

Nebert, D.W.; Adesnik, M; Coon, M.; (1987). The P- 450 gene
superfamily : recommended nomenclature. *DNA.*; 6 : 1 .

Neumann, S.; Hennrich, N.; Gunzer, G.; Lang, H.; (1983).

Enzyme- linked immunoassay for complexes of human granulocyte elastase with 1 OC - proteinase inhibitor in plasma. in : progress in clinical enzymology II. Goldberg, D.M. and werner. M. (Eds) , New york , Masson, Co. P : 293 .

Neville, A.M.; O'hare, M.; (1982). The human adrenal cortex . pathology and biology - an integrated approach. New york : springer - verlag .

Neville, A.M.; O'hare, M.; (1985). Histopathology of the adrenal cortex. J. clin. endocrinol. Metab.; 14 : 791 .

New, M.L.; Dupont, B.; Pang, S.; (1981). An update of congenital adrenal hyperphasia. Recent . prog. Horm. Res; 37 : 105. .

Nicholas G. guerina (1991). Bacterial and fungal infection. in manual of Neonatal Care, 3 rd edition. little, Brown and company. Boston, toronto, london. P : 150 .

Noel, G.L, Laufer, D.A; Edelson, P.J; (1988). Anaerobic bacteremia in a neonatal intensive care unit . an eighteen- year experience . pediatri . infect. dis . J.; 7 : 858 - 862 .

Noguchi, A.; Reynolds, J.W.; (1978). Serum cortisol and dehydroepiandrosterone sulfate responses to AcTH stimulation in premature infants . pediatri. Res.; 12 : 1057 - 61 .

Oberfield, S.E.; Bakshi, S.; Mayes, D.; Levine, L.S.; (1990).

Steroid response to adrenocorticotropin stimulation in children with human immunodeficiency virus infection. *J. clin. Endocrinol. Metab.*; 70 : 578.

Ohlsson, K. and Olsson, A.S; (1978). Immunoreactive granulocyte

elastase in human serum . *Hoppeseyler's Z physiol. chem*; 539 : 1531 .

Ohlsson. A; Bailey . T; Takiddine . F.; (1986). Changing etiology

and outcome of neonatal septicaemia in Riyadh, Saudi Arabia. *Acta. pediatr. scand*; 75 : 540 - 44 .

Okada, D.M; chow, A.W.; Bruce, V.T; (1977). Neonatal scalp abscesses

and foetal monitoring : factor associated with infections. *A . J. obst. gyn.*; 129 : 185 - 7 .

Olding, L.B; oldstone, M.G.; (1976). Thymus derived peripheral

lymphocyte from human newborn, inhibit division of their mother's lymphocytes. *J. immunol.* ; 116 : 682 .

Onishi, S.; Miyazawa, G.; nishimura, Y.; sugiyama, S.; (1983).

postnatal development of circadian rhythm in serum cortisol levels in children. *pediatr.*; 72 : 399 - 404 .

Overall, J.C; (1987). Infection of the newborn. In : Nelson textbook of

pediatrics . 13 th edition. Behrman, R.F. and vauan, V.C. (Eds) . W.B. saunders, Co. philadelphia, london, toronto, tokyo. P : 422 - 435 .

Padbury, J.F.; Roberman, B.; oddie, T.T.; hobel, C.J.; (1982).

Fetal catecholamine release in response to labor and delivery . obstet. Gynecol.; 60 : 607 - 611 .

Padmanbhan, J.; Risemery, H.M.; Rowe, D.; (1973). Howell- Jolly

bodies in the peripheral blood of full term and premature neonates . John. Hopkins. Med. J.; 132 : 146 - 50 .

Pang, S.; pollack, M.S.; Loo, M.; (1985). pitfalls of prenatal 21 -

hydroxylase deficiency in congenital adrenal hyperplasia . J. clin. endocrinol. Metab; 61 : 89 .

Parker, LN.; Sack, J.; Fisher, D.A.; (1978). The adrenarche : prolactin,

gonadotropins, adrenal androgen and cortisol . J. clin. endocrinol. Metab; 46 : 396 .

Parker, L.N.; Levin, E.R.; Lifrak, E.T.; (1985). Evidence for

adrenocortical adaptation to severe illness. J. clin. Endocrinol. Metab; 60 : 947.

Parker, C.R.; and wendel, G.D.; (1988). The effect of syphilis on

endocrine function of the fetoplacental unit . Am. J. obstet. Gynecol.; 159 : 1327 - 31 .

Parmelee, A.H.; wanner, W.H.; Schulz, H.R.; (1964). Infant sleep

patterns : from birth to 16 weeks of age . J. pediatr.; 65 : 576 - 82 .

Pavliuk, V.P.; (1988). Role of breast feeding in the development of the function of the hypophysial - adrenal system in newborn infants in the early neonatal period . Probl. Endokrinol (Mosk); 34 (5) : 35 - 8 .

Payne, N.R.; Burke, B.A; Day DL P.D; thompson, T.R Ferrieri, P; (1988). Correlation of clinical and pathologic finding in early onset neonatal group B streptococcal infection with disease severity and prediction of outcome. *pediatr. infect. Dis. J.* 7 (12) : 836 - 847 .

Pearce. R; Robertson. N.R; (1986). Sepsis neonatorum . in textbook of neonatology, 1 st edition, Robertson . N.R (Eds), Churchill Livingstone, London, Edinburgh and New York. P : 654 .

Philip, A.G.S; Hewitt, J.R; (1980). Early diagnosis of neonatal sepsis. *pediatr*; 65 (5) : 1036 - 1041 .

Philip, A.G.S; (1981). Decreased use of antibiotics using neonatal sepsis screening technique . *J. pediatr*; 98 : 795 - 799 .

Philip, A.G.S; (1984). Acute phase protein in neonatal infection *J. pediatr*; 105 : 940 .

Placzek M.M and Whitelaw. A (1983). Early and late neonatal septicaemia. *arch. dis. child*; 58 : 728 - 731 .

Plow, E.F.; (1982). Leukocyte elastase during blood coagulation . *J. clin. invest.*; 69 : 564 .

- Pohjavuori, M.; and Fyrhquist, F.; (1983). Vasopressin, AcTH and neonatal haemodynamics. *Acta. paediatr. scand*; 305 (suppl) : 79 - 83 .
- Pommier, C.G.; O'shea, J.; chused, T.; (1984). Studies on fibronectin receptors of human peripheral blood leucocytes. *J. exp. Med.*; 159 : 137 .
- Poskitt, T.R.; Poskitt, P.K.; (1985). Thrombocytopenia of sepsis. The role of circulating IgG containing immune complexes. *Arch. intern. Med.*; 145 : 891 - 4 .
- Pourcyrons, M; Korohees, S.B.; Bada, H.S.; patterson, T.; (1988). Indwelling umbilical arterial catheter : A preferred sampling site for blood culture . *paediatr.*; 81 (6) : 821 - 25 .
- Price, D.; Close, G.; Fielding, B.A.; (1983). Age of appearance of circadian rhythm in salivary cortisol values in infancy . *arch. dis. child.*; 58 : 454 - 456 .
- Procianoy, R.S.; Cecin, S.K.; Pinheiro, C.E.; (1983). Umbilical cord cortisol and prolactin levels in preterm infants . *Acta. paediatr. scand.*; 72 : 713 - 716 .
- Procianoy, R.S. and Cecin, K.G.; (1985). The influence of labor and delivery on preterm fetal adrenal function. *Acta. paediatr. scand*; 74 : 400 - 4 .
-

- Procianoy, R.S.; Giacomini, C.B.; oliveria, M.L.; (1988). Fetal and neonatal adrenal function in birth asphyxia. *Acta. pediatr. scand.*; 77 : 671 - 74 .
- Pryds, O.; Christensen, N.J.; Friis-hausen, B.; (1990). Increased cerebral Blood flow and plasma epinephrine in hypoglycemic preterm neonates. *Pediatr.*; 85 (2) : 172 - 176 .
- Puntis, J.W.; (1986). percutaneous insertion of central feeding catheters. *arch dis. child*; 61 : 1138 - 40 .
- Pyatl, S.P.; Pildes, R.S.; Jacobes, M.M.; Raval, D.S.; Amma.P.; Lillen. L.; (1983) .Penicillin in infants weighing two kilograms or less with early oncet group B streptococcal disease . *N. Engl. J. Med.*; 308 : 1383 - 9 .
- quie, P.G.; (1976). Neonatal septicemia. *Antibiotic. chemother*; 21 : 128 .
- Raff, H.V.; Siscoe, P.J.; wolf, E.A.; Maloney, G.; shuford, W.; (1988). Human monoclonal antibodies to GBS : reactivity and in vivo protection against multiple serotypes . *J. Exp. Med.*; 168 : 905 - 17 .
- Ramin, S.M.; Porter, J.C.; Gilstrap, L.C.; Rosenfeld, C.R.; (1991). Stress hormones and acid- base status of human fetuses at delivery *J. clin. endocrinol. Metab.*; 73 (1) . 182 - 186 .
-

- Rayan, K.; and Hopper, B.R.; (1974). placental biosynthesis and metabolism of steroid hormones in primates . contribution in perinatology; 3 : 258 .
- Reed, R.K.; larter. W.E.; sieber, D.F.; (1976). Peripheral nodular lesions in pseudomonas sepsis . J. pediatri; 88 : 97 .
- Remington, J.S. and klein, J.O; (1983). Infectious diseases of the fetus and newborn infant, 1 st edition . philadelphia, W.B. saunders, Co. P : 142.
- Riad- Fahmy, D.; Read, G.F.; Gaskell, S.J.; (1970). A simple direct radioimmnnassay for plasma cortisol, featuring a ¹²⁵ I- radioligand and a solid- phase separation technique. Clin. chem.; 25 : 665 - 8 .
- Rivero, N.C.; Chernow, B.; stoiko, M.A.; todres, D.; (1989). Hypocalcemia in critically ill children. J. pediatri; 114 : 946 - 51 .
- Robert, M.L; (1979). Neonatal bacteremia : diagnosis and mangement. Brit. Med. J; 6202 : 1385 - 6 .
- Roberton, N.R.C.; (1986). Neonatal infections. in textbook of neonatology. churchill livingstone, London, Melbourne, New york. P : 738.
- Robson, A.M.; (1992) . Sodium, Potassium and calcium. In : Nelson Textbook of pediatrics 14 th Edition, vaughan, V.C.; Nelson, W.E.; Behrman, R.E. (editors). W.B. Saunders, Co. London, Toronto, Tokyo. P : 182.

Rockicki, W.; Forest, M.; Bonnet, H.; Bertrand, J.; (1990). Free cortisol of human plasma in the first three months of life . Biol . Neonate; 57 (1) : 21 - 9 .

Rodwell, L.R.; Anton, L.L; David. T.; (1988). Early diagnosis of neonatal sepsis using a haematological scoring system . J. pediatr; 112 : 761 - 7 .

Roger, S.; and Gordon, B.; (1990). Morphology of the adrenal medulla and cortex . In : principles and practice of endocrinology and metabolism. by Becker. K.L. (Ed) . J.B. lippincott, Co. london, New york, sydney . P : 572.

Rokicki, w.; Krasnodebski, J.; cwik, A; lewandoski.; J.; (1981). Neonatal blood aldosterone . pediatr. pol.; 64 (6) : 395 - 400 .

Rozycki, H.J.; stahl, G.E; Baumgart.s; (1987). Impaired sensitivity of a single early leucocytic count in screening for neonatal sepsis . pediatr. inf. dis. J; 6 : 440 - 2 .

Sacchi. F., Rondini. G, Mingrat.g; stronati M and siccordi. A.G.; (1982). Different maturation of neutrophil chemotaxis in term and preterm newborn infants (clinical and laborotory observation) J. pediatr; 101 (2) : 273 - 274 .

Sanchez, G.J.; pryor, R.W.; parker, M.K.; Fry. H.D.; (1989).

Hypercalcitoninemia and hypocalcemia in acutely ill children : studies in serum clacium, blood ionized calcium, and calcium- regulating hormones . J. pediatri.; 114 : 952 - 6 .

Sanchez, P.; Siegel. J.D.; Cushion. N.B; Norma. T; (1990).

Significance of a positive urine group B streptococcal latex agglutination test in neonates . J. pediatri.; 116 : 601 - 6 .

Sann.;; Bienvenu, F; Bourgeois, J.; Bethenod. M; (1984). Evolution of serum prealbumin, CRP and orosomucoid in neonates with bacterial infection J. pediatri.; 105 : 977 - 81 .

Scheifele, DW.; Melton, P.; whitcelo, V; (1981). Evaluation of the limulus test for endotoxemia in neonates with suspected sepsis J. Pediatri; 98 : 899 .

Schiff, D.; Stern, L.; Leduc, J.; (1966). Chemical thermogenesis in newborn infants . Catecholamine excretion and the plasma non- esterified fatty acid response to cold exposure. Pediatri.; 37 : 577 .

Schmeling, D.J.; and Coran, A.G.; (1991). Hormonal and metabolic response to operative stress in the neonate .JPEN. parenter- enteral- Nutr.; 15 (2) : 215 - 38 .

- Schwab, K.O; Kruse, K.; Breitung, B.; Paulick, R.;(1989). Free and sulfoconjugated plasma catecholamines in premature infants and mature newborn infants after birth . *Monatsschr kinderheilkd*; 137 (1) : 28 - 32.
- Seely, J.R.; (1974). The fetal and neonatal adrenal cortex . In metabolic, endocrine, and genetic disorders of children . Kelley, V.E. (Ed). Harper & Row. New york, London . P : 225 .
- Sherman, M.P.; Goetzman, B.W.; Ahlfors, C.E.; weinberg, R.P; (1980). Tracheal aspiration and its clinical correlates in the diagnosis of congenital pneumonia . *pediatr*; 65 : 258 .
- Shibusawa, H.; Sano, y.; okinaga, S.; (1980). Studies on 11 β - hydroxylase of the human fetal adrenal gland. *J. steroid- biochem*; 13 : 881 - 887 .
- Shigeoka. A, Santos. J, Hill. H (1979). Functional analysis of neutrophil granulocytes from healthy infected and stressed newborns . *J. pediatr*; 95 : 454 .
- Shinefield, H.R (1983). Staphylococcal infections . in : infectious disease of the fetus and newborn infant, Remington . J.S and klein, J.O (Eds), philadelphia, W.B saunders, Co. P : 882 .
- Shohaib, S.; EL- far, N.; Zoarir, A.; (1989) . Blood culture and throat culture in neonates with respiratory tract infections . M. SC. Thesis, *pediatr*; tanta university.

Siegel, S.R.; Fisher, D.A.; william, o.H.; (1973). Renal function and serum aldosterone levels in infants with respiratory distress syndrome. J. pediatr.; 83 (50) : 854 - 858 .

Siegel, S.R.; Fisher, D.A.; williams, o.H.; (1974). Serum aldosterone concentrations related to sodium balance in the newborn infant . pediatr.; 53 (3) : 410 - 413 .

Siegel, J.D.; McCracken, G.H; (1981). Sepsis neonatorm. N. Engl. J. Med; 304 : 642 .

Siegel. J. D; (1985). Neonatal septicaemia. semin perinatol.; 9 : 20 - 28 .

Simon, C.; Schroder, H.; Beyer, C.; Zerbst, T.; (1991). Neonatal sepsis in an intensive care unit and results of treatment. infection ; 19 (3) : 146 - 9.

Simonian, M.; Capp, M.; (1984). Characterization of steroidogenesis cell cultures of the human fetal adrenal cortex : comparison of definitive zone, and fetal zone cells . J. clin. endocrinol. Metab; 59 : 643 .

Simpson, E.; Carr, B.; Parker, C.; Porter, J.; (1979). The role of serum lipoprotein in steroidogenesis by the human fetal cortex . J. clin endocrinol. Metab.; 49 : 146 .

Sippell, W.G.; Becker, H.; Versmold, H.; Knorr, D.; (1978).
longitudinal studies of plasma aldosterone, corticosterone, Deoxy
corticosterone, progesterone, 17 - hydroxyprogesterone, cortisol, and
cortisone determined simultaneously in mother and child at birth and during
the early neonatal period . J. clin . endocrinol. Metab; 46 : 971 - 82.

Sklavunu, T.; Gatzola, K.; Hatzioannidis, K.; (1991). Efficacy of
aztreonam in the treatment of neonatal sepsis . Rev. infect. dis.; 13 (suppl
7) : S591 - 3 .

Slagle, T.A.; Bifano, E.M.; wolf, J.W.; Gross, S.J.; (1989).
Routine endotracheal culture for the prediction of sepsis in ventilated
babies. arch. dis. child.; 64 : 34 - 38 .

Slotkin, T.A.; Seidler, F.J.; (1988). Adrenomedullary catecholamine
release in the fetus and newborn : secretory mechanisms and their role in
stress and survival . J. Dev. physiol.; 10 (1) : 1 - 16 .

Speck, W.T.; Fanaroff, A.A; klans. M; (1979). Neonatal infection . In :
Care of the high risk neonates, 2nd Edition, W.B. Saunders, Co.
philrdelphia, London, toronto. P : 267 - 293 .

Speer, C.P.; hauptmann, D.; Stubble, P.; Gahr, M.; (1985).
Neonatal septicaemia and meningitis in Göttingen, west germany . pediatr.
infect. dis; 4 : 36 - 14 .

Speer, C.P.; Ninjo, A.; Gahr. M; (1986). Elastase- α 1- proteinase inhibitor in early diagnosis of neonatal sepsis . J. pediatri; 108 (6) : 987 - 990.

Sperling, M.A.; (1980). Newborn adaptation : adrenocortical hormones and AcTH . In : maternal- fetal endocrinology . tulchinsky, D. and Ryan, K. (Eds) . philadelphia, W.B. saunders, Co. P : 387 .

spiegelblatt. L; Saintonge. J; chicoine. R; (1985). Changing pattern of neonatal streptococcal septicemia . pediatri. infect. dis; 4 : 56 .

Squire, E.; Favara, B.; Todd, J.; (1979). Diagnosis of neonatal bacterial infection : hematologic and pathologic findings in fatal and nonfatal cases . pediatri.; 64 (1) : 60 - 65 .

Stang, H.J.; Gunnar, M.R.; snellman, L.; Condon, L.M.; (1988). local anaesthesia for neonatal circumcision; effects on distress and cortisol response . JAMA; 259 (10) : 1507 - 11 .

Starr, S.E; (1985). Antimicrobial therapy of bacterial sepsis in the newborn infant . J. pediatri.; 106 (6) : 1043 - 1048 .

Stern, L.; lees, M.H.; Lednc, J.; (1965). Enviromental temperature, oxygen consumption, and Catecholamine excretion in newborn infants. pediatri.; 36 (3) : 367 - 372 .

- Storm, W.; (1985). Prognostic implication of functional hyposplenism in neonatal septicaemia . Acta. paediatr. scand; 74 : 611 - 12 .
- Strauss, R.G.; (1988). Granulopoiesis and neutrophil function in the neonate. In : stockman, JA; Pochedly, C. (Eds) . developmental and neonatal Hematology. New york : Raven press. P : 88 - 101 .
- Strauss, R.G.; (1991). Transfusion therapy in neonates. AJDC; 145 : 904 - 11 .
- Sulyok, E.; varga, F.; Gyory, E.; Jobst, L.; Gaba, I.; (1979). postnatal development of renal sodium handling in premature infants. J. paediatr.; 95 : 787 .
- Sulyok, E.; kovacs, L.; Lichardus, B.; varga, L.; Ertl, T.; (1985). late hyponatremia in premature infants : role of aldosterone and arginine vasopressin. J. paediatr.; 106 : 990 - 995 .
- Susan, Alicia A.; Kennard, L.; (1991). Factors associated with umbilical catheter - related sepsis in neonates. AJDC; 145 : 675 - 680 .
- Symington, T.; (1982). The adrenal cortex. In : endocrine pathology. general and surgery, by Bloodworth, J.M.B. (ED). Baltimore, wolliams & wilkins. P : 419 .
-

Takala A.K, pekkanen. E, Eskola. J (1991). Neonatal haemophilus influenzae infections. Arch. dis. child; 66 : 437 - 440 .

Talbert, L.M.; Kraybill, F.E.; potter, H.D.; (1976). Adrenal Cortical response to circumcision in the neonate. obstet. & gynecol.; 48 (2) : 208 - 10

Tate, D.y.; Carlton. G.T.; Johnson, D.; white. J.; (1981). Immune thrombocytopenia in severe neonatal infections. J. pediatr.; 98 (3) : 449 - 453.

Tennes, K.; Vernadakis, A.; (1977). Cortisol excretion levels and day time sleep in one- year - old infants . J. clin. endocrinol. Metab.; 44 : 175 .

Tietz, N.W.; (1970). Fundamentals of clinical chemistry. W.B. saunders, Co. philadelphia.

Togari, H.; Iwanaga, T.; Mastsumoto, A.; Hagusawa, T.; Ogino, R.; Kito, H.; (1983). Endotoxin clearance by exchange blood transfusion in septic shock neonates . Acta. pediatr. scaud.; 72 : 87 - 91 .

Togari, M.; Sugiyama, S.; OGino, T.; Suzuki, S.; Ito, T.; Kamiya, K.; (1986). Interactions of endotoxin with cortisol and acute phase proteins in septic shock neonates. Acta. pediatr. Scand.; 75 : 69 - 74 .

Tollner, U.; (1982). Early diagnosis of septicemia in the newborn : clinical studies and sepsis score. Eur. J. pediatr.; 138 : 331 - 7.

- Tullus, B.; Lundstrom, G.; Burman, G.; (1991).** Antibiotic susceptibility of 629 bacterial blood and CSF isolates from Swedish infants and the therapeutic implication. *acta. paediatr. scand.*; 80 : 205 - 212 .
- Tyrrell, J.B.; Aron, D.C.; Frosham, P.H.; (1991).** Glucocorticoids & adrenal androgens . In : basic and clinical endocrinology, 3rd Edition, by Greenspan, F.S. (Ed), Los Altos, California, Appleton & Lange . P : 323 .
- Ungar, A.; Phillip, J.H.; (1983).** Regulation of adrenal medulla . *physiol. Rev.*; 63 : 787 - 843 .
- Vain, N.E.; Mazlumian, J.R.; Swarner, O.W.; (1980).** Role of exchange transfusion in the treatment severe septicaemia. *paediatr.*; 66 - 693 .
- Van der heijden, A.J.; Wolff, E.D.; Nauta, J.; (1987).** Renal tubular function in preterm neonates . *Tijdschr kindergeneeskd*; 55 (5) : 186 - 91 .
- Vawter, G.E; (1981).** perinatal listeriosis. in : Rosenberg, H.S; Bernstein, J. (Eds), *pediatric pathology* 2nd Edition, Masson Publishing Co, USA. P : 153 .
- Vesikari, T.; Isolauri, E.; Tuppurainen, N; Renlund, M; Koivisto, M and Kunnas, M; (1989).** Neonatal septicaemia in Finland 1981 - 1985. *Acta. paediatr. scand.*; 78 : 44 - 50 .
- Visser, V.E.; Robert, T; Hall, M.; (1979).** Urine culture in the evaluation of suspected sepsis. *J. paediatr.*; 94 (4) : 635 - 639 .

- Voorhess, M.L.; (1974). Disorders of adrenal medulla. In : Metabolic, endocrine, and genetic disorders of children, Kelley, V.C. and Limbeck, G.A. (EDS) . London, San Francisco, New York. Harper & Row, Publisher, Inc. P : 357 .
- Weiner, D.; Smith, J.; Dahlem, S.; Berg, G.; (1987). Serum adrenal steroid levels in healthy full-term 3 - day old infants. J. pediatr.; 110 (1) : 122 - 124 .
- Weldon, V.V.; Kowarski, A.; Migeon, J.C.; (1967). Aldosterone secretion rate in normal subjects from infancy to adulthood. pediatr.; 39 (5) : 713 - 723 .
- Whitelaw, A.; (1990). Treatment of sepsis with IgG in very low birth weight infants . arch. dis. child.; 65 : 347 - 8 .
- Wilkinson, C.W.; (1989). The adrenal cortex . In : textbook of physiology, 21st Edition, by Patton, H.D; Fuchs, A.F.; Hille,.; Scher, A.; Steiner, R. (EDS), Philadelphia, Sydney, Tokyo, W.B. Saunders, Co. P : 1501 .
- William. A; Douglas. S; Jane. M; Narendra. K; (1991). Echerichia coli bacteremia in children. AJDC; 145 : 671 - 4 .
- Wilson, H.D.; Eichenwald, H.F.; (1974). Sepsis neonatorum. ped. clin. Nor. AM.; 21 : 571 - 582 .
-

- Wilson, J.; and Foster, D.W.; (1985). The adrenal cortex . In : williams textbook of endocrinology, 7 th Edition, philadelphia, toronto, tokyo, W.B. saunders, Co. P : 822 .
- Winston, D.J.; McGrattan, M.A.; Busuttil, R.W.; (1984). Imipenem therapy of pseudomonas aeruginosa and other serious bacterial infections . Antimicrob. agents. chemother.; 26 : 673 .
- Winter, J.S.D.; (1985). The adrenal cortex in fetus and neonate. In : adrenal cortex. Anderson, D.C. and winter, J.S.D (Eds), Butterworths, london, Boston, toronto. P : 32 .
- Wong, S.N; (1991). Haemophilus influenza in neonates. J. Pediatr. child. health; 27 (2) : 113 - 5.
- Xanthou, M.; Xypolyta, A.; Anagnostakis, D.; (1975). Exchange transfusion in severe neonatal infection with sclerema. arch. dis. child; 50 : 901 .
- Yashiro, Y.; kudo, T.; Kishimoto, Y.; (1985). Catecholamines in amniotic fluid as indicator of intrapartum fetal stress . Acta. Med. okayama.; 39 (4) : 253 - 63 .
- Yoder, P.R.; Gibbs, R.S.; Balanco, J.D.; (1983). A prospective, controlled study of maternal and perinatal outcome after intra - amniotic infection at term . Am. J. obstet . Gyn.; 145 : 695 - 8 .

- Yoder, P.C.; and polin, R.A.; (1986). Immunotherapy of neonatal septicemia. *ped. clin . Nor. Am.*; 33 (3) : 481 .
- Yoshida, N.; Sekiba, K.; Sano, Y.; Arai, K.; (1978). Biosynthetic pathways for corticoids and androgen formation in human fetal adrenal tissue in vivo. *Endocrinol. Japonica*; 25 : 191 .
- Yuen, B.; and Mincey, K.; (1987). Human chorionic gonadotropin, prolactin, estriol, and dehydroepiandrosterone sulfate concentration in cord blood of premature and term newborn infants : relationship to the sex of the neonate . *Am. J. obstet. gynecol.*; 156 : 396 - 400 .
- Zeitoun, M.M; hassan, A.I; Hussain, Z.M.; Fahmy, M.S.; hussein, M.; (1973). Adrenal glucocorticoid function in acute viral infections in children . *Acta. pediatri. scand.*; 62 : 608 - 14 .
- Zipursky, A.; Palko, J.; Milner. R.; Akenzua, G.; (1976). The heamatology of bacterial infections in premature infants. *pediatr.*; 57 (6) : 839 - 853.
- Zipursky, A.; and Jaber, H.; (1978). The Haematology of bacterial infection in newborn infants . *clin. Haematol*; 7 : 175 .