

CONTENTS

	Page
INTRODUCTION	1
REVIEW OF LITERATURE	7
MATERIAL AND METHODS	
1. Isolation of microorganisms	18
1.1 Collection of soil samples	18
1.2 Treatment of soil samples	18
2. Isolation of actinomycetes	18
3. Isolation and identification of dermatophytes fungi	20
3.1 Isolation of dermatophytes from patients	20
3.2 Identification of dermatophytes fungi	20
3.2.1 Mycological laboratory work	20
3.2.2 Other confirmatory tests were also performed for proper identification of species	21
4. Screening of the actinomycete isolates for antifungal activities	24
4.1 Taxonomically and biological studies of the selected actinomycete isolates.....	24
4.1.1 Identification of genera	24
4.1.1.1 Methods used for morphological studies	24
4.1.1.2 I.S.P. methods	25
4.1.1.3 The cover slip culture technique	25
4.2 The methods used for the classification of genera	26
4.2.1 Culture media	26
4.2.2 Additional characterization media used by Waksman (1961) .	26
4.2.3 Morphological studies	27
4.2.3.1 Morphology of spore chain and spore bearing hyphae ...	27
4.2.3.2 Electron- microscopy of spore surface	28
4.3 Physiological and biochemical properties	28

5. Optimization of environmental factors and nutritional requirements for antifungal production	33
6. Determination of growth rate	33
7. Determination of protein	33
8. Determination of inhibition percentages	34
9. Production of antifungal product	34
10. Effect of essential oils on dermatophyte fungal growth ...	35
11. Effect of propolis (bee glue) on dermatophyte fungal growth ...	35
12. Effect of commercial antifungal drugs and antifungal product of <i>Streptomyces kanamyceticus</i> EHE-68 on dermatophytes fungal growth	36
13. Index of experimental media	37

RESULTS

Chapter I

1. Isolation and identification of the most active isolate of actinomycetes producing antifungal compound	45
1.1 Isolation of actinomycete isolate from different soil samples and screening for production of antifungal	45
1.2 Identification of isolate 68 (produces of the highest antifungal compound).....	51
1.2.1 Diagnostic characteristics and identification of isolate number 68..	51
1.2.2 Physiological and biochemical properties of isolate number 68	54
i- Utilization of carbon sources	54
ii- Growth on Czapek's medium	54
iii- Other physiological and biochemical properties of isolate numbers 68	54
iv- Growth test	54
v- Resistance to certain antibiotics	55
2. Isolation and identification of dermatophyte fungi.....	63

Chapter II

Factors affecting the antifungal biosynthesis by *Streptomyces kanamyceticus* EHE-68

1- Environmental factors

- 1.1 Effect of different incubation periods on the extracellular protein, antifungal biosynthesis, growth rate and final pH of *Streptomyces kanamyceticus* EHE-68 70
- 1.1.1 Shaking condition at 180 rpm 70
- 1.1.2 Static condition 71
- 1.2 Effect of different incubation temperatures on the production of antifungal, extracellular protein, final pH and growth rate of *Streptomyces kanamyceticus* EHE-68 76
- 1.3 Effect of initial pH values on the antifungal production, extracellular protein, final pH and growth rate of *Streptomyces kanamyceticus* EHE-68..... 79

2- Nutritional Requirements

- 2.1 Effect of different carbon sources on antifungal production, extracellular protein, growth rate and final pH of *Streptomyces kanamyceticus* EHE-68 82
- 2.2 Effect of different nitrogen sources on antifungal production, extracellular protein, growth rate and final pH of *Streptomyces kanamyceticus* EHE-68 85
- 2.3 Effect of different phosphorous sources on antifungal production, extracellular protein, growth rate and final pH of *Streptomyces kanamyceticus* EHE-68 91
- 2.4 Effect of different metallic-ions on anti-microbial potentiality, extracellular protein, growth rate and final pH of *Streptomyces kanamyceticus* EHE-68 94

Chapter III

1- Effect of different concentration of essential oils on the growth rate of tested dermatophyte fungi	97
2- Effect of different concentration of propolis on the growth rate of tested dermatophyte fungi	114
3- Fungistatic and fungicidal activity of tested essential oils and propolis against <i>T. rubrum</i> , <i>T. mentagrophytes</i> and <i>M. canis</i> ...	119
4- Effect of different concentration of commercial drugs (griseofulvin, ketoconazole and nystatin) on the growth rate of tested dermatophyte fungi	122
5- Minimum fungal concentrations (MFCs) of some antifungal drugs, essential oils and antidermatophyte product by <i>Streptomyces kanamyceticus</i> EHE-68	125
DISCUSSION	128
SUMMARY	139
REFERENCES	142
ARABIC SUMMARY	