
REFERENCES

- Agate, A.D. and Bhat, J.V.(1963): A method for the preferential isolation of actinomycetes from soils. *Antonie van Leeuwenhock. J. Microbiol. and Serol.*, 29: 297.
- Ahmad, M.S. (1990): Production of some antibiotics by certain microorganisms, Ph. D. Thesis. Fac. Sci. All-Azhar Univ.
- Ajello, L. and Gerog, L.K. (1957): In vitro hair cultures for differentiating between typical isolates of *Trichophyton mentagrophytes* and *Trichophyton rubrum*. *Mycopathol. Mycol. Appl.*, 8: 3.
- Alderman, G.G. and Marth, E.H. (1976): Inhibition of growth and aflatoxin production of *Aspergillus parasiticus* by citrus oils. *Z. Lebensm. Unters. Forsch.*, 160: 353-358.
- Anker, J.N.; Popelevan, N.M.L. and Sauer, P.J.Z. (1995): Antifungal, antimicrob Agents *Chemother.*, 39: 1391-1397.
- Baratta, M.T.; Dorman, H.J.D.; Deans, S.G.; Figueiredo, A. C.; Barroso, J.G. and Ruberto, G. (1998): Antimicrobial and antioxidant properties of some commercial essential oils. *Flavour and Fragrance*, 13: 235-244.
- Barbara, J.; John, F.; Alice, S.; Thomas, F. and Richard, C. (1994): Clinical and pathogenic microbiology, Mosby- year book-2nd ed. pp.587.
- Bergey's Manual of Determination Bacteriology (1979): Determinative Bacteriology, 8th ed. Williams and Wilkin (Eds.) Co., Baltimore, London.

- Bergey's Manual of Systemic Bacteriology (1989): By Williams, T. Stanely and Harriet Flescher. Vol. 4. Williams & Wilkins Co., Bltimore., Hong kong, London, Sydney.
- Bergey's Manual of Determination Bacteriology Minth Edition (1994): By Holt, J.G.; Krieg, N.R.; Sneath, P.H.A.; Staley, J.T. and Williams, S.T.
- Bertasso, M.; Holzen Kampfer, M.; Zeeck, A.; Stacke-brandt, E.; Beil, W., Fiedler, HP. (2003): Ripromycin and other polycyclic macrolatams from *Streptomyces* sp. Tu6239: Taxonomy, fermentation, isolation and biological properties. J. Antibiot.(Tokyo). 56(4): 364-371.
- Bocobo, F.C. and Benham, R.W. (1949): Pigment production in the differentiation of *Trichophyton mentagrophytes* and *Trichophyton rubrum*.. Mycologia., 41: 291.
- Brewer, G.A., Jr. and Frazier, W.R. (1962): "Antimicrobial agents & Chemother- 1961" ed. Findland, M. & Savate, G. M., American society of Microbiology, Michigan, MSA, pp. 212-217.
- Cardile, V.; Panico, A.; Gentile, B.; Borrelli, F. and Russo, A.. (2003): Effect of propolis on human cartilage and chondrocytes. Life Sci. 11; 73 (8): 1027-1035.
- Chandler, F.W.; Kaplan, W. and Ajello, L. (1980): Histopathology of mycotic infections. Chicago: Year Book.
- Chattopadhyay, D. and Sen, S.K. (1997): Optimization of cultural conditions for antifungal antibiotic accumulation by *Streptomyces rochei* G164. Hind. Antibiot. Bull. 39(1-4): 64-71.
- Chu, M.; Truumees, I.; Mierzwa, R.; Terracciano, J.; Patel, M.;

- Loebenberg, D.; Kamminski, J.J.; Das, P. and Puar, M.S. (1997): A new polycyclic xanthone Sch54445 with highly potent antifungal activity produced by *Actinoplanes sp.* J-Nat-Prod. 60(5): 55-528.
- Cidaria, D.; Borgonovi, G. and Pirali, G. (1993): Novel polyene antibiotics AB023. I. Taxonomy of the producing organism, fermentation and antifungal activity. J. Antibiot. (Tokyo). 46(2): 251- 254.
- Conant, N.F. (1971): Manual of clinical mycology. 3rd ed. Philadelphia, Saunders.
- Conner, D.E. and Beuchat, L.R. (1984): Effects of essential oils from plants on growth of food spoilage yeast. J. Food Sci. 49(2) : 429-434.
- Cooper, R.; Horan, A.C.; Gentile, F.; Gullo, V.; Leobenberg, D.; Marquez, J.; Patel, M.; Puar, M.S. and Truumees, I. (1988): A novel antifungal compound produced by *Micromonospora sp.* Taxonomy, fermentation, isolation, structure and biological properties. J. Antibiot. (Tokyo). 41(1): 13-19.
- Cooper, R.; Truumees, I.; Gunnarsson, I.; Loebenberg, D.; Horan, A.; Marquez, J.; Patel, M.; Gullo, V.; Puar, M. and Das, P. (1992): A novel antifungal antibiotic from an *Actinoplanes sp.* Fermentation, isolation, structure and biological properties. J. Antibiot. (Tokyo). 45(4): 444-453.
- Cowan, S.T. (1974): Cowan and Steel's manual for the identification of medical bacteria; 2nd ed. Cambridge. Univ. press, London.
- Crawford, D.L. and McCay, E. (1972): Cellulose of *Thermomonospora fusca* and *Streptomyces thermocliastaticus*. Appl. Microbiol., (24): 150.

- Cross, T., Maciver, A. and Laoy, J. (1968): The thermophilic *Actinomyces* in mouldy hay *Microspolyspora foeni* sp. J. Gen. Microbiol., 50: 351-359.
- Cutler, J.E. (1991): Putative virulence factors of *Candida albicans*, Ann. Rev. Microbiol., 45: 187-218.
- Daouk, R.K.; Dagher, S.M. and Sattout, E.J. (1995): Antifungal activity of the essential oil of *Orignum syriacum* L.J. Food Prot. 54: 1147-1149.
- Deans, S.G. and Ritchie, G. (1987): Antibacterial properties of plant essential oils. Int. J. Food Microbiol., 5: 165-180.
- Delespaul, Q.; Billerbeck, V.G.D.; Roques, C.G.; Michel, G.; Vinuales, C.M. and Bessiere, J.M. (2000): The antifungal activity of essential oils as determined by different screening methods. J. Essential oil Res., 12: 256-566.
- Deshmukh, S.K.; Jain, P.C. and Agrawal, S.C. (1986): A note on mycotoxicity of some essential oils. Fitoterapia LVII 4, 295.
- Drago, L.; Mombelli, B.; De Vecchi, E.; Fassina, M.C.; Tocalli, L. and Gismonodo, M.R. (2000): In vitro antimicrobial activity of propolis dry extract. J. Chem., 12: 390-395.
- Dukić, N.M.; Božin, B.; Sokovic, M.; Mihajlovic, B. and Matavulj, M. (2003): Antimicrobial and antioxidant activities of three *Mentha* species essential oils. Planta Med., 69: 413-419.
- Dwivedi, S.K. and Dubey, N.K. (1993): Potential use of the essential oil of *Trachyspermum ammi* against seed-borne fungi of Guar (*Cyamopsis tetragonoloba* L. (Tab.). Mycopathologia., 121: 101-104.

- Egorov, N.S. (1985): Antibiotics, a scientific approach MIR publishers, Moscow .
- Ellaiah, P.; Venkateswara Rao, V.S.; Rama Krishna, S.V.; Ven Kateswarulu, Y.; Annapurna, J. and Venkatesam, U. (1998): A new *Streptomyces* producing diphenyl sulfone antibiotic-S. sulfonensis. J. Hind. Antibiot. Bull. 40: 31-37.
- Elliott, T.; Hastings, M. and Dessel berger, M. (1997): Medical microbiology. Antifungal Agents 3rd ed. 27: 194-195.
- Elwan, S.H.; Ammar, M.S. and EL-Moussalomy, M.K.(1986): Relation of the *Penicillium chrysogenum* Lipase to α - amylase biosynthesis and some factors affecting the crude lipase activity- Egypt. J. Microbiol., 21(2): 129-142.
- Emmons, C.W.; Binford, C.H. and Kwonchung, K.J. (1977): Medical mycology, Lea and Febiger. 3rd ed. Philadelphia.
- Farag, R.S.; Daw, Z.Y. and ABO. Raya, S.H. (1989): "Influence of some spice essential oils on *A. parasiticus* growth and Production of aflatoxins in synthetic medium. J. Food Sci., 54(1): 74-76.
- Farid, M.A.; El-Enshasy, H.A.; El-Diwany, A.I. and El-Sayed, S.A. (2000): Optimization of cultivation medium for natamycin production by *Streptomyces natalensis*. J. Basic-Microbiol., 40(3): 157-166.
- Frandberg, E.; Petersson, C.; Lundgren, L.N. and Schnurer, J. (2000): *Streptomyces halstedii* k122 produces the antifungal compounds bafilomycin B1 and C1. Can. J. Microbiol., Aug., 46(8): 753-758.
- French, R.C. (1985): The bioregulatory action of flavor compounds on fungal

- spore and other popagules. *Ann. Rev. Phytopathol.*, 23: 173-199.
- Fries, N. (1973): Effects of volatile organic compounds on the growth and development of fungi. *Trans. Br. Mycol. Soc.*, 60: 1.
- Furumai, T.; Hasegawa, T.; Kakushima, M.; Suzuki, K.; Yamamoto, S.; Hirano, M. and Oki, T. (1993): Pradimicins T1 and T2, new antifungal antibiotics produced by an actinomycete. I. Taxonomy, production, isolation, phsico-chemical and biological properties. *J. Antibiot. (Tokyo)*. 46(4): 589-597.
- Garg, R.C. and Siddiqui, N. (1992): Antifungal activity of some essential oils isolate. *Pharmazie*., 47: 467-468.
- Gautam, M.P.; Jain, P.C. and Singh, K.V. (1988): Activity of some essential oils against dermatophytes. *Indian Drugs* 17(9): 269.
- Geoff, H., Baker; Simon, E.; Blanch, F.; Roderick, J.J.; Dorgan, R.; Everett, B. R., Manger, C. R.; Reading, S. A.; Read, S. and Peter S. (1995): Further novel Milbemycin antibiotic from *Streptomyces* sp. 5217-02. *J. Antibiot. (Tokyo)*. 49(12): 1226-1231.
- Georgopapadakou, N.H and Walsh, T.J. (1996): Antifungal agents: Chemotherapeutic target and immunological strategies. *Antimicrob Agents Chem.*, 40: 279-291.
- Ghaly, M.F.E. (1985): Studies on some *Actinomycetes* producing amylase enzyme. Ph.D. Thesis Botany Department, Fac. Sci. Zagazig Univ. EL-Nakeeb, M.A. and Lechevalier, H.A. (1963). *Appl. Microbiol*, H: 75-77.
- Glazebrook, M.A.; Vining, L.C.; White, R.L.; Smith, K.C. and Chedrawy, E.G. (1993): Nutrient effects on growth and the production of 5-

- hydroxy-4-oxonorvaline by *Streptomyces akiyoshiensis*. Can. J. Microbiol., 39(5): 536-542.
- Goodfellow, M. and Orchard, V.A. (1974): Antibiotic sensitivity of some nocardioform bacteria and it's value as criterion for taxonomy. J.Gen. Microbiol., 83: 375-387.
- Gordon, R.E. (1966): Some criteria for the recognition of *Nocardia madura* (vincent) Blanchard. J. Gen. Microbiol., 45: 355-364.
- Gupte, T.E. and Naik, S.R. (1998): Optimization of nutritional requirements and process control parameters for the production of HA-91, a new Tetraene polyene Antibiotic Hind. Antibiot. Bull., 40: 5-13.
- Hacene, H.; Kebir, K.; Othmane, D.S. and Lefebvre, G. (1994): A new polyene antifungal antibiotic HM17 produced by a new strain of *Spirillospora*. J.Appl.Bacteriol., 77(5): 484-489.
- Hacene, H. and Lefebvre, G. (1995): a new non-polyenic antifungal antibiotic AH17 produced by a strain of *Spirillospora*. Microbiol., 83(336): 199-205.
- Hacene, H.; Boudjellal, F. and Lefebvre, G. (1998): A non-polyenic antifungal antibiotic AH7 produced by a new strain of *Streptosporangium roseum*. Microbiol., 96(384): 103-109.
- Hanafi, M.; Shibata, K.; Ueki, M. and TaKniguchi, M. (1996): Novel antifungal antibiotics UK-2A, B, C and D from *Streptomyces* sp. 5217-5302. J. Antibiot. (Tokyo). 49(12): 1226-1231.
- Harman, A.W. (1983): Hive products for therapeutic use. J. Amer. Bee., 39: 44.

- Hesseltine, C.W.; Bendict, R.G. and Pridham, T.G. (1954): Useful criteria for species differentiation in the genus *Streptomyces*. Ann. N. Y. Acad. Sci., 60: 163.
- Holtzel, A.; Kempter, C.; Metzger, J.W.; Jung, G.; Groth, I.; Fritz, T. and Fiedler, H.P. (1998): Spirofungin, a new antifungal antibiotic from *Streptomyces violaceus niger* Tu4113. J. Antibiot. (Tokyo). 51(8): 699-707.
- Hus, S.C. and Lockwood, J.L. (1975): Powder chitin agar as a selective medium for enumeration of *Actinomycetes* in water and soil. Appl. Microbiol., 29: 422-426.
- Igarashi, M.; Kinoshite, N.; Ikeda, T.; Nakagawa, E.; Hamada, M. and Takeuchi, T. (1997): Formamicin, a novel antifungal antibiotic produced by a strain of *Saccharothrix* sp. I. Taxonomy, production, isolation and biological properties. J. Antibiot. (Tokyo). Nov., 50(11): 926-931.
- Igarashi, M.; Kinoshite, N.; Ikeda, T.; Nakagawa, E.; Hamada, M. and Takeuchi, T. (1997): Resromycin, a novel herbicidal and antifungal antibiotic produced by a strain of *Streptomyces plantensis*. I. Taxonomy, production, isolation and biological properties. J. Antibiot. (Tokyo). Dec., 50(12):1020-1025.
- Ivanov, P. (1987): A study on some qualitative parameters of Bulgarian propolis. Apimondia (Warsaw) 31.
- Jacobson, M. (1958): Insecticides from plants. Agriculture Hand book No. 154, USDA.

- Jain, P.C. and Agrawal, S.C. (1978): Notes on the activity of some odoriferous organic compounds against some keratinophilic fungi Trans.Br. Mycol. Soc., 19:197.
- Janssen, A.M.; Scheffer, J.J. and Baerheim Svendsen, A. (1987): Antimicrobial activity of essential oils: a 1976-1986 Literature review. Aspects of the test methods. Planta Med., 53: 395-398.
- Jeuniaux, C. (1966): Chitinases. Methods enzymol. 8: 644-650.
- Johnson, M.J. (1955): Recent Advances in penicillin fermentation Bull. World Health org., 6: 99-121.
- Johnson, L.F.; Curl, E.A.; Bpnd, J.H.; and Fribourg, H.A. (1959): "Methods for studying soil microflora plant disease relationships" Burgess Minneapolis.
- Jones, K. (1949): Fresh isolates of *Actinomycetes* in which the presence of sporogenous aerial mycelia is a fluctuating characteristics. J. Bacteriol., 57: 141-145.
- Junya, N.; Mayum, I.; Yayol, I. and Iwao, I. (1998): Experimental penetration of *Trichophyton mentagrophytes* into human stratum corneum. Mycopathologia. 141: 153-157.
- Jurd, L. and Manners, G. (1980): Wood extractive as models for the development of new types of pest control agents. J. Agri. Food Chem., 28: 183-188.
- Kaneda, M.; Naid, T.; Kitahara, T.M.; Nakamura, S.; Hirata, T. and Suga, T. (1988): Carbazomycins G and H, novel carbazomycin-congeners containing a quinol moiety. J. Antibiot. (Tokyo). 41(5): 602-

608.

- Kawato, M. and Shinobu, R. (1959): *A Streptomyces herbicolor* nov. sp. supplement: A simple technique for the microbial observation Mem. of the Osaka Univ. Lib. Erts of Education, 8: 144.
- Ketoki, B and Majumdar, S.K. (1973): Utilization of carbon and nitrogen sources by *Streptomyces kanamyceticus* for kanamycin production Anti-microbial agents chem., 4(1): 6-10.
- Kishore, N.; Mishra, A.K. and Chansouria, J.P.N. (1993): Fungitoxicity of essential oils against dermatophytes. Mycoses. 36: 211-215.
- Koninklijke, N.V. (1960): Pimarcin atetraene polyene antibiotic. British Patent pp.844-289.
- Koo, H.; Gomes, B.P.; Rosalen, P.L.; Ambrosano, G.M.; Park, Y.K. and Cury, J.A. (2000): In vitro antimicrobial activity of propolis and *Arnica montana* against oral pathogens. Arch. Oral Biol., 45: 141-148.
- Kotake, C.; Yamasaki, T.; Moriyama, T.; Shinoda, M.; Komiyama, N.; Furumai, T.; Konishi, M., and Oki, T. (1992): Butyrolactols A and B, new antifungal antibiotics. Taxonomy, isolation, physico.chemical properties, structure and biological activity. J. Antibiot. (Tokyo). 45(9): 1442-1450.
- Krossilnikov, N.A. (1970): Pigmentation of *Actinomycetes* and its significance taxonomy. The *Actinomycetes* (Prauser Ed.) 123-131. Guster Ficher Verlag, Jena.
- Küster, E. (1959): Introductory remarks to the round table conference on *Streptomyces*. Int. Bull Bact. Nomencl. Taxon., 9: 57.

- Küster, E. (1959a): Outline of a comparative study of criteria used in characterization of the *Actinomycetes*. Intern. Bull. Bact. Nomen and Taxon., 98-104.
- Küster, E. and Williams, S.T. (1964): Production of hydrogen sulphide by *Streptomyces* and methods for its detection. App. Microbiol., 12: 46-52.
- Küster, E. (1972): Simple working key for the classification and identification of named taxa included in the International Streptomyces Project. Int. J. Syst. Bacteriol., 22: 139.
- Kützner, H.J. (1967): Methoden zur Untersuchung von Streptomyces und geeigneten anderen Actinomyceten: Darmstadt, Teilsammlung Darmstadt an Institute für Mikrobiologie der technischen Hochschule (cit. Williams, 1983).
- Kützner, H.J. (1981): The family *Streptomycetaceae* in the prokaryotes, A handbook of habitats; Isolation and identification of bacteria, vol. II. (M. Pstarr, H. Stolp, H.G. Truper, A. Balows and H.G. Schlegel "Eds") 2028-2090. Springer. Verlag Berlin.
- Kwon-Chung, K.J. and Bennett, J.E. (1992): Medical mycology. Philadelphia: Lea and Febiger.
- Lee, M.S.; Kojima, I. and Demian, A.L. (1997): Effect of nitrogen sources on biosynthesis of rapamycin by *Streptomyces hygroscopicus*. J. Ind. Microbiol. Biotech. 19(2): 83-86.
- Lincoff, G. and Mitchell, D.H. (1977): Toxic and hallucinogenic mushroom poisoning, New York, Van Nostrand Reinhold.
- Liu, C.M., McDaniel, L.E. and Schaffner, C.P. (1975): Antimicrob. Agents Chem., 7: 196-202.

- Lowry, O.H.; Rosebrough, N.J.; Farr, A.L. and Randall, R.J. (1951): Protein measurement with folin phenol reagent . J. Biol. Chem. 193: 265.
- Mac Ginnis, M.R. (1980): Laboratory hand book of medical mycology. Academic press, New York. pp.661.
- Maresca, B.I. and Kobayashi, G.S. (1996): HSPs in fungal infections. In Van Eden W, Young D, editors: stress proteins in medicine, New York, Marcel Dekker.
- Marquez, J.A.; Horan, A.C.; Kalyanpur, M.; Lee, B.K.; Loebenberg, D.; Miller, G.H.; Patel, M. and Waitz, J.A. (1983): The hazimicins, a new class of antibiotics. Taxonomy, fermentation, isolation, characterization and biological properties. J. Antibiot. (Tokyo). 36(9): 1101-1108.
- Martin, J.F. and McDaniel, L.E. (1977): Production of polyene macrolide antibiotics. Adv. Appl. Microbiol., 21:1-52.
- Martin, J. F. and Demian, A.L. (1980): Control of antibiotic synthesis. Microbiol. Rev., 44: 230-251.
- Meddof, G.; Brajtburg, J.; Kobayashi, G.S. and Bolard, J. (1983): Antifungal agents useful in therapy of systemic fungal infections. Ann. Rev. Pharma., 23:303-330.
- Mishra, A.K. and Dubey, N.K. (1994): Evaluation of some essential oils for their toxicity against fungi causing deterioration of stored food commodities. Appl. Environ. Microbiol., 60: 1101-1105.
- Montes, B.R. and Carvajal, M. (1998): Control of *Aspergillus flavus* in maize with plant essential oils and their components. J. Food Prot., 61(5): 616-619.

- Moselio, S.; Cary, E.; Barry, I. and Medoff, G. (1998): Microbial disease. Williams & Wilkins 3rd ed. 45: PP 419-421.
- Muller-Riebau, F.; Berger, B. and Yegen, O. (1995): Chemical composition and fungitoxic properties to phytopathogenic fungi of essential oils of selected aromatic plants growing wild in Turkey. J. Agric. Food. Chem. 43: 2262-2266.
- Musilikova, M. (1961): Folia Microbiol., 6: 175-178.
- Nair, M.G.; Putnam, A.P.; Mishra, S.K.; Mulks, M.H.; Taft, W.H.; Keller, J.R.; Zhu, P.P.; Meinhart, J.D. and Lyun, D.G. (1989). Faeriefungin: a new broad spectrum antibiotic from *Streptomyces griseus* var. autotrophicus. J. Nat.Prod. 52(4): 797-809.
- Nair, M.G.; Mishra, S.K.; Putnam, A.R. and Pandey R.C. (1992): Antifungal anthracycline antibiotics, spartanamicins A and B from *Micromonospora* sp. J. Antibiot. (Tokyo). 45: 1738-1745.
- Nonomura, H. (1972): Key for classification and identification of 458. species of the *Streptomyces* included in I.S.P. J. Fermentation Techn., 52: 7.
- Nüesch, J.(1965): Isolierung und selektionierung von Aktinomyceten In: Symposium anreicherung skulturund Mu-tenauslse "Gottungen April 1964" Zentralblatt für Bakteriologie. Parasitenkunde, Infektion skrnk Heiten und hygiene, Abt. I supp., 1: 234-252.
- Okami, Y. and Suzuki, M. (1958): A simple method for microscopical observation for *Streptomyces* and critiria of *Streptomyces* grouping with references to aerial structure. J. Antibiot., 11: 250-253.
- Owen, S.P. and Johnson, M.T. (1955): The effect of temperature changes on

- the production of penicillin by *Penicillium chrysogenum*. Microbiol., 3: 375-379.
- Oxoid Limited (1982): The oxoid manual of culture media, ingredients and other Laboratory services. Turnergraphic Ltd. England 4th Ed.
- Padhye, A.A.; Young, C.N. and Ajello, L. (1980): Hair perforation as a diagnostic criterion in the identification of *Epidermophyton*, *Microsporum*, and *Trichophyton* species. PAHO Sci. pub No 396, pp. 115.
- Patel, M.; Hegde, V.; Horan, A.C.; Gullo, V.P.; Loebenberg, D.; Marquez, J. A.; Miller, G.H.; Puar, M.S. and Waitz, J.A. (1984): A novel phenazine antifungal antibiotic, 1,6-dihydroxy-2-chlorophenazine. Fermentation, isolation, structure and biological properties. J. Antibiot. (Tokyo). Sep., 37(9): 943-948.
- Patrick, R.; Ken, S.; George, S. and Micheal, A. (1998): Medical microbiology, Mosby-year book. 3rd ed. 66: pp558.
- Pepeljnjak, S.; Magsinger, D. and Jalsenjak.I. (1982): Effect of propolis extracts on some fungi. Sci. Pharma., 50:165-167.
- Pepeljnjak, S.; Magsinger, D. and Jalsenjak.I. .; Magsinger, D. and Jalsenjak.I. (1985b): Inhibition of growth and biosynthesis of ochratoxin A in *A. sulphureus* 407 by propolis extract. Apic. Abstr., 293.
- Perfect, J.R. (1996): Fungal virulence genes as target for antifungal chemotherapy. Antimicrob Agents Chemo., 40: 1577-1583.
- Philpot, C. (1967): The differentiation of *Trichophyton mentagrophytes* from *Trichophyton rubrum* by a simple urease test. Sabouraudia. 5: 189.

- Pridham, T.G. and Gottlieb, D. (1948): The utilization of carbon compounds by some *actinomycetes* as an aid for species determination. J. Bacteriol., 56(1): 107-114.
- Pridham T.G.; Linder Felser, L.A.; Shotwell, O.L.; Stodola, F.A.; Benedict, R.G.; Foley, C. and Mitchell, P.W. (1956): Antibiotics against plant disease. (I) Laboratory and green house survey-Phytopathol., 46: 568-575.
- Pridham, T.G.; Anderson, P.; Foley, C.; Lindengelser, H.A.; Hesseltine, C.W. and Benedict, R.G. (1957): A selection of media for maintenance and taxonomic study of *Streptomyces*. Antibiotics. Ann. (1956 / 1957). Medical Encyclopedia. Inc. New York. 947-953.
- Pridham, T.G.; Hesseltine, C.W. and Benedict, R.G. (1958): A guide for the classification of *Streptomyces* according to selected groups placement of strains in morphological section. Appl. Microbiol., 6: 52-79.
- Pridham, T.G. and Lyons, A.J. (1961): *Streptomyces albus* (Rossi-Doria) Waksman et. Henrici: Taxonomic study of strain labeled *Streptomyces albus*. J. Bacteriol., 81: 431-441.
- Popova, J.N.; Wilhelm, J.J. and Tresner, H.D. (1960): Optimization of nutritional requirements and control parameters for production of nystatin. Appl. Microbiol., 8: 174-178.
- Porter, J.N.; Wilhelm, J.J. and Tresner, H.D. (1960): Method for the preferential isolation of *Actinomycetes* as an aid for species determination. J. Bacteriol., 56 (1): 107-114.

- Raytapadar, S. and Paul, A.K. (2001): Production of an antifungal antibiotic by *Streptomyces aburaviensis* 1DA-28. Microbiol Res. 155(4): 315-323.
- Rebell, G. and Taplin, D. (1970): Dermatophytes; their recognition and identification. University of Miami press, coral gables, florida.
- Rippon, J.W. (1974): Ascomycetes: The dermatophytes in fungi pathogenic for human and animals. Vol.3 Mareel Dekker, Inc. New York and Basal pp.113.
- Robert, F.B. (1995): Basic medical microbiology, The pathogenic fungi, Little Brown and company. 5th ed. 33: 481.
- Rook, A.; Wilkinson, D.S. and Ebling, F.J.G. (1979): Text book of dermatology. Vol. 1. Blackwell Scientific publication. 3rd ed. Pp. 783-789.
- Saithoh, K.; Temmyo, O.; Yamamoto, S.; Furumai, T. and Oki, T. (1993): Pradimicin S, a new pradimicin analog. I. Taxonomy, fermentation and biological activities. J. Antibiot. (Tokyo). 46(4): 580-588.
- Samoilov, P.; Kruglyak, M. and Aderonskay, L.M. (1967): Effect of culturing conditions of *Actinomyces lavendulae* on the composition of phytobacteriomycin. Prikl Biokhim Microbiol., 13 (3): 388-390.
- Sarhan, M.M. (1999): Antifungal implications of some selected essential oils on citrus fruit rot. J. Union Arab Biol. Cairo vol. 8(B) Microbiology & Viruses. Nov., 65-76.
- Schaffner, C.M.; Steinman, I.D.; Safferman, R.S. and Lechevalier, H.J. (1958): "Antibiotics Annual 1957-58". Ed. Welch, H. & Marti-Ibanez, F., Medical Encyclopaedia. Inc., New York, pp 869-873.

- Shash, S. (1992): Biological, chemical and numerical taxonomical studies on grey series of *Streptomyces* from Egyptian soils, ph.D. Thesis, Fac. Sci. Benha, Zagazig Univ.Egypt.
- Shimoni, M.; Putievsky, E.; Ravid, U. and Reuveni, R. (1993): Antifungal activity of volatile fractions of essential oils from four aromatic wild plants in Israel, J. chem. Ecol. 19 (6): 1129-1133.
- Shirling, E.B. and Gottlieb, D. (1966): Methods for characterization of *Streptomyces* sp. Intern. J. Syst. Bacteriol., 16: 313-340.
- Shirling, E.B. and Gottlieb, D. (1968a): Cooperative description of type culture of *Streptomyces* sp. II. Description from first study. Intern. J. Sys. Bacteriol., 18(2): 69-189.
- Shirling, E.B. and Gottlieb, D. (1968b): Cooperative description of type cultures of *Streptomyces* sp. III. Additional species description from first and second studies. Intern. J. Sys. Bacteriol., 18(4): 279-392.
- Shirling, E.B. and Gottlieb, D. (1969): Cooperative description of type cultures of *Streptomyces* sp. description from studies. Intern. J. Sys. Bacteriol., 19: 439-512.
- Shirling, E.B. and Gottlieb, D. (1972): Cooperative description of type strains of *Streptomyces*. Additional description, Int. J. Syst. Bacteriol., 22: 262-394.
- Singh, K.V. and Deshmukh, S.K. (1984): Volatiles constituents from members of *Liljaceae* and spore germination of *M. gypseum* complex. Fitoterapia., 55(5): 297.
- Sivropoulou, A.; Papanicolaou, E.; Nicolaou, C.; Kokkini, S.; Lanaras, T.

- and Arsenakis, M. (1996): Antimicrobial and cytotoxic activities of *Origanum* essential oils .J. Agric. Food. Chem., 44: 1202-1205.
- Smith, M.A.; Perry, G. and Pryor, W.A. (2002): Causes and consequences of oxidative stress in Alzheimer's disease (1,2). Free radical Biol. Med., 32:1049.
- Staib, F.; Mishra, S.K.; Formtling, R.A.; Blisse, A.; Kleinkauf, F.H.; Mahmutoglu, I.; Muller, K.D.; Salnikow, J.;6 Hohne, G. and Schwarz, H. (1983): A new polyene with antimycotic activity produced by *Streptomyces* from the rhizosphere of an indoor plant. Arzneimittelforschung,, 33(1): 84-87.
- Stanley, W.; Chapman, M.D. and Ralph Danid, C. (1994): Cutaneous manifestations of fungal infection. Infection Disease Clinics of North America. 8 (4): 879-910.
- Stuart, C.A.; Vanstratum, E. and Rustigian, R. (1945): Further studies on urease production by *Proteus* and related organisms. J.Bacteriol., 49: 437-444.
- Sutcliffe, J.and Georgopapadakou, N.H. (1992): Emerging targets in antibacterial and antifungal chemotherapy. New York: Chapman and Hall.
- Szabo, I.M.; Marton, M.; Buti, I. and Fernandez, C. (1975): A diagnostic key of the identification of " species" of *Streptomyces* and *Streptoverticillium* included in the International Streptomyces Project. Acta Botanica. Acad. Sci. Hungavica., 21: 387.
- Tadashi, A. (1975): Culture media for *Actinomycetes*. The society for

actinomycetes. Japan. Nat. Agricul. Library. 1: 1-31.

Takiguchi, Y.H.; Muramatsu, M.S.; Mishina, I.J.H. and Mibencin, M.
(1981): A new family of macrolides antibiotics, fermentation, isolation
and Physio-chemical properties of Mibenycins, D.E, F, G and H, J.
Antibiotics, 36: 502-508.

Tereshin, I.M. (1976): "polyene antibiotic-present and future". Univ. Tokyo
press, (Tokyo). pp. 7-62.

Thirumalachar, M.J.; Menon, M.V. and Bhatt, V.V. (1961): "Polyene
antibiotic- hamycin". Hind. Antibiot. Bull., 3: 136-138.

Thirumalachar, M.J. and Menon, S.K. (1962): "polyene antibiotic –
dermostatin". Hind. Antibiot. Bull., 4: 106-108.

Thirumalachar, M.J.; Rahalkar, P.W. and Gopalkrishnan, K.S. (1964):
Polyene antibiotic –aureofungin. Hind. Antibiot. Bull. 6: 108-111.

Thom, C. and Raper, K.B. (1945): An Annual of the Aspergilli williams and
williams Co. Baltimore Md.

Thompson, D.P. (1989): Fungitoxic activity of essential oil components on
food storage fungi. Mycologia. 81: 151-153.

Towers, G.H.N.; Spencer, P.A. and Rodriguez, E. (1989): Recent topics in
phytochemical ecology research. Phytochemical ecology;
Allelochemicals, mycotoxins and insect pheromones and allomonens
(C.H. Chou, G.R. Waller, Eds.). Institute of Botany. Academia Sinica
Monograph Series No. 9. Taipei, ROC.

Trease, E. G. and Evans, C.W. (1985): Pharmacognosy, 12th ed. ELBS,
London.

- Tresner, H.D.; Davies, M.C. and Backus, E.J. (1961): Electron Microscopy of *Streptomyces* spore morphology and its role in species differentiation- J.Bacteriol., 81: 70.
- Tsao, P.H.; Leben, C. and Keitt, G.W. (1960): An enrichment method for isolating *Actinomycetes* that produces diffusible antifungal antibiotics. Phytopathol., 50:88-89.
- Ueki, M.; Kusumoto, A.; Hanafi, M.; Shibata, K.; Tanaka, T. and Tangiuchi, M. (1997): A novel antifungal antibiotic UK-3A from *Streptomyces* sp. 517-02: Fermentation, isolation, structural elucidation and biological properties. J. Antibiot. (Tokyo). 50 (7): 551-555.
- Urakawa, A.; Otani, T.; Yoshida, K.; Nakayama, M.; Suzukake- Tsuchiya, K. and Hori, M. (1993): Isolation, structure determination and biological activities of a novel antifungal antibiotic, S-632-C, closely related to glutarimide antibiotics. J. Antibiot. (Tokyo). 46(12): 1827-1833.
- Van de Vijver, L.P.; Kardinaal, A.F.; Grobb, D.E.; Princen, H.M. and van Poppel, G. (1997): Lipoprotein oxidation, antioxidants and cardiovascular risk: epidemiologic evidence. Prostaglandins Leukot Essent Fatty Acids. 57: 479-487.
- Velikova, M.; Bankova, V.; Sorkun, K.; Houcine, S.; Tsvetkova, I. and Kujumgiev, A. (2000): Propolis from the Mediterrean region: Chemical composition and antimicrobial activity. Z Naturforsch (c) 55: 790-793.
- Vertesy, L.; Aretz, W.; Ehlers, E.; Hawser, S.; Isert, D.; Knauf, M.; Kurz,

M.; Schiell, M.; Vogel, M. and Wink, J. (1998): 3874H1 and H3, novel antifungal heptaene antibiotics produced by *Streptomyces* sp. HAG 003874. J. Antibiot. (Tokyo). 51(10): 921-928.

Wagman, G.H. and Weinstein, H.J. (1980): Antibiotics from *Micromonospora*. Ann. Rev. Microbiol. 34: 537-557.

Waksman, S.A. (1919a): Culture studies of species of *Actinomycetes*. Soil. Sci. 8: 71.

Waksman, S.A. (1919b): Cultural studies of species of *Actinomycetes*. Soil. Sci. 8: 71.

Waksman, S.A. (1961): The *Actinomycetes*, vol. II & III, classification, identification and description of genera and species, the Williams & Wilkins, Baltimore.

Waksman, S.A. and Lechevalier, H.A. (1961): The *Actinomycetes* vol. II, The Williams and Wilkins. Baltimore.

Williams, S.T. and Davies, F.L. (1965): Use of antibiotic for selective isolation and enumeration of *Actinomycetes* in soil. J. Gen. Microbiol., 38: 251-262.

Williams, S.T.; Goodfellow, M.; Alderson, G.; Wellington, E.M.H. and Sockin, M.J. (1983): Numerical classification of *Streptomyces* and related genera. J. Gen. Microbiol., 129: 1793-1813.

Yamaguchi, H.; Kobayashi, G.S. and Takahashi, H. (1992): Recent advances in antifungal chemotherapy, New York, Marcel Dekker.

Yano, K.; Yokoi, K.; Sato, J.; Oono, J.; Kouda, T.; Ogawa, Y. and

- Nakashima, T. (1986): Actinopyrones A, B and C, new physiologically active substances. J. Antibiot. (Tokyo). 39(1): 32-37.
- Yano, K.; Oono, J.; Mogi, K.; Asaoka, T. and Nakashima, T. (1987): Pyrroxamycin, a new antibiotic taxonomy, fermentation, isolation, structure determination and biological properties. J. Antibiot. (Tokyo). 40(7): 961-969.
- Yassin, M.H.(1998): Studies on a potential antifungal *Streptomyces violaceus*, Al-Azhar Bull. Sci. 9(2): 369-376.
- Yassin, M.H. and Galal, A.M. (1998): Antiphytoviral potentialities of some fungi and *Actinomycetes* isolates against kidney bean plants infected with T.N.V. 1st Conf. Protec. Egypt., pp. 156-161.