



REFERENCES

- * Aarden, L.A., Brunner, T.K., Certini, J.C., Dayer, J.M., DeWeek, A., Sabato, G.D., and Greg, L. (1979): Revised nomenclature for antigen-nonspecific T-cell proliferation and helper factors. *J. Immunology*, 123:2928.
- * Adachi, K., Yoshikawa, K., Halprin, K.M. and Levine, V. (1975): Prostaglandins and cyclic AMP in epidermis. *Br. J. Dermatol.*, 92:381.
- * Adachi, K., Lizuka, H., Halprin, K.M. and Levine V. (1980): Epidermal cyclic AMP is not decreased in psoriasis lesions. *J. Invest Dermatol.*, 74:74.
- * Adachi, K., Halprin, K.M. and Takeda, J. (1982): Epidermal surface receptors which link pharmacological mediators to the adenylate cyclase system. *Br. J. Dermatol.*, 107:111.
- * Adachi, K. and Aoki, T. (1989): IgE antibody, to sweat in atopic dermatitis. *Acta. Dermatol. Venereol (Stockh)*, 144:83.
- * Adams, D.H., Hubscher, S.G., Wang, L., Elias, E. and Neuberger, J.M. (1989): Soluble interleukin-2 receptors in serum and bile of liver transplant recipients. *Lancet*; 1 (8636):169.
- * Aggarwal, B., Hezel, W., Moffat, B., Kohr, W., and Harkins, R. (1985): Primary structure of human lymphotoxin derived from 1788 lymphoblastoid cell line. *J. Biol. Chem.*, 266:2334.
- * Altmann, A., Coggeshall, K. and Mustelin, T. (1990) : Molecular events mediating T cell activation. *Adv. Immunol.*, 48:227.
- * Amadori, A.P., Zanollo, P., Eozzi, E., Ciminale, V., Borghesan, F., Fagiolo, V. and crepladi, G. (1988): Study of some early immunological parameters in aging humans. *Gerontology*, 34:277.

-
- * Aoyagi, T., Umeda, K. and Kato, N. (1982): Cyclic AMP-dependent protein phosphorylation of pig epidermal homogenates. *J. Dermatol.*, 9:461.
 - * Arnold, W.P., van Hooijdonk, C.A., Mier, P.D. and van de-Kerkhof, P.C. (1990): Induction of ornithine decarboxylase following sellotape stripping in normal and psoriatic skin. *Br. J. Dermatol.*, 123-581.
 - * Asoa, H., Takeshita, T., Nakamura, M., Nagata, K. and Sugamura, K.C. (1990): Interleukin-2 (IL-2) induced tyrosine phosphorylation of IL-2 receptor p75. *J. Exp. Med.*, 172:637.
 - * Baadsgaard, O., Hansen, E. and Chan, L. (1988): Lesional psoriasis T cell cultures reflect the in situ subset distribution and demonstrate autoreactivity. *J. Invest. Dermatol.*, 90:545.
 - * Baadsgaard, O., Gupta, A.K. and Billings, A. (1989): Psoriasis epidermal cells demonstrate increased number and function of non Langerhans cell antigen presenting cells. *J. Invest. Dermatol.*, 92:190.
 - * Bacha, P., Williams, D.P., Waters, C., Williams, J.M., Murphy, J.R. and Strom, T.B. (1988): Interleukin 2 receptor-targeted cytotoxicity. Interleukin 2 receptor-mediated action of a diphtheria toxin-related interleukin 2 fusion protein. *J. Exp. Med.*, 167:612.
 - * Baden, H.P., Kubilus, J. and MacDonald, M.J. (1991): Normal and psoriatic keratinocytes and fibroblasts compared in culture. *J. Invest. Dermatol.*, 76:53.
 - * Baker, B.S., Swain, A.F., Fry, L. and Valdimarsson H. (1984): Epidermal T lymphocytes and HLA-DR expression in psoriasis. *Br. J. Dermatol.*, 110:555.
 - * Balkwill, F.R. (1989): Interferons. *Lancet*, 1:1060.
 - * Bancroft, G., Schreiber, R., Bosma, G., Bosma, M. and Unanue, E. (1987): A T cell-independent mechanism of macrophage activation by interferon-gamma. *J. Immunol.*, 139:1104.
-

- * Bani, D. and Moretti, D. (1987): Are Langerhans cells usual components of the dermal infiltrate of mycosis fungoides ? Arch. Dermatol., 279:561.
- * Barker, J.N.W.N., Sarma, V., Mitra, R.S., Dixit, V.M. and Nickoloff, B.J. (1990): Marked synergism between tumor necrosis factor-alpha and interferon-gamma in regulation of keratinocyte derived adhesion molecules and chemotactic factors. J. Clin. Invest., 85:605.
- * Barna, M., Snigdwint, F.G., Van-der-Hegden, F.L., Bos, J.D. and Kapsenberg, M.L. (1994): Characterization of lesional psoriatic skin T-lymphocyte clones. Acta-Dermatol Venereol (stockh.), 186:9.
- * Barr, R.M., Brain, S. and Camp, R.D.R. (1984): Human allergic and irritant contact dermatitis: levels of arachidonic acid and its metabolites in involved skin. Br. J. Dermatol., 111:23.
- * Benoist, C. and Mathis, D. (1989): Positive selection of the T cell repertoire : Where and when does it occur ? Cell, 58:1027.
- * Berridge, M.J. (1984): Inositol triphosphate and diacylglycerol as second messengers. Biochem. J., 220:345.
- * Berridge, M.J. and Irvine, R.F. (1984): Inositol triphosphate, a novel second messenger in cellular signal transduction. Nature, 312:315.
- * Beutler, B., Greenwald, D., Hulmes, J.D., Chang, M., Pan, Y.C. and Cerami, A. (1985): Identity of tumour necrosis factor and the macrophage-secreted factor cachectin. Nature, 316:552.
- * Beutler, B. and Cerami, A. (1987): Cachectin: more than a tumor necrosis factor. New. Engl. J. Med., 316:379.
- * Beverly, B., Kang, S., Lenardo, M., and Schwartz, R. (1992): Reversal of in vitro T cell clonal anergy by IL-2 stimulation. Int. Immunol., 4: 661.

- * Bhardway, N., Santhanam, V., Lau, L.L., Tatter, S.B., Ghrayeb, J., Rivelis, M., Steinman, R.M., Sehgal, P.B. and May, L.T. (1989): IL-6/IFN β 2 in synovial effusions of patients with rheumatoid arthritis and other arthritis. *J. Immunol.*, 143:2153.
- * Bjerke, J.R., Krogh, T.K. and Matre, J.D. (1978): Characterization of mononuclear cell infiltrates in psoriatic lesions, *J. Invest. Dermatol.*, 71:340.
- * Bjerke, J.R., Livden, J.K., Degre, M. and Matre, R. (1983): Interferon in suction blister fluid from psoriatic lesions. *Br. J. Dermatol.*, 108:295.
- * Bjerke, I.R. and Matre, R. (1983): Demonstration of Ia-like antigen on T lymphocytes in lesions of psoriasis lichen planus and discoid lupus erythematosis. *Acta Dermatol. Venereol (Stockh)*, 63:103.
- * Bluestone, J., and Matis, L. (1989): TCR gamma/delta cells minor redundant T cell subset or specialized immune system component. *J. Immunol.*, 142:1785.
- * Bocci, V. (1988): Roles of interferon produced in physiological conditions. A speculative review. *Immunology*, 64:1.
- * Bochner, B.S.; Landy, S.D.; Plaut, C.A.; Dinarello, C.A. and Schleimer, R.P. (1987): Interleukin γ production by human lung tissue. I. Identification and characterization. *J. Immunol.*, 139:2297.
- * Borowitz, M.J., Croker, B.P. and Burchette, J. (1982): Immunocytochemistry detection of lymphocyte surface antigens in fixed tissue sections. *J. Histochem. Cytochem.*, 30:171.
- * Bos, J.D. (1988): The pathomechanism of psoriasis, the skin immune system and cyclosporin. *Br. J. of Dermatol.*, 118:141.

- * Bos, J.D., Hulsebosch, H.J., Krieg, S.R., Bakker, P.M. and Cormane, R.H. (1983). Immunocompetent cells in psoriasis; In situ immunophenotyping by monoclonal antibodies. *Arch. Dermatol.*, 275:181.
- * Bos, J.D. and Krieg, S.R. (1985): Psoriasis infiltrating cell immunophenotype : changes induced by PUVA or corticosteroid treatment in T cell subsets . Langerhans cells and interdigitating cell. *Acta Dermatol Venereol (Stockh)*, 65:39.
- * Bos, J.D., Van Garderen, I.D., Krieg, S.R. and Poulter, I.W. (1986): Different in situ distribution patterns of dendritic cells having langerhans (T6+) and interdigitating cell immunophenotype in psoriasis, atopic dermatitis and other inflammatory dermatosis. *J. Invest., Dermatol.*, 87:358.
- * Bouclier, M.; El Baze, P., Milano, G., Shroot, B. and Ortonne, J.P. (1986): Hyperinduction of skin ornithine decarboxylase activity in psoriatic patients. *Br. J. Dermatol.*, 115:93.
- * Braun-Falco, O. and Christophers, E. (1974): Structural aspects of initial psoriatic lesions. *Arch. Dermatol.*, 251:95.
- * Braun-Falco, O., Plewig, G., Wolff, H.H. and Winkelmann, R.K. (1991) : Erythematous and erythemasquamous skin diseases. In dermatology. Ch. 4, P. 417. Springer Verlag. Berlin, Heidelberg, New York , London, Paris, Tokyo.
- * Braverman, I.M. and Sibley, J. (1982): Role of the microcirculation in the treatment and pathogenesis of Psoriasis. *J. Invest. Dermatol.*, 78:12.
- * Brenner, M., Strominger, J., and Krangle, M. (1988): The gamma delta T cell receptor. *Adv. Immunol.*, 43: 133.
- * Briggaman, R.A. and Wheeler, C.F. (1980): Nude mouse-human skin graft model III. studies on generalized psoriasis. *J. Invest. Dermatol.*, 74:262.

- * Brunish, R. and Sorensen, B. (1965): Urinary excretion of acid mucopolysaccharides and hydroxyproline in psoriasis. *Dermatologica*, 130:165.
- * Buchsbaum, M.E., Lewis, R.E., Kupper, T.S. and Murphy, G.F. (1989): Differential induction of intercellular adhesion molecule-1 in human skin by recombinant cytokines. *J. Invest. Dermatol.*, 92:408.
- * Bunjes, D., Hardt, C., Rllinghoff, M. and Wagner, H. (1981): Cyclosporin A mediates immunosuppression of primary cytotoxic T cell responses by impairing the release of interleukin1 and interleukin 2. *Eur. J. Immunol.*, 11:657.
- * Burns, G.F.; Triglia, T. and Werkmeister, J.A. (1984): In vitro generation of human activated lymphocyte killer cells: seperate precursors and modes of generation of NK-like cells and "anomalous" killer cells. *J. Immunol.*, 133:1656.
- * Camp, R.D.R. (1992): Psoriasis in : *Rook/Wilkinson/Ebling Textbook of Dermatology*, Ed. by Champion, R.H., Burton, J.L. and Ebling, F.J.G. Ch. 31, p. 1319. Oxford Blackwell Scientific Publications, London; Edinburgh, Boston Melbourne, Paris, Berlin, Vienna.
- * Campbell , M.J. and Machin D. (1993): *Medical statistics. Acommon sense approach*. John Wiley & Sons. New-York, Chichester, Brisbane, Toronto and Singapore.
- * Carswell, E.A., Old, L.J., Kasscl, R.L., Green, S. Fiore, N. and Williamson, B. (1975): An endotoxin-induced serum factor that causes necrosis of tumours. *Proc. Natl. Acad. Sci. USA.*, 72:3666.
- * Castells- Rodellas, A., Castell, J.V., Ramirez-Bosca, A., Nicolas, J.F., Valcuende-Cavero, F. and Thivolet, J. (1992): Interleukin-6 in normal skin and psoriasis. *Acta-Dermatol. Venereol. (Stockh.)*, 72:165.

- * Chafouleas, J.G., Lagace, L., Bolton, W.E., Boyd, A.E.III. and Means, A.R. (1984): Changes in calmodulin and its mRNA accompany reentry of quiescent (G₀) cells into the cell cycle. *Cell*, 36:73.
 - * Chin, Y.H., Falanga, V., and Cai, J.P. (1990): Lymphocyte adhesion to psoriatic dermal endothelium: mechanism and modulation. *J. Invest. Dermatol.*, 95:295.
 - * Chong, A.S.F., Scuderi, P., Grimes, W.J. and Hersh, E.M. (1989): Tumor targets stimulate IL-2 activated killer cells to produce interferon-gamma and tumor necrosis factor. *J. Immunol.*, 142:2133.
 - * Christophers, E. (1996): The immunopathology of psoriasis, *Int. Arch. Allergy- Immunol.*, 116:199.
 - * Clever, H., Alarcon, B., Wilemann, T., and Terhost, C. (1988): The T cell receptor /CD3 complex: A dynamic protein ensemble, *Annu. Rev. Immunol.*, 6:629.
 - * Collins, L., Tsien, W.H., Seals, C., Hakini, J., Weber, D., Bailon, P., Toome, V. and Ju, G. (1988): Identification of specific residues of human interleukin 2-that affect binding to the 70kDa subunit (p70) of the interleukin 2 receptor. *Proc. Natl. Acad. Sci. USA.*, 85:7709.
 - * Colvin, R.B., Fuller, T.C., Mckeen, L., Kung, P.C., Ip, S.H. and Cosin, A.B. (1987): Plasma interleukin-2 receptor levels in renal allograft recipient. *Clin. Immunol.*, 43:273.
 - * Combe, B. Pope, R.M., Fischbach, M., Darnell, B., Baron, S. and Talal, N. (1985): Interleukin 2 in rheumatoid arthritis : production of and response to interleukin 2 in rheumatoid synovial fluid, synovial tissue and peripheral blood. *Clin. Exp. Immunol.*, 59:520.
 - * Cooper, K.D. (1990): Psoriasis: leukocytes and cytokines. *Dermatol. Clinics*, 8 (4):737.
-

- * Cooper, K.D., Hammerberg, C., Baadsgaard, O., Elder, J.T., Chan, L.S. Sauder, D.N., Voorhees, J.J. and Fisher, G. (1990): IL-1 activity is reduced in psoriatic skin. Decreased IL-1 α and increased nonfunctional IL-1 β . *J. Immunol.*, 144:4603.
- * Cooper, K.D. and Baadsgaard, O. (1991): Immunological features of psoriasis In. *Immunologic diseases of the skin*, ed. by Jordon R.E. Ch. 51 P. 611. Appleton & lange. Norwalk, Connecticut. San Mateo, California.
- * Cox, A.J., Watson, W. (1972): Histologic variations in lesions of psoriasis. *Arch. Dermatol.*, 106:305.
- * Creaven, P.J. and Stoll, H.L. (1991): Response to tumor necrosis factor in two cases of psoriasis. *J. Am. Acad. Dermatol.*, 24:735.
- * Cross, S.L., Feinberg, M.B.; Wolf, J.B.; Holbrook, N.J.; Wong- Staal, F. and Leonard, W.J. (1987): Regulation of the human interleukin-2 receptor alpha chain promoter: activation of a nonfunctional promoter by the transactivator gene of HTLV-1. *Cell*, 49:47.
- * Czarnetzki, B.M. (1983): Increased monocyte chemotaxis towards leukotriene B₄ and platelet activating factor in patients with inflammatory dermatoses. *Clin. Exp. Immunol.*, 54:486.
- * Daniel, W.W. (1987): *Biostatistics, A foundation for analysis in The health sciences*. Johnwiley & sons. New-York; Chichester; Brisbane; Toronto & Singapore.
- * Das, N.S., Chowdary, T.N., Sobhanadri, C. and Rao, K.V. (1978): The effect of topical isoprenaline on psoriatic skin. *Br. J. Dermatol.*, 99:197.
- * Defrance, T., Vandervliet, B., Aubry, J.P. and Banchereau, J. (1988): Interleukin 4 inhibits the proliferation but not the differentiation of activated human B cells in response to interleukin 2. *J. Exp. Med.*, 168:1321.

- * Depper, J.M., Leonard, W.J., Kronke, M., Noguchi, P.D., Cunningham, R.E., Waldmann, T.A. and Greene, W.C. (1984): Regulation of interleukin-2 receptor expression : effects of phorbol diester, phospholipase-C, and reexposure to lectin and antigen. *J. Immunol.*, 133:3054.
- * Derynck, R. (1986): Transforming growth factor α : structure and biological activities. *J. Cell. Biochem.*, 32:2293.
- * Detmar, M., Brown, L.F., Claffey, K.P., Yeo, K.T., Kocher, O., Jackman, R.W., Berse B and Dvovak, H.F. (1994): Over expression of vascular permeability factor/vascular endothelial growth factor and its receptors in psoriasis. *J. Exp. Med.*, 180:1141.
- * De-Waal Malefyt, R., Abrams, J., Bennett, B., Figdor, C., and d. Vries, J. (1991): IL-10 inhibits cytokine synthesis by human monocytes. An autoregulatory role of IL-10 produced by monocytes, *J. Exp. Med.*, 174:1209.
- * DiGiovanna, J.J., Aoyagi, T., Taylor, J.R. and Halprin, K.M. (1981): Inhibition of epidermal adenylyl cyclase by lithium carbonate. *J. Invest. Dermatol.*, 76:259.
- * Djeu, J.Y., Kasahara, T., Balow, J.E. and Tsokos, G.(1986): Decreased interleukin-2 inhibitor in sera of patients with autoimmune disorders. *Clin. Exp. Immunol.*, 65:275.
- * Donahue, R., Yang, Y., and Clark, S. (1990): Human p40 T-cell growth factor (interleukin-9) supports erythroid colony formation. *Blood*, 75:2271.
- * Dowd, P.M., Camp, R.D.R., and Greaves, M.W. (1987): In vivo effects of human recombinant interleukin-1 alpha in normal human skin. *J. Invest. Dermatol.*, 119:955.

- * Ducan, J.I., Horroch, C., Drmerod, A.D., Powles, A.V., Whiting, P.H., Fry, L. and Thomson, A.W. (1991): Soluble IL-2 receptor and CD25 cells in psoriasis effects of cyclosporin A and PUVA therapy. *Clinical Exp. Immunol.*, 85 (2):293.
- * Duell, E.A., Fortune, J. and Petersen C. (1986): Eicosanoids quantitated simultaneously from keratomed epidermal strip of psoriatic skin. *J. Invest. Dermatol.*, 87:137.
- * Dukovich, M., Wano, Y., Bich Thuy, L.T., Katz, P., Cullen, B.R., Kehrl, J.H. and Greene, W.C. (1987): A second human interleukin-2 binding protein that may be a component of high-affinity interleukin-2 receptors. *Nature*, 327:518.
- * Eicheberger, M., Allan, W., Carding, S., Bottomly, K., and Doherty, P. (1991): Activation status of the CD4 gamma/ delta- T cells recovered from mice with influenza pneumonia. *J. Immunol.*, 147:2069.
- * Elder, J.T.; Tavakkol, A., Klein, S.B., Zeigher, M.E., Eicha, M. and Voorhees, J.J. (1990): Protooncogene expression in normal and psoriatic skin. *J. Invest. Dermatol.*, 94:19.
- * Elder, T.T., Sartor, C.T., Boman, D.K. and Fisher, G.J. (1992): Interleukin-6 in psoriasis. Expression and mitogenicity studies. *Arch. Dermatol Res.*, 284 (6). 324.
- * Enk, A., and Katz, S. (1992): Identification and induction of keratinocyte-derived IL-10. *J. Immunol.*, 149:92.
- * Essmann, J., Cooper, K.D., Voorhees, J.J. and Fisher, G.J. (1988): Membrane associated transglutaminase activity is increased in psoriatic involved in normal epidermis. *J. Invest. Dermatol.*, 90:556.
- * Esmann, J., Voorhees, J.J. and Fisher G.J. (1989): Increased membrane-associated transglutaminase activity in psoriasis. *Biochem. Biophys. Res. Commun.*, 164 (1):219.

- * Espevik, T., Figan, I.S., Shalaby, M. R., Lackides, G.A., Lowis, G.D. and Palladino M.A.Jr. (1987): Inhibition of cytokine production by cyclosporin A and transforming growth factor beta. *J. Exp. Med.*, 166:571.
- * Espevik, T., and Nissen Meyer, J. (1987): Tumour necrosis factor like activity on paraformaldehyde fixed monocyte monolayers. *Immunology*, 61:443.
- * Ezepechuk, Y.V., Leung, D.Y.M., Middleton, M.H., Bina, P., Reiser R and Norris, D.A. (1996): Staphylococcal toxins and protein A differentially induce cytotoxicity and release of tumour necrosis factor- α from human keratinocytes. *J. Invest. Dermatol.*, 107:603.
- * Ettehadi, P., Greaves, M.W., Camp R.D.R. and Wallach, D. (1994): TNF- α Immunoreactivity and Bioactivity in psoriasis. *J. Dermatol. Venercol. (Stockh)*, 186:14.
- * Farrar, W.L., Johnson, H.M. and Farrar, J.J. (1981): Regulation of the production of immune interferon and cytotoxic T lymphocytes by interleukin-2. *J. Immunol.*, 126:1120.
- * Fernandez-Botran, R., Sander, V.M. and Vitetta, E.S. (1989): Interactions between receptors for interleukin 2 and interleukin 4 on lines of helper T cells (HT-2) and B lymphoma cells (BCL1). *J. Exp. Med.*, 169:379.
- * Ferro, T.J., Johnson, A., Everitt, J. and Malik, A.B. (1989): IL-2 induces pulmonary edema and vasoconstriction independent of circulating lymphocytes. *J. Immunol.*, 142:1916.
- * Fidel, P.L. (1994): T-cell Mediated Lymphokines In *Immunochemistry* ed. Van Oss, C.J. and Van Regenmortel M.H.V., Ch. 37 P. 937. Marcel Dekker, Inc. NewYork.
- * Fiorentino, D., Bond, D. and Mosmann, T. (1989): Two types of mouse helper T cells. *J. Exp. Med.*, 176:2081.

- * Fiorentino, D., Zlotnik, A., Vieira, P., Mossmann, T. Moore, K. and O'Garra, A. (1991): IL-10 acts on the antigen-presenting cell to inhibit cytokine production by Th/cells. *J. Immunol.*, 146:3444.
- * Fisher, G.J.; Harris, V.A. and Voorhees, J.J. (1987): Purification and characterization of calcium phospholipid-dependant kinase from adult human epidermis. *J. Invest. Dermatol.*, 89:484.
- * Fisher, G.J.; Talwar, H.S.; Baldassare, J.J., Henderson, P.A. and Voorhees, J.J. (1990,a): Increased phospholipase C-Catalyzed hydrolysis of phosphatidylinositol 4,5 biphosphate and 1,2- Diacylglycerol content in psoriatic involved compared to uninvolved & normal epidermis. *J. Invest. Dermatol.*, 95 (4):428.
- * Fisher, G.J., Talwar, H.S., Tavakkol, A., Esmann, J., Baldassare, J.J., Edler, J.T., Griffiths, C.E.M., Baadsgaard, O., Cooperik, D. and Voorhees, J.J. (1990,b): Phosphorinositide-mediated signal transduction in normal and psoriatic epidermis. *J. Invest. Dermatol.*, 95(5); 15S.
- * Fleischer, A.B., Feldman, S.R., Rapp, S.R., Reboussin, D.M., Exum M.L., Clark, A.R. and Rajashkar, V. (1996): Disease severity measures in a population of psoriasis patients: the symptoms of psoriasis correlate with self-administered psoriasis area severity index score. *J. Invest. Dermatol.*, 107:26.
- * Forster, S., Ilderton, E., Summerly, R. and Yardley, H.J. (1983): The level of phospholipase A₂ activity is raised in the uninvolved epidermis of psoriasis. *Br. J. Dermatol.*, 108:103.
- * Fraker, D.L., Langstein, H.N. and Norton, J.A. (1989): Passive immunization against tumor necrosis factor partially abrogates interleukin 2 toxicity. *J. Exp. Med.*, 170:1015.
- * Fraki, J.E. (1983): Correlation of epidermal plasminogen activity with disease activity in psoriasis. *Br. J. Dermatol.*, 108:39.
- * Fraki, J.E. (1985): Proteinases and psoriasis. *Cutis*, 36:32.

- * Fraki, J.E., Briggaman, R.A. and Lazarus, G.S. (1982): Uninvolved skin from psoriatic patients develops signs of involved psoriatic skin after being grafted onto nude mice. *Science*, 215:685.
- * Fredriksson, T. and Pettersson, U. (1978): Severe psoriasis-oral therapy with a new retinoid. *Dermatologica*, 157(4):238.
- * Fung, M.R., Searce, R.M., Hoffman, J.A., Pepper, N.J., Hammes, S.R., Hosking, J.B., Schmandt, R., Kuziel, W.A., Haynes, B.F., Mills, G.B. and Greene, W.C. (1991): A tyrosine kinase physically associates with the β -subunit of the human IL-2 receptor. *J. Immunol.*, 147:1253.
- * Gajewski, T., Schell, S., Nau, G., and Fitch, F. (1989): Regulation of T-cell activation: Differences among T cell subsets. *Immunol. Rev.*, 111:79.
- * Galizzi, J., Zuber, C., Harada, N., Gorman, D., and Miyajima, A. (1990): Molecular cloning of a DNA encoding the human interleukin 4 receptor. *Int. Immunol.*, 2:669.
- * Gaspari, A., Jenkins, M. and Kutz, S.I. (1988): Induction of antigen specific T cell unresponsiveness by lat. Keratinocytes. *J. Invest. Dermatol.*, 90:562.
- * Gazzinelli, R., Oswald, I., James, S., and Sher, A. (1992): IL-10 inhibits parasite killing and nitric oxide production by IFN-gamma-activated macrophages. *J. Immunol.*, 148:1792.
- * Geerdink, J.F.M., de Mulder, P.H.M., Franzen, M.P., Gommans, J.M., van Rennes, H., van Erp, P.E.J., Bergers, M. and Mier, P.D. (1984): Monocyte function is normal in quiescent psoriasis. *J. Invest. Dermatol.*, 82:122.
- * Geiger, T., Aordus, T., Klapproth, J., Hirano, T., Kishimoto, T. and Herinrich, P.C. (1988): Induction rat aortic phase proteins by interleukin 6 in vivo. *Eur. J. Immunol.*, 18:717.

- * Gelfant, S. (1976): The cell cycle in psoriasis. A reappraisal. *Br. J. Dermatol.*, 95:577.
 - * Gibson, L.E. and Perry, H.O. (1992): Papulosquamous eruptions and exfoliative dermatitis. In: *Dermatology*. Ed. by Moschella, S.L. and Hurley, H.J. Ch 26. P. 607. W.B. Saunders Company, Philadelphia. London. Toronto.
 - * Gillis, S., Crabtree, C.R., and Smith, K.A. (1979): Glucocorticoids-induced inhibition of T-cell growth factor production. The effect on the in vitro generation of cytolytic T-cells. *J. Immunol.*, 123:1632.
 - * Gillitzer, R., Berger, R., MNieler, V., Muller, C., Wolff, K. and Sting, G. (1991): Upper keratinocytes of psoriatic skin lesions express high levels of NAP-1/IL-8 mRNA in situ. *J. Invest. Dermatol.*, 97:73.
 - * Gnarr, J.R., Otani, H., Wang, M.G., McBride, O.W., Sharon, M. and Leonard, W.J. (1990): Human interleukin-2 receptor beta-chain gene: chromosomal localization and identification of 5 regulatory sequences. *Proc. Natl. Acad. Sci. USA.*, 87:3440.
 - * Gomi, T., Shiohara, T., Munakata, T., Imanishi, K. and Nagashima, M. (1991): Interleukin 1 α , tumor necrosis factor α , and interferon γ in psoriasis. *Arch. Dermatol.*, 127:827.
 - * Gommans, J.M., Bergers, M.m Van Erp, P.E.J. (1979): Studies on the plasma membrane of normal and psoriatic keratinocytes. *Br. J. Dermatol.*, 101:413.
 - * Gommans, J.M., Vanden Hurk, J.J.M.A and Bergers, M. (1982): Studies on the plasma membrane of normal and psoriatic keratinocytes. 5. Lectin binding. *Br. J. Dermatol.*, 106:317.
 - * Goodwin, P., Hamilton, S. and Fry, L. (1973): A comparison between DNA synthesis and mitosis in uninvolved and involved psoriatic epidermis and normal epidermis. *Br. J. Dermatol.*, 106:317.
-

- * Goodwin, J.S., Bankhurst, A.D. and Meisner, R.P. (1977): Suppression of human T-cell mitogenesis by prostaglandin. *J. Exp. Med.*, 146:1719.
- * Gottlieb, A.B. (1988): Immunologic mechanisms in psoriasis. *J. Am. Acad. Dermatol.*, 18:1376.
- * Gottlieb, A.B., Lifschitz, B. and Fu, S.M. (1986): Expression of HLA-DR molecules by keratinocytes and HLA-DR expression in psoriasis. *Br. J. Dermatol.*, 110:555.
- * Gottlieb, A.B., Luster, A.D., Posnett, D.N. and Carter, D.M. (1988): Detection of a (gamma) interferon-induced protein IP-10 in psoriatic plaques. *J. Exp. Med.*, 168:941.
- * Gottlieb, S.L., Gilleaudeau, P., Johnson, R., Estes, L., Woodworth, T.G., Gottlieb, A.B. and Krueger, J.G. (1995): Response of psoriasis to a lymphocyte-selective toxin (DAB389 IL-2) suggests a primary immune, but not a keratinocyte, pathogenesis. *Nature. Med.*, 1:442.
- * Gray, P. and Goeddel, D. (1983): Cloning and expression of murine immune interferon DNA. *Proc. Natl. Acad. Sci. USA.*, 80:5842.
- * Greene, W., Robb, R., Depper, J., and Leonard, W. (1985): Stable expression of DNA encoding the human interleukin-2 receptor in eukaryotic cells. *J. Exp. Med.*, 162:363.
- * Griffiths, C.E.M., Voorhees, J.J. and Nicholoff, B.J. (1989): Characterization of intercellular adhesion molecule-1 and HLA-DR expression in normal and inflamed skin: Modulation by recombinant gamma interferon and tumor necrosis factor. *J. Am. Acad. Dermatol.*, 20:617.
- * Grimm, E., Mazumder, A., Zhang, H. and Rosenberg, S. (1982): Lymphokine- activated killer cell phenomenon. *J. Exp. Med.*, 155:1823.

- * Hatakeyama, M., Mori, H., Doi, T., and Taniguchi, T. (1989): A restricted cytoplasmic region of IL-2 receptor beta chain is essential for growth signal transduction but not for ligand binding and internalization. *Cell*, 59:837.
- * Haworth, C. and Feldmann, M. (1992): Applications of cytokines in human immunotherapy in : *The cytokine Handbook* Ed. by Thomson, A.W. Ch 15 p. 301. Academic Press. Harcourt Brace Jovanovich. London, San Diego, New York, Boston, Sydney, Tokyo, Toronto.
- * Heng, M.C.Y., Moy, R.L. and Lieberman, J. (1985): Alpha-1-antitrypsin deficiency in severe psoriasis. *Br. J. Dermatol.*, 112:129.
- * Henny, C., Kern, D., and Gillis, S. (1981): Interleukin-2 augments natural killer activity. *Nature*, 291:335.
- * Hershey, G. and Scheiber, R. (1989): Biosynthetic analysis of the human interferon-gamma-receptor. Identification of N-linked glycosylation intermediates. *J. Biol. Chem.*, 264:11988.
- * Higgs, J.B., Zeldes, W. and Kozarsky, K. (1988): A novel pathway of human T. Lymphocyte activation. Identification by a monoclonal antibody generated against a rheumatoid synovial T cell line. *J. Immunol.*, 140:3758.
- * Hirano, T., Yasukawa, K. and Harada, H. (1986): Complementary DNA for a novel human interleukin (BSF-2) that induces B lymphocytes to produce immunoglobulin. *Nature*, 324:73.
- * Ho, A.S.Y.; Lui, Y., Khan, T.A., Bazan, J.F. and Moore, K.W. (1993): A receptor for IL-10 is related to interferon receptors. *Proc. Natl. Acad. Sci. USA.*, 90:11267.
- * Hohmann, H., Remy, R., Brockhaus, M., and Vanloon, A. (1989): Two different cell types have different major receptors for human tumor necrosis factor. *J. Biol. Chem.*, 264:12927.

- * Horrocks, C., Ormerod, A.D., Duncan, J.I. and Thomson, A.W. (1989): Influence of systemic cyclosporin A on interleukin-2 and epidermal growth factor receptor expression in psoriatic skin lesions. *Clin. Exp. Immunol.*, 78:166.
- * Horrocks, C., Duncan, J.I., Sewell, H.F., Ormerod, D. and Thomson, A.W. (1990): Differential effects of cyclosporine A on Langerhans cells and regulatory T-cell populations in severe psoriasis: An immunohistochemical and flow cytometric analysis. *J. Autoimmunity*, 3:559.
- * Horvath, H. and Tobora E. (1972): Alterations of collagen in psoriatic skin. *Dermatologica*, 144:832.
- * Huang, Y.P., Pechere, J.C., Michel, M., Gauthey, L., Loreto, M., Gurran, J.A. and Michel, J.P. (1992): In vivo T cell activation, in vitro defective IL-2 secretion, and response to influenza vaccination in elderly women. *J. Immunol.*, 148:715.
- * Hunt, S.J. and Jegasothy, B.V. (1992): The function of cells in inflammatory disease. In: *Dermatology*. Ed. by Moschella, S.L. and Hurely, H.T. Ch 10 P. 250. W.B. Saunders Company. Harcourt. Brace Jovanovich, Inc., Philadelphia, London, Toronto, Montreal, Sydney, Tokyo.
- * Itoh, N., Yonehara, S., Schreurs, J., Gorman, D.M., Maruyama, K., Isii, A., Yahara, I., Avai, Kr and Miyajima, A. (1990): Cloning of an interleukin-3 receptor gene: a member of distinct receptor gene family. *Science*, 247:327.
- * Inn, H., Sato, S., Fugimoto, M., Kikuch, K. and Takchara, K. (1996): Clinical significance of serum level of soluble interleukin-2 receptor in patients with localized scleroderma *Br. J. Dermatol.*, 134:843.
- * Jelinek, D.F. and Lipsky, P.E. (1987): Regulation of human B lymphocyte activation, proliferation and differentiation *Adv. Immunol.*, 40:1.

- * Jensen, P.J., Baird, J., Belin, D., Vassalli, J.D., Busso, N., Gubler, P. and Lazarus, G.S. (1990): Tissue plasminogen activator in psoriasis. *J. Invest. Dermatol.*, 95:135.
 - * Johanneson, A., Hammur, H. and Sunqvist, K.G. (1982): The deposition of immunoglobulins and complement in stratum corneum in microscopic lesions in patients with active psoriasis the relationship to hyperproliferation. *Acta. Dermatol. Venereol. (Stockh)*, 62:21.
 - * Johnson, H.M. and Torres, B.A. (1985): Leukotrienes positive signals for regulation of γ -interferon production *J. Immunol.*, 132:413.
 - * Jung, V., Jones, C., Kumar, C., Stefanos, S., O'Connell, S. and Pestka, S. (1990): Expression and reconstitution of a biologically active human IFN-gamma receptor in hamster cells. *J. Biol. Chem.*, 265:1827.
 - * Kaplan, G., Kiessling, R., Teklemariam, S., Hancock, G., Sheftel, G., Job, C.K., Converse, P., Ottenhoff, T.H.M., Becx-Bleumink, M., Dietz, M. and Cohn, Z.A. (1989a): The reconstitution of cell-mediated immunity in the cutaneous lesions of lepromatous leprosy by recombinant interleukin-2. *J. Exp. Med.*, 169:893.
 - * Kaplan, G., Sampaio, E.P., Walsh, G.P., Burkhardt, R.A., Fajardo, T.T., Guido, L.S., Machado, A., Cellona, R.V., Abalos, R.M., Sarno, E.N. and Cohn, Z.A. (1989b): Influence of mycobacterium leprae and its soluble products on the cutaneous responsiveness of leprosy patients to antigen and recombinant interleukin 2. *Proc. Natl. Acad. Sci. U.S.A.*, 86:6269.
 - * Kapp, A. (1993): The role of cytokines for the pathogenesis of psoriasis. *Hautarzt*, 44(4):201.
 - * Kapp, A., Piskorski, A. and Schopf, E. (1988): Elevated levels of interleukin 2 receptor in sera of patients with atopic dermatitis and psoriasis *Br. J. Dermatol.*, 119:707.
-

- * Kapp, A., Textor, A., Krutmann, J. and Mller, A. (1990): Immunomodulating cytokines in atopic dermatitis and psoriasis: production of tumour necrosis factor and lymphotoxin by mononuclear cells in vitro. *Br. J. Dermatol.*, 122:587.
- * Kapp, A., Neuner, P., Krutmann, J., Luger, T.A. and Schopf, E. (1991): Production of interleukin-2 by mononuclear cells in vitro in patients with atopic dermatitis and psoriasis. *Acta. Dermatol. Venereol. (Stockh)*, 71:403.
- * Kappler, J.W., Roehm, N. and Marrack, P. (1987): T cell tolerance by clonal elimination in the thymus. *Cell*, 49:273.
- * Kaudewitz, P., Braun-Falco, O., Kind, P., Galosi, A., Rieber, P. and Riethmuller, G. (1984): Distribution of T-cell subsets as defined by monoclonal antibodies in skin lesions of psoriasis vulgaris. *Arch. Dermatol. Res.*, 276:33.
- * Kawakami, M. and Cerami, A. (1981): Studies of endotoxin induced decrease in lipoprotein lipase activity. *J. Exp. Med.*, 154:631.
- * Kehrl, J.H., Miller, A. and Fauci, A.S. (1987): Effect of tumor necrosis factor α on mitogen activated human B cells. *J. Exp. Med.*, 166:786.
- * Kehrl, J.H., Dukovich, M., Whalen, G. Katz, P., Fauci, A.S., and Greene, W.C. (1988): Novel interleukin 2 (IL-2) receptor appears to mediate IL-2 induced activation of natural killer cells. *J. Clin. Invest.*, 81:200.
- * Kemmett, D., Symons, A., Colver, B. and Duff, G.W. (1990): Serum soluble interleukin-2 receptor in psoriasis. *Acta. Dermatol. Venereol. (Stockh)*, 70:264.
- * Kimura, S. and Nishikawa, T. (1978): An immuno-histochemical analysis of the deposited immunoglobulins or fibrinogen in parakeratotic psoriatic horny layer and pemphigus skin lesions. *Arch. Dermatol.*, 261:55.

- * King, L.E., Gates, G.R., Stoscheck, C.M. and Nanney, L.B. (1990): Epidermal growth factor/Transforming growth factor alpha receptors and psoriasis. *J. Invest. Dermatol.*, 95:105.
- * Kisielow, P., Teh, H., Bluthmann, H., and von Boehmer, H. (1988): Positive selection of antigen-specific T cells in thymus by restricting MHC molecules. *Nature*, 335:730.
- * Kobayashi, H., Yasuda, H., Ohkawa, A., Dosaka, H. Oda, A., Ogiso, Y. and Kuzumaki, N. (1988): Enhanced expression of ras gene products in psoriatic epidermis. *Arch. Dermatol. Res.*, 280:259.
- * Kock, A., Urbanski, A. and Luger, T.A. (1989): mRNA expression and release of tumor necrosis factor alpha by human epidermal cells. *J. Invest. Dermatol.*, 92:462.
- * Kock, F.; Heufler, C., Kampgen, E., Schwneeweiss, D. Bock, G. and Schuler, G. (1990): Tumor necrosis factor alpha maintains the viability of murine epidermal langerhans cells in culture, but in contrast to granulocyte/macrophage colony-stimulating factor without inducing their functional maturation. *J. Exp. Med.*, 171:159.
- * Kohase, M., Henroksen-Destefano, D., May, L.T., Vilcek, J., and Sehgal, P.B. (1986): Induction of beta, 2. Interferon by tumour necrosis factor: A homeostatic mechanism in the control of cell proliferation. *Cell*, 45:655.
- * Kojima, J., Cromie, M.A., Fisher, G.J., Voorhees, J.J. and Elda, J.T. (1993): GRO α mRNA is selectively overexpressed in psoriatic epidermis and is reduced by cyclosporin A in vivo, but not in cultured keratinocytes. *J. Invest. Dermatol.*, 101:767.
- * Konnikov, N., Ree, H.J., Dinarello, C.A. and Pincus, S.H. (1987): Inteleukin-1 is present in psoriatic plaque. *J. Invest. Dermatol.*, 88:499.
- * Kozin, V.M., Adamenko, G.P., Bogdanovid, L.I. and Iakirevich, E.V. (1989): The functional properties of interleukins in psoriasis. *Acta. Dermatol. Venereol. (Stockh)*, 7:61.

- * Krane, J.F., Murphy, D.P., Gottlieb, A.B., Carter, D.M., Hart, C.E. and Krueger, J.G. (1991): Increased dermal expression of platelet-derived growth factor receptors in growth activated skin wounds and psoriasis. *J. Invest. Dermatol.*, 96:983.
- * Krueger, G.G. and Jederberg, G.W. (1980): Alteration of Hel a cell growth equilibrium by supernatants of peripheral blood mononuclear cells from normal and psoriatic subjects. *J. Invest. Dermatol.*, 74:148.
- * Krueger, J.G., Krane, J.F., Carter, D.M. and Gottlieb, A.B. (1990): Role of growth factors, cytokines, and their receptors in the pathogenesis of psoriasis. *J. Invest. Dermatol.*, 91:1355.
- * Kumar, S. and West, D.C. (1990): Psoriasis, angiogenesis and hyaluronic acid. *Lab. Invest.*, 62:664.
- * Kuo, L.M. and Robb, R.J. (1986): Structure-function relationships for the IL-2 receptor system. *J. Immunol.*, 137:1538.
- * Kupper, T.S., Chuna, A.O., Flood, P., McGuire, J. and Gubler, U. (1987): Interleukin-1 gene expression in culture keratinocytes is augmented by ultraviolet irradiation. *J. Clin. Invest.*, 90:430.
- * Kuziel, W.A., and Greene, W.C. (1992): Interleukin-2 In: *The cytokine handbook* ed. by Thomas, A Ch. 4 P 83. Academic press. Harcourt Brace Jovanovich pub. London. San Diego, New York, Boston, Sydney, Tokyo, Toronto.
- * Larsen, C.G., Anderson, A.O., Appella, E., Oppenheim, J.J. and Matsushima, K. (1989): The neutrophil activating protein (NAP-1) is also chemotactic for T lymphocytes. *Science*, 243:1464.
- * Lauharanta, J., Kousa, M., Kappalyo, K., Linnamaa, K. and Mustakallio, K. (1981): Reduction of increased polyamine levels in psoriatic skin by retinoid and PUVA treatments. *Br. J. Dermatol.*, 105:267.
- * Lazarus, G.S. (1992): New aspects of pathogenesis of psoriasis. *Seminars in Dermatology*, 11:2.

- * Lazarus, G.S., Yost, F.J. and Thomas, C.A. (1977): Polymorphnuclear leukocytes: possible mechanism of accumulation in psoriasis. *Science*, 198:1162.
- * Le, J. and Vilcek, J. (1987): Tumour necrosis factor and interleukin 1: Cytokines with multiple overlapping biological activities. *Lab. Invest.*, 56:234.
- * Lee, R.E., Gaspari, A.A., Lotze, M.T., Chang, A.E. and Rosenberg, S.A. (1988): Interleukin 2 and psoriasis. *Arch. Dermatol.*, 124:1811.
- * Lefrancois, L. and Goodman, T. (1989): In vivo modulation of cytolytic activity and Thy-7 expression in TCR-gamma delta intraepithelial lymphocytes. *Science*, 243:1716.
- * Lefrancois, L., Lecorre, R., Mayo, J., and Blueston, J. (1990): Extrathymic selection of TCR gamma delta cells by class II major histocompatibility complex molecules. *Cell*, 63:333.
- * Lelchuk, R. and playfair, J.H.K. (1985): Serum IL-2 inhibitor in mice 1 increase during infection. *Immunology*, 56:113.
- * Lenardo, M. (1991): Interleukin-2 programs mouse alpha/beta T-lymphocytes for apoptosis. *Nature.*, 353, 858.
- * Leung , D.Y., Walsh, P., Hiorno, R. and Norris, D.A. (1993): A potential role for superantigens in the pathogenesis of psoriasis. *J. Invest. Dermatol.*, 100:225.
- * Lever, W.F. and Schaumburg-Lever, G. (1990): Morphology of the cell in the dermal infiltrate In : *Histopathology of the skin*. Ch. 5. P. 58. J. B. Lippincott Co. Philadelphia, New York, St. Lous, London , Tokyo.
- * Lichtman, A.H., Chin, J., Schmidt, J.A. and Abbas, A.K. (1988): Role of interleukin 1 in the activation of T lymphocytes. *Proc. Natl. Acad. Sci. USA.*, 85:9699.

- * Littman, S., Faltynek, C. and Baglioni, C. (1985): Binding of human recombinant I. interferon gamma to receptors on human cells. *J. Biol. Chem.*, 260:1191.
- * Liu, Y., Wei, S.H.Y., de Waal Malefyt, R. and Moore K.W. (1994): Expression cloning and characterization of a human IL-10 receptor. *J. Immunol.*, 152:1821.
- * Livden, J.K., Bjerke, J.R., Degre, M. and Metre, R. (1986): The effect of Goeckerman therapy on interferon in serum and suction blister fluid from patients with psoriasis. *Br. J. Dermatol.*, 114:217.
- * Lizuka, H., Adachi, K., Halprin, K.M., Levine, V. (1978): Cyclic nucleotide-phosphodiesterase in the uninvolved and involved skin of psoriasis. *J. Invest Dermatol.*, 70:246.
- * Loetscher, H., Pan, Y.C., Lahm, H.W., Gentz, R., Tabuchi, H. and Lesslauer, W. (1990): Molecular cloning and expression of the human 55 kd tumour necrosis factor receptor. *Cell*, 61:351.
- * Lotze, M.T., Matory, Y., Ettinghausen, S.E. Rayner, A.A., Sharrow, S.O., Seipp, C.A., Custer, M.C. and Rosenberg, S.A. (1985): In vivo administration of purified human interleukin-2. *J. Immunol.*, 135:2865.
- * Lowe, N.J., Breeding, J. and Russell, D. (1982): Cutaneous polyamines in psoriasis. *Br. J. Dermatol.*, 107:21.
- * Lowenthal, J.W., Ballard, D.W., Bogerd, H., Bohnlein, E. and Greene, W.C. (1989): Tumor necrosis factor- α activation of IL-2 receptor- α gene involves the induction of κ β -specific DNA binding proteins. *J. Immunol.*, 142:3121.
- * Luger, T.A. (1989): Epidermal cytokines. *Acta. Dermatol. Venereol.*, 69:61.
- * Maca, R.D. (1983): The effects of prostaglandins on the proliferation of cultured human T lymphocytes. *Immunopharmacol.*, 6:267.

- * Mackay, C. (1991): Lymphocyte homing. Skin seeking memory T cell. *Nature.*, 549:737.
- * Majerus, P.W., Conlloy, T.M., Deckmyn, H., Ross, T.S., Bross, T.E., Ishii, H., Bassal, V.S. and Wilson, D.B. (1986): The metabolism of phosphoinositide-derived messenger molecules. *Science*, 234:1519.
- * Majewski, S., Tigalonna, M., Jablonska, S., Polakowski, I. and Janczura, E. (1987): Serum samples from patients with active psoriasis enhance lymphocyte-induced angiogenesis and modulate endothelial proliferation. *Arch. Dermatol.*, 123:221.
- * Male, D., Champion, B., Cook, A. and Owen, M. (1991): Cytokines. In *Advanced Immunology*, Ch. 11, P.11 J.B. Lippincott Co., Philadelphia, Gower Medical Publishing. London, New York.
- * Marcelo, C.L., Duell, E.A., Stawiski, M.A., Anderson, T.F. and Voorhees, J.J. (1979): Cyclic nucleotides in psoriatics and normal epidermis. *J. Invest. Dermatol.*, 72:20.
- * Marlin, S.D. and Springer, T.A. (1987): Purified intercellular adhesion molecule-1 (ICAM1) is a ligand for lymphocyte function associated antigen-1 (LFA-1). *Cell*, 51:813.
- * Marks, J.G., Jr., Zaino, R.J., Bressler, M.F. and Williams, J.V. (1987): Changes in lymphocytes and Langerhans cell populations in allergic and irritant contact dermatitis. *J. Immunol.*, 26:254.
- * Marrack, P. and Kappler, J. (1990): The staphylococcal enterotoxins and their relatives. *Science*, 248:705.
- * Matsushima, K., Yoshimura, T., Lavu, S., Kobayashi, Y., Lew, W., Appella, E., Kung, H.F., Leonard, E.J., and Oppenheim, J.J. (1988): Molecular cloning of a human monocyte-derived neutrophil chemotactic factor (MDNCF) and the induction of MDNCF mRNA by interleukin 1 and tumor necrosis factor. *J. Exp. Med.*, 167:1883.

- * Mckenzie, R.C. and Sauder, D.N. (1990): Keratinocyte cytokines and growth factors. *Dermatologic Clinics*, 8: 649.
- * McIntyre, B., and Allison, J. (1983): The mouse T-cell receptor: structural heterogeneity of molecules of normal T cells defined by xeno antiserum. *Cell*, 34:739.
- * Meager, A. (1991): Lymphocyte activation. In.: Cytokines Ed. by Meager, A. P. 105 Published by Prentic Hall. Englewood Clifs. New Jersey.
- * Metcalf, D. (1989): Haemopoietic growth factors. *Lancet*, I: 825.
- * Meuer, S.C., Hussey, R.E. and Cantrell, D.A. (1984): Triggering of the T3-T1 antigen receptor complex results in clonal T cell proliferation through an interleukin 2 dependent autocrine pathway. *Proc. Natl. Acad. Sci. USA.*, 81:1509.
- * Meuer, S.C.; Dumann, H., Meyer-Zum-Buschenfelde, K.H. and Kohler, H. (1989): Low dose interleukin-2 induces systemic immune responses against HBsAg in immunodeficient non-responders to hepatitis B vaccination. *Lancet*, I:15.
- * Mielke, V., Bauman, J.G.J., Sticherling, M., Ibs, T., Zomerschoe, A.G., Seligmann, K., Henneicke, H.H., Schror, J.M., Sterry, W. and Christophers, E. (1990): Detection of neutrophil-activating peptide NAP/IL-8 and NAP/IL-8 mRNA in human recombinant IL-1 alpha and human recombinant tumor necrosis factor-alpha-stimulated human dermal fibroblasts. *J. Immunol.*, 144:153.
- * Mita, S., Harada, N., Naomi, S., Hitoshi, Y. and Takats, K. (1988): Receptor for T cell-replacing factor/interleukin 5 : Specificity, quantitation, and its implication. *J. Exp. Med.*, 168:863.
- * Miyajima, A., Kitamura, T., Yokota, T., and Aral, K. (1992): Cytokine receptors and signal transduction. *Ann. Rev. Immunol.*, 10:295.

- * Mizumoto, T.; Hashimoto, Y., Hirokawa, M., Ohkuma, N., Lizuka, H. and Ohkawara A. (1985): Calmodulin activities are significantly increased in both uninvolved and involved epidermis in psoriasis. *J. Invest. Dermatol.*, 85:450.
- * Molinas, F., Wietzerbin, J., and Falcoff, E. (1987): Human platelets possess receptors for a lymphokine : Demonstration of high specific receptors for IFN-gamma. *J. Immunol.*, 138:802.
- * Mond, J., Carman, J., Sarma, C., Ohara, J. and Finkelman, F. (1986): Interferon-gamma Suppresses B cell stimulation factor induction of class II MHC. determinants on B cells. *J. Immunol.*, 137:3535.
- * Moore, K., Vieira, A., Fiorentino, D., Khon, T. and Mosmann, T. (1990): Homology of cytokine synthesis inhibitory factor (IL-10) to the Epstein-Barr virus gene. *Science*, 248:1230.
- * Morel, P., Revillard, J.P., Nicolas, J.F., Wijdenes, J., Rizova, H. and Therolet, J. (1992): Anti-CD4 monoclonal antibody therapy in severe psoriasis. *J. Autoimmun.*, 5(4):465.
- * Morgan, D.A., Ruscetti, F.W. and Gallo, R.C. (1976): Selective in vitro growth of T-lymphocytes from normal human bone marrow. *Science*, 193:1007.
- * Morhenn, V.B., Wastek, G.J., Cua, A.B. and Mansbridge, J.N. (1989): Effects of recombinant interleukin 1 and interleukin-2 on human keratinocytes. *J. Invest. Dermatol.*, 93:121.
- * Morimoto, C. Romain T. and Fox D.A.(1988): Abnormalities in CD4 T-lymphocytes subsets in inflammatory rheumatic diseases. *Am. J. Med.*, 84:817.
- * Morioka, S., Lazarus, G.S., Baird, J.L., and Jensen, P.J. (1987): Migrating keratinocytes express urokinase type plasminogen activator. *J. Invest. Dermatol.*, 88:418.

- * Morita, C., Verna, S., Aparicio, P. and Bernner, M. (1991): Functionally distinct subsets of human gamma/delta T cells. *Eur. J. Immunol.*, 21:2999.
- * Mosmann, T., and Coffman, R. (1989): Th 1 and Th2 cells: different patterns of lymphokine secretion lead to different functional properties. *Ann. Rev. Immunol.*, 7:145.
- * Mosley, B., Beckmann, P., March, C., Friend, D. and Pask, L. (1989): The murine interleukin-4 receptor Molecular Cloning and characterization of secreted and membrane bound form. *Cell*, 59:335.
- * Mozzanica, N., Cattaneo, A., Trabattoni, D., Finzi, A.F., Schmitt, E. and Ferrario, E. (1995): Production of type-1 and type 2 cytokine by peripheral blood mononuclear cells of psoriatic patients. *Immunology*, 86(3):422.
- * Mrowietz, V., Falsafi, M., Schrder, J.M. and Christophers, E. (1992): Inhibition of human monocyte functions by anthralin. *Br. J. Dermatol.*, 127:382.
- * Munk, M., Gatrill, A., and Karfmann, S. (1990). Target cell lysis and IL-2 secretion by gamma/delta T lymphocytes after activation with bacteria. *J. Immunol.*, 145:2424.
- * Muraguchi, A., Hirano, T., Tang, B., Matsuda T., Horis, Y. and Kishimoto, T. (1988): The essential role of β cell stimulatory factor 2 (BSF-2) IL-6 for the terminal differentiation of β cell. *J. Exp. Med.*, 167:332.
- * Murphy, G.F. (1995): Psoriasiform Dermatitis. In *Dermato pathology* Ch.4 p. 74. W.B. Saunders company. Philadelphia, London, Toronto, Montreal, sydney, Tokyo.

- * Nagel, J.E., Chopra, P.K., Chrest, F.J., McCay, M.T., Schneider, E.L., Holbrook, N.J. and Alder, Will (1988): Decreased proliferation, interleukin-2 synthesis, and interleukin 2 receptor expression are accompanied by decreased in RNA expression in phytohemagglutinin-stimulated cells from elderly donors. *J. Clin. Invest.*, 31:1096.
- * Nandi, D. and Allison, J. (1991): Phenotypic analysis and gamma delta T cell receptor repertoire of murine T-cells associated with the vaginal epithelium. *J. Immunol.*, 147:177.
- * Nanney, L.B., Stoscheck, C.M., Magid, M. and King, L.E. Jr. (1986): Altered epidermal growth factor binding and receptor distribution in psoriasis. *J. Invest. Dermatol.*, 86:260.
- * Nawroth, P.P., and Stern, D.M. (1986): Modulation of endothelial cell hemostatic properties by tumour necrosis factor. *J. Exp. Med.*, 163:740.
- * Nestle, F.O. and Nicholoff, B.J. (1995): Dermal dendritic cells are important members of the skin immune system. *Adv. Exp. Med. Biol.*, 378:111.
- * Neuner, P., Kapp, A., Kirnbauer, R., Schwartz, T., Krutmann, J. and Luger, T.A. (1989): Monocytes derived from patients with psoriasis synthesize and release increased levels of interleukin 6. *J. Invest. Dermatol.*, 92:490.
- * Neuner, P., Urbanki, A., Trautinger, F., Miller, A., Kirnbauer, R., Kapp, A., Schopf, E., Schwartz, T. and Luger, T. (1991): Increased IL-6 production by monocytes and keratinocytes patients with psoriasis. *J. Invest. Dermatol.*, 97:27.
- * Nguyen, D.T., Eskandari, M.K., Deforge, L.E., Raiford, C.L. Kunkel, S.L. and Remick, D.G. (1990): Cyclosporin a modulation of tumour necrosis factor gene expression and effects in vitro and vivo. *J. Immunol.*, 144: 3822.

- * Nicolas, J.F.; Kaiserlian, D., Daretienne, M., Faure, M. and Thivolet, J. (1987): Epidermal cell-derived lymphocyte differentiating factor (ELDIF) inhibits in vitro lymphoproliferative responses and interleukin-2 production. *J. Invest. Dermatol.*, 88:161.
- * Nikaein, A., Phillips, C., Gilbert, S.C., Savino, D., Silverman, A., Stone, M.J. and Menter, A. (1991): Characterization of skin-infiltrating lymphocytes in patients with psoriasis. *J. Invest. Dermatol.*, 96:3.
- * Nikolic-Zugic, J., and Bevan, M. (1990): Role of self peptides in positively selecting the T-cells repertoire. *Nature.*, 344:65.
- * Nickoloff, B.J. (1989): Role of interferon-gamma in cutaneous trafficking of lymphocytes with emphasis on molecular and cellular adhesion events. *Arch. Dermatol.*, 124:1835.
- * Nickoloff, B.J. (1991): The cytokine network in psoriasis. *Arch. Dermatol.*, 127:871.
- * Nickoloff, B.J., Riser, B.L., Mitra, R.S., Dixit, V. M. and Varani, J. (1988): Inhibitory effects of gamma interferon on cultured human keratinocyte thrombospondin production, distribution, and biologic activities. *J. Invest. Dermatol.*, 91:213.
- * Nickoloff, B.J., Barker, J., Griffiths, C.E.M., Elder, J.T., Kunkel, S. and Dixit, V. (1990): Molecular and cellular localization of IL-8 and its inducer-TNF in psoriasis. *J. Invest. Dermatol.*, 94:559 (abstr.).
- * Nickoloff, B.J., Fivenson, D.P., Griffiths, C. and Modlin, R. (1991): Preferential but not unique cutaneous endothelial cell expression of ELAM-1. *Clin. Res.*, 39:153.
- * Norris, D.A., Travers, J.B. and Leung D.Y.M. (1997): Lymphocyte activation in the pathogenesis of Psoriasis. *J. Invest. Dermatol.*, 109:1.
- * Nunberg, J.H., Doyle, M.V., York, S.M. and York, C.J. (1989): Interlekin-2 acts as an adjuvant to increase the potency of inactivated rabies virus vaccine. *Proc. Natl. Acad. Sci, USA.*, 86:4240.

- * O'Garra, A. (1989): Interleukins and the immune system 1. *Lancet*, 1:943.
- * O'Garra, A. Umland, S., France, T., and Christiansen, J. (1988): β cell factor are pleiotropic *Immunol. Today*, 9: 45.
- * O'Garra, A., Stapteton, G., Dhar, V., Pearce, M., Stall, A., Cupp, J., Mosmann, T. and Howard, M. (1990): Production of cytokines by mouse β cells; β lymphomas and normal β cells produce interleukin 10. *Int. Immunol.*, 2:821.
- * Ohashi, Y., Takeshita, T., Nagata, K., Mori, S. and Sugamura, K. (1989): Differential expression of IL-2 receptor subunits, p55 and p75 on various populations of primary peripheral blood mononuclear cells. *J. Immunol.*, 143:3548.
- * Oppenheim, J.J., Kovacs, E.J., Matsushima, K. and Durum, S.K. (1986): There is more than one interleukin-1. *Immunol. Today*, 7:45.
- * Oppenheim, J.J. Ruscetti, F.W. and Falynek, C.R. (1991): Cytokines. In: *Basic and clinical Immunology*. Ed. by: Stites, P. and Terr A.I. Appelton and Lange. Norwalk, Los Altos., California.
- * Orenberg, E.K., Wilkinson, D.I., Mansbridge, J.N. (1983): Localization of beta-adrenergic receptor sites in human epidermis with a fluorescent beta- antagonist. *J. Invest Dermatol.*, 80:311.
- * Orfanos, C.I., Schaumberg-Lever, G. (1973): Alterations of the cell surfaces as a pathogenetic factor in psoriasis. *Arch. Dermatol.*, 107:38.
- * Oxholm, A.; Oxholm, P., Staberg, B. and Bendtzen, K. (1989): Interleukin-6 in the epidermis of patients with psoriasis before and during PUVA treatment. *Acta. Dermatol. Venereol. (Stockh)*, 69:195.
- * Oxholm, A. (1992): Epidermal expression of interleukin-6 and tumor necrosis factor-alpha in normal and immunoinflammatory skin states in humans. *APMIS-Suppl.*, 24:1 (abstr).

- * Parent, D., Bernard, B.A., Deshas, C., Heenen, M., and Darmon, M.Y. (1990): Spreading of psoriatic plaques: alteration of epidermal differentiation precedes capillary leakiness and anomalies in vascular morphology. *J. Invest. Dermatol.*, 95:333.
- * Pennica, D., Nedwin, G.E.; Hayflick, J.S., Seeburg, P.H., Derynck, R., Palladino, M.A., Kohr, W.J., Aggarawal, B.B. and Goeddel, D.V. (1984): Human tumor necrosis factor: precursor structure, expression and homology to lymphotoxin. *Nature*, 312:724.
- * Pennica, D., Hayflick, J.S., Bringman, T.S., Palladino, M.A. and Goeddel, D.V. (1985): Cloning and expression in *Escherichia coli* of DNA for murine tumour necrosis factor. *Proc. Natl. Acad. Sci. USA.*, 82:6064.
- * Pietenpol, J.A., Stein, R.W., Moran, E., Yaciuk, P., Schlege, R., Pittelkow, M.R., Munger, K., Howley, P.M. and Moses, H.L. (1990): TGF- β 1 inhibition of c-myc transcription and growth in keratinocytes in abrogated by viral transforming proteins with pRB binding domains. *Cell*, 61:777.
- * Pigatto, P.D., Bersani, L., Altomare, G., Mozzanica, N. and Finzi, A. (1989): Effects of cyclosporin A on psoriatic monocyte functions and secretion. *Acta. Dermatol. Venereol. (Stockh)*, 146:165.
- * Pigatto, P.D., Pigatto, L.B., Bigardi, A., Altomare, G. and Finzi, A.F. (1990): Factors secreted by untreated psoriatic monocytes enhance neutrophil functions. *J. Invest. Dermatol.*, 94:372.
- * Piguet, P.F., Grau, G.E. and Vassalli, P. (1990): Subcutaneous perfusion of tumor necrosis factor induces local proliferation of fibroblasts, capillaries, and epidermal cells, or massive tissue necrosis. *Am. J. Pathol.*, 136:103.
- * Pike, M.C., Lee, C.S., Elder, J.T., Voorhees, J.J. and Fisher, G.J. (1989): Increased phosphatidylinositol kinase activity in psoriatic epidermis. *J. Invest. Dermatol.*, 92:791.

- * Pober, J., Gimbrone, M., Lapierre, L., Fiers, W., Rothlein, R. and Springer, T. (1986): Overlapping patterns of activation of human endothelial cells by interleukin 1, tumour necrosis factor and immune interferon. *J. Immunol.*, 137:1893.
- * Polla, B.S. (1988): A role for heat shock proteins in inflammation? *Immunol. Today*, 9:134.
- * Porcelli, S., Brenner, B. and Band, H. (1991): Biology of the human gamma/delta T cell receptor. *Immunol Rev.*, 120:137.
- * Poulter, L.W., Segmour, G.J. and Duke, O. (1982): Immunohistochemical analysis of delayed type hypersensitivity in man. *Cell Immunol.*, 74:358.
- * Prens, E.P., Van-Joost, J., Hegmans, J.P. Thooft-Benne, K. and Yesselmuider, D.E. (1995): Effects of cyclosporine on cytokines and cytokines receptors in psoriasis. *J. Am. Acad. Dermatol.*, 33(6):947.
- * Priestley, G.C. (1983): Hyperactivity of fibroblasts cultured from psoriatic skin. Faster proliferation and effect of serum withdrawal. *Br. J. Dermatol.*, 109:149.
- * Proctor, M.S., Wilkinson, D.I. and Orenberg, E.K. (1979): Lowered cutaneous and urinary levels of polyamines with clinical improvement in treated psoriasis. *Arch. Dermatol.*, 115:945.
- * Quesniaux, V. (1992): Interleukin 9,10,11 and 12 and kit ligand: A brief overview. *Res. Immunol.*, 143: 385.
- * Quesada, J.R. and Gutterman, J.U. (1986): Psoriasis and alpha-interferon. *Lancet*, i:1466.
- * Ragaz, A. and Ackerman, B. (1979): Evolution, maturation, and regression of lesions in psoriasis. *Am. J. Dermatopathol.*, 1:199.

- * Remick, D.G. (1994): Macrophage-Derived cytokines In: Immunochimistry ed. by Van Oss C.J. and Van Regenmortel, M.H.V. Ch. 38 P. 995. Marcel Dekker, Inc. New York.
- * Reusch, M., Studtmann, K., Schrder, M.J., Sticherling, M. and Christophers, E. (1990): NAP-1/IL-8 is a potent mitogen for human keratinocytes in vitro. J. Invest. Dermatol., 95:485.
- * Robb, R.J. (1984): Interleukin 2 the molecule and its function. Immunology Today, 5:203.
- * Robb, R.J., Munck, A. and Smith, K.A. (1981): T-cell growth factor receptors: Quantification, specificity, and biological relevance. J. Exp. Med., 154:1455.
- * Robb, R., and Greene, W. (1983): Drrcet demonstration of the identity of T-cell growth factor binding protein and the Tac antigen. J. Exp. Med., 158:1332.
- * Robb, R.J., Kurtny, R.M., Panico, M., Morris, H.R. and Chowdhry, V. (1984): Amino acid squence and post transtational modification of human interleukin 2. Proc. Natl. Acad. Sci. U.S.A., 81:6486.
- * Robb, R.J. and Green, W.C. (1987): Intenalization of IL-2 is mediated by the beta chain of the high affinity IL-2 receptor. J. Exp. Med., 165:1201.
- * Rosenberg, S.A., Lotze, M.T., Muul, L.M., Leitman, S., Chang, A.E., Ettinghausen, S.E., Matory, Y.L., Skibber, J.M., Shiloni, E. and Vetto, J.T. (1985): Observations on the systemnic administration of autologous lymphokine-activated killer cells and recombinant interleukin-2 to patients with metastatic cancer. N. Engl. J. Med., 313:1485.
- * Rosenberg, S.A. and Lotze, M.T. (1986): Cancer immunotherapy using interleukin-2 and interleukin-2 activated lymphocytes. Ann. Rev. Immunol., 4:681.

- * Rosenberg, S.A., Lotze, M.T., Muul, L.M., Chang, A.G., Avis, F.P., Leitman, S., Linehan, W.M., Robertson, C.N., Lee, R.E. and Rubin, J.T. (1987): A progress report on treatment of 157 patients with advanced cancer using lymphokine-activated killer cells and interleukin-2 or high dose interleukin-2 alone. *N. Engl. J. Med.*, 316:889.
- * Rosenstein, M., Ettinghausen, S.E. and Rosenberg, S.A. (1986): Extravasation of intravascular fluid mediated by the systemic administration of recombinant interleukin-2. *J. Immunol.*, 137:1735.
- * Rothe, M. and Falange, V. (1989): Growth factors. Their biology and promise in dermatologic diseases and tissue repair. *Arch. Dermatol.*, 125:1390.
- * Rubin, L.A., Kurman, C.C., Fritz, M.E., Biddison, W.E., Yarchoan, R. and Nelson, L.(1985): Soluble interleukin-2 receptors are released from activated human lymphoid cells in vitro. *J. Immunol.*, 135:3172.
- * Rubin, L.A., Jay, G. and Nelson, D.L. (1986): The released interleukin 2 receptor binds interleukin-2 efficiently *J. Immunol.*, 137:3841.
- * Ruddle, N. and Waksman, B. (1968): Cytotoxicity mediated by soluble antigen and lymphocytes in delayed hypersensitivity, *J. Exp. Med.*, 128:1255.
- * Rusk, C.Y., Neeper, M.P., Kuo, L.M., Kutny, R.M. and Robb, R.J. (1988): Structure function relationships for the IL-2 receptor system. V structure activity analysis of modified and truncated forms of the Tac receptor protein: site-specific mutagenesis of cysteine residues. *J. Immunol.*, 140:2249.
- * Russell, D.H., Cobest, W.L., Duell, E.A., Stawiski, H.A., Anderson, T.F. and Voorhees, J.J. (1978): Glucocorticoids inhibit polyamine biosynthesis in psoriasis. *J. Invest. Dermatol.*, 71:177.

- * Ruzicka, T., Simmet, T., Peskar, B.A. and Ring, J. (1986): Skin levels of arachidonic acid-derived inflammatory mediators and histamine in atopic dermatitis and psoriasis. *J. Invest. Dermatol.*, 86:105.
- * Sackstein, R., Falanga, V.m Streilein, W. and Chin, Y.H. (1988): Lymphocyte adhesion to psoriatic dermal endothelium is mediated by a tissue-specific receptor /ligand interaction. *J. Invest. Dermatol.*, 91:423.
- * Saiag, P., Coulomb, B., Lebreton, C., Bell, E. and Dubertret, L. (1985): Psoriatic fibroblasts induce hyperproliferation of normal keratinocytes in a skin equivalent model in vitro. *Science*, 230:669.
- * Salomon, D. and Halioua, B. (1990): Cyclosporin in dermatology. *Schweiz-Rundsch-Med. Prax.*, 16:1258 (Abstr.).
- * Sauve, K., Nachman, M., Spence, C., Bailon, P., Campbell, E., Tsien, W.H., Kondas, J.A., Hakimi, J. and Ju, C. (1991): Localization in human interleukin-2 of the binding site to the alpha chain (p55) of the interleukin-2 receptor. *Proc. Natl. Acad. Sci. USA.*, 88:4636.
- * Schlaak, J.F., Buslau, M., Jachum, W. Hermann, E., Cirndt, M. and Gallati H. (1994): T cell involved in psoriasis vulgaris belong to the Th 1 subset .. *J. Invest. Dermatol.*, 102 (2):145.
- * Schophf, E., Schwarz, T. and Luger, T.A. (1991): Increased IL-6 production by monocytes and keratinocytes in patients with psoriasis. *J. Invest. Dermatol.*, 37:27.
- * Schroder, J.M., Mrowietz, V., Morita, E. and Christophers, E. (1987): Purification and partial biochemical characterization of a human monocyte-derived, neutrophil-activating peptide that lacks interleukin 1 activity. *J. Immunol.*, 139:3473.
- * Schroder, J.M. and Christophers, E. (1989): Secretion of novel and homologous neutrophil-activating peptides by LPS-stimulated human endothelial cells. *J. Immunol.*, 142:244.

- * Schubert, C. and Christophers, E. (1985): Mast cells and macrophages in early relapsing psoriasis. *Arch. Dermatol. Res.*, 277:352.
- * Schuler, G., and Steinman, R. (1985): Murine epidermal Langerhans cells mature into potent immuno activating factor. *Lymphokines*, 11:87.
- * Sher, A., and Coffman, R.R. (1992): Regulation of immunity to parasites by T cells and T-cell derived cytokines. *Ann. Rev. Immunol.*, 10:385.
- * Shirai, I., Yamaguchi, H. Ito, H., Todd, C.W. and Wallace, R.B. (1985): Cloning and Expression in *Escherichia coli* of the gene for human tumor necrosis factors. *Nature.*, 313:803.
- * Smith, K.A. (1980): T- cell growth factor. *Immunol. Rev.*, 51:51.
- * Smith, K.A. (1984): Interleukin 2, *Annu Rev. Immunol*, 2:319.
- * Smith, K.A., Gillis, S., Baker, P.E. Mckerzie, D. and Ruscett, F.W. (1979): T cell growth factor mediated T cell proliferation. *Ann. Ny. Acad. Sci.*, 332:423.
- * Smith, K.A. (1988): Interleukin 2: Inception impact, and implications. *Science*, 240:1169.
- * Smith, C., Davis, T., Anderson, D., Solam, L., Jerzy, R., Power, S. and Goodwin, R. (1990): A receptor for tumour necrosis factor defines an unusual family of cellular and viral protein *Science*, 248:1019.
- * Soter, N.A., Kopelson, P., Karas, E. and Lim, H.W. (1989): Effect of cytokines on purified murine mast cells. *J. Invest. Dermatol.*, 92:521.
- * Spits, H., Yssel, H., Paliard, X., Kastelein, R., Figdor, C. and De Vries, J.E. (1988): IL-4 inhibits IL-2 mediated induction of human, lymphokine-activated killer cells, but not the generation of antigen-specific cytotoxic T lymphocytes in mixed leukocyte cultures. *J. Immunol.*, 141:29.

- * Sprent, J., Lo, D., Gao, E., and Ron, Y. (1988): T cell selection in the thymus. *Immunol Rev.*, 101:173.
- * Stawiski, M.A., Rusin, L.J., Burns, T.L. (1979): Ro 20-1724: An agent that significantly improves psoriatic lesions in double-blind clinical trials. *J. Invest. Dermatol.*, 73:261
- * Stethi, K.K. and Naher, H. (1986): Elevated titres of cell free interleukin of receptors in serum and cerebrospinal fluid specimen of patients with acquired immuno deficiency syndrome. *Immunol.*, 13:179.
- * Stoscheck, C.M. and King, L. E. Jr. (1986): Functional and structural characteristics of EGF and its receptor and their relationship to transforming proteins. *J. Cell. Biochem.*, 31:135.
- * Strange, P., Cooper, H.D., Hansen, E.R., Fisher, G., Larsen, J.K., Fox, D., Hrag, C., Voorhees, J.J. and Baadsgaard, O. (1993): T-lymphocyte clones initiated from lesional psoriatic skin release growth factor that induce keratinocyte proliferation. *J. Invest. Dermatol.*, 101:695.
- * Strieter, R.M., Kunhlee, S.L., Showell, H.J., Renick, D.G., Phan, S.H. and Marks, R.M. (1989): Endothelial cell gene expression of a neutrophil chemotactic factor by TNF-alpha, LPS, and IL-1beta. *Science*, 243:1467.
- * Swope, V., Abdel-Malek, Z. and Nordlund, J. (1990): IL-1 alpha, IL-6 and tumor necrosis factor-alpha are paracrine inhibitors of human melanocyte proliferation and melanogenesis. *J. Invest. Dermatol.*, 94:582.
- * Tagami, H.I., Jwatsuki, K. and Takematsu H. (1987): Psoriasis and leukocyte chemotaxis. *J. Invest. Dermatol.*, 88:18.
- * Taira, S., Matsui, M., Hayakawa, M., Yokoyama, T. and Naraichi, H. (1987): Interleukin secretion by cell lines and splenic β cells stimulated with calcium ionophore and phorbol ester. *J. Immunol.*, 139:2957.

- * Takahashi, H., Lizuka, H. and Katagiri, M. (1990): No evidence for the mutation of ras gene in psoriatic epidermis. *Arch. Dermatol. Res.*, 282:9.
- * Takematsu, H., Ohta, H. and Tagami, H. (1989): Absence of tumor necrosis factor- α in suction blister fluids and stratum corneum from patients with psoriasis. *Arch. Dermatol. Res.*, 281:398.
- * Takematsu, H. and Tagami, H. (1990): Interleukin-2, soluble interleukin-2 receptor, and interferon- γ in the suction blister fluids from psoriatic skin. *Arch. Dermatol. Res.*, 282:149.
- * Takeshita, T., Asoa, H., Tanaka, N. and Sugamura, K. (1992): Cloning of the gamma chain of the human IL-2 receptor. *Science*, 257:379.
- * Talwar, H.S., Fisher, G.J., Havris, V.A. and Voorhees, J.J. (1989): Agoniste-induced hydrolysis of phospho inositides and formation of 1,2-diacylglycerol in adult human keratinocytes. *J. Invest. Dermatol.*, 93:241.
- * Taniguchi, T., Matsui, H. and Hamuro, J. (1986): Molecular analysis of the interleukin-2 system, *Immunol Rev.*, 92:121.
- * Taniguchi, T. and Minami, Y. (1993): The IL-2 receptor system, a current overview. *Cell*, 73:5.
- * Teshigawara, K., Wang, H., Kato, K. and Smith, K. (1987): Interleukin-2 high-affinity receptor expression requires two distinct binding proteins *J. Exp. Med.*, 165:223.
- * Ternowitz, T. (1986): Monocyte and neutrophil chemotaxis in psoriasis. *J. Am. Acad. Dermatol.*, 15:1191.
- * Ternowitz, T. and Thestrup - Petersen, K. (1985): Neutrophil and monocyte chemotaxis in pustulosis palmo-plantaris and pustular psoriasis. *Br. J. Dermatol.*, 113:507.

- * Thoman, M.L. and Weigle, W.O. (1981): Lymphokines and aging: Interleukin-2 production and activity in aged animals. *J. Immunol.*, 127:2102.
- * Thompson-Snipes, L., Dhar, V., Bond, M., Mosmann, T., Moore, K. and Ronnick, D. (1991): Interleukin-10: A novel stimulatory factor for mast cells and their progenitors. *J. Exp. Med.*, 173:507.
- * Tigges, M.A., Casey, L.S. and Koshland, M.E. (1989): Mechanism of interleukin-2 signaling: mediation of different outcomes by a single receptor and transduction pathway. *Science*, 243:781.
- * Trotter, J.L., Clifford, D.B., and Anderson, C.B. (1988): Elevated serum interleukin-2. *N. Engl. J. Med.*, 318:1206.
- * Tsudo, M., Kitamura, F. and Miyasaka, M. (1989): Characterization of the interleukin-2 receptor β chain using three distinct monoclonal antibodies. *Proc. Natl. Acad. Sci. USA.*, 86:1982.
- * Tsujimoto, M., Yip, Y.K. and Vilcek, J. (1986): Interferon- γ enhances expression of cellular receptors for tumor necrosis factor. *J. Immunol.*, 136:2441.
- * Tucker, W.F.G., MacNeil, S., Bleehen, S.S. and Tomlinson, S. (1984): Biologically active calmodulin levels are elevated in both involved and uninvolved epidermis in psoriasis. *J. Invest. Dermatol.*, 82:298.
- * Tucker, W.F.G., MacNeil, S., Dawson, R.A., Tomlinson, S. and Bleehen, S.S. (1986): Calmodulin level in psoriasis: The effect of treatment. *Acta. Dermatol. venereol. (Stockh)*, 66:241.
- * Uchiyama, T., Broder, S. and Waldmann, T.A. (1981): A monoclonal antibody (anti Tac) reactive with activated and functionally mature human T cells. *J. Immunol.*, 126: 1293.

- * Ulich, T.R., Yin, S., Guo, K., Remich, D. and Delcastillo, J., (1991): Intratracheal injection of endotoxin and cytokines II. Interleukin-6 and transforming growth factor beta inhibit acute inflammation. *Am. J. Pathol.*, 138:1097.
- * Unchigama, T., Broder, S., and Waldmann, T. A. (1981): (A monoclonal antibody) (anti-Tac) reactive with activated and functionally mature human T. Cells, *J. Immunol.*, 126:1393.
- * Uyemura, K., Yamamamura, M., Fiverson, D.F., Modlin, R.L. and Nickoloff B.J. (1993): The cytokine network in lesional and lesion free psoriatic skin is characterized by a T-helper type 1 cell-mediated response. *J. Invest. Dermatol.*, 101:701.
- * Valdimarsson, H., Baker, B.S., Jonsdottir, I. and Fry, L. (1986): Psoriasis: a disease of abnormal keratinocyte proliferation induced by T-lymphocytes. *Immunology Today*, 7:256.
- * Valge, V.E., Wong, J.G.P., Datlof, B.M., Sinskey, A.J. and Roa, A. (1988): Protein Kinase C is required for responses to T cell receptor ligands but not to interleukin 2 in T cells. *Cell*, 55:101.
- * Vallat, V.P., Gilleaudeau, P., Battat, L., Wolfe, J., Nabeya, R., Heftler, N., Hodak, E., Gottlieb, A.B. and krueger J.G. (1994): PUVA bath therapy strongly suppresses immunological and epidermal activation in psoriasis: a possible cellular basis for remitties therapy. *J. Exp. Med.*, 180:283.
- * Van Damme, J., Van Beeumen, J., Opdenakker, G. and Billiau, A. (1988): NH₂- terminal sequence-characterized human monokine possessing neutrophil chemotactic, skin-reactive and granulocytosis promoting activity. *J. Exp. Med.*, 167:1364.
- * Van Snick, J. (1990): Interleukin-6 : An over view. *Ann. Rev. Immunol.*, 8:293.

- * Vermeer, M.H. and Streilein, J.W. (1990): Ultraviolet B and TNF-alpha cause similar alterations in intraepidermal Langerhans cells. *J. Invest. Dermatol.*, 94:588.
- * Vieira, D. Dang, M., Johnson, K., Kastelein, R., deVries, J. and Moore, K. (1991): Isolation and expression of human cytokine synthesis inhibitory factor (CSIF/IL10). Homology to Epstein-Barr virus open reading frame BCRF1. *Proc. Natl. Acad. Sci. USA.*, 88:1172.
- * Vissinga, C.S., Dirven, C.J.A.M., Steinmeyer, F.A., Benner, R. and Boersma, W.T.A. (1987): Deterioration of cellular immunity during aging. The relationship between age-dependent impairment of delayed -type hypersensitivity reactivity interleukin-2 production capacity. *Cell Immunol.*, 108:323.
- * Vollmer, S., Menssen, A., Trommler, P., Schendel, D. and Prinz, J.C. (1994): T lymphocytes derived from skin lesions of patients with psoriasis vulgaris express a novel cytokine pattern that is distinct from that of T helper type I and T helper type II cells. *Eur. J. Immunol.*, 24:2377.
- * Vonderheid, E.C., Kang, C.Ae., Kadin, M., Bigler, R.D., Griffin, T.D. and Rogers, T.J. (1990): Extracorporeal photopheresis in psoriasis vulgaris: Clinical and immunologic observations. *J. Am. Acad. Dermatol.*, 23:703.
- * Voorhess, J.J. (1983): Leukotrienes and other lipoxigenase products in the pathogenesis and therapy of psoriasis and other dermatoses. *Arch. Dermatol.*, 119:541.
- * Wagner, D.K., Kiwanuka, J., Edwards, B.K., Rubin, L.A., Nelson, D.L. and Margrath, I.T. (1987): Soluble interleukin-2 receptors levels in patients with undifferentiated and lymphoblastic lymphomas. Correlation with survival. *J. Clin. Oncol.*, 5:1262.
- * Wahba, A., Cohen, H.A., Bar-Eli, M. and Gallily, R. (1978): Enhanced chemotactic and phagocytic activity of leukocytes in psoriasis vulgaris. *J. Invest. Dermatol.*, 71:186.

- * Wakefield, P.E., James, W.D., Samlaska, C.P. and Meltzer, M.S. (1991): Tumor necrosis factor. *J. Am. Acad. Dermatol.*, 24:675.
- * Waldmann, T., Goldman, C. Robb, R. and Greene, W. (1984): Expression of interleukin 2 receptors on activated human β cells. *J. Exp. Med.*, 160:1450.
- * Waldmann, T. and Gobbold, S. (1993): The use of monoclonal antibodies to achieve immunological tolerance. *Immunol. Today*, 14:247.
- * Walz, A., Peveri, P., Aschauer, H. and Baggiolini, M. (1987): Purification and amino acid sequencing of NAF, a novel neutrophil-activating factor produced by monocytes. *Biochem. Biophys. Res. Commun.*, 149:755.
- * Wang, A.M., Creasey, A.A., Lander, M.B., Lin, L.S., Yamamoto, R., and Mark, D.F. (1985): Molecular cloning of the complementary DNA for human tumour necrosis factor. *Science*, 288:149.
- * Watson, J.D., and Mochizuti, D. (1980): IL-2 A class of lymphocyte growth factors. *Immunol. Rev.*, 512:257.
- * Weigel, V., Meyer, M. and Sebald, W. (1989): Mutant proteins of human interleukin-2: renaturation yield proliferative activity and receptor binding. *Eur. J. Biochem.*, 180:295.
- * Weinstein, G.D., McCullough, J. and Colton, A. (1983): Proliferative defects defects in psoriasis. *J. Invest. Dermatol.*, 80:360.
- * Weinstein, G.D., McCullough, J.L. and Ross, P.A. (1985): Cell kinetic basis for pathophysiology of psoriasis. *J. Invest. Dermatol.*, 95:579.
- * Wheelock, E. (1965): Interferon like virus-inhibitor induced in human leukocytes by phytohemagglutinin. *Science*, 149:310.
- * Wikner, N.E., Huff, J.C., Norris, D.A., Boyce, S.T., Cary, M., Kissinger, M. and Wesfon, W.L. (1986): Study of HLA-DR synthesis in cultured human keratinocytes. *J. Invest. Dermatol.*, 87:559.

- * Wiley, H.E. and Weinstein, G.D. (1979): Abnormal proliferation of uninvolved psoriatic epidermis: Differential induction by saline, propranolol and tape stripping in vivo. *J. Invest. Dermatol.*, 73:545.
- * Wofsy, D., Murphy, E.D., Roths, J.D., Daupinee, M.J., Kipper, S.B. and Tala, N. (1981): Deficient interleukin-2 activity in MRL/MP and C57 BL/6J mice bearing the Ipr. gene. *J. Exp. Med.*, 154:1671.
- * Wollina, V., Klinger, R., Wetzker, R. and Knopf, B. (1989): Increase of epidermal calmodulin precedes the formation of a psoriatic lesion. *Arch. Dermatol. Res.*, 281:73.
- * Wong, E., Camp, R.D. and Greaves, M.W. (1985): The response of normal and psoriatic skin to single and multiple topical applications of leukotriene. *Arch. Dermatol.*, 119:541.
- * Wong, G.H.W., Krowka, J.R., Stites, D.P. and Goeddel, D.V. (1988): In vitro anti-human immunodeficiency virus activities of tumour necrosis factor- α and interferon-gamma. *J. Immunol.*, 140:120.
- * Woorhees, J.J. (1983): Leukotrienes and other lipoxygenase products in the pathogenesis and therapy of psoriasis and other dermatoses. *Arch. Dermatol.*, 119:541.
- * Yagita, H., Nakata, M., Azuma, A., Nitta, T., Takeshita, T., Sugamura, K. and Okumura, K. (1989): Activation of peripheral blood T cells via the p75 interleukin 2 receptor. *J. Exp. Med.*, 170:1445.
- * Yamaski, K., Taga, T. and Hirata, Y. (1988): Cloning and expression of human IL-6 receptor. *Science*, 241: 825.
- * Yang, Y., Ricciardi, S., Ciarletta, A., Clavetti, J. and Clark, S. (1989): Expression cloning of DNA encoding a novel human hematopoietic growth factor: Human homologue of murine T-cell growth factor. *Blood*, 74:1880.

- * Yarden, Y., Escobedo, J.A., Kuang, W.J., Yang-Feng, T.L., Danie, T.O., Tremble, P.M., Chen, E.Y., Ando, M.E., Harkins, R.N., Francke, V., Fried, V.A., Ullrich, A. and Williams, L.T, 1986 : Structure of the receptor for platelet derived growth factor help define a family of closely related growth factor receptors. *Nature.*, 323:226.
- * Yasumoto, S. Imayama, S. and Hori, I. (1995): Increased serum level of interleukin-6 in patients with psoriatic arthritis and thrombocytosis. *J. Dermatol.*, 22(10):718.
- * Yokota, T., Arai, N., Spit, H. and Avai, K. (1988): Molecular biology of interleukin 4 and interleukin 5 genes *Immunol. Rev.*, 102:113.