

REFERENCES

REFERENCES.

- Bardin, B.C. and Paulsen, C. A. (1981):
The testis. In: Textbook of Endocrinology,
Williams, R.H. (ed). Philadelphia, W.B.
Saunders Company., P. 293.
- Barham, S.S.; Berlin, J.D. and Brackeen, R.B. (1976):
The fine structural localization of testi-
cular phosphatase in man: The control testis
Cell Tiss. Res. 166: 497.
- Bawa, S.R. (1963):
Fine structure of the Sertoli cell of the
human testis, J. Ultrastruct. Res. 9: 459.
- Bhalla, V.K. and Reichert, L.E.Jr. (1974):
Properties of follicle stimulating hormone-
receptor interactions. J. Biol. chem.
249: 43.
- Black, V.H. (1971):
Gonocytes in fetal guinea-pig testis phago-
cytosis of degenerating gonocytes by Sertoli
cells Am. J. Anat. 131: 415.
- Blanchard, W. G. and Josso, N. (1974):
Source of the anti-Mullerian hormone syn-
thesized by the fetal testis. Pediat.
Res. 8 : 968.

Blaquier, J.A.; Cameo, M.S. and Burgos, M.H. (1972):

The role of androgens in the maturation of epididymal spermatozoa in the guinea pig. *Endocrinol.* 90 : 839.

Borland, K.; Mita, M.; Oppenheimer, C.L. ; Blinderman, L.A.; Massaque, J.; Hall, P.F.; and Czech , M.P. (1984):

The actions of insulin-like growth factors I and II on cultured Sertoli cells. *Endocrinol* 114: 240.

Bouin , P. and Angel, P. (1903):

Sur la signification de la glande interstitielle du testicule embryonnaire. *Compl. Rend. Soc. Biol.*, 55: 1682. Quoted by Fawcett, D.W. (1975): Ultrastructure and function of the Sertoli cell. In: *Handbook of Physiology, Endocrinology*, Greep, R.O. and Astwood, E.B. (eds), Washington, American Physiological Society, P. 21.

Brokelmann, J. (1963):

Fine structure of germ cells and Sertoli cells during the cycle of the seminiferous epithelium in the rat. *Z. Zellforsch. Mikroskop. Anat.*, 59: 820. Quoted by Fawcett, D.W. (1975): Ultrastructure and function of

Sertoli cell. In: Handbook of Physiology; Endocrinology, Greep, R.O. and Astwood, E.B. (eds), Washington, American Physiological Society, P. 21.

Burgos, M.H. and Fawcett, D.W. (1955)

Studies on the fine structure of the mammalian testis. J. Biophys. Biochem. Cytol. 1 : 287.

Burgos, M.H. and Vitale-Calpe, R. (1967):

The mechanism of spermiation in the toad. Am. J. Anat. 120 : 227.

Burgos, M.H. and Vitale-Calpe, R. (1969):

Gonadotrophic control of spermiation. In: Progress in Endocrinology, Gual, C. (ed). Amsterdam, Excerpta Medica Foundation, P. 1030.

Chemes, H.E.; Dym, M. and Raj, H. G.M. (1979):

Hormonal regulation of Sertoli Cell differentiation. Biol. Reprod., 21 : 251.

Chowdhury, M.; Steinberger, A. And Steinberger, E. (1978)

Inhibition of denovo synthesis of FSH by the Sertoli cell factor. Endocrinol. 103 : 644.

Christensen, A.K. and Mason, N.R. (1965):

Comparative ability of seminiferous

tubules and interstitial tissue of rat testes to synthesize androgens from progesterone in vitro *Endocrinol* 76 : 646.

Christensen, A.K. (1965):

The fine structure of testicular interstitial cells in the guinea pig. *J. cell biol.* 26 : 911.

Christensen, A.K. and Fawcett, D.W. (1966):

The fine structure of the interstitial cells of the mouse testis. *Am. J. Anat.* 118 : 551.

Christensen, A.K. (1975):

Leydig cells, In: *Handbook of Physiology*, Hamilton, D. and Greep, R.O. (eds). Baltimore, Williams and Wilkins. P. 57.

Clegg, E.J. and MacMillan, E.W. (1965):

The uptake of vital dyes and particulate matter by the Sertoli cells of the rat testis, *J. Am.* 99 : 219.

Clermont, Y. (1966): Renewal of spermatogonia in man
Am. J. Anat., 118 : 509.

Clermont, Y.; McCoshen, J. and Hermo, L. (1980):

Evolution of the endoplasmic reticulum in the Sertoli cell cytoplasm encapsulating

the heads of late spermatids in the rat.

Anat. Rec. 196 : 83.

Cohen, P.; Burchell, A.; Foulkes, J.G.; Cohen, P.T.W.;

Vanaman, T.C. and Nairn, A.C. (1978):

Identification of the Ca^{+2} dependent
modulator protein as the fourth subunit
of rabbit skeletal muscle phosphorylase
kinase. F.E.B.S. Lett., 92 : 297.

Collin S.P. and Lacy, D. (1969): Studies on the struc-

ture and function of the mammalian testis.

Proc. Roy. Soc. London, Ser. 172 : 17.

Colman, O.D. and Stockert, J.C. (1978):

Observations on the nucleolar staining
by ethanolic phosphotungstic acid. Microsc.
Acta 81 (1) : 27.

Connell, C.J. (1978):

A freeze fracture and lanthanum tracer
study of the complex junction between
Sertoli: Cells of the canine testis.
J. Cell. Biol. 76 : 57.

Crew , F.A.E. (1922): A suggestion as to the cause

of the aspermatic condition of the imper-
fectly descended testis. J. Anat., 56:98.

Quoted by Hall, P. F.; Kew, D.; and Mita,
M. (1985): The influence of temperature on

primary culture. Endocrinol., 105: 1419.

de Kretser, D.M. and Burger, H.G. (1972):

Hormonal, histological and chromosomal studies in adult males with testicular disorders. J. Clin. Endocrinol. Metab., 35 : 392.

de Kretser, D.M.; Burger, H.G.; Hudson, B. and Keogh, E.J. (1975a):

The HCG stimulation test in men with testicular disorders. Clin. Endocrinol., 4:591.

de Kretser D.M.; Kerr, J.B. and Paulsen, C.A. (1975b):

The peritubular tissue in the normal and pathological human testis. Biol. Reprod., 12: 317.

Demoulin, A.; Koulischer, L.; Hustin, J.; Hazee, -

Hagelstein, M.T.; Lambotte, R. and Franchimont, P. (1979): Organ culture of mammalian testes. Horm. Res. 10 : 177.

Desjardins, C.; Zeleznik, A.J.; Midgley, A.R. and Reichert, L. E. Jr. (1974):

In vitro binding and autoradiographic localization of human chorionic gonadotrophin and follicle stimulating hormone in rat testis during development. In: Hormone Binding and Target Cell Activation in Testis,

Dufau, M.L. and Means, A.R. (eds) New York, Plenum Press. P. 221.

Donahoe, P.K.; Ito, Y.; Morikawa, Y. and Handren, W.H. (1977): Mullerian inhibiting substance in human testes after birth. J. Pediat Surg., 12 : 323.

Dufau, M.L.; Catt, K.J. and Tsurahara, T. (1971): Gonadotropin stimulation of testosterone production by the rat testis in vitro. Biochem. Biophys. Acta, 252 : 574.

Dym, M. and Fawcett, D.W. (1970): The blood-testis barrier in the rat and the physiological compartmentation of the seminiferous epithelium. Biol. Reprod., 3:308.

Dym, M. and Fawcett, D.W. (1971): Further observations on the numbers of spermatogonia, spermatocytes and spermatids joined by intercellular bridges in mammalian spermatogenesis. Biol. Reprod, 4:195.

Dym, M. (1977): The male reproductive system. In: Textbook of Histology, Weiss, L. and Greep, R.O. (eds), New York, McGraw-Hillbook Company A Blackiston Publication P. 979.

Dym, M. and Cavicchia, J.C. (1977):

Further observations on the blood testis barrier in monkeys. Biol. Reprod., 17:390.

Dym, M. and Madhwa Raj, H.G. (1977):

Response of adult rat Sertoli cells and Leydig cells to depletion of lutenizing hormone and testosterone. Biol. Reprod., 17 : 676.

Dyson, A.L.M.B. and Orgebin-Crist, M.C. (1973):

Effect of hypophysectomy, Castration and androgen replacement upon the Fertilizing ability of rat epididymal spermatozoa. Endocrinol., 93 : 391.

Edress, Z.; Raslan, N.; Badawy, Z. and Rashwan, S. (1977) : Phagocytic potentialities of Sertoli cells. Medical J. of Cairo University. 45 : 77.

Elkington, J.S.H.; Sanborn, B.M. and Martin, M.W.(1977):

Effects of testosterone propionate on ABP levels in rat hypophysectomized at different ages using individual sampling. Mol. cell Endocrinol., 6 : 203.

Ewing, L.L. and Chang, T.K. (1986):

The Testis , Epididymis, and Ductus Deferens. in Campbell's Urology, Cann, C. (ed) Philadelphia, W.B Saunders Company P. 200.

Fakunding, J.L. ; Tindall, D.J.; Dedman, J.R.; Mena, C.R. and Means, A.R. (1976): Biochemical actions of FSH in the Sertoli cell of the rat testis. *Endocrinol*, 98: 392.

Fawcett, D.W. and Burgos, M.H. (1956):

The fine structure of Sertoli cells in the human testis. *Anat. Rec.* 124: 401.

Fawcett, D.W.; Ito , S. and Slautterback, D.L. (1959):

The occurrence of intercellular bridges in groups of cells exhibiting synchronous differentiation. *J. Biophys. Biochem. Cytol.* 5 : 453.

Fawcett, D.W. and Phillips, D.M. (1969):

Observations on the release of spermatozoa and on changes in the head during passage through the epididymis *J. Reprod. Fertility (Suppl.)* 6 : 405.

Fawcett, D.W.; Long, J. A. and Jones, A.L. (1969):

The ultrastructure of the endocrine glands. *Recent progr. Hormone Res.* 25 : 315.

Fawcett, D.W.; Leak, L.V. and Heidger, P.M. (1970):

Electron microscopic observations on the structural components at the blood testis barrier. *J. Reprod. Fertil.* 10 : 105.

Fawcett, D.W. (1975): Ultrastructure and function of

the Sertoli cell. In: Handbook of Physiology; Endocrinology, Greep, R.O. and Astwood, E.B. (eds), Washington, American Physiological Society, P. 21.

Fawcett, D.W. (1979):

The cell biology of gametogenesis in the male. *Perspect. Biol. Med.* p. 56.

Fawcett, D.W. and Hoffer, A.P. (1979): Failure of exogenous androgen to prevent regression of the initial segments of the rat epididymis after efferent duct ligation or orchidectomy. *Biol. Reprod.* 20 : 162.

Flickinger, C.J. (1967): The postnatal development of the Sertoli cells of the mouse. *Z. Mikrosk. Anat. Forsch.* 78 : 92. Quoted by Tindall, D.J., Rowley, D.R. and Lipshultz, L.I.
(1983): Sertoli cell structure and function in vivo and in vitro. In: *Infertility in the male*, Lipshultz, L.I. and Howards, S.S. (eds). New York, Churchill Livingstone., P. 71.

Flint, A.P.F. and Armstrong, D. T. (1971): The compartmentation of nonesterified and esterified cholesterol in the superovulated rat ovary *Biochem. J.* 123: 143.

Fonkalsrud, E.W. and Mengel, W. (1981): Morphology of normal and cryptorchid testes. In: The Undescended Testis, Fonkalsrud, E.W. and Mengel, W. (eds). Chicago, Year Book Medical Publishers. p. 59.

Franke, W.W.; Grund, C. and Schmid, E. (1979): Intermediate-sized filaments present in Sertoli cells are of the vimentin type. Eur. J. Cell Biol. 19: 269.

French, F.S. and Ritzen, E.M. (1973): A high affinity androgen-binding protein (ABP) in rat testis. Endocrinol. 21: 611.

French, F.S.; Mclean, W.S. and Smith, A.R. (1974): Androgen transport and receptor mechanisms transport and receptor mechanisms in testis and epididymis. In: Hormone Binding and Target Cell Activation in the Testis, Dufau, M. L. and Means, A.R. (eds) New York, Plenum Press p. 311.

Fritz, I.B.; Kopee, B. ; Lam, K. and Vernon, R.G. (1974): Effects of FSH on levels of androgen binding protein in the testis. In: Hormonal binding and target cell activation in the testis, Means, A.R. and Dufau, M.L. (eds). New York, Plenum Publishing Corp. p. 156.

- Fritz, I.B.; Rommerts, F.G.; Louis, B. G. and Dorrington, J.H. (1976): Regulation by FSH and dibutyryl cyclic AMP of the formation of androgen binding protein in Sertoli cell enriched cultures J. Reprod. Fertil., 46 : 17.
- Fritz, I.B. (1978): Sites of action of androgens & F.S.H on cells of seminiferous tubules. In: Biochemistry of Hormones, Litwack, G. (ed) , New York, Academic Press, P. 249.
- Furuya, S.; Kumamoto, Y. and Sugrama, S. (1978): Fine structure and development of Sertoli junctions in human testis. Arch. Androl. 1:211.
- Gabrilove, J.L.; Freiberg, E.K.; Leiter, E. and Nicolis, G.L. (1980): Feminizing and non-feminizing Sertoli cell tumors J. Urol., 124: 757.
- Gilula, N.B.; Fawcett, D.W. and Aoki, A. (1976): The Sertoli cell occluding junction and gap junctions in mature and developing mammalian testis. Develop. Biol. 50: 142.
- Girgis, S.M. and Hafez, E.S.E. (1980): Basic concepts in andrology. In: Human Reproduction; Conception and Contraception, Hafez, E.S.E. (ed), New York, Harper and Row Publishers, p. 130.

- Gravis, C.J. (1979): Interrelationships between Sertoli cells and germ cells in the Syrian Hamster. *Z. Mikrosk. Anat. Forsch.* 93:321. Quoted by Tindall, D.J.; Rowley, D.R. and Lipshultz, L.I. (1983): Sertoli cell structure and function in vivo and in vitro. In: infertility in the Male, Lipshultz, L.I. and Howards, S.S. (eds) New York; Churchill Livingstone., P. 71.
- Griffiths, J. (1893): The structural changes in the testicle of the dog when it is replaced within the abdominal cavity. *J. Anat. Physiol.*, 27 : 483. Quoted by Hall, P. F.; Kew, D.; and Mita, M. (1985): The influence of temperature on the functions of cultured Sertoli cells. *Endocrinol.*, 116 : 1926.
- Griswold, M.D.; Mably, E.R. and Fritz, I.B. (1976): FSH stimulation of DNA synthesis in Sertoli cells in culture. *Mol. cell Endocrinol.*, 4:139.
- Guerriero, Jr. V.; Rowley, D.R. and Means, A.R. (1981): Production and characterization of an antibody to myosin light chain kinase and intracellular localization of the enzyme. *Cell* 27 : 449.
- Hadziselimovic, P. (1977):: Cryptorchidism. Ultrastructure of normal and cryptorchid testis development *Adv. Anat. Embryol. Cell Biol.*, 53 : 3.

- Hagenas, L.; Ritzen, E.M.; French, F.S. and Hansson, V.
(1974): Sertoli cell production of testicular
androgen-binding protein (ABP). J. Steroid
Biochem. 5:383.
- Hagenas, L. and Ritzen, E.M. (1975): Impaired Sertoli
cell function in secondary cryptorchidism
in the rat. Mol. cell. Endocrinol., 7:1.
- Hagenas, L.; Ritzen, E.M.; Ploen, L.; Hansson, V.; French, F.S. and Nayfeh, S.N. (1975): Sertoli
cell origin of testicular androgen binding
protein. (ABP). Mol. cell Endocrinol.,
2:339.
- Hagenas, L.; Ritzen, E.M.; Svensson, J.; Hansson, V. and
Purvis, K. (1978a): Temperature dependence
of Sertoli cell Function. Int. J. Androl,
2 : 449.
- Hagenas, L.; Ploen, L. and Ritzen, E.M. (1978b): The
effect of nitrofurazone on the endocrine
secretory and spermatogenic functions of
the rat testis. Int. J. Androl., 2: 107.
- Hallex, C.G. and Clermont, Y. (1964): Kinetics of the
germinal epithelium in man. Recent prog.
Horm. Res., 20 : 545.
- Hall, P.F. (1965): Influence of temperature upon the
biosynthesis of testosterone by rabbit

- testis in vitro. *Endocrinol.*, 76: 396.
- Hall, P.F.; Irby, D.C. and dekretser, D.M. (1969):
Conversion of cholesterol to androgens by
rat testes. *Endocrinol.*, 84: 488.
- Hall, P.E.; Kew, D. and Mita, M. (1985): The influence
of temperature on the functions of cultured
Sertoli cells. *Endocrinol.*, 116:1926.
- Hansson, V. (1972): Further characterization of the 5α -
dihydrotestosterone binding protein in the
epididymal cytosol fraction. In vitro stud-
ies, *Steroids*, 20 : 575.
- Hansson, V.; Reusch, E.; Trygstad, O.; Torgersen, O.;
French, F.S. and Ritzen, E.M. (1973): FSH
stimulation of testicular androgen binding
protein (ABP). *Nature New Biol.* 246:56.
- Hansson, V.; Trygstad, O.; French, F.S.; McLean, W.S.;
Smith, A.A.; Tindall, D.J.; Weddington, S.C.;
Petrusz, P.; Nayfeh, S.N. and Ritzen, E.M.
(1974): Androgen transport and receptor
mechanisms in testis and epididymis. *Nature*,
250 : 367.
- Hansson, V.; Ritzen, E.M.; French, F.S. and Nayfeh, S.N.
(1975): Androgen transport and receptor
mechanisms in the testis and epididymis. In:
Handbook of physiology, Section 7, Endocrinol.

Vol. 5, Greep, R. and Astwood, E.B. (eds).
Washington, D.C. Amer. Physiol Society,
p. 21.

Hansson, V.; Ritzen, E.M. and French, F.S. (1976): Androgen transport and receptor proteins in the testis. In: Human Semen and Fertility Regulation in Men, Hafez, E.S.E. (ed) Saint Louis The C.V. Mosby Company p. 51.

Hansson, V.; Purvis, K. and Attramadal, A. (1978): Sertoli cell function in the androgen insensitive (tfm) rat. Int. J. Androl., 1:96.

Heller, C.G.; Paulsen, C.A.; Mortimore, G. E.; Jungck, E.C. and Nelson, W.O. (1952): Urinary gonadotrophins, spermatogenic activity and classification of testicular morphology, Ann. NY Acad. Sci., 55 : 685.

Quoted by Bardin and Paulsen (1981)

Herbert, D.C. and Tindall, D.J. (1980): Epididymal androgen binding protein in protein-calorie malnourished rats. Endocrinol. Res. Commun. 7 : 61.

Hilscher, W. (1978): Beitrage den spermatogenese des menschen. Symposion "Prophylaktische Andrologie in Kindesalter". Remscheid 26:8. Quoted by Fonkalsrud, E.W. and Mengel, W. (1981):

Res., 5:510.

Josso, N. (1973): In vitro synthesis of Mullerian inhibiting substance hormone by seminiferous tubules isolated from the calf fetal testis. Endocrinol., 93: 829.

Josso, N. (1974): Mullerian-inhibiting activity of human fetal testicular tissue deprived of germ cells by in vitro irradiation. Pediat. Res. 8: 755.

Josso, N.; Picard, J.Y. ; Dacheux, J.L. and Courot, M. (1979): Detection of anti-Mullerian activity in boar rete testis fluid J. Reprod. Fertil. 57:397.

Jost, A. (1947): Recherches sur la differenciation sexuelle de l'embryon de lapin. Arch. Anat. Micro. Morpho. Exp. 36:271. Quoted by Price J.M. (1979).: The secretion of Mullerian inhibiting substance by cultured isolated Sertoli cells of the neonatal calf. Am. J. of Anat. 156:147.

Jost, A. (1953): Problems of fetal endocrinology: The gonadal and hypophyseal hormones. Rec. Prog. in Hor. Res. 8: 379.

Jost, A.; Vigier, B.; Prepin, J. and Perchellet, J.P.

- (1973): Studies on sex differentiation in mammals Recent Prog. Horm. Res. 29 : I.
- Karl, A.F. and Griswold, M.D. (1980): Actions of insulin and vitamine A on Sertoli cells. Biochem. J., 186:1001.
- Kerr, J.B. ; Rich, K.A. and de Kretser, D.M. (1979): Effects of experimental cryptorchidism on the ultrastructure and function of the Sertoli cell and paratubular tissue of the rat testis. Biol. Reprod., 21 : 823.
- Kierszenbaum, A.L.; Feldman, M. and Lea, O. (1980): Localization of androgen-binding protein in proliferating Sertoli-cells in culture. Proc. Natl. Acad. Sci. USA, 77: 5322.
- Kingsley-Smith, B.V. and Lacy, D. (1959): Residual bodies of seminiferous tubules of the rat. Nature, 184: 249.
- Kormano, M.; Koskimies, A.I.; and Hunter, R.L. (1971): The prescence of specific proteins, in the absence of many serum proteins in the rat seminiferous tubule fluid. Experientia, 27:1461.
- Koskimies, A.I.; Kormano, M.; and Alfthau, O. (1973): Proteins of the seminiferous tubule fluid in man-evidence for a blood-testis barrier. J. Reprod. Fertil. 32 : 79.

- Kotite, N.J.; Nayfeh, S.N. and French, F.S. (1978)
FSH and androgen regulation of Sertoli cell
function in the immature rat. Biol. Reprod;
18 : 65.
- Krebs, E.G.; Graves, D.J. and Fischer, E.H. (1959):
Factors affecting the activity of muscle
Phosphorylase b kinase. J. Biol. Chem.,
234 : 2867.
- Krebs, E.G., and Fischer, E.H. (1956): The phosphory-
lase b to a converting enzyme of a rabbit
skeletal muscle. Biochim. Biophys. Acta.,
20 : 150.
- Krimer, D.B. (1977): Ultrastructural and cytochemical
aspects of the nucleus of mouse Sertoli cells
Arch. Biol. 88 : 117.
- Krueger, P.M.; Hodgen, G.D. and Sherins, R.J. (1974):
New evidence for the role of the Sertoli
cell and spermatogonia in feedback control
of FSH secretion in male rats. Endocrinol.,
95 : 955.
- Lacy, D. (1960): Light and electron microscopy and
its use in the study of factors influencing
spermatogenesis in the rat. J. Royal Micro-
scop. Soc. 79 : 209.
- Lacy, D. (1962): Certain aspects of testis structure
and function. Brit. Med. Bull. 18 : 205.
- Lacy, D. and Pettitt, A.J. (1970): Sites of hormone

Morphology of normal and cryptorchid testes
In: The Undescended Testis, Fankalsrud, E.W.
and Mengel, W. (eds), London, Year Book
Medical Publishers, P. 59.

Hodgen, G.D. and Sherins, R.J. (1973): Enzymes as markers
of testicular growth and development in the
rat. Endocrinol. 93: 985.

Holmes, S.D.; Lipshultz, L.I. and Smith, R.G. (1984):
Regulation of Transferrin secretion by human
Sertoli cells cultured in the prescence or
absence of human peritubular cells. J.
Clin. Endocrinol. Metab., 59: 1058..

Howard, R.P. and Sniffen, R.C. (1950): Testicular defi-
ciency: Clinical and pathologic study. J.
clin. Endocrinol. Metab., 10:121.
Quoted by Bardin and Paulsen (1981).

Hsuch, A.J.W. ; Schreiber, J.R.; and Erickson, G.F.(1981):
Inhibitory effect of gonadotrophin releasing
hormone upon cultured testicular cells. Mol.
cell Endocrinol., 21 : 43.

Huggenvik, J. and Griswold, M.D. (1981): Retinol binding
protein in rat testicular cells. J. Reprod.
Fertil, 61: 403.

Huggins, C. and Moulder, P.V. (1945): Estrogen production
by Sertoli cell tumours of the testes. Cancer

production in the mammalian testis, and their significance in the control of male fertility. British Medical Bulletin, 26:87.

Levine, N. and Marsh, D.J. (1971): Micropuncture studies of the electrochemical aspects of fluid and electrolyte transport in individual seminiferous tubules, the epididymis and the vas-deferens in rats. J. Physiol. 213:557.

Lipshultz, L.I.,; Murthy, L. and Tindall, D.J. (1982): Characterization of human Sertoli cells in vitro. J. clin. Endocrinol. Metab., 55:228.

Louis, B.G. and Fritz, I.B. (1979): Follicle-stimulating hormone and testosterone independently increase the production of androgen binding protein by Sertoli cells in culture. Endocrinol, 104:454.

Lubarsch, O. (1896): Über des Vorkomen Kryptallinischer Und krystalloider Bildungen in den Zellen des menschlichen Hodens. Arch. Pathol. Anat. Physiol. 145:316.

Quoted by Tindall, D.J.; Rowley, D.R. and Lipshultz, L.I. (1983): Sertoli Cell structure and function in vivo and in vitro. In: Infertility in the Male, Lipshultz, L.I. and Howards, S.S. (eds). New York, Churchill Livingstone., p. 71.

Lu, C. and Steinberger, A. (1977): Gamma-glutamyl trans-peptidase activity in the developing rat testis. Biol. Reprod., 17:84.

Lyon, M.F.; Glenister, P.H. and Lamoreus, M.L. (1975): Normal spermatozoa from androgen resistant germ cells of chimeric mice and the role of androgen in spermatogenesis. Nature, 258:620.

Mather, J.P. ; Gunsalus, G.L.; Musto, N.A.; Cheng, C.Y.; Parvinen, M.; Wright, W.; Perez-Infant, V.; Margioris, A.; Liotta, A.; Becker, R.; Krieger, D.T. and Bardin, C.W. (1983): The hormonal and cellular control of Sertoli cell secretion. J. Steroid Biochem., 19:41.

Maximow, A.A. and Bloom, W. (1957): A textbook of Histology, Philadelphia, W.B. Saunders Co.
Quoted by Walsh, P.C. and Amelar, R.D. (1977): Embryology, anatomy and physiology of the male reproductive system. In: Male infertility, Amelar, R.D.; Dubin, L. and Walsh, P.C. (eds). Philadelphia, W.B. Saunders Company, P. 14.

McGinley, D.M.; Posalaky, Z.; Porvazink, M. and Russel, L. (1979): Gap junctions between Sertoli and germ cells of rat seminiferous tubules.

Tissue cell. 11 : 741.

- Means, A.R., and Vaitukaitis, J. (1972): Localization of follicle stimulating hormone and luteinizing hormone in the rat testis using immunological tests. *Endocrinol.*, 90:39.
- Means, A.R. and Tindall, D.J. (1975): FSH-induction of androgen binding protein in testis of Sertoli cell-only rats. In: *Hormonal Regulation of Spermatogenesis*, French, F.S.; Hansson, V.; Ritzen, E.M. and Nayfeh, S.N. (eds). New York. Plenum Press, P. 383.
- Means, A.R.; Fakunding, J.L.; Huckins, C.; Tindall, D.J. and Vitale, R. (1976): FSH, the Sertoli cell, and spermatogenesis. *Recent prog. Horm. Res.*, 32:477.
- Means, A.R.; Dedman, J.R.; Fakunding, J.L. and Tindall, D.J. (1978): Mechanism of action of FSH in the male rat. In: *Receptors and Hormone Action*, Birnbaumer, L. and O'Malley, B.W. (eds). New York, Academic Press, p. 363.
- Means, A.R.; Dedman, J.R.; Welsh, M.J. ; Marcum, M. and Brinkley, B.R. (1979): FSH and calcium as modulators of Sertoli cell differentiation and function. In: *Receptors and Reproductive Hormone Action*, Hamilton, T.; Clark, J.

and Sadler, W. (eds). New York, Raven Press, P. 207.

Means, A.R.; Dedman, J.R. and Tash, J.S. (1980): Regulation of the testis Sertoli cell by Follicle stimulating hormone. *Annu. Rev. Physiol.* 42:59.

Meyer, R.; Posalaky, Z. and McGinley, D. (1977): Inter cellular junction development in maturing rat seminiferous tubules. *J. Ultrastruct. Res.* 61:271.

Mita, M.; Borland, K.; Price, J.M.; and Hall, P.F. (1985): The influence of insulin and insulin-like growth factor-1 on hexose transport by Sertoli cells. *Endocrinol.* 116:987.

Moger, W.H. and Murphy, R.R. (1982): Reevaluation of effect of follicle stimulating hormone on the steroidogenic capacity of the testis. *Biol. Reprod.*, 26 : 422.

Moore, C.R. (1922): Cryptorchidism experimentally produced. *Anat. Rec.*, 24:383. Quoted by Hall, P.F.; Kew, D.; and Mita, M. (1985): The influence of temperature on the functions of cultured Sertoli cells. *Endocrinol.*, 116 : 1926.

Moore, C.R. and Chase, H.D. (1923): Heat application

and testicular degeneration. Anat. Rec., 26: 344. Quoted by Hall, P.F.; Kew, D.; and Mita, M. (1985): The influence of temperature on the functions of cultured Sertoli cells. Endocrinol., 116:1926.

Moore, C.R. and Quick, W.J. (1924): The scrotum as a temperature regulator for the testes. Am. J. Physiol., 68: 70. Quoted by Hall, P.F.; Kew, D.; and Mita, M. (1985): The influence of temperature on the functions of cultured Sertoli cells. Endocrinol., 116:1926.

Moore, C.R. (1944): Hormonal secretion by experimental cryptorchid testes. Yale J. Biol. Med., 17 : 203.

Morales, C. and Clermont, Y. (1982): Evolution of Sertoli cell processes invading the cytoplasm of rat spermatids. Anat. Rec. 203: 233.

Morse, M. J. and Whitmore, W.F. (1986): Neoplasms of the testis. In: Campbell's Urology, Cann, C. (ed), Philadelphia, Saunders Company, p. 1535.

Musto, N.A. and Bardin, C. (1974): FSH action and androgen binding protein in a rat with hereditary seminiferous tubular failure. Endocrinol. 94 : 108.

- Musto, N.A.; Santen, R.J. ; Huckins, C. and Bardin, C.W. (1978): A study of animals with an inherited disorder of seminiferous tubular and Leydig cell function. Biol. Reprod., 19 : 797.
- Nagano, T. (1966): Some observations on the fine structure of the Sertoli cells in the human testes. Z. Zellforsch. Mikroskop. Anat. 73:89. Quoted by Fawcett, D.W. (1975): Ultrastructure and function of the Sertoli cell. In: Handbook of Physiology, Endocrinology, Greep, R.O. and Astwood, E.B.(eds), Washington, American Physiological Society, p. 21.
- Nagano, T. and Suzuki, F. (1978): Cell to cell relationships in the seminiferous epithelium in the mouse embryo. Cell Tissue Res. 189:389.
- Neely, J.R. and Morgan, H.E. (1974): Relationship between carbohydrate and lipid metabolism and the energy balance of the heart. Annu. Rev. Physiol. 36: 413.
- Neumann, F. and Schenck, B. (1978): Influence of anti-androgens on Sertoli cell function and intratesticular androgen transport. International J. of Endocrinol., 1: 459.

- Nicander, L. (1963): Some ultrastructural features of mammalian Sertoli cells. J. Ultrastruct. Res. 8 : 190.
- Orth, J. and Christensen, A.K. (1977): Localisation of ^{125}I -labelled FSH in the testis of hypophysectomized rats by autoradiography at the light and electron microscope levels. Endocrinol. 101 : 262.
- Picard, J.Y; Tran, D. and Josso, N. (1978): Biosynthesis of labelled anti-Mullerian hormone by fetal testes: evidence for the glycoprotein nature of the hormone and for its disulfide-bonded structure. Mol and cell. Endocrinol. 58:1.
- Picon, R. (1969): Action du testicule foetal sur le development in vitro des canaux de Muller le rat. Arch. Micro. Morph. Exp. 58:1. Quoted by Price, J.M. (1979).
- Picon, R. (1970): Modifications, chez le rat, au cours du development du testicule, de son action inhibitrice sur les canaux de Muller in vitro. C.R. Acad. Sci. 272:2370. Quoted by Price, J.M. (1979).
- Price, J.M. (1979): The secretion of Mullerian inhibiting substance by cultured isolated Sertoli cells of the neonatal calf. Am. J. of Anatomy. 156 : 147.

- Purvis, K.; Clausen, O.D.F. , and Hansson, V. (1979):
LH contamination may explain FSH effects
on rat Leydig cells. *Reprod. Fert.*,56:657.
- Rabin, D. (1974): Binding of human FSH and its subunits
to rat testis. In: *Hormone Binding and
Target cell Activation in Testis*, Dufau,
M.L. and Means, A.R. (eds). New York,
Plenum Press, P. 193.
- Rambourg, A.; Clermont, Y. and Hermo, L. (1979): Three-
dimensional architecture of the golgi
apparatus in Sertoli cells of the rat. *Am.
J. Anat.* 154 : 455.
- Rich, K. and de Kretser, D.M. (1977): Effect of differ-
ent degrees of destruction of the rat semi-
niferous epithelium on the levels of serum
FSH and ABP. *Endocrinol.*,101: 593.
- Rich, K.A.; Bardin, C.W.; Gunsalus, G.L. and Mather,
J.P. (1983): Age-dependent pattern of and-
rogen-binding protein secretion from rat
Sertoli cells in primary culture. *Endocri-
nol.* 113 : 2284.
- Ritzen, E.M.; Hansson, V. and French, F.S. (1981): The
Sertoli cell. In: *The testis*, Burger, H.
and de Kretser, D.M(eds), New York, Raven.
Chapter 8. p. 59.

- Ross, M.H. and Reith, E.J. (1985): Male reproductive system. In: Histology A Text and Atlas, Ross, M.H. and Reith, E.J. (eds). New York J.B. Lippincott Company, p. 607.
- Rosvoll, R.V. and Woodart, J.R. (1968): Malignant Sertoli cell tumor of the testis. Cancer, 22 : 8.
- Russell, L.D.(1977): Desmosome-like junctions between Sertoli cells and germ cells in the rat testis. Am. J. Anat., 148 : 301.
- Russell, L.D. (1979): Further observations on tubulobulbar complexes formed by late spermatids and Sertoli cells in the rat testis. Anat. Rec. 194 : 213.
- Russell, L.D. (1980): Sertoli-germ cell interactions. Gamete Res. 3 : 179.
- Russell, L.D. (1981): Observations on the inter-relationships of Sertoli cells at the level of the blood-testis barrier. Am. J. Anat.155:259.
- Sairam, M.R.; Ranganathan, M.R. and Ramasharma, K. (1981): Binding of an inhibin like protein from bull seminal plasma to ovine pituitary membranes. Mol. cell Endocrinol 22:251.
- Sanborn, B.M.;Elkington, J.S.H. and Chowdhury, M. (1975):

Hormonal influences on the level of testicular androgen binding protein activity. *Endocrinol*, 96 : 304.

Sanborn, B.M.; Elkington, J.S.H. ; Steinberger, A.; Steinberger, E., and Meistrich, M.L. (1976): Androphilic protein in the testis In: *Regulatory Mechanism of Male Reproductive Physiology*, Spilman, C.H.; Lobol, T.J. and Kirton, K.T. (ed.) New York, Elsevier. p. 45.

Sanborn, B.M.; Steinberger, A. ; Tcholakian, R.K. and Steinberger, E. (1977): Direct measurement of androgen receptors in cultured Sertoli cells. *Steroids*, 29 : 493.

Sapsford, E.S. and Rae, C.A. (1968): Sertoli cell-spermatid relationship ultrastructural studies of the movements of mature spermatids into the lumen of the seminiferous tubule *J. Anat.* 103:214.

Sapsford, C.S.; Rae, C.A. and Cleland, K.W. (1969): Ultrastructural studies on maturing spermatids and on Sertoli cells in the bandicoot. *Australian J. Zool.* 17: 195.

Schulze, C. (1979): Giant nuclear bodies (Sphaerida) in

Sertoli cells of patients with testicular tumours. J. Ultrastruct. Res. 67 : 267.

Scott, R.S.; Burger, H. and Quigg, H. (1980): A simple and rapid in vitro assay for inhibin. Endocrinol. 107: 1536.

Seethalakshmi, L.; Steinberger, A. and Steinberger, E. (1984): Pituitary binding of ^3H -labelled Sertoli cell factor in vitro. Endocrinol. 115 : 1289.

Sertoli, E. (1865): De l'assistenza di particolari cellule ramificate nei canalicoli seminiferi dell' testicolo umano Morgagni, 7:31. Quoted by Fawcett, D.W. (1975): Ultrastructure and function of the Sertoli cell. In: Handbook of Physiology, Endocrinology, Greep, R.O. and Astwood, E.B. (eds), Washington, American Physiological Society, p.21.

Setchell, B.P.; Voglmayr, J.K. and Waites, G.M.H. (1969): A blood-testis barrier restricting passage from blood into rete testis fluid but not into lymph. J. Physiol. 200 : 73.

Setchell, E. (1970): Testicular blood supply, lymphatic drainage and secretion of fluid. In: The Testis, Johnson, A.D. and Comes, W.R. (eds). New York, Academic Press, p. 101.

Setchell, B.P. and Waites, G.M. H. (1975): The blood

testis barrier. In: Handbook of physiology, Section 7. Endocrinology, Greep, R.O. and Astwood, E.B. (eds). Baltimore, The Williams and Wilkins Co. , P. 143.

Setchell, B.P. (1978): The Mammalian Testis. Ithaca, Cornell University Press, P. 50.

Setchell, B.P. (1980): The functional significance of the blood testis barrier. J. Andrology, 1 : 3.

Sherins, R.J. and Hodgen, G.D. (1976): Testicular gamma glutamyl-transpeptidase. J. Reprod. Fertil., 48 : 191.

Skinner, M.K. and Griswold , M.D. (1980): Sertoli cells synthesize and secrete a transferrin-like protein. J. Biol. Chem. 255: 9523.

Skinner, M.K. and Griswold, M.D. (1982): Secretion of testicular transferrin by cultured Sertoli cells is regulated by hormones and retinoids. Biol. Reprod., 27 : 211.

Slaughter, G.R. and Means, A.R. (1982): Follicle-Stimulating hormone activation of glycogen phosphorylase in the Sertoli cell-enriched rat testis. Endocrinol , 113 : 1476.

Sniffen, R.C. and Howard, R.P. (1951): The testis. Absence of germ cells, sclerosing tubular degeneration, "Male climacteric." Arch. Pathol., 51:293.

Spangaro, S. (1902): Über die histologischen Veränderungen des Hodens, Nebenhodens, Und Samenleiters von Geburt an bis Zum Griesalter, Mit besondere Berücksichtigung der Hodenatrophie, des vorkommens von krystallen in Hoden. Anat. Rec., 18: 593. Quoted by Tindall, D.J.; Rowley, D.R. and Lipshultz, L.I. (1983): Sertoli cell structure and function in vivo and in vitro. In: Infertility in the Male, Lipshultz, L.I. and Howards, S.S. (eds), New York, Churchill Livingstone., P. 71.

Steinberger, E. (1971): Hormonal control of mammalian spermatogenesis. Physiol. Rev. 51 : 1.

Steinberger, E. and Steinberger, A. (1974): Hormonal Control of testicular function in mammals. In: Handbook of physiology; Knobil, W.H. and Sawyer H. (eds), Washington, American Physiological Society D.D. P. 325.

Steinberger, A.; Elkington, J.S.H.; Sanborn, B.M.; Steinberger, E.; Heindel, J.J. and Lindsey, J.N. (1975): Culture and F.S.H. responses of Sertoli cells isolated from sexually mature rat testis. In: Hormonal Regulation of Spermatogenesis, French, F.S.; Hansson, V.; Ritzen, E.M. and Nayfeh, S.N. (eds). New York, Plenum Press. p.399.

Steinberger, A., and Steinberger, E. (1976): Secretion of an FSH inhibiting factor by cultured Sertoli cells. *Endocrinol.* 99 : 918.

Steinberger, A. and Steinberger, E. (1977): The Sertoli cells. In: *The testis*, Johnson, A.A. and Comes, W.R. (eds), New York, Academic Press. Chapter 11.

Steinberger, E.; Tcholakian, R.K. and Steinberger, A. (1979): Steroidogenesis in testicular cells. *J. Steroid Biochem.*, 11: 185.

Steinberger, A. (1980): Factors affecting in vitro secretion of inhibin by isolated Sertoli cells. In: *Endocrinology*, Cuming, J.A.; Funder, J. W. and Mendelsohn, F.A.O. (eds), Canberra, Australian Academy of Sciences, P. 259.

Tabone, E.; Benahmed, M.; Reventos, J., and Salz, J.M. (1984): Interaction between immature porcine Leydig and Sertoli cells in Vitro. *cell Tissue Res.* 237 : 357.

Tash, J.S.; Dedman, J.R. and Means, A.R. (1979): Protein kinase inhibitor in Sertoli cell enriched rat testis. *J. Biol. chem.* 254 : 1241.

Tash, J.S.; Welsh, M.J. and Means, A.R. (1981): Regulation of protein kinase inhibitor by follicle-stimulating hormone in Sertoli cells in vitro. *Endocrinol.* , 108 : 427.

- Tcholakian, R.K. and Steinberger, A. (1979): In vitro metabolism of testosterone by cultured Sertoli cells and the effect of FSH. *Steroids*, 33:495.
- Tindall, D.J.; Schrader, W.I. and Means, A.R. (1974): The production of androgen binding protein by Sertoli cells. In: *Hormone Binding and Target cell Activation in Testis*, Dufau, M.L. and Means, A.R. (eds) New York, Plenum Press, p. 167.
- Tindall, D.J.; Vitale, R. and Means, A. R. (1975): Androgen binding protein as a biochemical marker of function of the blood-testis barrier. *Endocrinol.*, 97 : 636.
- Tindall, D.J. and Means, A.R. (1976): Concerning the hormonal regulation of androgen binding protein in rat testis. *Endocrinol.*, 99:809.
- Tindall, D.J.; Miller, D.A. and Means, A.R. (1977): Characterization of androgen receptors in Sertoli cell-enriched testis. *Endocrinol.*, 101 : 13.
- Tindall, D.J.; Means , C.R. and Means, A.R. (1978): Hormonal regulation of androgen binding protein in hypophysectomized rats. *Endocrinol.*, 103:589.
- Tindall, D.J.; Tash, J.S. and Means, A.R. (1981): Factors affecting Sertoli cell function in the testis. *Environ. Health perspect.* 38 : 5.

- Tindall, D.J.; Rowley, D.R. and Lipshultz, L.I. (1983):
Sertoli cell structure and function in vivo and
in vitro. In: Infertility in the Male,
Lipshultz, L.I. and Howards, S.S. (eds). New-
York, Churchill Livingstone, p. 88.
- Toyama, Y. (1976): Actin-like filaments in peritubular
myoid and Sertoli cells of the swine and mouse
testis. Anat. Rec. 186 : 477.
- Toyama, Y.; Obinata, T. and Holtzer, H. (1979): Crysta-
lloids of actin-like filaments in the Sertoli
cell of the swine testis. Anat. Rec. 195: 47.
- Tran, D. and Josso, N. (1982): Localization of anti-
Mullerian hormone in the rough endoplasmic re-
ticulum of the developing bovine Sertoli cell
using immunocytochemistry with a monoclonal
antibody. Endocrinol. 111:1562.
- Tuck, R.R. ; Setchell, B.P.; Waites, G.M.H. and Young,
J.A. (1970): The composition of fluid collected
by micropuncture and catheterization from the
seminiferous tubules and rete testes of rats.
Eur. J. physiol. 318 : 225.
- Tung, P.S.; Dorrington, J.H. and Fritz, I.B. (1975):
Structural changes included by follicle stimula-
ting hormone or dibutytyl cyclic AMP on presump-
tive Sertoli cells in culture. Proc. Natl. Acad.

Sci. USA, 72 : 1838.

Vanage, G.R.; Thakur, A.N.; Kadam, M.S. and Sheth, A.R.
(1980): Metabolic clearance rate of inhibin in
mature and immature male rats. Biol. Reprod.
23 : 606.

Vansickle , M.; Oberwetter, J.M.; Birnbaumer, L. and
Means, A. R. (1981): Developmental changes in
the hormonal regulation of rat testis Sertoli
cell adenylyl cyclase. Endocrinol. 109:1270.

Verhoeven, G.; Koninckx, P., and De Moor, P. (1982):
Androgen and progesterone production in cultured
interstitial cells derived from immature rat
testis. J. Steroid Biochem., 17 : 319.

Vernon, R.G., Kopec, B. and Fritz, I.B. (1974): Observat-
ions on the binding of androgens by rat testis
seminiferous tubules and testis extracts. Mol.
Cell Endocrinol. 1 : 167.

Vigier, B.; Tran, D.; du Mesnil du Buisson, F.; Heyman,
Y. and Josso, N. (1983): Use of monoclonal anti-
body techniques to study the ontogeny of bovine
anti-Mullerian hormone. J. Reprod. Fertil.
69 : 207.

Vigirsky, R.A.; Loriaux, D.L.; Lipsett, M.B. and Charm-
bach, A.C. (1975): Androgen binding proteins of
testis, epididymis and plasma in man and monkey,

J. Clin. Endocrinol . Metab. (In press.).

- Vilar, O. (1970): Histology of the human testis from neonatal to adolescence. In: The Human Testis, Rosenberg, E. and Paulsen, C.A. (eds). New York, Plenum Press. P : 95.
- Vitale, C.R.; Fawcett, D.W. and Dym, M. (1973): The normal development of the blood-testis barrier and the effect of clomiphene and estrogen treatment. Anat. Rec. 176 : 333.
- Waites, G.M.H. and Setchell, B.P. (1969): Some physiological aspects of the function of the testis. In: The Gonads, McKerns, K.W. (ed). New York, Appleton, P. 649.
- Waites, G.M.H. (1977): Fluid Secretion. In: The Testis, Johnson, A.D. and Lomes, W.R. (eds). New York, Academic Press, P. 91.
- Walsh, D.A.; Perkins, J.P.; Brostrom C.O.; Ho, E.S. and Krebs, E.G. (1971): Catalysis of the phosphorylase kinase activation reaction. J. Biol. Chem. 246 : 1968.
- Wartenberger, H. (1978): Human testicular development and the role of the mesonephros in the origin of a dual Sertoli cell system. Andrologia, 10:1.
- Weddington, S.C.; Hansson, V. and Ritzen, E.M. (1975):

Sertoli cell secretory function after hypophysectomy. Nature, 254 : 145.

Weddington, S.C. ; Hanssen, V. and Purvis, K. (1976):
Biphasic effect of testosterone propionate on
Sertoli cell secretory function. Mol. cell
Endocrinol. , 5 : 137.

Welsh, M.J. and Wiebe, J.P. (1975): Rat Sertoli cells a
rapid method for obtaining viable cells. Endo-
crinol. 96 : 618.

Welsh, M.J. and Wiebe, J.P. (1978): Sertoli cell capacity
to metabolize C 19 Steroids. Endocrinol 103:838.

Wright, W.W.; Parvinen, M.; Musto, N.A.; Gunsalus, G.L.;
Phillips, D.M.; Mather, J.P. and Bardin, C.W.
(1983): Identification of stage-specific proteins
synthesized by rat seminiferous tubules. Biol.
Reprod., 29 : 257.