

REFERENCES

REFERENCES

- 1- Bainbridge, D.R.; Whitlake, R. H. and Shephard, B.G. f. (1971): Balanitis xerotica obliterans and urinary obstruction. Br. J. of urology, 43:487.
- 2- Bharagava, R.K. and Thin, R. N.T. (1983): Subpreputial carriage of aerobic micro-organisms and balanitis. Br. J. Vener. dis. 59:131.
- 3- Brown, S.T. (1974): Tetracycline balanitis fixed drug reaction. JAMA. 277 No. 7:801.
- 4- Campus, G.V.; Ena, P. and Scuderi, N. (1984): Surgical treatment of balanitis xerotical obliterans. Plastic and reconstructive surgery. 73 No. 4:652.
- 5- Catterall, R.D. (1979): Balanitis. A short textbook of venereology. Second edition. The english language book society: 182.
- 6- Catterall, R.D. and Oates, J.K. (1979): Treatment of BXO with hydrocortisone injections. Br. J. vener. Dis. 38:75.
- 7- Chalmers, R. J. G.; Burton, P.A.; Bennett, R. and Goring, C. C. (1982): Lichen sclerosus et atrophicus - A distinctive and common cause of phimosis in boy, Br. J. dermatol. 107(Sup. 22) : 29.

- 8- Chakraborty, A.K. and Dutta, A. K. (1981): Herptic balanoposthitis. Indian Journal of dermatol Vol. 26 No. 1 : 15.
- 9- Chang, T. W. ; fiumara, N. J. and Weinstein, L. (1974): Genital herpes. JAMA. 229 : 544.
- 10- Cooke, R.A.; Rodrigue, R. B. (1964): Amoebic balanitis. Med. J. aust. 1 : 114. Cited in Rodrigue (1978)
- 11- Csonka, G.W. ; Rosedale, N. and Walkden, L. (1971): Balanitis due to fixed drug eruption associated with tetracycline therapy. Br. J. vener. Dis. 47:42.
- 12- Dodds, P.R. and Chi, T. N. (1985): Balanitis as fixed drug eruption to tetracyclines. The journal of urology. 133:1044.
- 13- Domonkos, A. N.; Arnold, H. L. and Odom, R. B. (1982): Balanitis plasma cellularis (Zoon). Andrews diseases of the skin. Seven Edetion. W. B. sau-nders company philadelphia : 833.
- 14- Dupré, A. (1976): Bull. Soc. fr. Derm. syph. 83:62. Cited in Wilkinson et al., (1979).

- 15- Farid, M. (1981): Balanitis. Sexually transmitted diseases. First Edition. AinShams University presspp. 557.
- 16- Follmann, J. (1931) orv. Hetil .,75 : 680 cited in Lejman and Starzycki, (1975).
- 17- Harkness, A. H. and Harber, H. (1960): Constricting meatopathy of the male urethra. Acta dermatovener (Stokh.) 3 : 1002.
- 18- Herschorn, S. and Colapinto, V. (1979): Balanitis xerotica obliterans involving anterior urethra. Urology 14,6 : 592.
- 19- Kanerva, L. ; Kousa, M. ; Niemi, K. M.; Lassus, A.; Juvakoski, T. and Lauharanta, J. (1982): Ultra-hisopathology of balanitis circinata. Br. J. Vener. Dis. 58 : 188.
- 20- Kinghorn, G. R.; Jones, B. M.; Chowdhury, f. H. and Geary, I. (1982). Balanoposthitis Associated with gardnerella vaginalis infection in men. Br. J. vener. Dis. 58 : 127.
- 21- Lay mon, C. W. and frecman, C. (1944) cited in campus et al., (1984).

- 22- Lejman, K. and Starzycki, Z. (1975): Syphilitic balanitis of follmann developing after the appearance of the primary chancre. Br. J. vener. Dis. 51 : 138.
- 23- Leonforte, J. f. (1981): Balanitis Circumscripta plasmacellularis. Acta dermato venereal (Stockholom) 62(4) : 352.
- 24- Lever, W. f. and Schaumburg-Lever, G. (1983): Balanitis circumscripta plasmocellularis. Histopathology of skin- 6 edition. J. B. Lippincott company Philadelphia. PP:498.
- 25- Lortat. Jacob, E. and Civatte, J. (1961): Balanitis pseudoepitheliomatous Keratosique et micaceé. Bull soc. fr. Dermatol Sphilol 68 : 164 Cited in Read and Abell (1981).
- 26- Mann, P. R. and Cowan, N. A. (1973). Ultrastructural changes in four cases of lichen sclerosus et atrophicus. Br. dermatol 89 : 223 cited in Lever and Schaumburg-Lever (1983).
- 27- Mann, R. J. and Allen. B. R. (1982): Keratotic and micaceous pseudoepitheliomatous balanitis. Br. J. of Urology 54 : 431.

- 28- Masfari, A. N.; Kinghorn, G.R. and Duerden ,B. I.
(1983): Anaerobes in genitourinary infections
in men. Br. J. vener. Dis. 59:255.
- 29- Masterton, G.; Sengupta, S. M. and Schofield (1975):
Natamycin in genital candidosis in men. Br. j.
vener. Dis. 51 : 210.
- 30- McKay, D. L.; Fugua, J. f. and Weinberg, A. G. (1975):
Balanitis xerotica obliterans in children. The
Journal of Urology, 114 : 773.
- 31- Pasieczny, T. A. (1977): The treatment of balanitis
xerotica obliterans with testosterone propionate
ointment. Acta dermatovener. (Stockholm) 57:275.
- 32- Pasricha, J. S. (1979): Drug causing fixed eruptions.
Br. J. of dermatology. 100 : 183.
- 33- Pattman, R. S. (1983): Two unusual manifestations of
infection with staphylococcus aureus presenting
to a clinic for sexually transmitted disease.
Br. J. of clinical practice : 228.
- 34- Petrozzi, J. W. and Erlich, A. (1977): Pseudomonal
balanitis. Arch dermatol. 113 : 952.

- 35- Peutherer, J. F.; Smith, I.W. and Robertson, D. H.
H. (1979): Necrotising balanitis due to a generalised primary infection with herpes simplex virus type 2. Br. J. of vener. Dis. 55 : 48.
- 36- Powers; R. D.; Rein, M. f. and Hayden, F. G. (1982):
Necrotizing balanitis due to herpes simplex type I. JAMA. 248 No. 2 : 215.
- 37- Read, S. I. and Abell, E. (1981): pseudoepitheliomatous, keratotic, and micaceous balanitis. Arch dermatol. 117 : 435.
- 38- Rodrigue, R. B. (1978): Amoebic balanitis. JAMA. 239, No. 2 : L09.
- 39- Rosemberg, S. K. and Jacobs, H. (1982): Continuous wave carbon dioxide treatment of balanitis xerotica obliterans. Urology XIX N. 5 : 539.
- 40- Rosemberg, S. K. (1985) : Carbon dioxide laser treatment of external genital lesions. Urology. Volume XXV, N. 6 : 555.
- 41- Schinella, R. and Miranda, D. (1974) : Posthitis xerotica obliterans in circumcision specimens. Urology. 3 : 348.

- 42- Schuermann, H. (1960): Plasmocytosis circumorificiolis.
Dt. Zaharzti. Z. 15 : 601 Cited in Toonstra,
1986.
- 43- Sonnex, T. S. ; Dawber, R. P. R.; Ryan, T. j. and
Ralfs, I. G. (1982): Zoon's (plasma - cell)
balanitis treatment by circumcision. Br. J.
of dermatol, 106 : 585.
- 44- Souteyrand, P. ; Wong, E. and Macdonald, D. M. (1981):
Zoon's balanitis (balanitis circumscripta plasma
cellularis). Br. J. of dermatol, 105 : 195.
- 45- Steigleder, G. K. and Raab, W. P. (1961) : Lichen
Sclerosus et atrophicus. Arch dermatol 84 :
219.
- 46- Stewart, W. D.; Danto, J. L. and Maddin, S. (1974):
Dermatology, diagnosis and treatment of cutaneous
disorder, 3rd Edition : 350. C. V. Mosby Company,
St. Louis. Cited in Sonnex et al., (1982).
- 47- Stuhmer, A. (1928) : Balanitis xerotica obliterans
(post-operationem) und ihre Beziehungen Zur
"Kraurosis glandis et praeputii penis. Arch.
f. dermat. Und Syph. 156 : 613. Cited in Hersch-
orn and Colapinto, (1979).

- 48- Toonstra, J. and Wichen, D. f. (1986) : immunohistochemical characterization of plasma cell in Zoon's balanoposthitis and (pre) malignant skin lesions. *Dermatologica* 172 : 77.
- 49- Wallin, J. and Forsgren, A. (1975) : Group B streptococci in venereal disease clinic patients. *Br. J. vener. Dis.* 51 : 401.
- 50- Waugh, M. A.; Evans, E. G. V.; Nayyar, K. C. and Fong, R. (1978). Clotrimazole (Canasten) in the treatment of candidal balanitis in men. *Br. J. of vener. Dis.* 54 : 184.
- 51- Weitzner, S. (1972) : posthitis xerotica obliterans in a 12 years old boy. *Amer. J. Dis. child.*, 123 : 68. Cited in McKay et al., (1975).
- 52- Wilkinson, D. S. (1979). Diseases of umbilical, perianal and genital regions. In textbook of dermatology. Edited by Rook, A; Wilkinson, D. S. and Ebling, f. j. G. page 1952. Blackwell Scintific publication Oxford.
- 53- Zoon, J. J. (1950) : Balanitis circumscripta chronica met plasma cellen-infiltraat. *Ned. Tijdschr. Geneesk.* 94 : 1528. Cited in Toonstra (1986).