


A decorative rectangular border with a dark, textured outer line and a white inner line. Four stylized floral motifs are placed at the corners: top-left, top-right, bottom-left, and bottom-right. Each flower has multiple pointed petals and a dark center.

References

REFERENCES

- Abd El-Kader G.; Abd El-Fataha F.; Nosair E. et al. (1996):*
Granulocyte macrophage colony stimulating factor in cord blood.
Thesis, Zagazig University.
- Abdul-Karim R. and Assali N. (1982):*
Pressor response to angiotensin in pregnant and non-pregnant women.
Am. J. Obstet. Gynecol., 182:246.
- Aglietta M.; Pasquino P.; Sanavio F. et al. (1993):*
GM-CSF and IL-3: Target cells and kinetics of response in vivo.
Stem cells-Dayt., 2:83-7.
- Akintorin S.M; Kamat M; Pildes R.S; et al.(1997):*
A prospective randomized trial of feeding methods in very low birth weight infants.
Pediatrics, 100: 234-9.
- Al-Banna E.; El-Dere S.; Mahrous S. et al.(2002):*
Granulocyte macrophage colony stimulating factor in the cord blood of premature neonates born to mothers with pregnancy induced hypertension.
Thesis, Zagazig University.
- Alderson M.; Armitage R.; Tough T. et al. (1993):*
CD40 expression by human monocytes: Regulation by cytokines and activation of monocytes by the ligand for cd40.
J. Exp. Med., 179:669-674.
- Alexander G.; De Caunes F.; Hulsey T. et al.(1992):*
Validity of postnatal assessments of gestational age: a comparison of the method of Ballard et al. and early ultrasonography.
Am. J. Obstet. Gynecol., 166:891-895.

Aman M.J.; Stckderher K.; Thews A. et al.(1996):

Regulation of immunomodulatory functions by granulocyte-macrophage colony factor and granulocyte-colony stimulating factor in vivo.

Ann.Hematol. 73:231-238.

American College of Obstetricians and Gynecologists (ACOG) (2001):

Chronic hypertension in pregnancy.

Obstet. Gynecol., 98:177-185.

Amiel-Tison C. (1968):

Neurological evaluation of maturity of newborn infants.

Arch. Dis. Child.,43:89-93.

Asuncion M; Silvestre A; Morbach C.A; et al.(1996):

A prospective randomized trial comparing continuous versus intermittent feeding methods in very low birth weight neonates.

J. Pediatr. 128:748.

Avery M. and First L.R. (1994):

Disorders of respiration ,hematology and neurology in neonates:

In Pediatric Medicine (2nd ed), Waverly company, New York.
pp:199-209.

Bagby M. and Segal L. (1995):

Growth factors and the control of hematopoiesis in Hematology
basic principles and practice, 2nd ed., by Hoffman R., p. 207.

Bailie K.; Irvine A.; Bridges J. et al.(1994):

Granulocyte and granulocyte-macrophage colony stimulating factors in cord and maternal serum at delivery.

Pediatr. Res.35:164.

Baldwin G.C.; Dipersio J.; Kaufman S. et al. (1987):

Characterization of human GM-CSF receptors on non hematopoietic cells.

Blood, 70: Suppl. 1: 519.

Baldwin, G.C. (1992):

The biology of granulocyte-macrophage-colony stimulating factor: Effects on hematopoietic and non hematopoietic cells.
Dev. Biol. 151, 352-367.

Baley J.E. and Fanaroff A.A. (1992):

Neonatal infection. In: Sinclair, J. , Bracken, M. eds. Effective care of the newborn. Oxford, Oxford University press. P 454-506

Ballard J.L.; Kizmaier K.; and Driver M. (1979):

A simplified score for assessment of fetal maturation of newly born infant.
J. Pediatr. 93:769.

Ballard J.L.; Khoury J.C.; Wedig K. et al. (1991):

New Ballard Score, expanded to include extremely premature infants.
J. Pediatr. 95:120.

Begley G.G.; Micola N. and Metcalf D. (1988):

Proliferation of human promyelocytes and myelocytes after a single pulse stimulation by purified GM-CSF and G-CSF.
Blood, 71: 640.

Beissert S.; Hosoi J.; Grabbe S. et al. (1995):

IL-10 inhibits tumor antigen presentation by epidermal antigen-presenting cells.
J. Immunol. 145:1280-1286.

Bertil E.; and Michael D. (1994):

Hemostatic disorders: In Schaffer and Avery (2nd ed) the newborn disease. W.B. Saunders company .Philadelphia . pp:777.

Bifano E.M. and Ehrenkran Z. (1995):

Perinatal hematology.
Clin. Perinatol., 23: 3.

Bilgin K.; Yaramis A.; Haspolat K. et al. (2001):

A randoized trial of GM-CSF in neonates with sepsis and neutropenia.

Pediatrics, 107:36-41.

Blanchard D.K.; Michelini N. and Norris M.B. (1991):

Production of granulocyte macrophage colony-stimulating factor by large granular lymphocytes stimulated with candida albicans: Role in activation of human neutrophil function.

Blood, 77: 2259.

Blaymore B.J.; Ferguson A.E.; Morales Y. et al. (1997):

Breast feeding in infants who were extremely low birth weight.

Pediatrics., 100: 268-71.

Bober L.A.; Grace M.J.; Pugliese-Silvo C. et al. (1995):

The effect of GM-CSF and G-CSF on human neutrophil function.

Immunopharmacol., 29: 111-119.

Bracho F.; Goldman S.; and Cairo M.S.; (1998):

Potential use of granulocyte colony stimulating factor and granulocyte macrophage colony stimulating factor in neonates

Curr. Opin. Hematol. 5:215-220.

Brazy J.E.; Grimm J.K. and Little V.A.(1982):

Neonatal manifestations of sever maternal hypertension occurring before the thirty-sixth week of pregnancy.

J. Pediatr. 100: 265-271.

Broudy V.C. (1997):

Stem cell factor and hematopoiesis.

Blood, 90: 1345-1364.

Brugnara C. and Platt O.S. (1998):

The neonatal erythrocyte and its disorders. In: Nathan DG, Orkin SH (eds): Nathan and Oski's Hematology of infancy and childhood. 4th ed. Philadelphia, WB Saunders, PP: 17-52.

Burgess A.W.; Begely C.G.; Johnson G.R. et al. (1987) :

Purification and properties of bacterially synthesized human granulocyte macrophage colony stimulating factor.
Blood, 69: 43.

Cadnapaphornchai M. and Faix R.G.(1992):

Increased nosocomial infection in neutropenic low birth weight (2000 grams or less) infants of hypertensive mothers.
J. Pediatr. 121: 956-961.

Cairo M.S. (1989):

Neonatal neutrophil host defense: Prospects for immunologic enhancement during neonatal sepsis.
Am. J. Dis. Child. 143:40-46.

Cairo M.S.; Suen Y.; Knoppel E. et al. (1991):

Decreased stimulated GM-CSF production and GM-CSF gene expression in human preterm newborns compared with adults.
Pediatr. Res., 30: 362-7.

Cairo M.S.; Suen Y.; Knoppel E. et al. (1992):

Decreased G-CSF and IL-3 production and gene expression from mononuclear cells of newborn infants.
Pediatr. Res., 31:574-8.

Cairo M.S.; Christensen R.; Sender Y. et al. (1995):

Results of phase I/II of recombinant human granulocyte macrophage colony stimulating factor in very low birth weight neonates.
Blood, 86: 2509-2515.

Calhoun D. and Christensen R.D. (1995):

Commonly asked questions regarding infections in neonates.
Pediatr. Ann., 24:583-91.

Calhoun D.; Lunoe M.; Du Y. et al.(2000a):

Granulocyte colony stimulating factor (G-CSF) is present in human milk and its receptor (G-CSF-R) is present in human fetal intestine.
Pediatrics, 105:E7.

Calhoun D.; Lunoe M.; Du Y. et al.(2000b):

Granulocyte colony stimulating factor (G-CSF) serum and urine concentration in neutropenic neonates before and following the intravenous administration of recombinant G-CSF.

Pediatrics, 105:392.

Cannistra S.; Groshek P.; Garlick R. et al. (1990):

Regulation of surface expression of the granulocyte macrophage colony stimulating factor in normal human myeloid cells.

Proc. Natl. Acad. Sci., USA; 87: 93.

Caplen M.S.; Sun X. and Hsueh W.(1990):

Role of platelet-activating factor and tumor necrosis factor alpha in neonatal necrotizing enterocolitis.

J. Pediat. Gastroenterol. Nutr. 11:317-322.

Carr R. and Modi N. (1997):

Hemopoietic colony stimulating factors for preterm neonates.

Arch. Dis. Child. Fetal Neonatal. 76: 128-133.

Carrolli G.; Duley L. and Belzan V. (1994):

Calcium supplementations during pregnancy: A systematic review of randomized controlled trials.

Br. J. Obstet.Gynecol.;101:753-758.

Cebon J.; Nicola N.; Ward M.; et al.(1990):

Granulocyte-macrophage colony-stimulating factor from human lymphocytes. The effect of glycosylation on receptor binding and biological activity.

J. Biol. Chem.,265:4483.

Charles L.; Anderson J. and Stewart I.E. (1998):

Retinopathy of prematurity: In Cloherty J and Stark A (Eds), Manual of Neonatal Care 4th ed., Philadelphia, Lippincott-Raven, P: 643-648.

Chatson K.; Fant M.E. and Cloherty J.(1998):

Temperature control: In Cloherty J. and Stark A. (ed.) Manual of Neonatal Care.4th.ed.Lippincott-Raven Philadelphia p-p 139.

Chesly L.C.; Annitto J.E. and Cosgrove R. (1986):

The familial factor in toxemia of pregnancy.
J. Obstet. Gynecol.; 32:203-11.

Chesly L.C. and Cooper D.W. (1986):

Genetics of hypertension in pregnancy: possible single gene control of preeclampsia and eclampsia in the descendants of eclamptic women.
Br. J. Obstet. Gynaecol. ; 93: 898-908.

Chirico G.; Ciardelli L. and Cecchi P. et al.(1997):

Serum concentration of granulocyte colony stimulating factor in term and preterm infants.
Eur. J. Pediatr 156:269-71.

Christensen P.D. (1989):

Neutrophil kinetics in the fetus and neonate.
Am. J. Pediatr. Hematol. 11: 215-223.

Christiane M.; Witte J.; Hildmenn A. et al. (1999) :

Neonatal screening for hearing disorders in infants at risk : incidence, risk factors, and follow up.
Pediatrics, 104: 900-904.

Christin M.F. (2001):

Granulocyte colony stimulating factors: how different are they?
How to make a decision?
Anticancer- Drugs., 12(3): 185-91.

Christopher W.G.; Redman M.D.; Gavin P. et al. (1996):

Preeclampsia :An excessive maternal inflammatory response to pregnancy.
Am. Obstet. Gyn., 180: 449-5060.

Christou H.A. and Rowitch D.W. (1998):

Anemia in neonates: In Cloherty, J. and Stark, A. eds. Manual of Neonatal Care, 4th ed. Lippincott-Raven, Philadelphia. PP:453-640.

Ciotti R.; Rosti V.; Lucotti C. et al. (1995):

Treatment with hematopoietic growth factors in myelodysplastic syndromes.
Br. J. Hematol. 91: 773.

Cockell A.P. and Poston L. (1997):

Flow mediated vasodilatation is enhanced in normal pregnancy but reduced in preeclampsia.
Hypertension; 30:247-251.

Cornblath M. (1991):

Disorders of carbohydrate metabolism in infancy (3rd ed.).
Blackwell Scientific, Carbridge, MA.

Cornblath M. (1993):

Hypoglycemia in the neonate.
J. Pediatr. Endocrinol., 6: 113.

Coskun M.; Kardelen F.; Oygur N.; et al. (1997):

Granulocyte and granulocyte macrophage colony stimulating factors, their receptors and interleuken-3 in newborns.
Turk. J. Pediatr., 39: 295-301.

Cosmi E.V.; Torre R.L. and Oiazzi J. (1996):

New treatment of neonatal RDS. 17th international symposium of neonatal intensive care, Sanremoo Italy .

Couser R. J.; Ferrara T.B. and Wright G. (1996):

Prophylactic indomethacin therapy in the first 24hr. of life in the prevention of patent ductus arteriosus (PDA).
J. Pediatr. 128:631-7.

Cowett R.M. (1992):

Hypoglycemia and hyperglycemia in the newborn. In: Polin R and Fox W (Eds.), fetal and neonatal Physiology (1st ed.) Saunders Philadelphia, P. 406.

Creasy R.K. (1994):

Preterm labor and delivery. In: Creasy RK, Resmik R (editors). Maternal-fetal medicine, principles and practice (2nd ed.). WB. Saunders, Philadelphia, PP: 494-520.

Cunningham F. G.; Fernandez O. and Hernandez C. (1995):

Blindness associated with preeclampsia and eclampsia. Am. J. Obstet. Gynecol.; 172:1292-8.

Cunningham F. G.; Macdonald C. and Gant N.F.(1997):

Hypertensive disorders in pregnancy: In Williams Obstetrics 20th ed. Prentice-Hall International, New York p-p:693-744.

Danis V.A.; Frank G.M.; Rathjen D.A. et al.(1991):

Effects of granulocyte macrophage- colony stimulating factor (GM-CSF), IL-2, interferon-gamma (IFN-gamma), tumor necrosis factor-alpha(TNF-alpha)and IL-6 on the production of IL-1 and TNF-alpha by human monocytes. Clin. Exp. Immunol. 85:143-150.

David A.P.; Stepron A.; Mark F. et al. (1999):

Cranial sonography in very low birth weight infants: Do all need to be screened. Clin. Ped. ;503.

David D. (1999):

Left to right shunt lesion .
Pediatr. Clin. of North Am. 46:355-68.

DeGroot C.J.; Davidge S.T. and Friedman S. (1995):

Plasma from preeclamptic women increases human endothelial cell prostacyclin production without changes in cellular enzymatic activity or mass.
Am. J. Obstet. and Gynecol., 172, 976-85.

Demerti G.; Zenzic B.; Rheinwald J. et al.(1989):

Expression of colony stimulating factor genes by normal human mesothelial cells and malignant mesothelioma cell lines in vitro. Blood, 74: 940.

Demetri G. and Griffin J.(1991):

Granulocyte-colony-stimulating factor and its receptor.
J. Am. Soc. Hematol.78:2791.

Denzlinger W.; Gerhartz H.T. Sagebiel S.P. et al.(1993):

Differential activation of the endogenous leukotriene biosynthesis
by two different preparation of granulocyte macrophage- colony
stimulating factor in healthy volunteers.
Blood, 81: 2007-13.

Deresinsk S.D. and Kemper C.A. (1998):

The potential role of GM-CSF and G-CSF in infectious diseases.
Infect. Med., 15(2): 856-870.

Dexter T.M. and Testa N.G.(1993):

Hematopoietic growth factors: Review of Biology and clinical
potential . Macclesfield: Gardiner-Caldwell communications.

Diane D.M.; Miltovk and Thomas C. (1999) :

Risk factor for chronic lung disease in surfactant era : A north
carolina population based study of VLBW infants.
Pediatrics, 104. 1345-54.

Dipersio J.; Hedvat C.; Ford C.F. et al., (1991) :

Characterization of the soluble human granulocyte-macrophage
colony stimulating factor receptor complex.
J. Biol. Chem. 266: 279.

Doron M.W.; Makhlof R.A.; Katz V.L.; et al. (1994):

Increased incidence of sepsis at birth in neutropenic infants of
mothers with preeclampsia.
J. Pediatr. 125: 452-458.

Driscoll J.M. (1987):

Gestational age assessment : In Fanaroff A. and Martin J. (eds.),
Neonatal -Perinatal Medicine 4th ed. Mosby, St. Louis. P. 122.

Dubois C.; Schlageter M.H. and Gentile A. (1994):

Hematopoietic growth factor expression and ATRA sensitivity in acute promyelocytic blast cells.

Blood, 83(11): 3364.

Dubowitz L.M.; Dubowitz V. and Goldberg C.(1977):

Clinical assessment of gestational age in the newborn infant.

J. Pediatr., 77:1-10.

Egyptian National Neonatal Care Program (ENNCP) (2001):

Neonatal care: protocols for physicians (1st ed.), Egyptian Ministry of Health and Population.

El Beshlawy A. (1993):

Recombinant human erythropoietin in the anemia of prematurity.

Annual Scientific Meeting of the Egyptian Society of Hematology
Cairo University.

Eldaher S. and Chevalier R.L. (1990):

Special needs of the newborn infant in fluid therapy :

Pediatr. Clin. North. Am. 37(2):323.

Emersen S.G.; Yang Y.G.; Clark S.C. et al. (1989):

Recombinant granulocyte macrophage- colony stimulating factor and interleukin-3 have overlapping but distinct hematopoietic activities.

J. Clin. Invest., 82: 1282.

Emerson S.G. (1998) :

Hematopoiesis : In Fred. J. (ed). Hematologic Pathophysiology (1st ed.). Lippincott. Raven. Philadelphia. Pp. 7-11.

Engle W.D. and Rosenfield C.R. (1984):

Neutropenia in high risk neonates.

J. Pediatr. 105: 982-6.

Eregie C.O.(1993):

Determination of maturity at birth: further observations on a maturity scoring system for head circumference and mid-arm circumference.

East Afr. Med. J. 70:48-50.

Evans N. (1993):

Diagnosis of patent ductus arteriosus.

Arch. Dis. Child., 68:58-63.

Fanaroff A.A.; Wright L.L.; Stevenson D.K. et al.(1995):

Very low birth weight outcomes of National Institute of Child Health and Human Development. Neonatal Research Network, May 1991 through December 1992.

Am. J. Obstet. Gynecol., 173:1423.

Farr V.; Mitchell R.G.; Neligan G. et al.(1966):

The definition of some external characteristics used in the assessment of gestational age in the newborn infant.

Dev. Med. Child Neurol., 8: 507.

Feinberg B.B. and Repake J.K. (1998):

Preeclampsia and related conditions: In Cloherty, J. Stark, A. eds. Manual of Neonatal Care 4th ed., Lippincott-Raven, Philadelphia. pp:26-29.

Fibbe W.E.; Kuck P.M.C.; Duinkerken N. et al. (1988):

Factors influencing release of granulocyte macrophage colony stimulating factor activity from human mononuclear phagocytes.

Eur. J. Haematol., 41:352.

Fielder H.G. Suci E.; Wittlief C. et al. (1990):

Mechanisms of growth factor expression in acute myeloid leukemia.

Leukemia, 4: 459.

Finnstorm O.(1971):

Studies on maturity in newborn infant: Postnatal radiological examination of epiphyseal centers.

Neurpediatr., 3:119-128.

Finnstrom O.; Gothefors L. and Zetterlund B.(1975a):

Studies on maturity in newborn infants. VII. Fetal hemoglobin.
Acta. Pediatr. Scand. 64:404-408.

Finnstorn O.; Karlsson B. and Zetterlund B.(1975b):

Studies on maturity in newborn infants. VIII. Alpha-foetoprotein
and alburnin.
Acta. Pediatr. Scand. 64:409-412.

Fletcher M.A. (1997):

Assessment of gestational age: In Physical Diagnosis in
Neonatology, 1st ed., Lippincott-Raven, Philadelphia, P: 55-66.

Fraser S.H and Tudehope D.I. (1996):

Neonatal neutropenia and thrombocytopenia following maternal
hypertension.
J. Pediatr. Child Health., 32: 31-34.

Forestier F. Daffos F.; Catherine N. et al. (1991):

Developmental hematopoiesis in normal human fetal blood.
Blood, 77: 2360.

Fuke A.; Berner R.; Traichel B. et al. (2000):

Frequency, natural course and outcome of neonatal neutropenia.
Pediatrics, 106: 45-51.

Gabrilove J. (2000):

Novel cytokine strategies: improving the treatment hematological
and malignant diseases. Abstracts of the 42 Annual Meeting of
the American Society of Hematology, California.

Garland J.; Buck R. and Leviton A.(1995):

Effect of maternal glucocorticoid exposure on risk of severe
intraventricular hemorrhage in surfactant-treated preterm infants.
J. Pediatr., 126:273.

Gasson J. (1991):

Molecular physiology of GM-CSF.
Blood, 77: 1131.

- Gerhartz H.H.; Marcus R. and Delmer A. (1990) :**
Randomized phase II study with GM-CSF and low dose ARAC in patients with high risk myelodysplastic syndrome.
Blood, 76 (Supplement 1) : 337a.
- Gertner J. (1990):**
Disorders of calcium and phosphorus hemostasis .
Pediatr. Clin. North. Am. 37(6): 1441.
- Gessler P.; Kirchmann N.; Engel R. et al. (1993):**
Serum concentrations of granulocyte colony stimulating factor in healthy term and preterm neonates and in those with various diseases including bacterial sepsis.
Blood, 3177-3182.
- Goldman D.J.; Goldman S.L. and Hirata T. (1996) :**
Prematurity: In Jackson P. and Vesseny J. (ed.) Primary care of the child with a chronic condition. 2nd ed. Mosby, New York, PP: 650-669.
- Goldman S.; Ellis R.; Dhar V. et al. (1998):**
Rationale and potential use of cytokines in the prevention and treatment of neonatal sepsis.
Clin. Perinatol., 25:699-710.
- Gomella T. L. (Ed.) (1999):**
Neonatology, 4th. ed. Appelton and Lange, Stamford.
- Gotoff S.P. (2000) :**
Infections in the neonatal infant : In Behrman, Kliegman and Jenson (eds.). Textbook of pediatrics 16th ed., W.B. Saunders. Philadelphia. Pp. 538-552.
- Graeber J.E. and Schwartz T.L. (1993):**
Retinopathy of prematurity: diagnosis, treatment and outcome. In: Witter, F. and Keith L. (eds): Textbook of Prematurity, Little Brown, New York, 212-217.

Gray P.H.; Rodwell R. and Rogers Y.(1996):

Risk of neutropenia and nosocomial infection associated with maternal hypertension.
Pediatr. Res.40: 351A.

Green D.M.; Hendon B. and Minouni F.B.(1995):

Nucleated erythrocytes and intraventricular hemorrhage in preterm neonates.
Pediatrics, 96:475

Hamilton J. (1993):

“ CSF, Cytokines and monocytes-macrophages. Some controversies”.
Immunol. Today; 14:18-24.

Hammerman C. (1995):

Patent ductus arteriosus.
Clin. Perinatol., 22:457.

Hansen T. and Cobet A.(1991):

Hyaline membrane disease, Bronco-pulmonary dysplasia, In: Taeusch, H; Ballard, R; Avery, M;(eds.) Disease of the Newborn, 6th ed . W. B. Saunders Philadelphia. Pp 498-519.

Hansens M.; Keirse J.; Spitz B. et al.(1991):

Angiotensin II levels in hypertensive and normotensive pregnancies.
Br. J. Obstet. Gynaecol. ;98:155-159.

Hercus T.R.; Bagley C.G.; Cambarera B. et al. (1994):

“Specific human GM-CSF antagonists”.
Nalt. Acad.Sci. USA; 13:5838-42.

Hittner H.M.; Hirsch N.J. and Rudolph A.J.(1977):

Assessment of gestational age by examination of the anterior vascular capsule of the lens.
J. Pediatr. 91:455.

Hoglund M.; Simonsson B.; Oberg G. et al. (1994):

The effect of rhGM-CSF on neutrophil and esinophil regeneration after ABMT as monitored by circulating levels of granules proteins.

Br. J. Hematol., 86: 709.

Hume R.(1992):

Neonatal metabolic disorders: In Campbel A. and McIntosh N. (eds.), Forfar and Arnelils Textbook of Pediatrics, 4th ed. Churchill Livingstone, London p-p322.

Huttner K.M. (1998):

Hypocalcemia, hypercalcemia and metabolic bone disease of prematurity: In Cloherty, J. and Stark, A. (eds.). Manual of Neonatal Care (4th ed.) Lippincott-Raven, Philadelphia., P-P: 553-564.

John H.(1991):

Intracranial hemorrhage In: Taeusch, H; Ballard, A; and Avery, M; (eds.). Diseases of the newborn ,6th ed. W. B. Saunders. Philadelphia. pp:422-425.

Kalkon S.C. (1997):

Metabolic and endocrine disorder. In: Fanaroff A and Martin P (eds.), Neonatal Perinatal Medicine, 6th ed. St. Louis, Mosby, P. 1439.

Kaushanksy K.; O'Hara P.J.; Berkener K. et al. (1986):

Genomic cloning, characterization, multilineage growth-promoting activity of human granulocyte-macrophage colony stimulating factor.

Proc. Natl. Acad. Sci. USA, 83: 3101.

Keens T. and Ward S.L. (1993):

Apnea spells sudden death and role of apnea monitor. Pediatr. Clin. North Am. 40 897- 905.

Kingdom J.C.; Macara L.M. and Whittle M.J.(1994):

Fetoplacental circulation in health and disease. Arch. Dis. Child; 70:161-163.

Kitagawa S.; You A.; Souza L.M. et al. (1987):

Recombinant human granulocyte colony-stimulating factor enhances superoxide release in human granulocytes stimulated by chemotactic peptide.

Biochem. Biophys. Res. Commun., 144: 1143-1146.

Kliegman R. (1996):

Fetus and neonate. In: Behram, R; Kliegman, R;(eds.). Nelson Textbook of Pediatrics (15th ed.). W B Saunders, Philadelphia. pp: 431.

Kliegman R. (1998):

Fetal and neonatal medicine: In Behrman, R; Kliegman, R.(eds.). Nelson Essentials of Pediatrics. (3rd ed.). W.B. Saunders, Philadelphia. pp 167.

Kloster B.E. (1996):

Cord blood stem cells: In Henry J. (ed.) Clinical Diagnosis and Management by Laboratory Methods., 19th ed., W.B. Saunders, Philadelphia, pp 837-845.

Knock G.A. and Poston L. (1990) :

Bradykinin-mediated relaxation of isolated maternal resistance arteries in normal pregnancy and preeclampsia. Am. J. Obstet. Gynecol. 175: 1668-1671.

Kocherlakota P. and LaGamma E.(1998):

RhG-CSF may reduce the incidence of neonatal sepsis in prolonged preeclampsia-associated neutropenia. Pediatrics, 102:1107-1111.

Koenig J.M. and Christensen R.D. (1989):

Incidence, neutrophil kinetics, and natural history of neonatal neutropenia associated with maternal hypertension. N. Eng. J. Med. 321:557-562.

Koenig J.M. and Christensen R.D. (1991):

The mechanism responsible for diminished neutrophil production in neonates delivered to women with pregnancy-induced hypertension.

Am. J. Obstet. Gynecol. 165:467-473.

Koo W.(1992):

Parenteral nutrition-related bone disease.

J. Parent. Nutr., 16:386.

Kuban K.C. (1998):

Intraventricular hemorrhage: In Cloherty, J. and Stark, A. (eds.), Manual of Neonatal Care, 4th ed. Lippincott-Raven, Philadelphia. p-p: 505-515.

Kumppel R.A. and Drukker J.(1993):

Hypertention in pregnancy: In High risk pregnancy.

A team Approach, 2nd ed. 468-517.

LaGamma E.; Alpan O. and Kocherlakota P. (1995) :

Effect of granulocyte colony- stimulating factor on pre-eclampsia-associated neonatal neutropenia.

J. Pediatr. 126: 457-459.

Lau C.; Sheena H.; Shulman R. et al. (1997):

Oral feeding in low birth weight infants.

J. Pediatr., 130:561-567.

Laver J.; Duncan E.; Abbound N. et al.(1990):

High levels of granulocyte and granulocyte- macrophage colony-stimulating factors in cord blood of normal full term neonates.

J. Pediatr. 116:627-32.

Liechty K.W.; Schibler K.R.; Ohls R.K. et al. (1993):

The failure of newborn mice infected with Escherichia Coli to accelerate neutrophil production correlates with their failure to increase transcripts for granulocyte colony-stimulating factor and interleukin-6.

Biol. Neonate., 64: 31.

Lieschke G.J. and Burgess A.W. (1992):

Granulocyte colony-stimulating factor and granulocyte-macrophage colony-stimulating factor (1 and 2).
N. Eng. J. Med., 327: 28-35.

Lieschke G.J. (1994):

Granulocyte colony stimulating factor (G-CSF).
Aust. Prescr., 17: 96-98.

Liley H.G. and Stark A.R. (1998):

Respiratory distress syndrome/ Hyaline Membrane Disease: In
Cloherty J and Stark A. (Eds) Manual of Neonatal Care 4th ed.
Lippincott-Raven, Philadelphia, PP. 329-336.

Lowenberg B. and Touw I.P. (1993):

"HGFS and their receptors in acute leukemia".
Blood, 81: 281-292.

Mackendrick W. and Caplan M. (1993):

Necrotizing enterocolitis: New thoughts about pathogenesis and
potential treatments.
Pediatr. Clin. North Am. 40: 1047-59.

Mahrous S. El-Dere S. and Yousef A. (1996):

Circulating progenitors in neonates delivered to mothers with
pregnancy induced hypertension.
Zag. Medical J., Vol. II, 234-241.

Manroe B.L.; Winberg A.G. Rosenfeld C.R. et al. (1979):

The neonatal blood count in health and disease. Reference values
for neonatal neutrophilic cells.
J. Pediatr., 95: 89-98.

Marland G.; Bakker A.B.; Adema G.J. et al. (1996):

Dendritic cells in immune response induction.
Stem Cells. 14:501-507.

Masson J.A. and Herrman K.R. (1998) :

Universal infant screening by automated auditory response measurement.

Pediatrics, 101: 221-228.

McAlman K.R. (1998):

Necrotizing Enterocolitis. In Cloherty, J. and Stark, A. (Eds.), Manual of Neonatal Care (4th ed.), Lippincott-Raven, Philadelphia, PP. 609-615.

McNice I.K.; McGroth H. and Quesenberry P. (1988):

Granulocyte-colony stimulating factor augments in vitro megakaryocyte colony formation by interleukin-3.

Exp. Heme., 16: 807.

Meager A. (1990):

Cytokines, Meager A. (ed). Publ. Redwood Press Lim., 139.

Meister B.; Herold M.; Mayr A. et al. (1993):

Interleukin-3, interleukin-6, granulocyte macrophage- colony stimulating factors and erythropoietin cord blood levels of preterm and term neonates.

Eur. J. Pediatr. 152:569-73.

Ment L.R.; Vohr B. and Whitelaw A.H. (1994):

Low dose of indomethacin and prevention of intraventricular hemorrhage: a multicenter randomized trial.

Pediatrics, 93: 543.

Metcalf D. (1986):

The molecular biology and functions of the granulocyte-macrophage colony stimulating factors.

Blood, 67: 257.

Metcalf D. and Micola N.A. (1995):

The hematopoietic colony stimulating factor: In from biology to clinical Applications. Cambridge University Press. Cambridge, 1995 PP. 44-55.

Miller G.; Heckmatt J.Z.; Dubowitz M.S. et al.(1983):

Use of nerve conduction velocity to determine gestational age in infants at risk and in very low birth weight infants.
J. Pediatr., 103: 109-112.

Modrowski D.; Lomri P.J. and Mariel P.J.(1997):

Endogenous GM-CSF is involved as an autocrine growth factor for human osteoblastic cells.
J. Cell. Physiol. 170:35-46.

Morita M.; Lamkhioued B.; Gounni A.S. et al.(1996):

Induction by interferons of human esinophil apoptosis and regulation by interleukin-3, granulocyte/macrophage-colony stimulating factor and interleukin-5.
Eur. Cytokin. Network. 7:725-732.

Mouzinho A.; Rosenfield C.R.; Sanchez P.J. et al. (1992):

Effect of maternal hypertension on neonatal neutropenia and risk of nosocomial infection.
Pediatrics, 94:76-82.

Nagata S.; Tsuchinya M.; Asano S. et al. (1986):

Molecular cloning and expression of cDNA for human granulocyte-colony stimulating factor.
Nature; 319: 415.

Nakamura H.; Ueki Y.; Sakito S. et al. (2000):

High serum and synovial fluid colony stimulating factor concentration in patient with rheumatoid arthritis.
Clin. Exp. Rheumatol., 18(6): 713-8.

Narayan I.; Dua K.; Gujral. V.V. et al. (1982):

A simple method of assessment of gestational age in newborn infants.
Pediatrics, 69: 27-32.

National Maternal Mortality Study (NMMS) (2000) :

Egyptian Ministry of Health and Population.

Nelson K.B. and Grether J.K. (1995):

Can magnesium sulphate reduce the risk of cerebral palsy in very low birth weight infants?
Pediatrics, 95: 263.

Nemunaitis D. (1993) :

GM-CSF : A review from preclinical development to clinical application.
Transfusion, 33: 70-83.

Nicola N.; Metcalf D.; Matsumoto M. et al., (1983):

Purification of a factor inducing differentiation in murine myelomonocytic leukemia cells: Identification as granulocyte colony stimulating factor (G-CSF).
J. Biol. Chem. 258:9017-22.

Nicola N.; Begley C. and Metcalf D. (1985):

Identification of human analogue of a regulator that induces differentiation in murine leukemic cells.
Nature 314:625-631.

Ogawa M. (1993):

Differentiation and proliferation of hematopoietic stem cell.
Blood, 81: 2844-2853.

Ohls R.K.; Liy Abdel-Mageed A.; et al.(1995):

Neutrophil pool size and granulocyte colony stimulating factor production in human mid-trimester fetuses.
Pediatr. Res. 37:806.

Ohls R.K. and Christensen R.D. (2000):

Development of the hematopoietic system. In: Behrman, Kliegman and Jenson (eds), Nelson Textbook of Pediatrics, 16th ed., WB. Saunders, Philadelphia, PP: 1456-1460.

Oski F.K.(1991):

Physiologic jaundice : In Teausch, H; Ballard, R; and Avery, M; (eds.).Diseases of The Newborn (6th ed.). W.B. Saunders, Philadelphia. 754.

Overturf F.G.; Scherman M. and Scheifele D.(1990):

Neonatal necrotizing enterocolitis associated with deltatoxin-producing methicillin-resistant staph.aureus.
Pediatr. Infect. Dis. 9 . 88.

Ozer H.; Armitage J.O. and Bennett C.L. (2000) :

Update of recommendations for the use of hematopoietic colony-stimulating factors: evidence-based, clinical practice guidelines.
J. Clin. Oncol. 18: 3558-3585.

Padbury J.F.; Gore E.M.; and Polk D.H.(1996):

Extrapulmonary effects of antenatally administered steroids.
J. Pediatr. 128:167-172.

Parad R.B. and Berger T.M. (1998):

Chronic lung disease: In Cloherty J and Stark A (eds.) Manual of Neonatal Care 4th ed. Lippincott-Raven, Philadelphia., P-P: 378-388.

Parkin J.M.; Hey E.N. and Clowes J.S. (1976):

Rapid assessment of gestational age at birth.
Arch. Dis. Child., 51: 259-263.

Patil R.R. and Borch R.F. (1995):

Granulocyte macrophage colony stimulating factor expression by human fibroblasts is both upregulated and subsequently downregulated by interleukin-1.
Blood, 80(1): 85.

Peter G.B. and Johnson M.B. (1994):

Preterm infants: In Vulliamy"s The Newborn Child 17th ed.
W.B. Saunders, Philadelphia. pp: 93.

Phelps D.L. (1993):

Retinopathy of prematurity.
Pediatrics, 91: 224.

Robberts J.M.(1995):

Preeclampsia-not simply induced hypertention.
Hosp. Pract.; 30: 25-28.

Robberts J.M. and Redman C.W.(1993):

Preeclampsia more than pregnancy-induced hypertention.
Lancet. 341:1447-1451.

Robinson R.J.(1966):

Assessment of gestational age by neurological examination.
Arch. Dis. Child.41:437.

Ross R. and Anthomy W.(1991):

Growth factors and their receptors: Genetic control and rational application.
J. Nat. Cancer Institute,82:874.

Rudolph N. (1998):

Low birth weigh: In Finberg, L. ed. Saunders manual of Pediatric Practice (1st ed) WB Saunders Philadelphia, PP: 64-69.

Ryan C.A. and Finer N.N.(1995):

Antenatal corticosteroid therapy to prevent respiratory distress syndrome.
J. Pediatr., 126: 317-319.

Salah G.; Ashraf M.; Affifi H. et al. (2002) :

Granulocyte- colony stimulating factor in cord blood in preterm neonates born to mothers with pregnancy induced hypertension.
Theisis, Ain Shams University

Sanders M.; Allen M.; Alexander G.R. et al.(1991):

Gestational age assessment in preterm neonates weighting less than 1500 grams.
Pediatrics, 88: 542-546.

Saugasted O.D.(1992):

Neonatal oxygen radical disease.
Recent Adv. Pediatr. 10: 173.

Sieff C.A. (1987):

Hematopoietic growth factors.
J. Clin. Invest., 79: 1549.

Simmers R.M.; Webber L.M.; Shannon M.F. et al. (1987):

Localization of the G-CSF gene on chromosome 17 proximal to the breakpoint in the acute promyelocytic leukemia.
Blood, 70: 330.

Slayton W.B.; Juul S.; Calhoun D.A. et al. (1998):

Hematopoiesis in the liver and marrow of human fetus at 5 to 16 weeks post conception: Quantitative assessment of macrophage and neutrophil populations.
Pediatr. Res., 43: 774.

Souza L.M.; Boone T.C.; Gabrilove J. et al. (1986):

Recombinant human granulocyte-colony stimulating factor: effects on normal and leukemic myeloid cells.
Science, 232: 61.

Spiekerman K.; Roesler J.; Emmendoerffer A. (1997):

Functional features of neutrophils induced by G-CSF and GM-CSF treatment: differential effects and clinical implications.
Leukemia, 11: 466-478.

Stark A.R. (1998):

Apnea. In: Cloherty J and Stark A (Eds), Manual of Neonatal Care, 4th ed. Lippincot-Raven, Philadelphia, P: 374-378.

Steel R.G.D. and Torrie J.H. (1960):

Principles and procedures of statistics, with special reference to the biological sciences. McGraw-Hill Book Company, New York.

Stocking C. (1997):

Gene transfer into hemopoietic cells.
Bailleres Clinc. Haemat. 10: 445-466.

Stoll B. and Kliegman R. (2000):

The fetus and the neonatal infants, non-infectious disorders. In: Behrman, Kliegman, and Jenosen (eds.) Textbook of Pediatrics 16th ed., WB Saunders, Philadelphia. pp: 451-538.

Stolz J.W. (1998):

Hearing loss in neonatal intensive care unit graduates. In: Cloherty J and Stark A. (eds), Manual of Neonatal Care 4th ed. Philadelphia, Lippincott-Raven, P. 648-649.

Strauss R.G. (1995):

Erythropoietin and neonatal anemia.
N. Engl. J. Med., 330: 1227.

Strife A.; Labek C.; Wisniewski D. et al. (1987):

Activities of four purified growth factors on highly enriched human hematopoietic progenitor cells.
Blood., 69: 1508.

Sun Y.; Awnetwant E.L.; Colliers S. et al. (1998):

Nutrition in neonates: In Clohertry, J. and Stak, A. Manual of Neonatal Care 4th ed. Lippincott-Raven, Philadelphia. pp: 101-142.

Surbeck D.V.; Danzer E.; Steinmann C. et al. (2001) :

Effect of preeclampsia on umbilical cord blood hematopoietic progenitor-stem cells.
Am. J. Obstet. Gynecol. 158(3) : 191-200.

Testa N.G. and Dexter T.M. (1999):

The regulation of hemopoietic cell production. In: Hoffbrand Lewis and Tuddenham (eds), postgraduate hematology (4th ed.). Butterworth/ Heinemann, Oxford, Boston, PP: 1-12.

Tobler A.; Meier R.; Seitz M. et al. (1992):

Glucocorticoids downregulate gene expression of GM-CSF, IL-8 and IL-6 but not of M-CSF in human fibroblast.
Blood, 79: 45.

Torehenriksen, (1997):

Hypertension in pregnancy: Use of antihypertensive drugs.
Acta. Obstet. Gynecol. Scand., 76:96-106.

Tsang R.C. (1995):

Calcium, phosphorus and magnesium metabolism, section XXV.
In Pollin R and Fox W (eds.), fetal and neonatal physiology.
Saunders, Philadelphia.

Tsao P.; Teng R.; Tang J. et al. (1999):

Granulocyte colony-stimulating factor in cord blood of premature neonates born to mothers with pregnancy induced hypertension.
J. Pediatr. 135:56.

Tsuchiya M.; Asano S.; Kazioro Y. et al. (1986):

Isolation and characterization of the cDNA for murine granulocyte colony stimulating factor.
Proc. Natl. Acad. Sci. USA, 83: 7633.

Udall J. N.(1990):

Gastrointestinal host defence and necrotizing enterocolitis.
J. Pediatr. 117: 533.

Van B.O.; Follens I.; Hartmann S. et al. (1997):

Treatment of patent ductus arteriosus with ibuprofen.
Arch. Dis. Child., 77: F147.

Van-Marter L.J.; Pagano M. and Allerd G.(1992):

Rate of bronchopulmonary dysplasia as a function of neonatal intensive care practices.
J. Pediatr. 120: 938-946.

Varlotta L. and Allen J.(1994):

Apnea: In Schidlow D. and Smith D.(eds) A Practical Guide To Pediatric Respiratory Diseases; Saunders, Philadelphia. p 149.

Vellenga E.; Dokter W.; Wolf I.M. et al. (1991):

Interleukin-4 prevents the induction of G-CSF mRNA in human adherent monocytes in response to endotoxin and IL-1 stimulation.

Br. J. Haematol., 79: 22.

Veukateswaran L.; Wilimas J.A.; Dancy R. et al. (2000):

Granulocyte-macrophage colony stimulating factor in the treatment of neonates with neutropenia and sepsis.

Pediatr. Hemat. Oncol. 17:(6) 469-73.

Victor Y.H.(1994):

Prematurity and low birth weight: In Robinson M. Robertson D. and Pllan P. (eds.) Practical Pediatrics, 3rd. ed. Chirchul-Livingstone London. P-p237.

Volpe J. J. (ed.) (1996):

Intraventricular hemorrhage of premature infants: In Neurology of the newborn. 3rd. ed. Saunders, Philladelphia. p-p 403-460.

Voher B. and Ment L.R. (1996):

Intraventricular hemorrhage in the preterm infant.

Early Hum. Dev., 44: 1-9.

Wada H.; Noguchi Y.; Marino M.W. et al. (1997):

T cells function in granulocyte-macrophage colony-stimulating factor deficient mice.

Proc. Natl. Acad. Sci. USA. 94:12557-12561.

Walker J.J.(1998):

Current thoughts on the pathophysiology of preeclampsia\ eclampsia. In: Progress In Obstetrics and Gynecology. Vol.13, Studd, J;(ed.). Churchill Livingston. London . pp:177-190.

Walker J.J. (2000):

Preeclampsia.

Lancet, 356:1260-1265.

Walsh S.W. (1985) :

Preeclampsia: An imbalance in placental prostacyclin and thromboxane A₂ production.
Am. J. Obstet. Gynecol. 152: 1690-1694.

Ward K.; Hata A.; Jeunemaitre A. et al. (1994):

Amelior variant of angiotensinogen associated with preeclampsia (comment).
Nat. Genet.;4: 59-63.

Ward R.M.(1994):

Pharmacological enhancement of fetal lung maturation.
Pediatr. Clin. North Am. 21:523.

Watchko J.F. (1992):

Kernicterus in preterm newborn: Past, present, and future.
Pediatrics, 90: 707.

Wechsler S.B. and Wernovsky (1998):

Patent ductus arteriosus: In: Cloherty J and Stark A. (eds). Manual of Neonatal Care 4th ed. Lippincott-Raven, Philadelphia, PP: 430-31.

Weinmann E.; Rutkowski S. and Reisbach G. (1998):

G-CSF, GM-CSF, and IL-6 levels in cord blood : Diminished increase of G-CSF and IL-6 in preterms with perinatal infection compared to term neonates.
J. Pediatr. Med. 26:211.

Weiss L.P. (1995a):

The structure of hematopoietic tissues. In: Handin RI, Lux SE, Stossel TP, eds. Blood: Principles and Practice of Hematology. Philadelphia: JB Lippincott: 155—169.

Weiss L.P. (1995b):

Functional organization of hematopoietic tissue. In: Hoffman RH, et al., eds. Hematology: Basic Principles and Practice, 2nd ed. New York: Churchill Livingstone; 193-206.

Welte K.; Platzer E.; Lu L. et al. (1985):

Purification and biochemical characterization of human pluripotent hematopoietic colony-stimulating factor.
Proc. Natl. Acad. Sci. USA, 82: 1526.

Wickenhauser C.; Lorenzen J.; Thiele H. et al. (1995) :

Secretion of cytokines (IL-1 alpha, IL-3, IL-6 and GM-CSF) by normal human megakaryocytes.
Blood. 85: 685.

Wilker R.E. (1998):

Hypoglycemia and hyperglycemia. In: Cloherty, J. and Stark, A. (eds.). Manual of Neonatal Care 4th ed. Lippincott-Raven, Philadelphia, PP: 545-553.

Wilimas J.A.; Wall J.E.; Fairclough D.L. et al. (1995) :

A longitudinal study of granulocyte colony-stimulating factor levels and neutrophil counts in newborn infants.
J. Pediatr. Hematol. Oncol. 17(2): 176-9.

Williams D.A. (1995):

The stem cell model of hematopoiesis. In: Hoffman R, Benz E, Shattil S, Furie B, Cohen H, Silberstein L, eds. Hematology: Basic Principles and Practice. Williams & Wilkins, Baltimore. pp. 180-192.

Wimperis J.Z.; Niemeyer C.M.; Sieff C.A. et al. (1989):

Granulocyte macrophage colony stimulating factor and interleukin-3 mRNA are produced by a small fraction of blood mononuclear cells.
Blood, 74: 1525.

Whetton A.D. ed.(1997):

Molecular Hemopoiesis.
Clinical Haemat. 10: 429-619.

Wolach B.; Gavrieli R. and Pomeranz A. (2000):

Effect GM-CSF and G-CSF on neonatal neutrophil functions.
Pediatr. Res., 48: 369-373.

Wong G.G.; Witek J.S. and Temple P.A. (1985) :

Human GM-CSF : Molecular cloning of complementary DNA and purifications of the natural and recombinant proteins.
Science, 228: 810.

Wright D.; Kenney R.; Oette D. et al. (1994) :

Contrasting effects of recombinant human granulocyte macrophage colony stimulating factor and granulocyte colony stimulating factor treatment on the cycling of blood elements in childhood-onset cyclic neutropenia.
Blood. 84: 1257.

Xanthou M. (1970) :

Leucocyte blood picture in healthy full-term and premature babies during neonatal period.
Arch. Dis. Child. 45: 242-249.

Yong K.L. and Linch D.C. (1992):

Differential effects of granulocyte-and granulocyte macrophage. Colony stimulating factors (G- and GM-CSF) on neutrophil adhesion in vitro and in vivo.
Eur. J. Haematol., 49: 251-259.

Zsebo S.N.; Cohen A.M.; Murdock D.C. et al. (1986):

Recombinant human granulocyte colony stimulating factor: Molecular and biological characterization.
Immunobiol., 172: 175.

Zsebo S.N.; Yushenkoff V.; Schiffer S. et al. (1988):

Vascular endothelial cells granulopoiesis IL-1 stimulate release of G-CSF and GM-CSF.
Blood, 71: 103.