

REFERENCE

- Adcok LM, Yamashita Y, Gooddard -Finegold J, Smith CV, (1996):** Cerebral hypoxia-ischemia increases microsomal iron in newborn piglets. *Metab Brain Dis.*; 11: 3 59-367.
- Akhter W, Ashraf QM, Zanelli SA, Mishra OP, Delivoria-Papadopoulos M, (2001):** Effect of graded hypoxia on cerebral cortical genomic DNA fragmentation in newborn piglets. *Biol Neonate*, 79: 187-93.
- Al Nakib L, Pennock J Hajnal JV, et at, (1999):** Diffusion weighted MRI in neonatal stroke-timing of onset of lesions and prediction of lesion size. *Eur. J Pediatr Neurol*; 3: A124-125.
- Al Naqeeb N, Edwards AD, Cowan FM et at., (1999):** Assessment of neonatal encephalopathy by amplitude-integrated electroencephalography. *Pediatrics*; 103: 1263-71.
- Alejandro G, Jenik, Jose M, Ceriani Cernadas, Adriana Gorenstein, Jose A, Ramirez, Nestor Vain, Macelo Armadans, and Jorge R, Ferraris, (2000):** A randomized, Double-blind, placebo-controlled trial of the effects of prophylactic theophylline on renal function in term neonates with perinatal asphyxia. *Pediatrics*. 105 (4).
- Alfy A, Carroll JE, Devarajan LV, Moussa MAA, (1990):** Term infant asphyxia in Kuwait. *Ann Trop Paediatr*; 10: 355-361.
- American Academy of pediatrics Committee on Fetus and Newborn (2000).** Use of inhaled nitric oxide. *Pediatrics*; 106: 344-5.

- American College of obstetricians and Gynecolgists (1995) :** Umbilical artery blood acid-base analysis. Washington : The college: 1-5. Technical Bulletin No 216.
- American Psychiatric Association. (2000):** Diagnostic and statistical manual of mental disorders, text revision, 4th ed. Washington: American Psychiatric Association; 41-49.
- Amiel-Tison and Ellison. (1986):** Birth asphyxia in the full term newborn early assessment and outcome: Dev Med Child Neur 128: 671.
- Ana Martin-Ancel, Alfredo Garcia Alix, Francisco Gaya Fernanado Cabanas, Margarita Burgueros Jose Quero, (1995):** Multiple organ invovlement in perinatal asphyxia. J Pediatr 127: 786-93.
- Ansari, M.Q, Chincanchan, C'.A and Armstrong, DL (1990.):** Brain calcification in hypoxic- ischemic lesions: An autopsy review. Pediatr. Neurol 6:94-101.
- Arebelaez A, Castillo M, and mukherji SK (1999):** Diffusion-weighted MR imaging of global cerebral anoxia. Am J Neuroradiol; 20: 999-1007.
- Ashwal S, (1989):** Brain death in the newborn. Clin Perinatol 16:501-518.
- Badawi N, Kurinczuk JJ, Keogh JM et al., (1998):** Antepartum risk factors for newborn encephalopathy: the Western Australian case control study. Br Med. J: 317: 1549-53.
- Baschat AA, Gembruch U, Reiss I, Gortner L, Harman CR, Weiner CP, (1999):** Neonatal neuclated red blood cell counts in growth – restriced fetuses: relationship to arterial and venous Doppler studies. Am J Obstet Gynecol; 181: 190-195.

- Basher A, Itoo, Zakaria M, Al-Hawsawi, DCH, CARP, Anwar H, Khan (2003):** Hypoxic ischemic encephalopathy incidence and risk factors in north western Saudi Arabia. *Saudi Med J*; Vol. 24(2): 147-153.
- Bauer M, Feucht K, Ziegenfuss I, Marzi I, (1995):** Attenuation of heat shock -induced hepatic microcirculatory disturbances by the use of starch -deferoxamine conjugate for resuscitation. *Crit Care Med*. 23: 316-322.
- Bax MC, (1964):** Terminology and classification of cerebral palsy. *Developmental Medicine and Child Neurology* 6: 295-307.
- Beckman JS, (1991):** The double-edged role of nitric oxide in brain function and superoxide-mediated injury. *J Develop Physiol* 15: 5 3-9.
- Belai Y, Goodwin TM, Durand M, Greenspoon IS, Paul RH, and Walther FJ (1998):** Umbilical arteriovenous Po₂ differences and neonatal morbidity in term infants with severe acidosis *Am J Obstet Gynecol*; 178: 13-19.
- Berger R, Garnier Y (1999):** Pathophysiology of perinatal brain damage. *Brain Res Rev*; 30: 107-134.
- Berseth CL, McCoy HH, (1992):** Birth asphyxia alters neonatal intestinal motility in term neonates. *Pediatrics*, 90: 669-673.
- Biagioni E, Bartalena L, Boldrini A, pieri R, Cioni G, (1999):** Constantly discontinuous EEG patterns in full-term neonates with hypoxic ischaemic encephalopathy. *Clin. Neurophysiol*: 110: 1510- 1515.
- Biomond P. Swaak AJG, Beindorff CM, Koster JF, (1986):** Superoxide dependent and independent mechanisms of iron mobilisation from ferritin by xanthine oxidase. *Biochem J* 239: 169-73.

- Bittigau P, Sifringer M, Genz K, Reid, E, Pospischil D, Govindarajalu S, Dzieko M, Pesditschek S, Mai I, Dikranian K Olney JIV, Ikonomidou C, (2002):** Antipileptic drugs and apoptotic neruodegeneration in the developing brain. *Proc. Natl Acad. Sci., USA.* 99: 15089-15094.
- Bjerre I, Hellstrom —Westas L, Rosen I, Svenningen NW, (1983):** Monitoring of cerebral function after sever asphyxia in infancy. *Arch. Dis. child* 58: 997-1002.
- Bloom SL, Swindle RG, McIntrie DD, Leveno KJ, (1999):** Fetal pulse oximetry: Duration of desaturation and integration outcome. *Obstet Gynecol*; 93: 1036.
- Boda V, Finck B, Durken M, Comments J, Hell wege HH, Kohlschutter A, (1998):** Monitoring erythrocyte free radical resistance in neonatal blood microsamples using a peroxy radical -medicated haemolysis test. *Scand J Clin Lab Invest*; 58: 317- 322.
- Boo NY, Lye MS, (1991):** Factors associated with clinically significant perinatal asphyxia in Malaysian neonates: a cases control study. *J. Trop Pediatric*, 38: 284-289.
- Bracci R, (1998): Free radials in neonatal brain disorders. Possible benefits of antioxidants.** In: Cosmi EV. Marinoni E, Di Lorio R. editors. *New technoliges in reproductive medicine. Neoantoology and , gynecology.* New York. Parthenon; 131-9.
- Bracci R, (1998): Oxygen toxicity in Kurjak A (ed): Text-book of perinatal medicine.** London, Parthenon, pp. 78-89.
- Bracci R, Perrone S, Buonocore G, (2001):** Red blood cell involvement in fetal neonatal hypoxia. *Biol Neonate* 79: 210-12.

- Bromont C, Marie C, Bralet J, (1989):** Increased lipid peroxidation in vulnerable brain regions after transient forebrain ischemia in rats. *Stroke* 20: 918-24.
- Buonocor G, Perrone S, Longini M, Vezzosi P, Marzocchi B, Paffetti P et al., (2002a):** Oxidative stress in preterm neonates at birth and on the seventh day of life. *Pediatr Res.* 52: 46-9.
- Buonocore G, Perrone S, Carlini M, Bagnoli F, Gatti MG, Paffetti P et al., (2001):** Non protein bound iron plasma levels in critically ill newborns (abstract), *Biol Neonate*; 80: 93-4.
- Buonocore G, Perrone S, Gioia D, Gatti MG, Massafra C, Agosta R, et al., (1999):** Nucleated red blood cell count at birth as an index of perinatal brain damage. *Am J Obstet Gynecol*; 181: 1500-5.
- Buonocore G, Perrone S, Longini M, Paffetti P, Marzocchi B, Vezzosi P, Bracci R, (2002):** Oxidative stress in preterm babies at birth and on the seventh day of life. *Pediatric Res*; in press.
- Buonocore G, Perrone S, Longini M, Terzuoli L, Bracci R, (2000):** Total hydroperoxide and advanced oxidation protein products in preterm hypoxic babies. *Pediatr. Res*; 47: 221-224.
- Buonocore G, Zani S, Sargenitin I, Gioia D, Signorini C, Bracci R, (1998):** Hypoxia-induced NPBI released in the red cells of newborn infants. *Acta Paediatr*; 87: 77-81.
- Buonocore G, Zani S, Sargentini I, Gioia D, Signorini C, Bracci R. (1998):** Hypoxia-induced free iron release in the red cells of newborn infants. *Acta Paediatr* 87: 77-81.

- Busza AL, Allen KL, King MD et al., (1992):** Diffusion weighted imaging studies of cerebral ischemia in gerbils: potential relevance to energy failure. *Stroke*; 23: 1602-1612.
- Cabaas, F., Pellicer, A Prez-Higueras, A., et al (1991):** Ultrasonographic findings in thalamus and basal ganglia in term asphyxiated infants. *Pediatr. Neurol* 7:211-215.
- Caroline A, Dorrepaal, Howard M, Beger, Manon J. N.L. Benders, (1996):** Non protein-bound iron in postasphyxial reperfusion injury of the newborn. *Pediatrics*. Nov; 98:5.
- Carson S.C., Hertzberg, B.S., Bowie, J.D., et al (1990):** Value of sonography in the diagnosis of intracranial hemorrhage and periventricular leukomalacia : A postmortem study of 35 cases *Am. J. Neuroradiol* 11:677-683.
- Casey BM, McIntrie DD, Leveno KJ, (2001):** The continuing value of the Apgar score for the assessment of newborn infants. *N Engl J Med*. 344: 467-71.
- Chandra S, Ram JI, Thirupuram S, (1997):** Perinatal asphyxia multivariate analysis of risk factors in hospital birth. *Indian Pediatr*; 34: 206-212.
- Charlotte van Kesteren, Manon J. N. L. Benders, Floris Groenendaal, Frank van Bel F. F. Tessa Ververs, and Carin M. A. Rademaker, (2006):** Population Pharmacokinetics of Allopurinol in full-Term Neonates With Perinatal Asphyxia. *Ther Drug Monit* ;28:339-344.
- Cheung PY, Robertson CMT and Finer NN (1996):** Plasma lactate as a predictor of early childhood neurodevelopmental outcome of

neonates with severe hypoxemia requiring extracorporeal membrane oxygenation . Arch Dis Child ; 74: F47 —F50.

Chiappa KH, (1990): Brainstem auditory evoked potentials: methodology. In: chaippa KH, editor, Evoked potentials in clinical medicine. New York: Raven Press: p. 173-221.

Churchal PC, Bidani AK, (1982): Hypothesis: adenosine mediates hemodynamic changes in renal failure. Med Hypotheses: 8: 275-285.

Ciccoli L, Rossi V, Signorini C, Leoncini S, Buonocore G, Paffetti P, et al, (2001): Iron release in newborn and adult erythrocytes exposed to hypoxia-reoxygenation Biol Neoante 80: 102.

Ciccoli L, Signorini C, Alessandrini C, Ferrali M, Comporti M, (1994): Iron release lipid peroxidation and morphological alterations of erythrocytes exposed to acrolein and phenylhydrazine. Exp Mol. Pathol 60: 108-118.

Ciccoli L, Signorini C, Scarano C, Rossi V, Bambaginoi S, Ferrali M, Comporti M, (1999): Iron release in erythrocytes from patients with B-thalassemic erythrocytes. Free Radic Res. 30: 401-413.

Clark SL, Gimovsky ML, Miller FC, (1984): The scalp stimulation test: A clinical alternative to fetal scalp blood sampling. Am J Obstet Gynecol; 148: 274.

Corneth L, Levene MI, (2001): Postoperative management of the asphyxiated term and preterm infant. Semin Neonatol. 627-82.

Coskum A; Lequin M; Vigneron D et al., (2001): Quantitative analysis of MR images in asphyxiated neonates: Correlation with

neurodevelopmental outcome. *AJNR Am.J. Neurodiol* ; 22 (2) 400-5.

Cowan FM, Pennick J, Hanrahan JD, et al., (1994): Early detection of cerebral infarction and hypoxic ischemic encephalopathy in neonates using diffusion-weighted magnetic resonance imaging. *Neuropediatrics*; 25: 172-175.

Curran JS and Barness LA (2000): Nutrition. In : Nilson textbook of pediatrics. Behrman RE, Kliegman RM and Jenson HB (eds) 16th edition, WB Saunders Co, Philadelphia, p:4

Dammann O, Leviton A, (1997): Maternal intrauterine infection, cytokines and brain damage in the preterm newborn. *Pediatr Res*; 42: 1-8.

Dauber IM, Krauss AN, Symchych PS, Auld PA, (1976): Renal failure following perinatal anoxia. *J pediatr*. 88(5):851-5.

De Menezes MS (2002): Hypoxic-ischemic Brain Injury In Newborn. *eMedicine J* [Serial online]; May 13 [33 screens]. Available from: <http://www.eMedicine.com>.

De Vries LS (1993): Somatosensory-evoked potentials in term neonates with postasphyxial encephalopathy. *Clin Perinatol*; 20: 463-482.

De Vries LS, Eken P, Groenendaal F, Rademaker KJ, Hoogervorst B, and Bruinse HW (1998): Antenatal onset of hemorrhagic and/or ischemic lesions in preterm infants: prevalence and associated obstetric variables. *Arch Dis Child Fetal Neonatal Ed*; 78: F51-F56.

De Vries, L.S., Wigglesworth, J.S., Regev R. et al., (1988): Evolution of periventricular leukomalacia during the neonatal period and

infancy Correlation of imaging and Postmortem findings Early Hum Dev 17:205-2 19.

Devoe- LD (1990,): The non-stress test. Obstet. Gynecol.Clin. North Am. 17: 111-128

Diego Gazzolo, Emanuela Marinoni, Romolo Di Iorio, Matteo Bruschetti, MD, Maria Koruacka, Mario Lituania, Urszula Majewska, MD, Giovanni Sera, Fabrizio Michetti, (2004): Urinary S 100B protein measurements: a tool for the early identification of hypoxic-ischemic encephalopathy in asphyxiated full-term infants; Crit Care, Med; 32: 131-136.

Dietrich RB, Bradley WG, (1998): Iron accumulation in the basal ganglia following severe ischemic-anoxic insults in children Radiology: 68: 203-206.

Dinah S Reddihough and Kevin J, Collins (2003): The epidemiology and causes of cerebral palsy, Australian Journal of Physiotherapy. 49: 7-12.

DiseSsa TG, Leitner M, Ti CC, et al., (1981): The cardiovascular effects of dopamine in the severely asphyxiated neonate. J Pediatr; 99: 772-6.

Dorrepaal CA, Berger HM, Benders Benders MJN, Van Zoeren-Grobbe D, Van De Bor M, Van Bel F, (1996): Non-protein bound iron in postasphyxial reperfusion injury of the newborn. Pediatrics: 98: 883-889.

Dubowitz L, Mercuri E, Dubowitz V, (1998): An optimality score for the neurologic examination of the term newborn J Pediatr. 133: 406-416.

- Edersheim TG, Hutson JM, Druzin ML, Kogut EA, (1987):** Fetal heart rate response to vibratory acoustic stimulation predicts fetal pH in labor. *Am. J Obstet Gynecol*; 157-1557.
- Ellis M, Mandandhar N, Mandandhar D, Costellos AM del. (2000):** Risk factors for neonatal ecephalopathy in kathmandu. Nepal. a developing country: unmatched case-control study. *BMJ*, 320: 1229-1236.
- Esther Zeinstra, Johanna M Fock, Jacobus H Begeer Tiemen W Van Weerden, Natash M, Maurits, Marianna J, Zweens, (2001):** The prognostic value of serial EEG recordings following acute neonatal asphyxia in full-term infants. *European journal of paediatric neurology*; 5: 155-160.
- Fawer, C.L. Calame, A, perentes, E. E. et al., (1985):** Periventricular leukomalacia: ultasound in the neonate. A correlation study between real- time autopsy findings. *Periventricular leukomalacia in the neonate Neurodiology* 27: 292-300.
- Fee SC, Malee K, Deddish R, (1990):** Sever acidosis and subsequent neurologic status. *Am J. Obstet Gynecol*; 162: 802-6.
- Fellman V, Raivio KO, (1997):** Reperfusion injury as the mechanism of brain damage after perinatal asphyxia. *Pediatric Res.* 41: 599-607.
- Fenichel GM (2001):** Psychomotor retardation and regression. In: *Clinical pediatric neurology a signs and symptoms approach*, 4th ed. Philadelphia: WB Saunders 117-147.

- Ferrali M, Signorini , Ciccoli L, Comporti M, (1993):** Iron released from an erythrocyte lysate by oxidative stress is diffusible and in redox active form, *FEBS Lett*; 319: 40-44.
- Ferrali M, Signorini C, Ciccoli L, Comporti M, (1992):** Iron release and membrane damage in erythrocytes exposed to oxidizing agents. phenylhydrazine, divicine and isouramil. *Biochem. J*; 285: 295-301.
- Finer NN, Barrington KJ, (2001):** Nitric oxide for respiratory failure in infants born at or near term. *Cochrane library Disk Issue*.
- Fischer A,Q., Anderson, J.C, and Shuman R.M. (1998):** The evolution of ischemic cerebral infarction in infancy: A sonographic evaluation *J. Child. Neurol.*, 3:105-109.
- Fitzhardinge P.M Flodmark O, Fiz, C.R., et al (1981):** The prognostic value of computed tomography as an adjunct to assessment of the term infant with postasphyxial encephalopathy *J. pediatr* 99:777- 781.
- Flodmark, O., Roland, E.H. Hill A. et al (1987):** Periventricular leukomalacia: Radiologic diagnosis *Radiology* 162:119-124 1987.
- France Cowan, (2000):** Outcome after in utero asphyxia in term infants. *Semin Neonatol*; 5: 127-140.
- Frankenburg WK, Dodds JB, (1992):** Training manual, Denver II. Denver Development Materials.
- Frankenburg WK, Doods JB, (1967):** The Denver Development Screening Test *J Pediatr*; 71(2): 181-91.

- Freeman RK, Garite TH, Nageotte MP, (1991):** Fetal heart rate monitoring 2nd ed. Baltimore, Williams and Wilins.
- Freeman S, Sohmer H, Silver S. (1991):** The effect of stimulus repetition rate on the diagnostic efficacy of the auditory nerve-brain-stem evoked response. *Electroencephalogr Clin neurophysiol* ; 78: 284-90.
- Frey B, Pfenninger J, Bachman D, (1995):** Reply. *J. Pediatric* 127: 334.
- Fullerton AJ, Ditelberg JS, Chen SV et al., (1998):** Copper/zinc superoxide dimutase transgenic brain accumulates hydrogen peroxide after perinatal hypoxic-ischemia. *Ann Neurol*; 44: 357-364.
- Gaffney G, Flavell V, Johnston A, Squier M, and Sellars S, (1994):** Cerebral palsy and neonatal encephalopathy. *Archives of disease in Childhood*, 70: F195 - F20.
- Ganong WF (1999):** Energy balance , Metabolism and nutrition. In: Ganong WF (ed), 19th edition, Appleton and Lange Co, California ; pp: 267
- Gary F, Cunningham, Norman F, Grant Kenneth J, Leveno, Larry C, Gilstre, John C, Heut, Kather D, Wenstro (2001):** William Textbook of obstetric 21st Edition.
- Gary F, Cunningham, Norman F, Grant Kenneth J, Leveno, Larry C, Gilstre, John C, Heut, Kather D, Wenstro (1903):** William Textbook of obstetric 1st Edition.
- Gathwala Geeta, Khera Atul, Singh Ishwar, Rohtak, Rohtak, (2006):** Mangesium therapy in birth asphyxia. *Indian J pediater* ; 73: 209-212.

- Ghosh B, Mittal S, Kumar S, Dadhwal V, (2003):** Prediction of perinatal asphyxia with nucleated red blood cells in cord blood of newborns. *International Journal of Gynecology and Obstetrics* 81: 267-271.
- Giuseppe Buonocore, Serafina Perrone, Mariangela Longini, Patrizia (2003):** Non protein bound iron as early predictive marker of neonatal brain damage. *Brain*, 126, 1224-1230.
- Glenys Dixon, BPsych, Nadia BAdawi, PhD, Jennifer J, Kuzuk, John M, (2002):** Early developmental outcomes After newborn encephalopathy. *Pediatrics* Vol., 109 No.1.
- Gomella TL, Cunningham MD, Eysel FG, eds. (1994):** Neonatology: Management, procedures, on-Call problems, Disease and Drugs. 3rd ed. Norwalk, Connecticut: Appleton and Lange.
- Gouyon JB, Guignard JP, (1988):** Theophyllin prevents the hypoxemia induced renal hemodynamic changes in rabbits. *Kidney. Int.* 33: 1078-1083.
- Greinberg L, Rachmilewitz EA, Kitrossky N, Chevion M, (1995):** Hydroxyl radical generation in B-thalassemic red blood cells. *Free radic Biol Med*; 18: 611-615.
- Greisen G, (1997):** Tape-recorded EEG and the cerebral function monitor: amplitude-integrated, time compressed EEG. *J. Perinat Med.* 22: 541-6.
- Grether JK, Nelson KB, (1997): Maternal infection and** cerebral palsy in infants of normal birth weight. *JAMA* 278-207-11.
- Groenendaal F, Shadid M, McGowan JE, Mishra OP, Van Bel F, (2000):** Effects of deferoxamine, a chelator of free iron, on Na⁺,

K⁺ ATPase activity of cortical brain cell membrane during early reperfusion after hypoxia ischemia in newborn, lams. *Pediatr Res.* 48: 560-564.

Gull I, Jaffa AJ, Oren M, Grisaru D, Peyser MR, Lessing JB, (1996): Acid accumulation during end-stage bradcardia in term fetuses: How long is too long? *Br J Obstet Gynecol* 103: 1096.

Gunn AJ, Gluckman PD, Gunn TR, (1998): Selective head cooling in newborn infants after perinatal asphyxia: a safety study. *pediatrics.* 102: 885-892.

Guoguang Fan, Zhenhua Wu, Liying Chen qiyon Guog, Binbin Ye, Jian Mao (2003): Hypoxic-ischemic encephalopathy in full-term neonate: correlation proten MR spectroscopy with MR imaging. Received 7 November 2001; *European journal of radiology.* 45: 9 1-98.

Gururaj A; Sztrihá L; Hass D et al. (2002): CT and MR patterns of hypoxic-ischemic brain damage following perinatal asphyxia *J. Trop Paediatr.*, 48(1) 5-9.

Hallioglu O, Topaloglu AK Zenciroglu A, Duzovali O, Yilgor E, and Saribas S (2001): DDST II for early identification of the infants who will develop major neurological sequale of HIE. *Pediatrics International*; 43: 400-404.

Halliwell B, (1989): Protection against tissue damage in vivo by desferrioxamine: what is its mechanism of action? *Free Rad Biol Med* 7: 645-51.

Halliwell B, (1992): Reactive oxygen species and the central nervous system. *J Neurocbemi* 59: 1609-23.

- Hamilton R J, Hodgett SG and O' Brien PM (1996):** Near infrared Spectroscopy applied to intrapartum fetal monitoring. *Baillieres clinical Obstet Gyncol*; 10(2): 307-24.
- Han B.K, Towbin R.B., De Courten — Myers, G. et al (1989.):** Reversal sign on CT : Effect of anoxic / ischemic cerebral injury in children *Am. J. Neuroradiol* 10:1191-1198.
- Handrahan JD, Cox IJ, Edwards AD, Cowan FM, Sargentoni J, Bell JD, Bryant DJ, Rutherford MA and Azzapordi D (1998):** Persistent increases in cerebral lactate concentration after birth asphyxia *pediatr Res*: 44(3): 304-11.
- Hankins GD, Koen S, Gei AF et al., (2002):** Neonatal organ system injury in actue birth asphyxia sufficient to result in neonatal encephalopathy. *Obstet Gynecol*: 99: 688-91.
- Hanlon Lundberg KM, Kirby RS, (1999):** Nucteated red blood cells as a marker of acidemia in term neonates. *Am J Obstet Gynecol*; 181: 196-201.
- Hanrahan JD, Cox IJ, Edwards AD, et al., (1998):** Persistent Increases in cerebral lactate concentration after birth asphyxia. *Pediatric Res*. 44: 304-311.
- Harachovy, R.A Mizrahi, E.M. and Kellaway, P (1990):** Electroencephalography of the newborn in Daly D.D. and Pedley T.A. (eds): *Current Practice of Clinical Electroencephalography*, 2nd ed. New York, Raven press 1990.
- Hein HA. (1993):** The use of sodium bicarbonate in neonatal resuscitation: help or harm? *Pediatrics* 91: 496-7.

- Hendrik J, Ter Horst, Constanze Sommer I, Klasien A, Bergman, Johanna M, Fock, Tiemen W, Van Weerden, and Arend F, Bos, (2004):** Prognostic significance of amplitude —integrated EEG during the first 72 hours after birth in severely asphyxiated neonates. *Pediatr Res.* 55: 18.
- Henrik Hagberg and Carina Mallard, (2000):** Antenatal brain injury: aetiology and possibilities of prevention. *Semin Neonatal*; 5: 41-51.
- Hill A (1989):** Clinical assessment of perinatal asphyxia in the newborn Canadian Medical Association J., 141 (Suppl) 53-58.
- Hiroyuki Ichiba, Hiroshi Tamai, Hirokuni Negishi Toru, Ueda Tue-Jung Kim Yutaka Sumida, Yukihiro Takahashi Hideshi Fujinaga and Hirotaka Minami for the Kansai Magnesium Study Group (2002):** *Pediatrics International*, 44, 505-509.
- Holmes GL and Kull LL (1990):** Neonatal seizures. *Am J EEG Technol*; 30: 281-308.
- Holmes GL, (2002):** Seizure-induced neuronal injury- animal data neurology: 59: S3-S6.
- Holt S, Guiderson M, Joyce K, Nayini NR, Eyster GI, Garitano AM, Zonia C, Krause GS, Aust SD, White BC, (1986):** Myocardial tissue: iron delocalization and evidence for lipid peroxidation after two hours of ischemia. *Ann Emerg Med* 15: 1155-9.
- Hope FL, Gould SJ, Howard S, et al., (1988):** Precision of ultrasound diagnosis of pathologically verified lesions in the brains of very preterm infants. *Dev. Med. Child neurol.*, 30: 457-471.

- Hwang J, Krebs C, Huynh BH, Edmondson DE, Theil EC, Penner-Han JE, (2000):** A short Fe-Fe distance in peroxidferric ferritin: Control of Fe substrate versus cofactor decay? *Science* 287: 122-125.
- Ikida T, Choi BH, Yee S, (1999,):** Oxidative stress, white matter damage and intrauterine asphyxia in fetal lambs. *Int J Dev. Neurosci*; 17: 1-14.
- International guidelines for neonatal resuscitation.** An excerpt from the guidelines 2000 for cardiopulmonary resuscitation and emergency cardiovascular care: International Consensus on Science. *Pediatrics*; 106: 29.
- Isabelle A. W, and Richard LB (1991):** Assessment of fetal well being. In *schauffers disease of the newborn*, Avery and Tacusch (eds) 6th ed., P81.
- James A Low, (2004):** Determining the contribution of asphyxia to brain damage in the neonate. *J Obstet Gynaecol. Res.* Vol. 30, No. 4: 276-286.
- James A, Low, Constadina Panagiotopoulos, and Jane Derrick BA, (1995):** Newborn complications after intrapartum asphyxia with metabolic acxidosiis in the preterm fetus. *Am J. Obstet Gynecol*; 172: 805-10.
- James A, Low, Heather Pickersgill, Helen Killen and Jane E, Derric BA, (2000):** The prediction and prevention of intrapartum fetal asphyxia in term pregnancies. *Am J Obstet Gynecol*: 184: 724-30.

- Jan Milson, Lars Ladfors Klara Hiringer, Aimon Nikiasson, Antlers Odeback and Eva Thornberg, (2002):** Influence of maternal, obstetric and fetal risk factors on the prevalence of birth asphyxia at term in a Swedish urban population. *Acta Obstet Gynecol scand*; 81: 909-917.
- Jefferey M. perlman, MB, ChB (2006):** Intervetnion Strategies for Neonatal Hypoxic-Ischemic Cerebral Injury. *Clin Ther*; 28: 1353-1365.
- Jenkin G, Ward J, Hooper SB, O'Connor AE, De Krester DM, Wallace EM, (2001):** Feto-Placental hypoxemia regulates the release of activin A and postaglandin E2. *endocrinology*; 142: 963-6.
- Jensen FE, Applegate CD, Holtzman D, Belin T, and Burchfiel J (1991):** Epileptogenic effects of hypoxia on immature rodent brain. *Ann neurol*; 29: 629-637.
- Johnson JH and Goldman J (1990):** Developmental assessment in clinical child psychology. New York: Pergamon Press.
- Karlowicz MG, Adelman RD. (1995):** Non oliguric and oliguric acute renal failure in asphyxiated term neonates *pediatr. Nephrol.* 718-22.
- Kaya H, Oral B, Ditrich R, Ozkaya O, (2000):** Lipid peroxidation in umbilical arterial blood at birth: The effect breech delivery. *Br .J Obstet Gynecol*; 107: 982-986.
- Ken-Dror A, Pratt H, Zeltzer M, Sujov P, Katzir J, Benderley A. (1995):** Auditory brainstem evoked potentials to clicks at different presentation rates: estimating maturation of preterm and full-term neonates. *Electro-encephalogr Clin Neurophysiol*: 68: 209-18.

- Kime R Gibson A, Yong W, Hider R, Powers H, (1996):** Chromatographic method for the determination of non-transferrin-bound iron suitable for use on the plasma and bronchoalveolar lavage fluid of preterm babies. *Clin Sci (Colch)*; 91: 633-638.
- Kinsbourne M and Graf WD (2001):** Disorders of mental development. In : Menkes JH, Sarnat HB, eds. *Child neurology*, 6th ed. Philadelphia: Lippincott Williams & Wilkins; 1155-1211.
- Klawitter, P. Morales, D. Bustamante, M. Goiny, and M. Herrera-Marschitz (2006):** Plasticity of the central nervous system (CNS) following perinatal asphyxia: Does nicotinamide provide neuroprotection? *Amino acids* 31: 377-384.
- Klawson TF, Vannucci SJ, Wang GM, et al., (1999):** Hypoxia-ischemia. Induced apoptotic cell death correlates with IGF- I mRNA Decreases in neonatal rat brain. *Biol Signals Recept.* 8: 281- 293.
- Klebermass K, Kuhle S, Kohlhauser Vollmuth C, Pollak A, Veninger M, (2001):** Evaluation of the cerebral function monitor as a tool for neurophysiological surveillance in neonatal intensive care patients. *Childs. Nerv Syst.* 17 :544-550.
- Kohlhauser C, Mosgoeller W, Hoeger H, et al., (1999):** Cholinergic. monoaminergic and glutamatergic changes following perinatal asphyxia in the rat. *Cell Mol Life Sci*, 55: 1491-1501.
- Komara JS, Nayini NR, Bialick HA, Indnieri RJ, Evans A T, Garritano AM, Hoehner TJ, Jacobs WA, Huang RR, Krause GS, white BC, Aust SD, (1986):** Brain iron delocalization and lipid peroxidation following cardiac arrest. *Ann Emerg Med* 15: 384-8.

- Kotagal S., Toce S.S., Rotagal P., et al (1983):** Symmetric bithalamic and striatal hemorrhage following perinatal hypoxic in a term infant. *Comput. Assist. Tomogr.* 7:353- 355.
- Kowaltowski AJ, Vercesi AE, (1999):** Mitochondrial damage induced by conditions of oxidative stress. *Free Rad Biol Med*: 26: 463-71.
- Kraig R, Pulsinelli W and Plum F (1986):** Carbonic acid buffer changes during complete brain ischemia *Am J Physiol* ; 250: R 348
- Krebs HB, Petres RE, Dunn U, (1981):** Intrapartum fetal heart rate monitoring, 5 Fetal heart rate patterns in the second stage of labor. *Am J Obstet Gynecol*; 140: 435.
- Kumar F; Katariya S; Sridhar K, (2001):** Postasphyxial encephalopathy in preterm neonates. *Indian J. Pediatr*, 68 (12) 1121-5.
- Lam BCC, Yeung CY, (1999):** Surfactant lavage for meconium aspiration syndrome: a pilot study. *pediatrics*; 103-1014-8.
- Laurel M, Bear, (2004):** Early identification of infants at risk for developmental disabilities. *Pediatr Clin N Am.* 51: 685-701.
- Leikin E, Verma U, Klein S, Tejani N, (1996):** Relationship between neonatal nucleated red blood cell counts and hypoxic-ischemic injury. *Obstet Gynecol*; 87: 439-43.
- Leth H, Toft F, Pryds O, (1995):** Brain lactate in preterm and growth retarded neonates. *Acta pediatr*; 84: 495-9.
- Leven MI, Sands C, Grindulis H, Moore JR, (1986):** Comparison of two methods of predicting outcome in perinatal asphyxia. *Lancet* 67: 68.
- Levene, MI, Evans DJ, Mason S, Brown J, (1999):** An international network for evaluating neuroprotective therapy after severe birth asphyxia. *Semin Perinatol*, 23: 226-233.

- Lindeman JH, Lenijes EG, van Zoeren-Grobbe D, Berger HM, (2000):** Postnatal changes in plasma ceruloplasmin and transferrin antioxidant activities in preterm babies. *Biol Neonate*; 78: 73-76.
- Lipp-Zwahlen AE, Deonna T, Chrzanowski R, et al, (1985):** Temporal evolution of hypoxic —ischemic brain lesion in asphyxiated full-term newborns as assessed by computerized tomography. *Neuroradiology*; 27: 138-44.
- Luckasson RL, Coulter DL, Polloway EA, et al. (1992):** Mental retardation definition, classification and systems of supports. Washington: AAMR.
- Lupton A, Lupton MB, Hill A, Roland EH, Whitfield MF, and Flodmark O (1988):** Brain swelling in the asphyxiated term newborn: Pathogenesis and outcome. *pediatrics*; 82: 139-46.
- Madle C, Kramer L, Domanovits H, et al., (2000):** Improved outcome prediction in unconscious cardiac arrest survivors with sensory evoked potentials compared with clinical assessment. *Crit Care Med*; 28: 721-6.
- Maier RF, Bohme K, Dudenhausen JW, Obladen M, (1993):** Cord blood erythropoietin in relation to different makers of fetal hypoxia. *Obstet Gynecol*; 81: 575-80.
- Mallard EC, Williams CE, Gun AJ et al., (1993):** Frequent episodes of brief ischemia sensitize the fetal sheep brain to neuronal loss and induce striatal injury *Pediatr Res*. 33: 61-5.
- Manning FA, Harman C'R, Morrison I, Menticogluo SM, Lange JR, Johnson JM, et al (1984):** Fetal assessment based on fetal

biophysical profile scoring. An Analysis of perinatal morbidity and mortality. Am J Obstet Gynecol; 150: 245.

Manniro, F.I and Trauner, D.A (1983): Stroke in neonates J. Pediatr. 102:605 -610.

Mari K, Zarifi, Loukas G, Astrakas, Ti, Young Poussaint, (2002): Prediction of adverse outcome with cerebral lactate level and apparent diffusion coefficient in infants with perinatal asphyxia. Radiology; 225-859-870.

Maria Gieron-Korthals and Jose Colon, (2005): Hypoxic ischemic encephalopathy in infants: new challenges. Fetal and pediatric pathology; 24: 105-120.

Marianne Thoresen and Andrew Whitelaw (2005): Therapeutic hypothermia for hypoxic- ischemic encephalopathy in the newborn infants current opinion in neurology, 18: 111-116.

Marin –Padilla M (1997): Pathology and pathogenesis of secondary epilepsy to hypoxic- ischemic encephalopathies. Rev Neuro; May 1,25 (141): 673-82.

Mark Scher, (2001): Perinatal asphyxia: Timing and mechanism of injury in neonatal encephalopathy. Current Neurology and neuroscience Reports; 1: 175-184.

Marret, S. Parain D, Jeannot , E, et al (1992): Positive rolandic sharp waves in the EEG of the premature newborn- a five - year prospective study . Arch Dis Child 67:948-951.

Martin -Ancel A, Garacia-Alix A, Cabanas FGF, et al., (1995): Multiple organ involvement in perinatal asphyxia. J Pediatr: 127-786-93.

- Marzocchi B, Paffetti F, Longini M, Gioia D, Magi B, Pallini V, et al., (2001):** Biomarkers of oxidative stress in preterm newborns. *Biol Neonate*; 80: 104A.
- Matute Sanchez-Gomez MV, Martinez-Millen L, (1997):** Glutamate receptor-mediated toxicity in optic nerve oligodendrocytes. *Proc Natl Acad Sci USA*; 94: 8830-8835.
- Mayes PA (2000):** Glycolysis and the oxidation of pyruvate In : Harper's biochemistry. Murray RK, Granner DK, Mayes PA and Podwell VW (eds) 25th edition, Appleton and Lange California. pp: 190.
- McBride MC, Laroia N, Guillet R, (2000):** Electrographic seizures in neonates correlate with poor neurodevelopment outcome. *Neurology*; 55: 506-513.
- McCardle CB, Richardson CJ, Hayden CK et al., (1987):** Abnormalities of the neonatal brain: MR imaging 11 hypoxic ischemia brain injury. *Radiology*; 163: 395-403.
- Meek JH, Elwell CE, McCormick DC, et al., (1999):** Abnormal cerebral hemodynamics in perinatally asphyxiated neonates related to outcome. *Arch Dis Child Fetal Neonatal Ed*; 81: F110-F115.
- Mehmet H, Yue X, Squire MV, et al., (1994):** Increased apoptosis in the cingulate sulcus of newborn piglets following transient hypoxia-ischemia is related to the degree of high energy phosphate depletion during the insult. *Neurosci Lett*; 181: 121-125.
- Meisun, Wei-Lin Wang, Wei Wang, De-Liang Wen, Hid Zhang and Yu Kun Han, (2001):** Gastroesophageal manometry and 24-hour double pH monitoring in neonates with birth asphyxia. *World J gastroenterol*; 7 (5): 695-697.

- Melltis E, Holden K, Freeman J, (1982):** Neonatal seizures 11.A multivariate analysis of factors associated with outcome. *pediatric* 70: 177-85.
- Mercuri E, Guzzett A, Haataja L, et al., (1999):** Neonatal neurological examination in infants with hypoxic-ischemic encephalopathy: correlation with MRI findings. *Neuropediatrics*; 30: 83-89.
- Mercuri E, Ricci D, Cowan FM, et al., (2000):** Head growth in infants with hypoxic-ischaemic encephalopathy; correlation with neonatal magnetic resonance imaging. *Pediatrics*; (In press).
- Michelle de Haan, John S. Wyatt, Simon Roth, (2006):** Brain development after asphyxia at term birth. *Developmental Science* 9: 4,pp350-358.
- Mireles LC, Lum MA, Dennery PA, (1999):** Antioxidant and cytotoxic effects of bilirubin on neonatal erythrocytes. *Pediatr. Res*; 45: 355-362.
- Mizock BA and Falk JL, (1992):** Lactic acidosis in Critical illness *critic Core Med*, 20: 80-93.
- Moison RWM, Palinckx JJS, Roest M, Houdkamp E, Berger HM, (1993):** Induction of lipid peroxidation of pulmonary surfactant by plasma of preterm babies. *Lancet* 341: 79-82.
- Nakai A, Asakura H, Tanichi Y, et al., (2000,):** Effect of a phenol-N-etr-Burtyl nitron of fetal cerebral energy metabolism during intrauterine ischemia and reperfusion in rats. *Pediatr Res*: 47: 451-456.
- Nath, KR; Nork M; Hass D et al (2002):** CT and MR patterns of hypoxic ischemic brain damage following perinatal asphyxia. *J. Trop. Paediatr.*, 48 (1): 5-9.

- Nathan L, Leveno KJ, Carmody TJ, Kelly MA, Sherman ML, Meconium, (1994):** A perspective on an old obstetric hazard. *Obstet Gynecol*; 83: 328.
- Nathoo KJ, Chimbra THK, Mtimavalye LAR, (1990):** Mortality and immediate morbidity in term babies with low Apgar scores in Zinbabwe. *Ann Trop Paediatrc*; 10: 239-244.
- Natsume J, Watanabe K, Kuno K, Hayakawa F, Hashizume Y, (1995):** Clinical, neurophysiologic and neuropathological features of an infant with brain damage of total asphyxia type (myers). *Pediatric Neurol*; 13: 6 1-4.
- Neefjes VME, Evelo CTA, Baars LGM, Blanco CE, (1999):** Erythrocyte glutathione S transferase as a marker of oxidative stress at birth, *Arch Dis Child Fetal Neonatol Ed*, 81: F130-133.
- Nelson K and leviton A, (1991):** How much of neonatal encephalopathy is due to birth asphyxia? *American Journal of disease in childhood* 145: 1325-1331.
- Nelson KB, Ellenberg JH, (1981):** Apgar scores as predictors of chronic neurologic disability. *Pediatrics* 68: 36-44.
- Nicole Nagdyman, Ingrid Grimmer, Tristess Scholz,. Chirstian Muller, (2003):** Predictive value of Brain-specific proteins in Serum for neurodevelopmental outcome after birth asphyxia. *Pediatr. Res* 54: 270-275.
- Nicole Nagdyman, Wolfgang Komen, Hae-Kyung Ko, (2001):** Early Biochemical Indicators of hypoxic-ischemic encephalopathy after birth asphyxia. *Pediatr. Res.* 49: 502-506.

- Nordstrom L, Ingemarsson I and Westgren M, (1996):** Fetal monitoring with lactate Baillieres clinical Obstet Gynecol., 10 (2) 225-42.
- Nordstrom L, Shuas, Roy A and Arulkumaran S, (1998):** Quality assessment of two lactate test strip methods suitable for Obstetric use. J. Perinatal Medicine, 26(2): 83-8.
- Nwaesei, CG., Pape K.E Martin D.J et al (1984):** Periventricular infarction diagnosed by ultrasound A postmortem correlation J. Pediatr 105:106-110.
- Palmer C, Menzies SL, Roberts RL, Pavlick G, Connor JR, (1999):** Changes in iron histochemistry after hypoxic-ischemic brain injury in the neonatal rat. J Neurosci Res; 56: 60-71.
- Palmer C, Roberts RL, Bero C, (1994):** Deferoxamine post treatment reduces ischemic brain injury in neonatal rats, stroke 25: 1039-45.
- Panter SS, Braugher JM, Hall ED, (1992):** Dextran coupled deferoxamine improves outcome in a murine model of head injury. J Neurotrauma 9: 47-53.
- Pasternak, J.F. (1987):** Parasagittal infarction in neonatal asphyxia. Ann. Neurol 21:202-204.
- Perlman JM, Tack ED, Martin T, Shackelford G, Amon E, (1989):** Acute systemic organ injury in term infants after asphyxia. Arch Pediatr Adolesc Med (Am J Dis Child); 143: 6 17-20.
- Perlman JM, Adcock L, dewitt S, et al., (1999):** Early identification of infants at highest risk for abnormal outcome secondary to intrapartum hypoxia ischemia (HI)- Texas Regional Survey. Pediatr Res; 45: 218A (Abstract).

- Perlman JM, Risser R, (1996):** Severe fetal acidemia: Neonatal neurologic features and short-term outcome. *Pediatr. Neurol.* 9: 277-82.
- Perrone S, Bracci R, and Buonocore G, (2002):** New biomarkers of fetal neonatal hyoxic stress. *Acta paediatr Supp*; 438: 135-138.
- Perrone S, De Marco-L, Ciccoli L, Tanganelli D, Rossi V, Bracci R, et al., (2001):** Intraerythrocyte non protein bound iron release and diffusion after in vitro hypoxic reoxygenation. *Biol Neonate* 80: 325.
- Perrone S, Florio F, Longini M, Tanganelli D, Luisi S, Petraglia F, (2002):** Activin A in newborn with and without hypoxia at birth *Pedatric Res*; 51: 255A.
- Petraglia F, Sawchenko P, Lim ATW, Vale W, (1987):** Loclaization. secretion and action of inhibin in human placenta. *Science*; 237: 189-92.
- Pezzani, C. Radvanyi-Bouvet M.F, Relier J,P, et al (1986):** Neonatal electroencephalography during the first twenty four hours of life in full-term newborn infants. *Neuropediatrics* 1 7: 11- 1 8
- Phelan, JP(1988):** Antepartum fetal evaluation. In: critical care obstetrics clark, S. L, Phelan JP, and Collon DB (eds) Medical Economics book.
- Picquard F, Hsiung R, Mattauer M, Schaefer A, Haberey P, (1988):** The validity of fetal heart rate monitoring during the second stage of labor. *Obstet Gynecol* 72: 746.
- Pierrat V, Haourari N, Liska A, Thomas, D, Subtil P, Truffert, (2003):** Prevalence, causes, and outcome at 2 years of age of newborn

encephalopathy: population based study. Arch Disc Child Fetal Neonatal Ed, 90 F 257-F261.

Pinto LC, Giliberti P, (2001): Neonatal seizures: background EEG activity and the electroclinical correlation in full-term neonates with hypoxic-ischemic encephalopathy. Analysis by computer — synchronized long-term polygraphic video-EEG monitoring. Epileptic Disorder 3: 125-132.

Polnad RL and Freeman RK (1992): Guidelines for perinatal care: Relationship between perinatal factors and neurologic outcome. am Acad of Pediatr: PP. 221.

Preman JM, Tack ED, (1988): Renal injury in the asphyxiated newborn infant. Relationship to Neurologic outcome. J. Pediatric; 113 (5): 875-9.

Radi R, Beckman JS, Bush K, Freeman BA, (1991): Peroxynitrite oxidation of sulfhydryl: the cytotoxic potential of superoxide and nitric oxide. J Biol Chem 266: 4244-50.

Radolfo Bracci, Serafina Perrone, Giuseppe Buonocore, (2001): Red blood cell involvement in Fetal/neonatal hypoxia. Biol Neonate 79: 210-212.

Raju TN(1991): Cerebral Doppler studies in the fetus and newborn infant. J. Pediatr., 119: 165-174.

Ralph Heinz, In Re: Barkovich AJ, MR and CT (1992): Evaluation of profound neonatal and infantile asphyxia: A JNR Am J Neuroradiol; 13: 959-972.

Ramin KD, Leveno KJ, Kelly MS, (1996): Amnionic fluid meconium: a fetal environmental hazard Ostet Gynecol; 87: 181.

- Rao R, De Ungria M, Sullivan D, Wu P, Wobken JD, Nelson CA, Georgieff MK, (1999):** Perinatal brain iron deficiency increases the vulnerability of rat hippocampus to hypoxic ischemia insult. *J Nutr.* 129: 199-206.
- Read JA, Miller FC, (1977):** Fetal heart rate acceleration in response to acoustic stimulation as a measure of fetal well-being. *Am J Obstet Gynecol*; 129: 512.
- Rebertson CM and Finer NN (1985):** Term infants with hypoxic-ischemic encephalopathy: Outcome of 3-5 Years. *Dev Med Child Neurol*; 27: 473-484.
- Repka T, Shalev O, Reddy R, Yuan J, Abraomov A, Racmilewitz EA, Low PS, Hebbed R, (1993):** Non random association of free iron with membrane of sickle and B. Thalassemic erythrocytes. *Blood*; 82: 3204-3210.
- Resenbloom L, (1999):** Dyskinetic cerebral pulsy and birth asphyxia. *Dev Med Child Neurol*; 36: 285-289.
- Rickman DW, Nacker RE, Rickman CB, (1999):** Characterization of the cell death promoter, BAD in the developing rat retina and forebrain, *Dev Brain Res* 115: 41-47.
- Rivkin MJ and Volpe JJ (1996):** Aasphyxia and brain injury : In Spitzer AR (Ed.) *Intensive Care of Foetus and Newborn*; Mosby, St. Louis: 59 : 685 - 95
- Robertson CM, Finer NN, (1993):** Long term follow up of term neonates with perinatal asphyxia. *Clin Perinatol.* 20 (20): 483-500.

- Robertson CM, Finer NN, Grace MG (1989):** School Performance of survivors of neonatal encephalopathy associated with birth asphyxia at term. *J pediatric* 114: 753-760.
- Robertson RL, Ben Sira L, Barnes PD, et al., (1997):** MR lin-scan diffusion-weighted imaging in term neonates with perinatal ischemia *AJNR*, 20: 1658-1670.
- Rodriguez E, Poskitt K, Roland EH et al. (1995):** Acute hypoxic ischemic basal ganglia injury in the term Neonborn: computed tomography and clinical syndrome. *Ann Neurol* 138: 544A.
- Roland EH and Hill A (1995):** Clinical aspects of perinatal hypoxic-ischemic brain injury. *Sem pediatr Neurol*; 2: 57.
- Rosenthal RE, Charderbhan R, Marchal G, Fiskum G, (1992):** Prevention of post-ischemia brain lipid conjugated diene production and neurological injury by hydroxyethyl starch-conjugated deferoxamin. *Free Rad Biol. Med* 12: 29-33.
- Roth, S.C, Baudin, .J., MocCormick, D.C., et al (1993):** Relation between ultrasound appearance of the brain of very preterm infants and neurodevelopmental impairment at eight years. *Dev. Med. Cud. Neurol.* 35:755-768.
- Rutherford MA, Pennock JM, Counsell SJ, et al., (1998):** Abnormal magnetic resonance imaging in the internal capsule predicts poor neurodevelopmental outcome in infants with HIE. *Pediatrics*; 102: 323-8.
- Rutherford MA, Pennock, I Schwieso J, et al., (1996):** Hypoxic ischaemic encephalopathy: Early and late magnetic resonance imaging findings in relation to outcome. *Arch Dis. Child*: 75: F145-151.

- Ruthrford MA, Pennock JM, Counsel! SJ et al., (1998):** Abnormal magnetic resonance signal in the internal capsule predicts poor neurodevelopmental outcome in infants with hypoxic-ischemic encephalopathy. *Pediatrics*; 102: 323-8.
- Sachin Shah, Mark Tracy Fracy John Smyth, (2004):** Postnatal lactate as an early predictor of short-term outcome after intrapartum asphyxia. *Journal of perinatology*, 24: 16-20.
- Sadovsky E, Ohel G, and Havazelet H, (1983):** The significance of decreased fetal movements. *Acta Obstet. Gynecol. Scand.*, 62:409.
- Sarnat HB, Sarnat MS, (1976):** Neonatal encephalopathy following fetal distress : a clinical and electroencephalographic study. *Arch. Neurol.* 33: 696-705.
- Schafer FQ, Qian SY, Buettner GR, (2000):** Iron and free radical oxidations in cell membranes. *Cell Mol Biol (Noisy Le-Grand)*: 46: 657-662.
- Scher M.S. and Beggarly M (1989):** Clinical significance of focal periodic discharges in neonates *J Child Neurol* 4:175-185.
- Scherjon N, Samolders-Dettas H and Kok JH (1993):** The (Brain sparing) effect: antenatal cerebral Doppler finding in relation to neurologic outcome in very preterm infants. *Am. J. Obstet. Gynecol.*, 169 (1): 169-175.
- Schmidt H, Grune T, Muller R, Siems WG, Wauer RR, (1996):** Increased levels of lipid peroxidation products malondialdehyde and 4-hydroxynonanal after perinatal hypoxia. *Pediatr. Res.*; 40: 15-20.

- Scott H, (1976):** Outcome of very severe birth asphyxia. Arch Dis Child 51: 712-6.
- Seetha Shankaran Eunice Woldt, Thomas Koepke, (1991):** Acute neonatal morbidity and long-term central nervous system sequelae of perinatal asphyxia in term infants. Early Human development, 25: 135-148.
- Seetha Shankaran, MD, (2002):** The postnatal management of the asphyxiated term infant. Clin Perinatol 29: 675-692.
- Seri I, (1995): Cardiovascular, renal and endocrine actions of dopamine in neonates and children.** J Pediatr. 126: 333-44.
- Shah F, Riphagen S, Beyene J, Penman M, (2004):** Multiorgan dysfunction in infants with post-asphyxial hypoxic-ischaemic encephalopathy. Arch Dis Child Fetal neonatal Ed:89: F 152- F 155.
- Shah PS, Raju NV, Beyene J and Perlman M, (2003):** Recovery of metabolic acidosis in term infants with postasphyxial hypoxic ischemic encephalopathy. Acta Paediatr. 92: 941-947.
- Shah S, Fernandez AR, and Chirila D (2001):** Role of brain SPECT in neonates with HIE and its correlation with neurodevelopmental outcome. Indian Pediatr Jul; 38(7): 705-13.
- Sheldon RA, Partidge JC, Formal DM, (1992):** Postischemic hyperglycemia is not protective to the neonatal rat brain. pediatr; Res. 32: 489-93.
- Siesjo BK, (1988):** Acidosis and ischemic brain damage. Neurochem Pathol 9:31-88.

- Sigel, M.J. Shachkelford, G.D., Penman, J.M., et al., (1984):** Hypoxic ischemic ultrasound Radiology 152:395 -399.
- Silva SD, Hennebert N, Denis R, Wayenberg JL, (2000):** Clinical value of single postnatal lactate measurement after asphyxia. Act Paediatr; 89: 320-3.
- Silverstein FS, Barks JDE, Hagan P, Luix, (1997):** Cytokines and perinatal brain injury. Neurochem Int; 30: 375-383.
- Simeonsson RJ and Simeonsson NW (2001):** Developmental surveillance and intervention. In: Hoekelman RA, Adam HM, Nelson NM, Weitzman ML, Wilson MH, eds. Primary pediatric care, 4th ed. St. Louis: Mosby, 274-282.
- Smith ME, Van Der Maesen K, Somera FP, (1998):** Macrophage and microglia response to cytokine in vitro. Phagocytic activity. Proteolytic enzymes release and free radical production. J Neurosci Res; 54: 68-78.
- Snyder EY and Cloherty JP (1998):** Neurology: Perinatal Asphyxia In: John P Cloherty and Ann Stark (Eds.): Manual of neonatal care: 4th edition; Little Brown and White Co; Boston ; 27: 522 -530.
- Sohmer H, Freeman S, Gafni M, Goitein K, (1986):** The depression of the auditory nerve-brainstem evoked response in hypoxemia mechanism and site of effect. Electroencephalogr. Clin Neurophysiol; 64: 334-8.
- Sohmer H, Gold S, Cahani ill, Attias J, (1989):** Effects of hypothermia on auditory brain-stem and somatosensory evoked response. A model of a synaptic and axonal lesion. Electroencephalogr Clin Neurophysio; 74: 50-7.

- Soll RF, Dargaville F, (2000):** Surfactant for meconium aspiration syndrome in full term infants. Cochrane Library Disk. Issue.
- Sorg O, Ham TFW, Yu NC, Gruol DL, Bloom FE, (1997):** Inhibition of astrocyte glutamate uptake by reactive oxygen species : role of antioxidant enzyme. *Mol Med*; 3: 431-40.
- Suehiro E, Fujisawa H, Ito H, et al., (1999):** Brain temperature modifeis glutamate neurotoxicity in vivo. *J Neurotrauma* 16: 285-97..
- Sutton HC, Winterbourn CC, (1989):** On the participation of higher oxidative states of iron and copper in Fenton reactions. *Free Radic Biol Med*; 6: 53-60.
- Tao Yu MD, Department of pediatrics, The second university Hospital. Sichuan University, Chengdu, Liao Qung Kui; Qian Zhong Ming, (2003):** Effect of asphyxia on non-protein-bound iron. *Developmental medicine Child Neurology*, 45: 24-27.
- Taylor and Francis Ltd (2002):** Early predictors of adverse outcome in term infants with post-asphyxial HIE. *Acta Paediatrica*; 89: 343-347.
- Temple CJ, Dennis J, Carney (1995):** Neoantal seizures: long-term outcome and cognitive development among normal survivors. *Dev Med Child Neurol*; 37: 109-118.
- Throp JA, Dalidy GA, Yeomans ER, Meyer BA, and Parisi N(1996):** Umbilical cord blood gas analysis at delivery. *Am. J. Obstet. Gynecol.*, 175 (3): 5 17-522.
- Toat ML, Hellstrom —Westas L, Groenendaal F, Eken F, de Vries LS, (1999):** Amplitude integrated EEG 3 and 6 hours after birth full

term neonates with hypoxic —ischemic encephalopathy. Arch Dis Child Fetal neonatal ED 81: F19-F23.

Toet Mc, Van der Mei W, de Vries LS, Uiterwaal CSPM, van, Huffelen AC, (2002): Comparison between simultaneously recorded amplitude integrated electroencephalogram (cerebral function monitor) and standard electroencephalogram in neonates. Pediatrics; 109: 772-779.

Toffaletti JC (1996): Electrolytes In : Clinical Chemistry : Principles procedures, correlations Bishop ML Duben — Englirk JL and Fody EP (eds) 3rd edition, Lippincott, Boston; 12:272-274

Tohru Ogihara Kazurya Hirano, Hiromi Ogihara, Kiryo Misaski, (2003): Non-protein bound transition metals and hydroxyl radical generation in cerebrospinal fluid of newborn Infants with hypoxic ischemic encephalopathy. Pediatr. Res 53: 594-599.

Toth-Heyn P, Drukker A, Guignard JP, (2000): The stressed neonatal kidney: from pathophysiology to clinical management of neonatal vasomotor nephropathy. Pediatr. Nephrol; 14: 227-39.

Tropper PJ, and Fox HE (1982): Evaluation of antepartum fetal well-being by measuring growth. Clin. Perinatol, 9: 27 1-284.

Van Bel F, Shadid M, Moison RM, Dorrepaal CIA, Fontijn J, Monteiro L Van De Bor M, Berger HM, (1998): Effect of allopurinol on postasphyxial free radical formation, cerebral hemodynamics. and electrical brain activity. Pediatrics 101: 185-93.

Van Rooij MC, Toel MC, Ostredkar D, Van Huffelen A, Groenendaal F, de Vries S (2005): Recovery of amplitude integrated electroencephalographic background patterns within 24 hours of

perinatal asphyxia. *Arc Dis child Fetal Neonatal Ed.* 90: 245-251.

Venta M, Asensi M, Sastre J, et al., (2001): Resuscitation with room air instead of 100% oxygen prevents oxidative stress in moderately asphyxiated term neonates; *pediatric*; 107: 642-7.

Volpe J.J. and Pasternak, JF: parasagittal cerebral injury in neonatal hypoxic ischemic encephalopathy. (1977): Clinical and Neurological features. *J. pediatr*; 91: 472.

Volpe JJ (1995): Hypoxic-ischemic encephalopathy. In. Volpe J.J. (ed). *Neurology of the newborn.* 3rd edition. W.B. Saunders Company P: 211-364.

Volpe JJ (1995): *Neurology of the newborn,* 3rd ed philadellphia WB Saunders.

Volpe JJ (2001): *Neurology of the newborn* 4th edition. WB Saunders Co., Philadelphia

Volpe JJ, (1981): Hypoxic ischemia encephalopathy: Neuropathology and clinical aspects, in Schaffer AJ, Markowitz M (eds): *Neurology of the Newborn.* Philadelphia, WB Saunders CO. Vol. 22: 180-238.

Volpe JJ, (2001): Editor. Hypoxic-isehemic encephalopathy: clinical aspects. In: *Neurology of the newborn.* 4th edition. Philadelphia: WB Saunders Co.; 331-94.

Volpe JJ, (2001): Hypoxic ischemic encephalopathy: clinical aspects. In: *Neurology of the newborn* 4th ed. philadelphia: WB Saunders Company; 331-94..

- Volpe JJ, (2001):** Neurology of the newborn. WB. Saunders. Company. Philadelphia. pp. 201-202.
- Volpe JJ, (2001):** Specialized studies in the neurological evaluation. In: Volpe JJ. Editor. Neurology of the newborn. Philadelphia. PA: WB Saunders: p. 134-77.
- Volpe, J.J. (1987):** Brain death determination in the newborn . Pediatrics 80: 293 -297.
- Warash S, Gaa J, Siewert B, et al., (1995):** Acute human stroke studies by whole brain echo planar diffusion weighted magnetic resonance imaging Am Neurol; 37: 231-241.
- Wayenberg JL, Vemeylen D, Bormans J, Magrez P, Muller MF, and Pardou A (1994):** Diagnosis of severe birth asphyxia and early prediction of neonatal neurological outcome in term asphyxiated newborns. J Perinat Med 22:129-136.
- Whittle M, (2000): Is it time to abandon** cardiotocographic ECG analysis? Lancet 355: 422.
- Williams JW, (1993):** Willams Obstetrics, 1st ed. New York. Appleton.
- Willis F, summers J, Minutillo C, Hewitt I, (1997):** Indices of renal tubular function in perinatal asphyxia. Arch Dis Child Fetal Neonatal Ed., 77: F57:F60.
- Wiswell TE, (2001):** Handling the meconium —stained infant. Semin Neonatal 6: 225-31.
- Wolff F, (1997):** Obligatory monitoring methods during delivery. Z Geburtshilfe Neonatol; 201: 1-5.
- Wu DD, Lai M, Hughes PE, Sirimanne E, Gluckmann PD, Williams CE, (1999):** Expression of activin axis and neuronal rescue effects of

recombination actin A following hypoxic-ischemic brain injury in the infant rat. *Brain Res*; 835-369-78.

Wyatt JS (1997): Near-infrared spectroscopy in asphyxial brain injury. *Clin perinatol*; 20: 369-378.

Yagel JY Heitjan DF, Towgighi J and Vannucci RC (1992): Effect of Insulin-Induced and fasting hypoglycemia on perinatal hypoxic-ischemic brain damage. *Pediatr. Res.*, 31: 138-142.

Yeargin-Allsopp M, Murphy CC, Cordero JF, et al., (1997): Reported biomedical causes and associated medical conditions for mental retardation among 10 year old children, metropolitan Atlanta, 1985 to 1987. *Dev Med Child Neurol*; 39: 142-149.

Yu J, Shao L, Vaughan J, Valle W, Yu AL, (1989): Characterization of the potentiation effect of activin on human erythroid colony formation in vitro. *Blood*; 73: 952-60.

Zeinstra E, Fock JM, Begeer JH, van Weerden TW, Mauritis NM, Zweens MJ (2002): The prognostic value of serial EEG recordings following acute neonatal asphyxia in full-term infants. *Eur J Paediatr Neurol*; 5: 155-160.