

REFERENCES

References

- **Abraham, A.S.; Eylath, U. and Weinstein, M. (1977):**
Serum magnesium levels in patients with acute myocardial infarction.
N. Engl. J. Med., 296: 862-863.
- **Aggett, P.J. (1985):**
Trace elements in human nutrition.
J. Trop. Ped., 26 (2): 43.
- **Aicardi, J. and Chevrie, J.J. (1976):**
Febrile convulsions. Neurological sequelae and Mental retardation. In: Brain dysfunction in infantile febrile convulsions.
Brazier, M.A. and Coceani, F. (eds)., 247.
- **Andrews, J.; Boutroy, M.J.; Bianchetti, G.; Vert, P. and Morselli, P.L. (1991):**
Clonazepam in Neonatal Seizures: Dose Regimens and Therapeutic Efficacy.
Eur. J. Clin. Pharmacol., 40: 193-195.
- **Anonymous, (1975):**
Scientific and technical assessment report on manganese.
Environmental Protection Agency. EPA-600/6-74-002.
United States Government Publishing Office, Washington, D.C. 564-620.

- **Antony, J.H. and Hawke, S.H. (1983):**
Phenobarbital Compared with Carbamazepine in Prevention of Recurrent Febrile Convulsions: A Double-Blind Study. *Am. J. Dis. Childr.*, 137: 892-895.

- **Armand, V.; Rundfeldt, C. and Heinemann, U. (2000):**
Effects of retigabine (D-23129) on different patterns of epileptiform activity induced by low magnesium in rat entorhinal cortex hippocampal slices. *Epilepsia.*, 41 (1): 28-33.

- **Arteaga, R.; Herranz, J.L.; Valdizan, E.M. and Armijo, J.A. (1992):**
Y-Vinyl GABA (Vigabatrin): Relationship between dose, plasma concentrations, platelet, GABA-transaminase inhibition, and seizure seduction in epileptic children *Epilepsia.*, 33: 923-931.

- **Aschner, M. and Aschner, J.L. (1991):**
Mercury neurotoxicity: mechanisms of blood-brain barrier. *Neurosci. Bioch. Rev.*, 14: 169-176.

- **Ashraf, W.; Jaffar, M.; Mohammed, D. and Iqbal, J. (1995):**
Utilization of scalp hair for evaluating epilepsy in male and female groups in the Pakistan population. *Sci. Total Environ.*, 164 (1): 69-73.

- **Aubby, D.H.; Saggu, K.; Henner, N. and Quinn, P. (1988):**
Leukocyte glutamate dehydrogenase activity in patients with degenerative neurological disorders. *J. Neurosurg. And Psychia.*, 51: 893-902.

- **Barlow, P.J. and Kapel, M. (1979):**
Hair metal analysis and its significance to certain disease

conditions.

Second Annual Trace Minerals Health Seminar, Boston.,
12: 67-75.

- **Ben-Ari, Y. (1985):**

Limbic seizure and brain damage produced by kainic acid; mechanisms and relevance to human temporal lobe epilepsy.

Neuroscience., 14: 375-403.

- **Ben-Ari, Y.; Trembley, E.; Riche, D.; Ghilini, G. and Nequet, R. (1981):**

Electrographic, clinical and pathological alterations following systemic administration of kainic acid, bicuculline or pentetrazole: metabolic mapping using the deoxyglucose method with special reference to the pathology of epilepsy.

Neuroscience., 6: 1361-1391.

- **Benassi, E.; Besio, G.; Bo, G.P.; Cocito, L.; Maffini, M.; Mainardi, P.; Morselli, P.L. and Loeb, C.P. (1988):**

Preliminary observations on the activity of progabide administered as monotherapy in complex partial seizures.

Int. J. Clin. Pharmacol. Res., 8: 353-361.

- **Benninger, C.; Matthis, P. and Scheffner, D. (1986):**

Changes of the EEG in children with primary generalized epilepsy during the introduction of antiepileptic therapy with sodium valproate.

Epilepsia., 27: 587.

- **Berkovic, S.F.; Andermann, F.; Guberman, A.; Hipola, D. and Bladin, P.F. (1989):**

Valproate prevents the recurrence of absence status.

Neurology., 39: 1294-1297.

- **Bertz, R.J. and Howrie, D.L. (1993):**
Diazepam by continuous intravenous infusion for status epilepticus in Anticonvulsant Hypersensitivity Syndrome. *Ann. Pharmacoth.*, 27: 298-301.

- **Booker, H.E. (1989):**
Trimethadione. In: R. Levy,; R. Mattson,; B. Meldrum, J.K. Penry, and F.E. Dreifuss (eds).
Antiepileptic Drugs. 3rd ed. New York. Raven Press.

- **Bradford, H.F. (1993):**
Current views of the pathobiochemistry of epilepsy. *Mol. Aspects. Med.*, 9: 119-172.

- **Browne, T.R.; Dreifuss, F.E.; Dyken, P.R.; Goode, D.J.; Penry, J.K.; Porter, R.J.; White, B.G. and White, P.T. (1975):**
Ethosuximide in the treatment of absence (petit mal) seizures.
Neurology., 25: 515-524.

- **Browne, T.R.; Feldman, R.G.; Buchanan, R.A.; Allen, B.C.; Fawcettvickers, L.; Szabo, G.K.; Mattson, G.F.; Norman, S.E. and Greenblatt, D.J. (1983):**
Ethosuximide for complex partial seizures: efficacy, toxicity, clinical pharmacology and drug interactions.
Neurology., 33: 414-418.

- **Browne, T.R.; Mattson, R.H. and Penry, J.K. (1987):**
Vigabatrin for refractory complex partial seizures: Multicenter Single-Blind Study with Long-Term Follow-Up. *Neurology.*, 37: 184-189.

- **Burch, R.E. and Hahn, K.J. (1979):**
Tace elements in human nutrition.

Med. Clin. North. Am., 63 (5): 1057.

- **Burch, R.E. and Sullivan, J.E. (1976):**
Diagnosis of zinc, copper, and manganese abnormalities in man.
Med. Clin. North. Am., 60 (4): 655.
- **Butler, E.J. and Newman, G.E. (1956):**
The urinary excretion of copper and its concentration in blood of normal human adults.
J. Clin. Pathol., 9: 157.
- **Callaghan, N.; Kemy, K.S. and O'Neill, B. (1985):**
A prospective study between carbamazepine, phenytoin and sodium valproate as monotherapy in previously untreated and recent diagnosed patients with epilepsy.
J. Neurol. Neurosurg. and Psychiatr., 48: 639-644.
- **Canadian Clobazam Cooperative Group. (1991):**
Clobazam in treatment of refractory epilepsy: The Canadian Experience. A Retrospective Study.
Epilepsia., 32: 407-416.
- **Casey, C.E.; Hambidge, K.M. and Neville, M.C. (1985):**
Studies in human lactation: Zinc, copper, manganese, and chromium in human milk in the first month of lactation.
Am. J. Clin. Nutr., 41: 1193-1200.
- **Cavazzuti, G.B. (1980):**
Epidemiology of different types of epilepsy in school-age children of Modena, Italy.
Epilepsia., 21: 57-62.
- **Chadwick, D. (1992):**
Gabapentin.

Recent Advances in Epilepsy., 5: 211-222.

- **Chadwick, D. (1993):**
Seizures, epilepsy and other episodic disorders. In: Walton, J.: Brain's diseases of the nervous system, 10th edition. New York. Oxford University Press. 1: 702-727.
- **Chandra, E. (1988):**
Epilepsy in developing countries. In: A textbook of Epilepsy (eds). Lailaw, J.; Richens, A. and Oxley, J. . Churchill Livingstone. London and New York.
- **Chandra, S.V.; Shukla, G.S. and Saxena, K. (1985):**
Manganese-induced behavioral dysfunction and its neurochemical mechanism in growing mice. J. Neurochem., 33: 1217-1221.
- **Chawla, L.S. (1983):**
Study of trace elements (Zinc, Iron, Copper, and Magnesium) in normal healthy populations. J. Assoc. Physicians India., 31: 141-145.
- **Chittleborough, G.A. (1980):**
A chemist's view of the analysis of human hair for trace elements. Sci. Total Environ., 14: 53-75.
- **Claudio, O.I.; Ferchmin, P; Velisek, L. Sperber, E.F.; Moshe, S.L. and Ortiz, J.G. (2000):**
Plasticity of excitatory amino acid transporters in experimental epilepsy. Epilepsia., 41 (Suppl. 6): 104-110.
- **Collaborative Study Group. (1987):**
Monotherapy with valproate in primary generalized

epilepsies.
Epilepsia., 28: 8-11.

- **Collingridge, G.L. and Bliss, T.V. (1987):**
NMDA receptors: their role in long-term potentiation.
Trends. Neuroscience., 10: 288-293.
- **Commission, (1981):**
Commision on Classification and Terminology of the
International League Against Epilepsy: Proposal for revised
clinical and electroencephalographic classification of
epileptic seizures.
Epilepsia., 22: 489-499.
- **Commission, (1989):**
Commission on Classification and Terminology of the
International League Against Epilepsy. Proposal for revised
clinical and electroencephalographic classification of
epileptic seizures.
Epilepsia., 30: 268-278.
- **Cousins, R.J. and Failla, M.L. (1980):**
Cellular and Molecular aspects of mammalian zinc
metabolism and haemostasis. In: Zinc in the Environment.
Part II: Health effects. Niriagu, J.O. (eds)., John Wiley &
sons, New York. Chichester. Brisbane. Toronto., 121-135.
- **Cowan, L.D.; Bodensteiner, J.B.; Leviton, A. and
Doberty, L. (1989):**
Prevalence of the epilepsies in children and adolescents.
Epilepsia., 30 (1): 94-106.
- **Crawford, P.; Ghadiali, E.; Lane, R.; Blumhardt, L. and
Chadwick, D. (1987):**
Gabapentin as an Antiepileptic Drug in Man.
J. Neurol. Neurosurg. and Psychiatr., 50: 682-686.

- **Danks, D.J.; Campbell, B.E. and Steven, B.J. (1972):**
Menke's kinky hair syndrome: an inherited defect in copper absorption with widespread effects.
Pediat., 50: 188.

- **Daugbjerg, P.; Brems, M.; Mai, J.; Ankerhus, J. and Knudsen, F.U. (1990):**
Intermittent prophylaxis in febrile convulsions: Diazepam or valproic acid ?
Acta. Neurol. Scand., 82: 17-20.

- **Delgado-Escueta, A.V. and Enrile-Bascal, F.E. (1984):**
Juvenile myoclonic epilepsy of Janz.
Neurology., 34: 285.

- **Di Chiara, G. and Gessa, G.L. (1986):**
Glutamate as a neurotransmitter.
Advances in neurology. New York. Raven Press., 44: 1011-1031.

- **Donaldson, J. (1987):**
The physiological significance of manganese in brain: It's relation to schizophrenia and neurodegenerative disorders.
Neurotoxicology., 8: 451-462.

- **Dost, R. and Rundfeldt, C. (2000):**
The anticonvulsant retigabine potently suppresses epileptiform discharges in the low calcium and low magnesium model in the hippocampal slice preparation.
Epilepsy. Res., 38 (1): 53-66.

- **Dreifuss, E. (1990):**
Clinical Implications of the International Classification.
Epilepsia., 31: S 3 - S 10.

- **Duncan, J.S. (1987):**
Antiepileptic drugs and the electroencephalogram.
Epilepsia., 28: 259-266.

- **Dunlap, W.M.; James, J.W. and Hume, D.S. (1974):**
Anaemia and neutropenia caused by copper deficiency.
Ann. Intern. Med., 88: 478.

- **El-Afify, H. and Mostafa, M. (1981):**
A study of epilepsy in general hospital population. M. Sc.
Thesis. Faculty of Medicine. Ain Shams University. Coted
from Nasr, A.A., Cognitive impairment in epileptic children.
M.D. Thesis; Cairo University, 1993.

- **Elwan, O. (1991):**
Principles of Anticonvulsant Therapy.
Egyptian Epilepsy Bulletin., 1: 15-23.

- **Eriksson, H.; Morath, C. and Heilbronn, E. (1984):**
Effects of manganese on the nervous system.
Acta. Neurol. Scand., 70: 89-93.

- **Erwin, H.; Seidl, R.; Freilinger, M.; Mali, C and Herkner,
K. (1996):**
Haematologic manifestations and impaired liver synthetic
function during valproate monotherapy.
Brain. 18: 105-109.

- **Familusi, J.B. (1985):**
Preliminary Nigerian Experience in the use of
Carbamazepine in Children with Intractable Seizures.
Epilepsia., 26: 10-14.

- **Farrell, K. (1986):**
Benzodiazepines in Children with Epilepsy.

Epilepsia., 27: 45-51.

- **Faught, R.N.; Sachdeo, R.C.; Remler, M.P.; Chayasirisobhen, S.; Iragui-Madoz, V.J.; Ramsay, R.E.; Sutula, T.P.; Kanner, A.; Harner, R.N.; Kuzniecky, R.; Kramer, L.D.; Kamin, M. and Rosenberg, A. (1993):**
Felbamate Monotherapy for Partial-Onset Seizures: An Active-Control Trial.
Neurology., 43: 688-692.
- **Fiatarone, M.A.; Marks, E.C. and Meredith, C.N. (1989):**
High intensity strength training in nonagenarians effects on skeletal muscles.
Clin. Res., 37: 330.
- **Finocchiaro, J.; Taroni, F. and Di-Donato, S. (1986):**
Glutamate dehydrogenase. In: olivo-ponto-cerebral atrophy in leukocytes, fibroblasts, and muscle mitochondria.
Neurology., 36: 550-553.
- **Flink, E.B. (1976):**
Magnesium deficiency and magnesium toxicity in man. In: Trace Elements In Human Health And Disease.
Prasad, A.S. (ed)., Academic Press. New York., 2: 1-21.
- **Floren, K.L.; Graves, N.M.; Leppik, I.E.; Remmel, R.P.; Margul, B.L. and Doose, D.R. (1989):**
Pharmacokinetics of Topiramate in Patients with Partial Epilepsy Receiving Phenytoin or Valproate.
Epilepsia., 30: 646.
- **Fogg, G.E. and Foster, A.C. (1983):**
Amino acid neurotransmitters and their pathways in the mammalian central nervous system.
Neuroscience. 9: 701-719.

- **Friis, M.L.; Kristensen, O.; Boas, J.; Dalby, M.; Deth, S.H.; Gram, L.; Mikkelsen, M.; Pedersen, B.; Sabers, A.; Worm-Petersen, J.; Andersen, D. and Jensen, P.K. (1993):**
Therapeutic Experiences with 947 Epileptic Out-Patients in Oxcarbazepine Treatment.
Acta. Nneuro. Scand., 87: 224-227.

- **Gastaut, H.; Gonclaves, E.; Silva, G.E. and Fernandez, G.R. (1975):**
Relative frequency of different types of epilepsy: A study employing the Classification of the International League Against Epilepsy.
Epilepsia., 16: 457.

- **Gella, F.G. (1985):**
Quantitative determination of aspartate aminotransferase in serum and leukocytes.
Clin. Chim. Acta., 153: 241-247.

- **Gerstle de Pasquet, E.; Scaramelli, A.; Pineyrua de Caceres, M.; L'Heritier, C.; Feldman, S.; Santana, R.; Aguilar, J.; Musch, B. and Morselli, P.L. (1991):**
Double-Blind, Placebo-Controlled, Cross-Over Trial of Progabide as Add-on Therapy in Epileptic Patients.
Epilepsia., 32: 133-139.

- **Goddard, G.V. (1967):**
Development of epileptic seizures through brain stimulation at low intensity. In: Pedly, T.A.; Meldrum, B.S. (eds).
Nature., 204: 1020-1021.

- **Goddard, G.V.; Mc Intyre, D.C. and Leech, C.K. (1969):**
A permanent change in brain function resulting from daily electrical stimulation.

Exp. Neurol., 25: 295-330.

- **Gram, L. (1994):**
Clinical Experience with Oxcarbazepine Treatment.
Epilepsia., 35: 23-25.

- **Gram, L.; Sabers, A. and Dulace, O. (1992):**
Treatment Of Pediatric Epilepsy with Y-Vinyl GABA
(Vigabatrin).
Epilepsia., 33: S 26 - S 29.

- **Guthrie, H.A. (1983):**
Introductory Nutrition, 5th edition, 119-127.

- **Halsted, J.A.; Ronaghy, H.A.; Abadi, P.; Haghshenass, M.; Amirhakeni, G.H.; Barakat, P.M. and Reinhold, J.C. (1974):**
Zinc deficiency in man. The shiraz experiment.
Am. J. Med., 53: 277-284.

- **Hauser, W.A. and Kurland, L.T. (1975):**
The epidemiology of epilepsy in Rochester, Minnesota, 1935 through 1967.
Epilepsia., 16: 1.

- **Heijbel, J. and Bergfors, P.G. (1975):**
Benign epilepsy of children with centrotemporal EEG foci. A study of incidence rate in outpatient care.
Epilepsia., 16: 657-664.

- **Holmes, G.L. (1986):**
Partial seizures in children.
Pediatrics., 77: 725-742.

- **Hoogenroad, T.U. and Van-den Hamer, C.J. (1983):**
3 years of continuous oral zinc therapy in 4 patients with

Wilson's disease.
Acta. Neurol. Scand., 87 (6): 356-364.

- **Horton, R.W. (1989):**
Amino acid neurotransmitters.
Neurotransmitters: Drugs And Disease.
Webster, R.A. and Jordon, C.C. (eds): 165-181.

- **Houtkooper, M.A.; Van Oorschot, C.A.; Rentmeester, T.W.; Hoppener, P.J. and Onkelinx, C. (1986):**
Double-Blind Study of Milacemide in Hospitalized Therapy-Resistant Patients with Epilepsy.
Epilepsia., 27: 255-262.

- **Hu, R.Q. and Davies, J.A. (1997):**
Glutamate receptor antagonists reduce spontaneous epileptiform activity in cortical wedges prepared from DBA/2 mice.
Exp. Brain. Res., 115 (2): 311-318.

- **Ikeda, T.; Higashi, A.; Matsukura, M. and Matsuda, I. (1983):**
Hair copper and zinc concentrations in handicapped children treated with anticonvulsants.
Dev. Pharmacol. Ther., 6: 381-387.

- **Ilhan, A.; Uz, E.; Kali, S.; Var, A. and Akyol, O. (1999):**
Serum and hair trace element levels in patients with epilepsy and healthy subjects: does the antiepileptic therapy affect the element concentrations of hair?
Eur. J. Neurol., 6 (6): 705-709.

- **Isokawa, M. and Mathern, G.W. (1999):**
Developmental changes in NMDA-induced intrinsic optical signals in the hippocampal dentate gyrus of children with medically intractable seizures.
Dev. Neurosci., 21(3-5): 215-222.

- **Janjua, N.A.; Itano, T.; Kugoh, T.; Hosokawa, K.; Nakano, M.; Matsui, H. and Hatase, O. (1992):**
Familial increase in plasma glutamic acid in epilepsy.
Epilepsy. Res., 11 (1): 37-44.

- **Jawad, S.; Oxley, J.R.; Yuen, W.C. and Richens, A. (1986):**
The Effects of Lamotrigine (A Novel Anticonvulsant) on Interictal Spikes in Patients with Epilepsy.
Br. J. Clin. Pharmacol., 22: 191-193.

- **Jawad, S.; Yuen, W.C.; Peck, A.W.; Hamulton, M.I.; Oxley, J.R.; and Richens, A. (1987):**
Lamotrigines: Single Dose Pharmacokinetics and Initial one week Experience in Refractory Seizures.
Epilepsy. Res., 1: 194-201.

- **John Cawte, M. (1985):**
Psychiatric sequelae of manganese exposure in the adult, fetal and neonatal nervous system.
J. of Psychiatry. Australian., 19: 211-217.

- **Johnson, J.W. and Ascher, P. (1987):**
Glycine potentiates the NMDA response in cultured mouse neurons.
Nature; 325: 529-533.

- **Johnston, G.A. (1987):**
Neuropharmacology of amino acid inhibitory transmitters.
Ann. Rev. Pharmacol., 28: 1705-1712.

- **Kahyama, K.; Ueno, S.; Tatsumi, T.; Yorifum, S.; Takahashi, M. and Tarul, S. (1988):**
Decreased glutamate dehydrogenase protein in spinocerebellar degeneration.

J. Neurosurg. And Psychia., 54: 1078-1080.

- **Kairiss, E.W.; Racine, R.J. and Smith, G.K. (1984):**
Development of interictal spike during kindling in the rat.
Brain. Res., 322: 101-110.
- **Kari, E.M. (1979):**
A conspectus of research on copper metabolism and requirements in man.
J. Nutr., 109 (11): 1979.
- **Kay, R.G. and Tasman-Jones, C. (1975):**
Zinc deficiency and intravenous feeding.
Lancet., 2: 605-606.
- **Kleckner, N. and Dingledine, R. (1988):**
Requirement for glycine in activation of NMDA receptors expressed in *Xenopus* oocytes.
Science; 241: 835-837.
- **Klevay, L.M.; Bistrian, B.R.; Fleming, C.R. and Neumann, C.G. (1987):**
Hair analysis in clinical and experimental medicine.
Am. J. Clin. Nutr., 46: 233-236.
- **Koch, H.J.; Smith, E.R. and Moneely, J. (1957):**
Analysis of trace elements in human tissue. In: the lymphomatous disease.
Cancer., 10: 151.
- **Koo, S.I. and Ramlet, J.S. (1983):**
Dietary cholesterol decreases the serum level of zinc: further evidence for the positive relationship between zinc and high density lipoproteins.
Am. J. Clin. Nutr., 37: 918-923.

- **Kosteljanetz, M.; Christiansen, J.; Mouritzen Dam, A.; Stensgaard Hansen, B.; Lyon, B.B.; Pederson, H. and Dam, M. (1979):**
Carbamazepine Versus Phenytoin: A Controlled Clinical Trial in Focal Motor and Generalized Epilepsy.
Arc. Neurol., 36: 22-24.

- **Kurtz, Z.; Tookey, P. and Ross, E. (1988):**
The epidemiology of epilepsy in childhood. In: Roger, J.; Dravet, C. and Bureau, M. (eds): Epileptic Syndromes in Infancy, Childhood and Adolescence. London, John Libbey, 68-72.

- **Kuzniecky, R.; Rubin, Z.K.; Faught, E. and Morawetz, R. (1992):**
Antiepileptic Treatment After Temporal Lobe Epilepsy Surgery: A Randomized Study Comparing Carbamazepine and Polytherapy.
Epilepsia., 33: 908-912.

- **Leach, M.J.; Baxter, M.G. and Critchley, M.A. (1991):**
Neurochemical and Behavioral Aspects of Lamotrigine.
Epilepsia., 32: 4-8.

- **Li, T.K. (1966):**
In: Zinc Metabolism (eds).
Prasad, A.S. and Thomas, T., 48.

- **Limberge, N.; Spath, L. and Starke, K. (1986):**
Release of previously incorporated aminobutyric acid in rabbit caudate nucleus slices.
J. Neurochem., 46: 1102-1108.

- **Linderman, R.D. (1987):**
Mineral in medical practice. In: Quick Reference to Clinical Nutrition.

J.B. Lippincott Company, Philadelphia, New York., (ed. Halpern, S.L.): 295-323.

- **Liu, C.S.; Wu, H.M.; Kao, S.H. and Wei, y.h. (1998):**
Serum trace elements, glutathione, copper/zinc superoxide dismutase, and lipid peroxidation in epileptic patients with phenytoin or carbamazepine monotherapy.
Clin. Neuropharmacol., 21 (1): 62-64.

- **Lucke, A.; Kohling, R. and Speckmann, E.J. (1996):**
Effects of glutamate application on the rhythm of low magnesium-induced epileptiform activity in hippocampal slices of guinea-pigs.
Eur. J. Neurosci., 8 (10): 2137-2148.

- **Lukaski, H.C.; Bolonchuk, W.W.; Klevay, L.M.; Milne, D.B. and Sandstead, H.H. (1983):**
Maximal oxygen consumption as related to magnesium, copper and zinc nutrition.
Am. J. Clin. Nutr., 37: 407-415.

- **Luna, D.; Dulac, O.; Pajot, N. and Deaumont, D. (1989):**
Vigabatrin in the Treatment of Childhood Epilepsy: A Single-Blind Placebo-Controlled Study.
Epilepsia., 30: 430-437.

- **Lund, M. and Trolle, E. (1973):**
Clonazepam (Ro 5-4023) in the Treatment of Epilepsy.
Acta. Neurol. Scand., 49: 82-90.

- **Maritin, W.D.; Mayes, A.P. and Rodwell, W.V. (1985):**
Harper's review of biochemistry., 19th ed.: 579.

- **Martinez-Lage, J.M.; Bossi, L.; Morales, G.; Martinez-Vila, E.; Orofiamma, B. and Viteri, C. (1984):**
Progabide Treatment in Severe Epilepsy: A Double-Blind

Cross-Over Trial Versus Placebo.
Epilepsia., 25: 586-593.

- **Martins da Silva, A.; Aarts, J.H. and Binnie, C.D. (1984):**
The Circadian Distribution of Interictal Epileptiform EEG Activity.
Electroenceph. Clin. Neurophysiol., 58: 1-13.

- **Matsura, S.; Hirayama, K. and Murata, R. (1993):**
Enhancement of synaptic facilitation during the progression of kindling epilepsy by amygdala stimulations.
J. Neurophysiol., 70 (2): 602-609.

- **Maytal, J.; Novak, G.P. and King, K.C. (1991):**
Lorazepam in the Treatment of Refractory Neonatal Seizures.
J. Child. Neurol., 6: 319-323.

- **McLean, M.J.; Schmutz, M.; Wamil, A.W.; Olpe, H.R. and Porter, C. (1994):**
Oxcarbazepine: Mechanism of Action.
Epilepsia., 35: 5-9.

- **Mekky, M.K.; Mostafa, M. and Abu-Zeid, Y.A. (1981):**
Prevalence of epilepsy in rural Egypt.
M. Sc. Thesis. Faculty of Medicine. Ain Shams University.
Coted from Nasr, A.A., Cognitive impairment in epileptic children. M.D. Thesis; Cairo University, 1993.

- **Meldrum, B.S. (1985):**
Amino acid neurotransmitters and new approaches to anticonvulsant drug action.
Epilepsia. 25: 140-149.

- **Meldrum, B.S. (1994):**
The role of glutamate in epilepsy and other CNS disorders.

Neurology; 44(suppl.8): 14-23.

- **Messenheimer, J.; Ramsay, R.E.; Willmore, L.J.; Leroy, R.F.; Zielinski, J.J.; Mattson, R.; Pellock, J.M.; Clakas, A.M.; Womble, G. and Risner, M. (1994):**
Lamotrigine Therapy for Partial Seizures: A Multicenter, Placebo-Controlled, Double-Blind, Cross-Over Trial.
Epilepsia., 35: 113-121.

- **Milgram, N.W.; Green, I.; Liberman, M.; Riexinger, K. and Petit, T.L. (1985):**
Establishment of status epilepticus by limbic system stimulation in previously unstimulated rats.
Exp. Neurol., 88: 253-264.

- **Milligan, N.; Dhillon, S.; Oxley, J. and Richens, A. (1982):** Absorption of Diazepam from the Rectum and its Effect on Interictal Spikes in the EEG.
Epilepsia., 23: 323-331.

- **Mitchell, W.G. and Chavez, J.M. (1987):**
Carbamazepine Versus Phenobarbital for Partial Onset Seizure in Children.
Epilepsia., 28: 56-60.

- **Mori, N. and Wada, J.A. (1989):**
Bidirectional transhemispheric interaction between amygdaloid kindling induced by excitatory amino acids and electrical stimulation.
Epilepsia., 30 (3): 271-275.

- **Morimoto, K. and Goddard, G.V. (1987):**
The substantia nigra is an important site for the containment of seizure generalization in the kindling model of epilepsy.
Epilepsia., 28: 1-10.

- **Moynahan, E.J. (1974):**
Acrodermatitis enteropathica: A lethal inherited human zinc deficiency disorder.
Lancet., 2: 399.

- **NAS. (1989):**
Recommended Dietary Allowance (RDA), 10th Revised Edition, Food and Nutritional Board, National Academy of Sciences, Washington, D.C., 231-235.

- **Nielsen, J.A. and Thaysen, E.H. (1971):**
Acute and chronic magnesium deficiency following extensive small gut resection.
Scand. J. Gastroenterol., 6: 663-666.

- **North, J.B.; Haniel, A.; Challen, R.G.; Penhall, R.R.; Hann, C.S. and Frewin, D.B. (1980):**
Postoperative Epilepsy: A Double-Blind Trial of Phenytoin after Craniotomy.
Lancet., 1: 384-386.

- **Ohmori, O.; Hirano, H. and Mita, T. (1999):**
Shifted cytosolic NADP-dependent isocitrate dehydrogenase on 2-D gel in the brain of genetically epileptic E1 mice.
Neurochem. Res., 24 (3): 365-369.

- **Ohtsuka, Y.; Amano, R.; Mizukawa, M.; Oka, E. and Ohtahara, S. (1992):**
Treatment of Intractable Childhood Epilepsy with High-Dose Valproate.
Epilepsia., 33: 158-164.

- **Okasha, A. (1988):**
Behavior therapy. In: Okasha's clinical psychiatry, published

by: Angelo Egyptian Bookshop. Cairo. A.R.E.

- **Olney, J.W. (1982):**
Excitotoxic amino acids and Huntington's disease.
Chase, T.N.; Wexler, N.S. and Barbeau, A. (eds)., New
York. Raven Press., 609-624.

- **Otoharu, S.; Ishida, S. and Oka, E. (1981):**
Epilepsy and febrile convulsions: A Neuroepidemiological
Study. In: International symposium on developmental
disabilities, (eds)., Fukuyama, Y. and Arima, M. Tokyo,
Amsterdam. Coted from Nasr, A.A., Cognitive impairment in
epileptic children. M.D. Thesis; Cairo University, 1993.

- **Overweg, J.; Binnie, C.D.; Meijer, J.W.; Meinardi, H.;
Neijten, S.T.; Schamltz, S. and Wauquier, A. (1984):**
Double-Blind Placebo-Controlled Trial of Flunarizine as
Add-on Therapy in Epilepsy.
Epilepsia., 25: 217-222.

- **Panayiotopoulos, C.P. (1989):**
Benign childhood epilepsy with occipital paroxysms: a 15-
year prospective study.
Ann. Neurol., 26: 51-56.

- **Parisi, A.F. and Vallee, B.J. (1970):**
Isolation of Zinc and Macroglobulin.
J. Biochem., 9: 2421.

- **Pineal, J.P. and Ronver, L.I. (1978):**
Electrode placement and kindling-induced experimental
epilepsy.
Exp. Neurol., 58: 335-340.

- **Pippenger, C.E. (1989):**
Pharmacodynamics of Antiepileptic Drugs. In: R.M. Post;

M.R. Trimble and C.E. Pippenger (eds). Clinical Use of Anticonvulsants in Psychiatric Disorders. New York. Demos.

- **Porter, J. (1989):**
Mechanism of Action of New Antiepileptic Drugs.
Epilepsia., 30: 29-34.

- **Post, R.M. and Weiss, S.R. (1989):**
Kindling and manic-depressive illness. In: Bolwig, T.G. and Trimble, M.R. (eds), *The Clinical Relevance Of Kindling*. Wiley. Chichester., 209-230.

- **Prasad, A.S. (1966):**
Metabolism of zinc and its deficiency in human subjects. In: *Zinc Metabolism*.
Archives of Internal Medicine., 111: 407-428.

- **Prasad, A.S. (1983):**
Clinical, endocrinologic, and biochemical effects of zinc deficiency.
Spc. Top Endocrinol. Metabol., 7: 45-76.

- **Prasad, A.S.(1985):**
Essential trace elements in human health and disease.
J. Am. Coll. Nutr., 4 (1): 1-2.

- **Prasad, A.S.; Dardenne, M.; Abdallah, J.; Meftah, S.; Brewer, G.J. and Bach, J.F. (1988):**
Serum thymulin and zinc deficiency in humans.
Ann. Rev. Pharmacol. Toxicol., 20: 393-426.

- **Ramsay, R.E. (1993):**
Treatment of Status Epilepticus.
Epilepsia., 34: 71-81.

- **Reynolds, E.H. (1992):**
Y-Vinyl GABA (Vigabatrin): Clinical Experience in Adult and Adolescent Patients with Intractable Epilepsy.
Epilepsia., 33: 30-35.

- **Richens, A. (1992):**
Lamotrigine.
Recent. Adv. Epilepsy., 5: 197-210.

- **Riekinen, P.J.; Pitkanen, A.; Ylinen, A.; Sivenius, J. and Halonen, T. (1989):**
Specificity of Vigabatrin for the GABA-Ergic System in Human Epilepsy.
Epilepsia., 30: 180-222.

- **Rimmer, E.M. and Richens, A. (1988):**
Clinical Pharmacology and Medical Treatment. In: W.J. Laidla; A. Richens and J. Oxley (eds). *Textbook of Epilepsy*.
New York.: Churchill Livingstone.

- **Robertson, M.M. (1986):**
Current Status of 1,4 and 1,5 Benzodiazepines in the Treatment of Epilepsy: The Place of Clobazam.
Epilepsia., 27: 27-41.

- **Rock, D.M.; Macdonald, R.L. and Taylor, C.P. (1989):**
Blockade of Sustained Repetitive Action Potentials in Cultured Spinal Cord Neurons by Zonisamide (AD 810, CI 912): A Novel Anticonvulsant.
Epilepsy. Res., 3: 138-143.

- **Rodin, E.A.; Rim, C.S. and Rennick, P.M. (1974):**
The Effect of Carbamazepine on Patients with Psychomotor Epilepsy: Results of a Double-Blind Study.
Epilepsia., 15: 547-561.

- **Rogawski, M.A. and Porter, R.J. (1990):**
Antiepileptic Drugs: Pharmacological Mechanisms and Clinical Efficacy with Consideration of Promising Developmental Stage Compounds.
Pharmacol. Rev., 42: 223-386.

- **Rose, S.E.; Penry, J.K.; Markush, R.E.; Radloff, L.A. and Putnam, P.L. (1973):**
Prevalence of epilepsy in children.
Epilepsia., 14: 133-152.

- **Ross, E.M. and Peckham, C.S. (1983):**
Seizure disorder in the National Child Development Study.
In: Research progress in epilepsy, (eds). Rose, F.C. New York. Raven Press.

- **Rowan, A.J.; Meijer, W.A.; Beer-Pawlikowski, N.; Van Geest, P. and Meinardi, H. (1983):**
Valproate, Ethosuximide Combination Therapy for Refractory Absence Seizures.
Arc. Neurol., 40: 797-802.

- **Rowan, A.J. and French, J.A. (1988):**
The Role of Electroencephalogram in the Diagnosis and Management of Epilepsy.
Recent. Adv. Epilepsy., 4: 63-92.

- **Rowan, A.J.; Schear, M.J.; Wiener, J.A. and Luciano, D. (1989):**
Intensive Monitoring and Pharmacokinetic Studies of Gabapentin in Patients with Generalized Spike-Wave Discharges.
Epilepsia., 30: 661.

- **Sachdeo, R. and Chokrovertty, S. (1985):**
Increasing Epileptiform Activities in EEG in Presence of

Decreasing Clinical Seizures after Carbamazepine.
Epilepsia., 26: 522.

- **Sandstead, H.H.; Burch, R.E.; Booth, G.H. and Darby, W.J. (1970):**
Current concepts on trace elements; Clinical consideration.
Med. Clin. North. Am., 54 (6): 1509.
- **Saner, G.; Dagoglu, T. and Ozden, T. (1985):**
Hair manganese concentrations in newborns and their mothers.
Am. J. Clin. Nutr., 41: 1042-1044.
- **Sangrador, C.O. and Luaces, R.L. (1991):**
Study of the prevalence of epilepsy among school children in Valladolid, Spain.
Epilepsia., 32: 791-797.
- **Sato, S.; White, B.G.; Penry, J.K.; Dreifuss, F.E.; Sackellares, J.C. and Kupferberg, H.J. (1982):**
Valproic Acid Versus Ethosuximide in the Treatment of Absence Seizures.
Neurology., 32: 157-163.
- **Saunders, J.W. and Shorvon, S.D. (1987):**
Incidence and prevalence studies in epilepsy and their methodological problems : A Review.
J. Neurol. Neurosurg. Psychiat., 50: 829.
- **Schauf, C.L. (1987):**
Zonisamide Enhances Slow Sodium Inactivation in Myxicola.
Brain. Res., 413: 185-188.
- **Scheuhammer, A.M. and Cherian, M.G. (1981):**
The influence of manganese on the distribution of essential

trace elements. Regional distribution of Mn, Na, K, Mg, Zn and Cu in rat brain after chronic Mn exposure.

Toxicol. Appl. Pharmacol., 61: 227-233.

- **Schroeder, H.A. (1966):**
Cadmium as a factor in hypertension.
J. Chronic. Dis., 18: 647.
- **Shakir, R.A.; Johnson, R.H.; Lambie, D.G.; Melville, I.D. and Nanda, R.N. (1981):**
Comparison of sodium valproate and phenytoin as Single Drug Treatment in Epilepsy.
Epilepsia., 22: 27-33.
- **Shammansky, S.L. and Glaster, G.H. (1979):**
Socioeconomic characteristics of childhood seizure disorders in New Haven area: An epidemiological study.
Epilepsia., 20: 457-474.
- **Sherwin, A.; Quesney, F.; Gauthier, S.; Olivier, A.; Robitaille, A.; McQuaid, P.; Harvey, C. and VanGelder, N. (1984):**
Enzyme changes in actively spiking areas of human epileptic cerebral cortex.
Neurology. 34: 927-933.
- **Sherwin, A.L. (1999):**
Neuroactive aminoacids in focally epileptic human brain: A review.
Neurochem. Res., 24 (11): 1387-1395.
- **Shrestha, K.P. and Oswaldo, A. (1987):**
Trace elements in hair of epileptic and normal subjects.
Sci. Total Environ., 67 (2-3): 215-225.

- **Shukla, G.S. and Shandra, S.V. (1982):**
Effect of manganese on carbohydrate metabolism in mitochondrial enzymes in rats.
Acta. Pharmacol. Toxicol., 51: 209-216.

- **Smith, K.T.; Cousins, R.J.; Silbon, B.L. and Failla, M.L. (1978):**
Zinc absorption and metabolism by isolated vascularly perfused rat intestine.
J. Nutr., 108: 1848-1857.

- **Solomons, N.W.; Pineda, O.; Viteri, F. and Sandstead, H.H. (1983):**
Studies in the bioavailability of zinc in humans: mechanism of the intestinal interactions of non-heme iron and zinc.
J. Nutr., 113: 339-349.

- **Staedt, J.; Stoppe, G.; Kogler, A.; Munz, D.L.; Hajak, G.; Rieman, H. and Ruther, E. (1994):**
Nightly myoclonus syndrome (NMS) and restless legs syndrome (RLS); Review and Case Report.
Neurol. Psychiatr., 62 (3): 88-93.

- **Stahl, Y.; Persson, A.A.; Petters, I.; Rane, A.; Theorell, K. and Walson, P. (1983):**
Kinetics of Clonazepam in Relation to Electroencephalographic and Clinical Effects.
Epilepsia., 24: 225-231.

- **Stasheff, S.F.; Mott, D.D. and Wilson, W.A. (1994):**
Axon terminal hyperexcitability associated with epileptogenesis in vitro. Pharmacological regulation by NMDA and GABA-A receptors.
J. Neurophysiol., 70 (3): 976-984.

- **Steidl, L.; Tolde, I. and Svomova, V. (1987):**
Metabolism of magnesium and zinc in patients treated with antiepileptic drugs and with magnesium lactate.
Magnesium. 6: 284-295.

- **Stelzer, A.; Slater, N.T. and Bruggencate, G. (1987):**
Activation of NMDA receptors blocks GABAergic inhibition in an in vitro model of epilepsy.
Nature., 326: 688-701.

- **Suzuki, T.; Koizumi, J.; Moroji, T.; Shiraishi, H.; Hori, T.; Baba, A.; Kawai, N. and Tada, K. (1992):**
Effects of long-term anticonvulsant therapy on copper, zinc, and magnesium in hair and serum of epileptics.
Biol. Psychia., 31: 571-581.

- **Suzuki, S.; Aso, K.; Ariyoshi, C. and Ishimaru, M. (1994):**
Acute Intermittent Porphyria and Epilepsy: Safety of Clonazepam.
Epilepsia., 33: 108-111.

- **Tartara, A.; Manni, R.; Galimberti, C.A.; Morini, R.; Mumford, J.P.; Iudice, A. and Perucca, E. (1992):**
Six-year Follow-Up Study on the Efficacy and Safety of Vigabatrin in Patients with Epilepsy.
Acta. Neurol. Scand., 86: 247-251.

- **Telfeian, A.E. and Connors, B.W. (1999):**
Epileptiform propagation patterns mediated by NMDA and non-NMDA receptors in rat neocortex.
Epilepsia., 40 (11): 1499-1506.

- **Thomas, L.; Perry, M.D. and Shirley Hansen, B.A. (1981):**
Amino acid abnormalities in epileptogenic foci.

Neurology. 31: 872-876.

- **Tietz, N.W. (1976):**
Isocitrate dehydrogenase. In: Fundamentals Of Clinical Chemistry. Saunders. Co, W.B. Philadelphia. London. Toronto., 635.
- **Tietz, N.W. (1976):**
Fundamentals of clinical chemistry.
Saunders Co., W.B., 849-865.
- **Treiman, D.M.; Wilensky, A.J.; Ben-Menachem, E.; Ojemann, L.; Yerby, M.; Barber, K.O.; McCormick, K.B.; Cereghino, J.J.; White, B.G. and Swisher, K. (1985):**
Efficacy of Nafimidone in the Treatment of Intractable Partial Seizures: Report of a Two-Center Pilot Study.
Epilepsia., 26: 607-611.
- **Trimble, M.R. (1989):**
Psychiatric Problems in Epilepsy and their Management. In: R.M. Post; M.R. Trimble and C.E. Pippenger (eds.), Clinical use of Anticonvulsants in Psychiatric Disorders. New York. Demos.
- **Troupin, A.S.; Ojemann, L.M.; Halpern, L.; Dodrill, C.; Wilkus, R.; Friel, P. and Feigl, P. (1977):**
Carbamazepine: A Double-Blind Comparison with Phenytoin.
Neurology., 27: 511-519.
- **Tuff, L.P.; Racine, R.F. and Mishia, R.K. (1989):**
The effects of kindling on GABA mediated in the dentate gyrus of the rat. II-Receptor binding.
Brain. Res., 277: 91-98.

- **Villareal, H.J.; Wilder, B.J.; Willmore, L.J.; Bauman, A.W.; Hammond, E.J. and Bruni, J. (1978):**
Effect of Valproic Acid on Spike-Wave Discharges in Patients with Absence Seizures.
Neurology., 28: 886-891.

- **Wada, J.A.; Osawa, T.; Wake, A. and Corcoran, M.E. (1975):**
Effects of taurine on kindled amygdaloid seizures in rats, cats, and photosensitive baboons.
Epilepsia., 16: 229-234.

- **Wada, A.S.; Sato, M.; Wake, A.; Green, G.R. and Troupin, A.S. (1976):**
Prophylactic effects of phenytoin, phenobarbital and carbamazepine examined kindling cat preparation.
Arch. Neurol., 33: 426-434.

- **Walshe, J.M. (1967):**
The physiology of copper in man and its relation to Wilson's disease.
Brain., 90: 109.

- **Wasterlain, C.G.; Morin, A.M.; FujiKawa, D.G. and Bronstein, J.M. (1989):**
Chemical kindling. In: Bolwig, T.G. and Trimble, M.R. (eds), *The Clinical Relevance Of Kindling*. Wiley. Chichester., 35-53.

- **Wenthold, R.J.; Altschuler, R.A.; Skaggs, K.K. and Reeks, K.A. (1987):**
Immunochemical characterization of glutamate dehydrogenase in the cerebellum of the rat.
J. Neurochem., 48: 636-643.

- **Whitehouse, R.C.; Prasad, A.S.; Rabbani, P.I. and Cossack, Z.T. (1982):**
Zinc in plasma, neutrophils, lymphocytes and erythrocytes as determined by flameless atomic absorption spectrophotometry.
N. Engl. J. Med., 298 (1): 6-10.

- **Wilensky, A.J.; Ojemann, L.M.; Chmelir, T.; Margul, B.L. and Doose, D.R. (1989):**
Topiramate Pharmacokinetics in Epileptic Patients Receiving Carbamazepine.
Epilepsia., 30: 645.

- **Wilkinson, R.J. and Nicholls, D.G. (1989):**
Compartmentation of glutamate and aspartate within cerebral cortical synaptosomes: evidence for a non cytoplasmic origin for the calcium releasable pool of glutamate.
Neurochem. Int., 15 (2): 191-197.

- **Wilkus, R.J.; Dodrill, C.B. and Troupin, A.S. (1978):**
Carbamazepine and the Electroencephalogram of Epileptics: A Double-Blind Study in Comparison to Phenytoin.
Epilepsia., 19: 283-291.

- **William, S.R. (1985):**
Mineral, foundation of nutrition. In: Nutrition and Diet therapy, Roberson, N.K. (eds)., 5th. Times Mirror. Masby. College Publishing., chap. 8: 165-211.

- **Wingrove, D.E. and Gunter, T.E. (1986):**
Kinetics of mitochondrial Ca efflux mechanism of liver mitochondria.
J. Biol. Chem., 261: 15159-15165.

- **Woodbury, D.M. and Kemp, J.W. (1989):**
Other Antiepileptic Drugs: Acetazolamide. In: R. Levy; R. Mattson; B. Meldrum; J.K. Penry and F.E. Dreifuss (eds). Antiepileptic Drugs 3rd ed. New York: Raven Press.
- **Yeh, G.C.; Bonhaus, D.W.; Nadler, J.V. and Mac Namara, J.O. (1989):**
N-Methyl-D-Aspartate receptor plasticity in kindling: quantitative-qualitative alterations in N-Methyl-D-Aspartate receptor-channel complex.
Proc. Natl. Acad. Sci., USA. 86: 8157-8160.
- **Young, C.; Gean, P.W.; Chiou, L.C. and Shen, Y.Z. (2000):**
Docosahexanoic acid inhibits synaptic transmission and epileptiform activity in the rat hippocampus.
Synapse., 37 (2): 90-94.
- **Zielinski, J.J. (1988):**
Epidemiology, in a textbook of Epilepsy (eds): Laidlaw, J.; Richens, A. and Oxley, J. . Churchill Livingstone Edinburgh, London and New York. Coted from Nasr, A.A., Cognitive impairment in epileptic children. M.D. Thesis; Cairo University, 1993.