

REFERENCES

REFERENCES

Abilgaard, M. (1978):

Purification of two progressive antithrombins of human plasma.

Scand. J. Clin. Lab Invest. 19, 190-195.

Aitchenson C.T.; Pebles. C. and Josling F. (1992):

Characteristics of antinuclear antibodies in rheumatoid arthritis.

Arth. Rheum, 26: 562.

Albrechtsen O.K. (1957):

The fibrinolytic activity of human tissue.

Br. J. Haematol., 3. 284.

Altman, R.D. (1991):

Criteria for classification of clinical osteoarthritis.

J. Rheumatol., 18 (Supl. 27): 10.

Anderassay K., Sherz. M., Ritz. E., Walter. E., Walter. E, Hasper. B., Starch. H. and Vomel. W. (1976):

Penicillin induced coagulation disorder.

Lancet., 2: 1039-1041.

Anderson, J. (ed.) (1986):

Muir's Textbook of pathology 11th Edition E.L.B.S. and Edward Arnold.

Arnett, F.C.; Edworthy, S. and Block, D.A. (1988):

The 1987 Revised ACR criteria for classification of RA.

Arth. Rheum., 31: 315.

Astrup T. (1966):

Tissue activator of plasminogen.

Fed Proc., 25: 42.

Axelrod, R.L. (1996):

Glucocorticoid therapy, Medicine, 55: 39.

Bacon, P. A. (1991):

Diagnosis of rheumatic disorders.
Amer. J. Med., 66: 337.

Bacon, P.A. (1993):

Rheumatoid arthritis in: Cecil textbook of medicine, 19th edition, PP., 1508. 1515 W.B. Saunders Company, Philadelphia, Lea and Febiger, PP. 811-840.

Bacon, P.A., (1993):

Rheumatoid arthritis. In McCarty DJ. Koopman, WJ. (eds): Arthritis and Allied conditions, 12th ed. Philadelphia. PP. 811-840.

Badley, E.M. and Wood, P.H. (1987):

Impairment, Disability an International Classification of Impairment, Disabilities and handicaps: The nature of the underlying conditions and patients of impairment.
Int. Rehabil Med., 8: 118.

Bagge, E.; Bjelle, A.; Eden, S. and Saunborg, A. (1991):

Osteoarthritis in the elderly: Clinical and radiological findings.
Ann. Rheum. Dis. 50: 535-539.

Ballord, T.R., Balow, J.E. and Dale, D.C. (1995):

Glucorticoid therapy: Mechanisms of action and clinical considerations.
Ann. Intern. Med., 44; 304-315.

Barlow, G.H. and Robbins R.C. (1982):

Fibrinolytic agents in factor VIII in health and disease.
Clin. Pathol. 11084.

Barnett, E.V., Condemi, J. and Atwater E.C. (1995):

The significance of antinuclear factors in rheumatoid arthritis.
Arth., Rheum, 8: 1080.

Belhon, I.R., and Hess E. V. (1993):

Erosive osteoarthritis.
Semin. Arth. Rheum., 22: 298.

Beller F.K.; Nachtigall L., Rosenberg M., (1972):

Coagulation studies of menopausal women taking estrogen replacement.
Obstet. Gynecol; 39 (5): 775-778.

Bert J.M. (1993):

Role of abrasion arthroplasty and debridement in the management of osteoarthritis of the knee.
Rheum. Dis. Clin. North Am. Aug.; 19 (3): 725-739.

Bick R.L., Bick M.D., Feketete L.F. (1980):

Antithrombin III patterns in disseminated intravascular coagulation.
American Journal Clin. Pathol. (4), 73: 557-583.

Bigg S.R. (1976):

Human blood coagulation. Haemostasis and thrombosis,
Blookwell scientific publication, 36.

Bodmer, J., and Schnebl, M. (1984):

Plasma proteinase inhibitors.
Schweiz Med. Wschr.
Research Department Ciba-Geigy, (114) (1359-1363).

Borrenstein G, Ping C, David L. and Craig M. K. (1995):

Coagulant proteins and thrombin generation in synovial fluids.
American Journal of Hematology (50); 79-83.

Braat E.A.M., Jie, A.F.H., Ronday H.K., Beekman. B. and Rijken D.C.; (2000):

Urokinase mediated fibrinolysis in the synovial fluid of rheumatoid arthritis patients may be affected by the inactivation of single chain urokinase type plasminogen activator by thrombin.
Ann rheum. Dis. (59): 315-318.

Brooze GJ. Jr., Warren La and Novotny WF. (1988):

The lipoprotein associated coagulation inhibitor that inhibits the factor VII tissue factor complex also inhibits factor Xa: insight into its possible mechanism of action.
Blood; 71: 335-343.

Camerer E., Kolsto AB. And Prydz H. (1996):

Cell biology of tissue factor, the principal initiator of blood coagulation.

Thromb. Res. Jan.; 81 (1): 1-14.

Capell H.A., Lewis D. and Careg J. (1996):

A three year follow up of patients to placebo, or oral-or injectable gold therapy for rheumatoid arthritis.

Ann. Rheum. Dis., 45: 705-711.

Carmichael J. and Shankel S.W. (1996):

Effect of nonsteroidal anti-inflammatory drugs on prostaglandin and renal function.

Am. J. med., 78; 992-1000.

Carson, D.A. (1997):

Antiglobin antibodies. In: Panayji, GS. (ed) the scientific basis of rheumatology.

Chruchill Livingstone, Edinburgh, P. 114.

Castillo B.A., El-Sallab R.A. and Scott J.T. (1994):

Physical activity, cystic erosions and osteoporesis in rheumatoid arthritis.

Ann. Rheum. Dis., 29: 522-527.

Cathcart E.S.; Hayes. K.C. and Franz Blau, (1986):

Experimental arthritis in a non human primate. I. Induction of bovine type II collagen.

Lab. Invest. 54: 26.

Chan, V., and Chan, T. (1979):

Antithrombin III in fresh and cultured human endothelial cells. A natural anticoagulant from the vascular endothelium.

Thrombosis Research, (15): 209-13.

Chandnani, V.; HOC.; Gerharter J., Neumann, C., Kursunoglu-Brahme S., Sartoris D.J., Resnick D., (1992):

MRI findings in asymptomatic shoulders: a blind analysis using symptomatic shoulders as controls.

Clin. Imaging Jun-Mar; 16 (1): 25-30.

Ciennensen I., Holund B., and Andersen RB. (1983):

Fibrin and fibronectin in rheumatoid synovial membrane and rheumatoid synovial fluid.
Arthritis Rheum Apr., 26 (4): 479-485.

Clements P.J., Paulus H.E. (1997):

Nonsteroidal antirheumatic drugs. in Kelly WN. et al., (eds) text book of Rheumatology, 5th edition Philadelphia, WB, Saunders, PP. 707-740.

Collen D, Wiman B, and Verstrate M. (1979):

The physiological inhibitors of blood coagulation and fibrinolysis proceeding of a round-table conference held at the university of Leuvan, Belgium July 2223, Amsterdam: Elsevier.

Colman RW, Jameson B. and Lin Y. and Conard J. (2000):

Domain 5 of high molecular weight kininogen down regulates endothelial cell proliferation and migration and inhibit angiogenesis.
Journal Haematol, 45: 543-550.

Colman RW, Pixely RA, Najamunnisa S. Comp PC and Horellon MH. (1997):

Binding of high molecular kininogen to human endothelial cells is mediated via a site within domains 2 + 3 of urokinase receptor.
Haemostasis. 100: 1481-1487.

Conard, J. and Samama, M. (1978):

L'antithrombin III.
La Revue de Medicine, 19: 2343-2351.

Condemi, J.; Barnett, E.V. and Atwater, E.C. (1995):

The significance of antinuclear factors in rheumatoid arthritis.
Arth. Rheum. 8: 1080.

Cooke, T.D. and Dwosh, I.L. (1986):

Clinical features of osteoarthritis in the elderly.
Rheum. Dis. Clin., 12: 155-159.

Copper, N.S. (1993):

Pathology of rheumatoid arthritis.
Med. Clin. North American, 52: 3.

Cosgriff TM, Bishop DT, Mershgold EJ, Skolnick MH, Martin BA, Baty BJ, Carlson KS (1983):

Familial Antithrombin III deficiency; Its natural history, genetics, diagnosis and treatment.
Medicine (Baltimore) 1983 Jul., 62 (4): 209-220.

Crey R.G.; Tenn Baum J, Gottlieb NL. (1993):

Local steroid injection treatment in rheumatic disorders.
Semin Arthritis Rheum. 10L: 231-254.

Dacie, J.V. and Lewis, S.M. (1975):

Practical haematology, 3rd edition, New York., Grune and Stratton Inc.

Deighton C.M. and Walker, D.J. (1991):

The familial nature of rheumatoid arthritis.
Ann. Rheum. Dis., 50: PP 62-65.

Dennis A., Carson, and Salvatore Albani, (1997):

Etiology and pathogenesis. In Arthritis and Allied condition
Ed. By Hollander, J. and Mc. Carty D.J. 12th Edition, Philadelphia, PP. 1041-1070.

Doherty, M. (1994):

Color atlas and text of osteoarthritis Wolfe, London.

Doolittle LR., Doalittle RF and Goldbaum RM (1978):

Designation of sequences involved in the coiled-coil interdominal connections in fibrinogen concentration of an atomic scale model.
J. Mol. Biol. 20; 311-325.

Duthie J., Brown P.E., Truelove L.H., Barager F.D., and Lawrie A.J. (1994):

Course and prognosis in rheumatoid arthritis.
Ann Rheum. Dis. 23; 193-195.

Dumonde, D., And Stewart, M. (1978):

The role of microbial infection in rheumatic disease. In Compeman's Text book of Rheumatic Disease Fifth edition. Livingstone. London. P. 222.

Dunne, J.U.; Carson, D.A. and Spiegelberge, H.L. (1995):

IgA rheumatoid factor in serum of patients with rheumatoid arthritis.

Ann. Rheum. Dis., 38: 161.

Edmonds, J.P. (1997):

Rheumatoid arthritis. In: Howe H.S., Feng P.H. (eds). Text book of clinical Rheumatology 1st ed. Singapore, PP. 54-111.

Edward and Harris (1997):

Clinical features of rheumatoid arthritis. In: Kelly W.N.; Harris E.D., Ruddy S., Sledge C.B. (eds): Text book of Rheumatology 5th ed Saunders Company, Philadelphia.

Edward, (2001):

Clinical picture of rheumatoid arthritis. In Kelly W.N.; Harris E.D., Ruddy S., Sledge C.B. (eds): Text book of Rheumatology 6th ed. Saunders Company Philadelphia. PP. 967-1000.

Edwards PH.; Grana W.A. (2001):

Anterior cruciate ligament reconstruction in the articular reconstruction.

Am. J. Knee Surg.; 14 (4): 232-237.

Edwards, R.L. and Rickles F.R. (1984):

Macrophage procoagulants.

Prog. Hemost. Thromb.; 7: 183-209.

Ehrlich M.G.; Houle, P.A.; Vigliani, G. and Mankin, H.J. (1978):

Correlation between articular cartilage collagenase activity and osteoarthritis

Arth. Rheum., 21: 761.

Emery S.E. and Bohlman H. (1992):

Osteoarthritis of the cervical spine. Osteoarthritis: Diagnosis and Management. Philadelphia, W.B Saunders company, PP. 255-263.

Fagerhol, M.K. and Abildgaard, U. (1970):

Immunological studies on human AT III. Influence of age, sex and use of oral contraceptives on serum concentration. Scand. Journal of Haematology, (7): 10-17.

Fagerhol, M.K. Korvaid, E.; Godal, H.C.; and Abildgaard, U. (1971):

Antithrombin III in patients with acute changes of the haemostatic mechanism, 2nd congress of International society of thrombosis and Haemostasis. Advances in Protein chemistry: Annual publication Academic Press. New York., London.

Fair, D.; and Bahnak, D. (1984):

Human hepatoma cells secrete single chain factor X, prothrombin and AT III. Blood, 64: 194-214.

Fair DS.; Marlar RA. (1986):

Biosynthesis and secretion of factor VII, protein C, protein S, and the protein C inhibitor from a human hepatoma cell line Blood; 67: 64-70.

Felson, D.T. (1996):

Osteoarthritis. Rheum. Dis. Clin., 16: 499.

Fenton J.W., Harpel, P.C. (1986):

Novel immunoenzyme assays for thrombin-antithrombin III and thrombin α_2 macroglobulin complexes in plasma. Acad Sci.; 485: 184-198.

Finazzi G., Caccia R. and Barbir T. (1987):

Different prevalence of thrombolism in the subtypes of congenital antithrombin III deficiency. Blood; 58: 109.

Firestein, G.S. (2001):

Etiology and Pathogenesis of rheumatoid arthritis. In Textbook of Rheumatology 6th ed. Saunders Company Philadelphia. PP. 851-897.

Fisher, R.A. (1964):

Statistical methods of research workers.
10th ed., Endenburg, Dilver and Boyed.

Fong, S.; Carson, D. and Vanghan, J. (1985):

Rheumatoid factor in "Immunology of Rheumatic disease".
Ed. By Gupta, S and Talal, N. 2nd pp. 167:197 New York,
Plenum Medical Book Comp.

Forsgren M, Raden B, Bannan S, and Sticklan M. (1987):

Molecular cloning and characterization of a full. Length
cDNA clone for human plasminogen.
FEBS lett; 213: 251-260.

Frank A. Wollheim (1997):

Clinical picture of rheumatoid arthritis. In: Oxford text book
of Rheumatology. 2nd edition Oxford, New York, Tokyo,
PP. 1004-1026.

**Furmaniak –Kazmierczak E., Cooke TD., Manuel R., Seudamore A.,
Hoogendorn H., Giles AR. and Nesheim M. (1994):**

Studies of thrombin-induced proteoglycan release in the
degradation of human and bovine cartilage.
J. Clin. Invest. Aug.; 94 (2): 472-480.

Fye, K.H.; Moutsopoulos S, H.H. and Talat, N. (1992):

Rheumatic disease in "Basic and clinical Immunology" Ed J.
4TH Ed. Los Angeles, California, PP. 422.

Gabazza EC, Osamu T, Yamakami T (1994):

Correlation between clotting and collagen markers in
rheumatoid arthritis.
Thromb Haemost; 71: 1199-2002.

Garber E.K., Targoff C. and Paulus H.E. (1987):

Corticosteroid in the rheumatic disease. In: Drug for
rheumatic disease. New York: Churchill Livingstone, 443.

Gault S.J. and Spyker, J.M. (1994):

Beneficial effect of immobilization of joint in rheumatoid
arthritis.
Arthritis Rheum. 91: 34-44.

Gettins P., Harten B. (1988):

Properties of thrombin and elastase modified human AT III.
Biochemistry May 17: 27 (10): 3634-3639.

Ghosh S., Mackie M.J., McVerry B.A., Galloway M., Ellis A. and Mckay J. (1983):

Chronic inflammatory bowel disease, deep venous thrombosis and antithrombin activity.
Acta. Haematol, 70 (1): 50-53.

Giercksly, K.E. (1996):

Piroxican and gastrointestinal bleeding.
Am. J. Med. 81: 2-5.

Goldenberg, D.L.: Egan, M.S. and Cohen (1982):

Inflammatory synovitis in degenerative joint disease
J. Rheumatol., 9: 204-209.

Golding, D.N. (1989):

Rheumatoid arthritis in A synopsis of rheumatic disease 5th ed, Wright, London, PP. 51-888.

Grigor'eva VD.; Fedorova NE., Suzdal'niskii DV., Dashina TA. and Kapinos EN. (1992):

Cryoelectrotherapy in the treatment of osteoarthritis patients.
Clin. Orthop. Jan. Feb.; (1): 16-20.

Guillaume D.N., Bernard P. and Alain B. (2001):

Disease modifying antiosteoarthritic drugs current therapies and new prospects around protease inhibition.
Elsevier Science IL Farmaco (56); 107-112.

Hardisty R.M.; (1974):

The haemostatic mechanism in blood and its disorders
Blackwell Scientific Publication, Edinburg, Melbourne, 5: 23.

Harfenist, E.J.; and Murray R.K.; (1993):

Plasma protein, Immunoglobulins and clotting factors
Harper's Biochemistry (59); 681, 684-685.

- Haslam R.J.; Davidson MM. and Fox JE. (1978):**
Cyclic nucleotides in platelet function blood; 40: 232-240.
- Hathaway, W.E. (1991):**
Seminars in Hematology: 28: 19.
- Hathway W. and Bonar J. (1978):**
Prenatal coagulation Grune and stratton. New York.
- Heinmoller E, Schroppt, Kisker O, Simon B, Seitz R and Weinel R.J. (1995):**
Tumor cell induced platelet aggregation in vitro by human pancreatic cancer cell lines.
Scand. J. Gastroenterol. Oct., 30 (10): 1008-16.
- Hender, U. and Nilson I.M. (1973):**
AT III in a clinical material.
Thromb. R. es., 8: 631.
- Herzog R., Lutz S., Blin N, Marasa JC., Blinder MA. and Tollefsen DM (1991):**
Complete nucleotide sequence of the gene for human heparin cofactor II and mapping to chromosomal band 22q11.
Biochemistry; 30: 1350-1357.
- High Smith R.F. and Rosenberg, R.D. (1974):**
The inhibition of human plasma by antithrombin heparin cofactor.
J. Biol. Chem., 249: 4335.
- Hiroshi Shin, Isao Kitajima, Toshihiro Nakajima, Qing Shao, Takeshi Tokioka, Ikukotaka Saki, Naohiro Hanya, Toshikazu Kubo and Ikuro Maruyama (1999):**
"Thrombin receptor mediated signals induce expression of interleukin 6 and granulocyte colony stimulating factor via NF-KB activation in synovial fibroblasts".
Ann. Rheum. Dis. (58): 55-60.
- Hiroyasu K.; Esteban C, Gabazza., Taguch O, Wada H. (1988):**
Protein C anticoagulant system in patients with interstitial lung disease.
Am. J. Respire Crit. Care Med., 157: 1850-1854.

Hirsh. J., Ofsu F., and Cairns J. (1985):

Advances in antithrombotic therapy. In: Recent Advances in Haematology. Churchill, Livingstone.

Hogg PJ and Jackson CM. (1989):

Fibrin monomer protects thrombin from inactivation by heparin-antithrombin III implications for heparin efficiency Proc. Natl. Acad. Sci. USA 86; 3619-3623.

Hoogendorn, H., Giles, A.R. and Nesheim, M. (1994):

Studies of thrombin induced proteoglycan release in the degradation of human and bovine cartilage. Clin. Invest. Journal (94) 2: 472-480.

Howard A. Fuchs and John S. Sergent (1997):

In clinical picture of rheumatoid arthritis. In: Arthritis and Allied conditions. 12th Edition Lea and Febiger, Philadelphia. PP. 1041-1070.

Howie P.W., Evans K.; Forbes, C.D. and Prentice, C.R.M. (1975):

The effects of stilboestrol and quineestrol upon coagulation and fibrinolysis during the puerperium. British Journal of Obstetrics Gynaecology, 82, 975-988.

Hoyer LW., Colman RW, Hirsh J. and Marder VJ. (1993):

Hemostasis and Thrombosis: Basic principles and clinical practice. 3rd edition Philadelphia: JB lippincott, 103-133.

Hsiao GH and Chiu HC. (1997):

Low-dose danazol in the treatment of livedoid vasculitis; Ann. Rheum. Dis., 19 (3): 251-255.

Hung DT., Vu T-KH, Nelten NA, Coughlin Sr (1992):

Thrombin-Induced events in non platelet cells are mediated by the unique proteolytic mechanism established for the cloned thrombin receptor. J. Cell. Biol. 116: 827-832.

Ilona K, Ewa Emich-W., Elzbieta M (1995):

Biochemical aspects of brain vascular diseases in children on the base of up to date literature Med. Sci. Monit, 1999; 5 (4): 809-813.

Inamo Y, Pemberton S., E.G.D. Tuddenham and P.Woo (1995):

Increase of activated factor VIII and haemostatic molecular markers in juvenile chronic arthritis.

Br. Journal for Rheumatology; (34): 466-469.

Ingrid Marty, Veronique Peclat, Gailate Kirdaite, Roberto Salvi, Alexander So and Nathalie Busso (2001):

"Amelioration of collagen induced arthritis by thrombin inhibition".

Journal of clinical investigation (107): 631-640.

Jackson. C.M. and Nemerson Y. (1980):

Blood coagulation.

Ann. Rev. Biochem.; 49: 765-811.

Jarrett M.P., Green D. and Taao, C.H. (1983):

Relation between AT III and clinical and serological parameters in systemic lupus erythematosus.

Thromb. Haemost, 77: 180-191

Jayson and Dixon, (1990):

Valvular mechanisms in Juxta-articular cysts.

Ann. Rheum. Dis., 29: 415.

Jones H.W., Bailey. R., Zhang Z., Dunne. A.K., Blake D.R.; Cox N.L., Morris C.J. and Winyard P.G. (1998):

Inactivation of antithrombin III in synovial fluid from patients with rheumatoid arthritis.

Ann. Rheum. Dis. (57): 162-165.

Juillet Y., Fiessinger J. and Alech M. (1978):

Thromboses vases viscerales au cours des deficiences congenitales et hereditaires en AT III.

Nouvelle Press Medicale, 7; 367.

Kaplan KL., Broekman MI. and Chernoff A. (1979):

Platelet alpha-granule proteins studies on release and subcellular localization.

Blood; 53: 604-18.

Keele, C.; Neil E. and Joils, N. (1984):

Samson Wright Applied Physiology.

Oxford University Press London. New York.

Kernoff, P.B.A., and McNicol, G.P. (1977):

Normal and abnormal fibrinolysis.
Br. Med. Bull. 33 (3): 239.

Klippel J.H. (1997):

Treatment of RA in Primer on the rheumatic disease.
Arthritis Foundation, Atlanta, GA, 93.

Koide T., Odani S., Takahashi K, One T. and Sakuragawa N. (1984):

Antithrombin III Toyama: replacement of arginine-47 by cysteine in hereditary abnormal antithrombin III that lacks heparin-binding ability.
Blood; 81: 289-293.

Koie K, Kamiya T. and Ogata K. (1978):

Alpha 2- plasmin inhibitor deficiency.
Thromb Haemost.; 2: 1334-1336.

Krans, S.M. (1993):

Articular and systemic manifestation of osteoarthritis. In
Arthritis and Allied conditions, 12th ed. Malvern, Lea and
Febiger PP. 763-765.

Kual M.P. (1994):

Superficial heat and cold. How to maximize benefits.
Phys. Sports,. Med., 22 (12): 66-74.

Kushner, I. (1991):

The phenomenon of the acute phase response.
Ann. New. York Acad. Sci., 389-348.

Lan DA, Bayston T, Olds RJ., Takahashi K. and Odani S. (1997):

Antithrombin mutation data base: 2nd up date. For the plasma
coagulation Inhibitors subcommittee of the Scientific and
Standardization Committee of the International society on
thrombosis and haemostasis.
Thromb Haemost, 77: 197-211.

Laposta M., Dohnarsky DK., Shin HS. (1983):

Thrombin-induced gap formation in confluent endothelial
cell monolayers in vitro.
Blood; 62: 549-556.

Laroche, P., Plassart V., Amiral J. (1989):

Rapid quantitative latex immuno assays for diagnosis of thrombotic disorders.

Thromb. Haemostas, 62, 379.

Laura P. Hale and Barton F, Hynes (1997):

Pathology of rheumatoid arthritis and associated disorders.

In Arthritis and Allied conditions 12th edition, Lea and Febiger Philadelphia, PP. 993-1016.

Lightfoot (1989):

Treatment of rheumatoid arthritis in Text book of Rheumatology 11th ed. Lea and Febiger, Philadelphia, 772-781.

Lin, Y. Harris RB.; Yan W. and Black. C. (1997):

High molecular weight kininogen peptides inhibit the formation of Kallikrein on endothelial cell surface and subsequent urokinase dependent plasmin formation.

Thromb. Haemost; 90: 690-697.

Louis Soloman: (2001):

Clinical picture of osteoarthritis in: Text book of Rheumatology 6th ed. Saunders Company, Philadelphia. PP. 1409-1418.

Handbook of Rheumatology, 2nd ed. Vol. 1. Arthritis and Allied Conditions. 1409-1418.

Mandle B, Jr., Colman R.W. and Kaplan AP. (1976):

Identification of prekalikrein and high molecular weight kininogen as a complex in human plasma.

Thromb. Haemost; 73: 4179-4183.

Mankin, Henry J. and Brandt, Kenneth D. (2001):

Pathogenesis of osteoarthritis. Text book of Rheumatology 6th ed. Saunders Company, Philadelphia. PP. 1391-1406.

Mankin H.J. (1983):

Orthopaedics in 2013: a prospect.

J. Bone Joint Surg. Am. Oct.; 65 (8): 1190-1194.

Mark and Harlod (1993):

Corticosteroid. In *Arthritis and Allied conditions: A textbook of rheumatology* 12th ed. Philadelphia, Lea and Febiger, PP. 665-682.

Maruyama Naoki, Hirano Fuminori, Yoshirawa Noritaday Migita Kiyoshi, Katsumi Eguchi and Hirotoshi Tanaka (2000):

Thrombin stimulates cell proliferation in human fibroblast-like synoviocytes in NF-KB factor activation and protein kinase mediated pathway.
Journal of Rheumatology (27): 2777-2785.

Mazzuca S.A., Brandt KD. (1999):

Plain radiography as an outcome. Measure in clinical trials involving patients with knee osteoarthritis.
Rheum. Dis. Clin. North Am. May; 25 (2): 467-480.

McAlindon T.E., Cooper C., Lirwan J.R. and Dieppe P.A. (1992):

Knee pain and disability in the community.
Br. J. Rheumatol. 31: 189-92.

McCarty, D.J. (1997):

Synovial fluid in *Arthritis and Allied Conditions* 12th Edition, P. 56.

McClelland, J.; Fail, B. and Walker, D. (1996):

E.S.R., Hb level, plasma and serum viscosity as measure of disease activity in rheumatoid arthritis.
Br. J. Rheumatol., 36: 156-158.

Mills J.A. (1991):

Value of bed rest in patients with rheumatoid arthritis.
N. Engl. J. Med., 289: 453-458.

Moll J.M. (1987):

Osteoarthritis. In: *Rheumatology in Clinical Practice* Edited Blackwell Scientific Publications.
 Oxford, P. 342-372.

Moncada S., and Vane JR. (1979):

The role of prostacyclin in vascular tissue.
Fed Proc; 38: 66-71.

Morris R, Morris CJ., Winyard PG., Martin JE (1993):

The proinflammatory action of intra-articular thrombin
virchows. Arch A.
pathol Anat (in Abstract). 8: 90

Morris D.T., Wilder R.L., Cupps, T.R., Balow, J.E. (1990):

Glucocorticoid therapy for immune-mediated diseases. Basic
and clinical correlates.
Ann. Intern. Med., 119: 1198-1208.

**Mosher DF.; Hurley R.W., Skorstengaard K. and Ohkubo I and
Asakura S. (1992):**

Inhibition of cell adhesion by high molecular weight
Kininogen.
J. cell. Biol. Jan; 116 (2): 465-476.

Moskowitz, R.W. (1992):

Osteoarthritis, signs and symptoms. In: Osteoarthritis.:
Diagnosis and Management. Philadelphia, W.B. Saunders
Company, PP. 225-263.

Murray, R.K., Granner, D.K., Mayes, P.A. and Rodwell. V.M. (1988):

Harper's biochemistry, 21st Edition.
Appleton and Lange.

Nakajima T., Shao Q, Tokioka T., Shin H and Kitajima I (1999):

Thrombin receptor mediated signals induce expression of IL-
6 and granulocyte colony stimulating factor via NF-KB
activation in synovial fibroblasts.
Ann. Rheum.Dis. 58: 55-60.

**Nakajima T.; Kitajima I, Shin H. Takasaki I, Shigeta K and Abeyama
K. (1994):**

Involvement of NF-KB activation in thrombin-induced
human vascular smooth muscle cell proliferation.
Biochem Biophys. Res. Commun.; 204: 950-955.

Nakano, K.K. (1995):

Neurological complications of rheumatoid arthritis.
Orthp. Clin. North Am., 8: 712.

Nakano, S., Ikata, T., Kinoshita. I., Kanemats, U.J. and Yasuoka. S. (1999):

Characteristics of the protease activity in the synovial fluid from patients with rheumatoid arthritis and osteoarthritis
Clinical and Experimental Rheumatology (17): 161 – 170.

Naoki Maruyama, Fuminori Hirano, Noritada Yoshikawa, Kiyoshi Migita, Katsumi Eguchi and Hirotoshi Tanaka (2000):

Thrombin stimulates cell proliferation in human fibroblast like synoviocytes in Nuclear factor-KB activation and protein kinase C mediated pathway.
Journal of Rheumatology; 27: 12.

Nemerson M. and Nossel H. (1982):

The biology of thrombosis.
Ann. Rev. Med. 33: 479.

Nesheim M.G., Gid., S., and Mann, K.G. (1981):

Assembly of the prothrombinase complex in the absence of prothrombin.
J. Biol. Chem., 256: 9874.

Nichols, P.J.R. (1976):

Rehabilitation medicine, the management of physical disabilities, Butter Worth, London, Boston., P. 146.

Nienaber, A.R.M. Gasper and A.W.H. Neitz (2000):

Partial isolation of a potent thrombin inhibitor from the salivary glands of the tick ornithodoros savignyi (Abstract)
Scand. J. Clin. Lab Invest. (20): 115-120

Nilson W., Vaughan, V. and Behram, R.E. (1988):

In: Nelson Textbook of pediatrics. 13th edition. Saunders Company. Philadelphia.

Odegard, M.K. and Lie, M. (1976):

Heparin cofactor activity and AT III concentration in plasma related to age and sex.
Scand. J. Haematol. 17: 258.

Ofusu, F. and Modi G. (1982):

Heparin with a low affinity to AT III.
Thrombosis Res., 28: 487-497.

- Paley D., Bhatnagar J., Herzenberg JE and Bhavé, A. (1994):**
A new procedures for tightening knee collateral ligaments in conjugation with knee realignment osteotomy
Clin. Orthop. North. Am. Jul; 25 (3): 533-555.
- Parrillo, J.E.; Faucci, A.S. (1996):**
Mechanism of glucocorticoid action on immune processes.
Ann. Rev. Pharmacol. Toxicol., 19: 179.
- Penner JA and Hunter MJ. (1978):**
Antithrombins: clinical aspects chemical and biological properties.
Prog. Clin. Biol. Res. (5): 277-300.
- Peyron, J.G. and Altman, R.D. (1992):**
The epidemiology of osteoarthritis. In: Osteoarthritis: Diagnosis and Management. Saunders Company Philadelphia. PP. 15-19.
- Pickup M.E, Dixon J.S., Hallett G., Bird, H.A. and Wright, V. (1991):**
Plasma viscosity a new appraisal of its use as index of disease activity in RA.
Ann. Rheum. Dis., 40: 272.
- Pincus T. and Callahan, L.F. (1992):**
Mortality in rheumatoid arthritis over 15 years (Abstr).
Arth. Rheum., 33: 140.
- Pope R.M., Mannix M., Grillo B.E. and Teller D.C. (1995):**
The hyperviscosity syndrome in rheumatic arthritis
Arth. Rheum., 18: 97.
- Radioux M., Furst D.E. and Cupps T.R. (1993):**
Glucocorticoid therapy for immune-mediated disease: Basic and Clinical correlates.
Ann. Intern. Med., 119: 1198-1208.
- Rashad S., Low F., Revell P., Hemingway A., Rainsford K. and Walker F. (1989):**
Effect of non-steroidal anti inflammatory drugs on course of osteoarthritis.
Lancet. Nov. 2 (8672): 1149.

Ravinder N Maini and Mar Feldmann (1988):

Immunopathogenesis of rheumatoid arthritis" In: 2nd edition
Oxford, New York. PP. 983-1000.

Resnick, D. and Niwayama, G. (1991):

Rheumatoid arthritis and sero-negative spondyloarthropathies. In Diagnosis of bone and joint disorders Vol. 1,
Saunders comp. Philadelphia PP. 852-1001.

Ritchie D., Boyle J.M. and Clnnes J.E. (1988):

Clinical studies with articular index for the assessment of
joint tendernes patients with Rheumatoid arthritis.
Quart, J. Med., 37: 393-406.

Robbins, K.C.; (1982):

The plasminogen-plasmin enzyme system. In text book of
haemostasis and thrombosis Philadelphia., J.B., Lippincott
company, 450.

Rodin E.L., Paul I.L. and Rose R.M. (1984):

Role of mechanical factors on pathogenesis of primary
osteoarthritis.
Lancet, 11: 519.

Rosenberg R. (1976):

Actions and Intreraction of antithrombin and heparin.
N. Engl. J. Med., 292: 146-151.

Rothchild BM, Thompson LD., Pifer DD, Chesnut CM. (1985):

Pertubation of protease inhibitors and substrates in
inflammatory arthritis.
Semin Thromb. Hemost. 11: 394.

Ruggeri Zm, Dent JA. and Saldivar E. (1999):

Contribution of distinct adhesive interactions to platelet
aggregation inflowing blood.
Blood; 94: 172-178.

Ruth Morris, Paul G Winyard, Lawrence F Brass, David R Blake, Christopher J Morris (1996):

Thrombin receptor expression in rheumatoid and
osteoarthritic synovial tissue.
Ann Rheum Dis 55: 841-843.

Salem H. (1986):

The natural anticoagulants.
Clinics in Haematology, vol. 15, No. 2.

Salem M., Abdellaal R., and El-Mahdi E. (1988):

The diagnostic relevance of AT III. An overview.
Gaz. Egypt. Ped. Ass., Vol. 36, No. 1-2 (Jan-June).

Samuels B., Lee, J.C., Engleman E.P. and Hooper J. Jr. (1997):

Membranous nephropathy in patients with rheumatoid arthritis: Relationship to gold therapy.
Medicine; 57: 319.

Schapira M., Scott. C.F. and Colman R.W (1981):

Protection of human plasma kallikrein from inactivation by C1 inhibitors and other protease inhibitors. The role of high molecular weight kininogen.
J. Hamatol; 20: 2738-2743.

Schumacher H.R., Klipple J.H. and Koopman W.J. (1993):

Rheumatoid arthritis. In: Primer on Rheumatic disease
Atlanta, Arthritis Foundation.

Schumacher, H.R.; Klipple, J.H. and Koopman, W.J. (1994):

Beneficial effect of immobilization of joint in rheumatoid arthritis.
Arthritis Rheum. 91: 34-44.

Schumacher. H.R.; Klippel, J.H. and Robinson, D.R. (1988):

Osteoarthritis in "Primer on the Rheumatic Diseases", 9th ed.,
ed. Arthritis Foundation, Atlanta, G.A. P. 171.

Scott CF, Solver LD. And Purdon AD. (1985):

Cleavage of human high molecular weight kininogen by factor XIa in vitro.
Blood coagulation Fibrino L. 260: 10856-10863.

Scott, D.L.; Grindulis, K.A.; Coulton, B.L.; Popert, A.I. and Bacon, P.A. (1984):

Progression of radiological changes in rheumatoid arthritis.
Ann. Rheum. Dis., 43: 8-17.

Shankar R, Delamotte CA, Dicorleto PE. (1992):

3-deazaadenosine inhibits thrombin stimulated in derived growth factor production and endothelial leucocyte adhesion molecule-1 mediated monocyte cell adhesion in human aortic endothelial cells.

J. Biol. Chem; 276: 9376-82.

Shapiro S.S., and Anderson D.B.; (1977):

Thrombin inhibition in normal plasma chemistry and biology of thrombin.

Ann Arbor., Science publishing, 361.

Shattil, S.J.; Hoxie JA. And Cunningham M. (1985):

Changes in the platelet membrane glycoprotein. During platelet activation.

Blood; 260: 11107-11114.

Shinmei M., Kobayashi T., Yoshihara Y., Samura (1995):

A significance of the levels of carboxy terminal type II procollagen peptide, chondroitin sulfate isomers, tissue inhibitor of metalloproteinases in osteoarthritis joint fluid.

J. Rheumatol. Suppl. Feb., 43: 78-81.

Siegel, D.B.; Gelberman, R.H. and Smith, R. (1992):

Osteoarthritis of the hand and wrist. In: Osteoarthritis: Diagnosis and Management Saunders Company, Philadelphia, PP. 547-560.

Sokoloff L. (1985):

Pathology of rheumatoid arthritis. In Arthritis and Allied disease, 9th ed PP. 309-331. Philadelphia, Lea and Febiger.

Sox HC Jr and Liang MH. (1986):

The erythrocyte sedimentation rate. Guidelines for rational use.

Ann. Intern. Med. Apr; (104 (4): 515-523.

Stasek JE and Garcia J.G.N. (1992):

Role of protein kinase C in thrombin-mediated endothelial cell contraction and barrier dysfunction.

Semin Thromb Hemost; 18: 117-125.

Steinbroker, O, Traeger, C.H. and Batterman R.C. (1949):

Therapeutic criteria in rheumatoid arthritis JAMA, 140: 659-662.

Stephan. I., Baretton GB., Menauer F., Blasenbru and Vogts., Lohrsu. (2000):

Sjogren's syndrome and MALT lymphomas of salivary glands: a DNA-cytometric and interphase-cytogenetic study Mod. Pathol Jan, 13 (1): 4-12.

Steultjens MP.; Dekker J., Van Baar ME.; Oostendorp RA. and Bijlsma JW. (2001):

Muscle strength, Pain and disability in patients with osteoarthritis. Clin Rehabil Jun; 15 (3): 331-341.

Stewart HD. and Newton G. (1992):

Long term results of the Manchester Knee. Surface arthroplasty of the tibiofemoral joint. Clin. Orthop. May; (278): 138-146.

Swezey R.L. (1993):

Rehabilitation medicine and arthritis in: Arthritis and Allied condition, 12th ed., Philadelphia, Lea and Febiger, PP. 887-917.

Takashi Ohba, Yuko Takase, Morihiro Ohhara and Rej Kasukawa (1995):

Thrombin in the synovial fluid of patients with rheumatoid arthritis mediates proliferation of synovial fibroblast-like cells by induction of platelet derived growth factor. The Journal of Rheumatology (23): 1505-1511.

Teitz, N. (1986):

Reference range and laboratory values of clinical importance.

In Cecil textbook of Medicine. 17th edition Saunders Company, Philadelphia.

Thaler, E., and Lechner, K. (1981):

Antithrombin deficiency and thromboembolism. Thrombosis Research (20): 200-205.

Travis J., and Salvesen G. (1983):

Plasma Protein inhibitors.
Ann. Rev. bioch., 52, 655.

Trentham D.E., Dynesius, R.A. and David, J.R. (1978):

Passive transfer by cells of type II collagen induced arthritis
in rats.
J. Clin. Invest., 62: 359.

Von Kaulla E., Kurt M. and Von Kaulla M. (1976):

Antithrombin III and disease.
American J. of Clin. Path., Vol. 48, No.1, 69-80.

Walker I.D., Davidson J.F., Young, P. and Conki J.A., (1975):

Effect of anabolic steroids on plasma antithrombin III, α_2
macroglobulin and α_1 antitrypsin levels.
Thromb. Diath. Haemorrh. Sept. 30; 34 (1); 106-114.

Watt I. and Rozman B. (1996):

The role of MRI in the assessment of the rheumatological
disease.
Br. J. Rheum., 35: Suppl. 3: 1-47.

Walter, J.B. and Israel, M.S. (1987):

General Pathology. 6th Edition.
Churchill Livingstone, Edinburgh, London.

Watson, R.G. (1965):

A more specific method for detecting and quantitating
rheumatoid factors using a simple modification of the
rheumatoid arthritis test.
Am. J. Clin. Path., 43: 152.

Weinberg JB, Pippen AMM and Greenberg C.S. (1991):

Extravascular fibrin formation and dissolution in synovial
tissue of patient with osteoarthritis and rheumatoid arthritis.
Arthritis Rheum; 34: 996-1005.

Weinblatt Trentham D.E. and Fraster P.A. (1995):

Long term prospective trial of low dose methotrexate in
rheumatoid arthritis. 31: 147-175.

Weksler BB., Ley CW. Jaffe EA. (1978):

Stimulation of endothelial cell PGI₂ production by thrombin, trypsin, and ionophore A.
J. Clin Invest; 62: 923-38.

Weltz JI, Hudoba M, Massel D. and Senecal JL. (1990):

Clot-band thrombin is protected from inhibition by heparin-antithrombin II but is susceptible to inactivation by antithrombin III-independent inhibitors.
Blood 86: 385-391.

Wilson, M.E., and Poss R. (1992):

Osteoarthritis of the hip. In: Osteoarthritis: Diagnosis and Management. Saunders Company, Philadelphia, PP. 621-650.

Wintrobe, M.; Ritchard, J.; Boggs, D. and Bithell (1981):

Clinical Haematology Text book, 7th Edition. Lea Febiger. Philadelphia.

Zacharski LP., Brown FE., Memoli VA., Kisiel W., Kurdryk B.J. Rousseau SM., Hunt. J.A., Dunwiddie C. and Nutt EM. (1992):

Pathways of coagulation activation in situ in rheumatoid synovial tissue.
Clin. Immunol. Immunopathol. May; 63 (2): 155-162.

Ziff, M. and Baum, J. (1993):

Laboratory finding in rheumatoid arthritis. In "Arthritis and Allied conditions. 12th Ed., Lea and Febiger, Philadelphia, P. 367.

Zimmerman GA., Melintyre TM. (1985):

Thrombin stimulates adhesion of neutrophils in human endothelial cells in vitro.
J. Clin Invest; 76: 2235-2246.

Zvaifler, (1995):

Immunopathology of joint inflammation in arthritis.
J. Adv. Immunol. 16: 265.