

Debut

REFERENCE

- Abildgaard, N.; Nielsen, J.L. and Heickendorff, L. (1994):*
Connective tissue components in serum in multiple myeloma: analyses of propeptides of type I and type III procollagens, type I collagen telopeptide, and hyaluronan. *Am. J. Hematol.*, 46, 173-178.
- Akimoto, S.; Akokuro, K. and Shimosaki, J. (1996):*
Clinical usefulness of serum carboxy terminal propeptide of type I procollagen and pyridinoline cross-linked carboxy terminal telopeptide of type I collagen in patients with prostate cancer. *J. Pn. J. Clin. Oncol.*, 26: 157.
- Aloia, J.F.; Cohn, S.M.; Vaswani, A.; James, K. and Ellis, K. (1985):*
Risk factors of postmenopausal osteoporosis. *Obstet. Gynecol. Survey*, 40: 689.
- Anderson, J.J. (1996):*
Calcium, phosphorus and human bone development. *J. Nutr. Apr.*; 126 (4 suppl.): 11536-85.
- Azria, M. (1989):*
The value of biomarkers in detecting alterations in bone metabolism. *Calcif. Tissue Int.*, 45, 7-11.
- Bailey, A.J.; Wotton, S.F.; Sims, T.J. and Thompson, P.W. (1992):*
Post-translational modification in the collagen of human osteoporotic femoral head. *Biochem. Biophys. Res. Commun* 185: 801-805.
- Bain, S.D.; Bailey, M.C.; Celino, D.L.; Lantry, M.M. and Edwards, M.W. (1993):*
High dose estrogen inhibits bone resorption and stimulates bone formation in the ovariectomized mouse. *J. Bone Miner Res.* 8 : 435-42.
- Barham, D. and Trinder, P.; Bartel, H. and Bohmer, M. (1972):*
Clin. Chem. Acta 37: 193.

- Batge, B.; Diebold, J.; Stein, H.; Bodo, M. and Muller, P.K. (1992):**
Compositional analysis of collagenous bone matrix-A study
on adult normal and osteopenic bone tissue.
Eur. J. Clin. Invest. 22: 805-812.
- Beardsworth, L.J.; Eyre, D.R. and Dickson, I.R. (1990):**
Changes in urinary excretion of lysyl and hydroxylysyl.
Pyridinoline, two new markers of bone collagen turnover.
J. Bone Mineral Rws.; Vol. 5: 671.
- Bissell, D.M. and Roll, J. (1990):**
Connective tissue metabolism and hepatic fibrosis.
Hepatology. 1: 424.
- Bjarnason, N.H.; Christiansan, C.(2000):**
Early response in biochemical markers predicts long-term
response in bone mass during hormone replacement therapy
in early postmenopausal women.
Bone, 26: 561-9.
- Blomqviste, Risteli, L. Risteli, J. et al. (1996) :**
Markers of type I collagen degradation and synthesis in the
monitoring of treatment response in bone metastases from
breast carcinoma.
Br. J. Cancer, 73 : 1074 – 1079.
- Bodanksy, O. (1972):**
Acid phosphatase.
Clin. Chem., 15: 44.
- Body, J.J.; Clearen, A.; Pot, M. and Borkowsky, A. (1986):**
Serum osteocalcin (BGP) in tumor-associated
hypercalcemia.
J. Bone. Mineral. Res.; Vol. 1: 523.
- Boskey, A. L.; and Posner, A. S. (1984):**
Bone structure, composition and mineralization.
Orthop. Coen. North Am 15: 597.
- Braham, H.; Mallmin, H.; Michaelsson, K. and Strom, H. (1998):**
Relationship between bone mass and physical activity.
Clacif.
Tissue Int. May. 62 (5): 400-12.

Brown, J.P. and Josse, R.C. (2002) :

Clinical practice guide lines for the diagnosis and management of osteoporosis society of Canada.
CMAJ. November, 12 : 1-83.

Calvo, M.S.; Eyre, D.R. and Gundberg, C.M. (1996):

Molecular basis and clinical application of biological markers of bone turnover.
Endocrinol Rev 17: 333-68.

Chailurkit, L.O.; Saetung, S. Piaseu, N. et al. (2001) :

Biochemical markers of bone turnover and response of bone mineral density to intervention in early post menopausal women : An experience in a clinical laboratory.
Clin Chem, 47 : 1083 –1088.

Charhon, S.A. Chapyu, M.C.; Delvin, E.E.; Valentin-Opran, A. Edouard, C.M. and Meunier, P.J. (1985) :

Histomorphometric analysis of sclerotic bone metastases from prostate carcinoma with special refence to osteomalacia. Cancer, 51, 918-924.

Charles, P.; Mosekilde, L.; Risteli, L.; Risteli, J. and Eriksen, E.F. (1994):

Assessment of bone remodeling using biochemical indicators of type I collagen synthesis and degradation: relation to calcium kinetics.
Bone Miner, 24, 81-94.

Cheung, C.K.; Panesar, N.S.; Lau, E.; Woo, J. and Swaminathan, R. (1995):

Increased bone resorption and decreased bone formation in chinese patients with hip fracture.
Calcif Tissue int 56: 374-9.

Clifford J Rosen, and Alan Tenenhouse (1998):

Biochemical markers of bone turnover postgraduate medicine, Vol. 104 No.4.

Courvoisier, P.B. (1985):

Clinical diagnosis of osteoporosis.
Med. Hyg. Gen. 43: 3091-3095.

Cuttman, S. and Burger, H.G. (1997):

Use of androgens in postmenopausal women.
Current Opin Obstet Gynecol. 9 : 177-180.

Defots, L.J. (1991):

Bone protein and peptide assays in the diagnosis and management of Skeletal diseases (Review).
Clin. Chem. 37: 1143.

De la Piedra, C.; Diaz, M.M.; Diaz, D.E.; Lopez, G.E.; Gonzalez, P.E.; Caramelo, C. and Rapado, A. (1994):

Serum concentrations of carboxyterminal cross-linked telopeptide of type I collagen (ICTP), serum tartrate resistant acid phosphatase, and serum levels of intact parathyroid hormone in parathyroid hyperfunction.
Scand. J. Clin. Lab. Invest., 54: 11-15.

Delmas, P.D. (1988):

Biochemical markers of bone turnover in osteoporosis: Riggs BL, Melton J, editors. Osteoporosis: etiology, diagnosis and management.
New York: Raven Press: p. 297-316.

Delmas, P.D. (1990):

Biochemical markers of bone turnover for clinical assessment of metabolic bone disease.
Endocrinol. Metab. Clin. North Am. 19: 1-18.

Delmas, P.D. (1993):

Biochemical markers of bone turnover.
J. bone Miner. Res., 8 (suppl. 2) S549-S555.

Delmas, P.D. (1995):

Biochemical markers for the assessment of bone turnover.
In: Riggs BL, Melton III J, editors. Osteoporosis: etiology, diagnosis and management. 2nd edn.
Philadelphia: lippincott-Raven; 1995. P. 319-33.

Demers, L.M. (1997):

Markers of bone degradation and formation.
J. Clin. Lab. Invest. Suppl. 227: 12-20.

- Devine, A.; Dick, I.M.; Heal, S.J. and Criddle, R.A. (1997):*
Follow up study of the effect of calcium supplementation on bone density in elderly postmenopausal women.
Osteoporosis Int., 7(2): 23-28.
- Doumas, P.T.; Watson, W.A. and Biggs, H.G. (1971):*
Determination of S. albumin.
- Earnshaw, S.A.; Worley, A. and Hosking, D.J. (1997):*
Diet does not relate to bone mineral density after menopause.
Br. J. Nutr. July; 78 (1): 65-72.
- Eastell, R.; Delmas, P.D.; Hodgson, S.F. and Erikson, F.E. (1988):*
Bone formation rate in older normal women.
J. Clin. Endocrinol. Metab. 67: 741.
- Ebeling, R.P.; Peterson, J.M. and Riggs, B.L. (1992):*
Utility of type I procollagen propeptide assay for assessing abnormalities in metastatic bone disease.
J. Bone Miner. Res.; 7: 1243-50.
- Ebeling, P.R.; Atley, L.M.; Cuthrie, R.H. (1996):*
Bone turnover markers and bone density across the menopausal transition.
J. Clin. Endocrinol. Metab. 81(9): 3366.
- Elomaa, P.; Virkkunen, L.; Risteli, L. and Risteli, J. (1992):*
Serum concentration of cross-linked carboxy-terminal telopeptide of type I collagen (ICTP) is useful prognostic indicator in multiple myeloma.
Br. J. Cancer. 66: 337-341.
- Eriksen, E.F.; Charles, P.; Mosekilde, L.; Risteli, L. and Risteli, J. (1992):*
Serum markers of type I collagen formation and degradation in metabolic bone disease: correlation to bone histomorphometry.
J. bon. Min. Res. (in Press).

- Eriksen, E.F.; Charles, P.; Melren, F.; Mosekilde, L.; Risteli, L.; Resteli, J. (1993):**
Serum markers of type I collagen formation and degradation in metabolic bone disease: correlation with bone histomorphometry.
J. bone miner. Res. 8: 127-32.
- Eriksen, E.F.; Charles, P.; Melren, F. et al. (1998):**
Serum markers of type I collagen formation and degradation in metabolic bone disease: correlation with bone histomorphometry.
J. bone miner. Res. 8: 127-132.
- Eyre, D.R (1980):** Collagen molecular diversity in the body's protein scaffold science; 207: 1315.
- Eyre, D.T. (1995):**
The specificity of collagen cross-links as markers of bone and connective tissue degradation.
Acata Orthop. Scand. (Suppl. 266) 66: 166-170.
- Eyre, D.R.; Dichson, I.R. and Van Ness, K.P. (1988):**
Collagen cross-linking in human bone and articular cartilage: age- related changes in content of mature hydroxypridinium biochem.
J. 495-500.
- Filipponip, Pedettim, Beghe, F.; Giovagnini, B.; Miam, M. and Cristallini, S. (1994):**
Effects of two different bisphosphonates on Paget's disease of bone: ICTP.
Assessed. Bone, 15,261-267.
- Fink, E.; Cromier, C.; Steinmetz, P.; Kindermans, C.; LeBouc, Y.; Souberbielle, J.C.(2000):**
Differences in the capacity of several biochemical bone markers to assess high bone turnover in early menopause and response to alendronate therapy.
Osteoporosis Int: 11:295-303.

- Frevert, E.U. Biester, A.; Muller, M.J. Schmidt, G.H. Von Zur muhlen A. and Brabant, G. (1994):**
Markers of bone metabolism during short-term administration of thyroxine in healthy volunteers.
Eur. J. Endocrinol., 131,145-149.
- Fukushima, O.; Bekker. P.J.; and Gay. C.V. (1991):**
Ultrastructural localiazation of tartarate-resistant acid phosphatase (purple acide posphatase) activity in chicken cartilage and bone.
Am. J. Anat. 191(3): 228.
- Gallagher, J.C.; Riggs, B.L.; Eisman, J.; Hamster, A.; Arnard, S.B.; and Deluce, H.F. (1979):**
Intestinal calcium absorption and serum vitamin D metabolities in normal subjects and osteoporotic patients.
J. Clin. Invest. 64:729.
- Gallagher, J.C.; Riggs, B.L.; and Jerpback, G.M. (1980):**
The effect of age on serum immunoreactive parathyroid hormone in normal and osteoporotic women.
J. Lab. Clin. Med. 95:373.
- Garnero, P.; and Delmas, P.D. (1997):**
Bone markers.
Baillieres clin. Rheumatol. 11(3):517-37.
- Garnero, P.; Delmas, P.D.(1998) :**
Biochemical markers of bone turnover.
Endocrinol Metab Clin North Am 27:303-23.
- Glimcher, M.J.(1975):**
Studies of the structure organization and reactivity of bone collagen. In Gibson T (ed): proceeding of the International sumposium on wound healing.
Montreaux. Foundation of International Cooperative Medical science. 253.
- Gordan, S.; (1976):**
In: Clinical management of osteoporosis.
Gordan, S. and Vaughan, C. (eds). H.M.+M. publisher
England

Greenspan, S.L.; Rosan, H.N. and Parker, R.A. (2000):

Early changes in serum N-telopeptide and C-telopeptide cross-linked collagen type I predict long-term response to alendronate therapy in elderly women,
J. Clin Endocrinol Metab 85:3537-40.

Gundberg, C.M. (2000):

Biochemical markers of bone formation.
Clin Lab Med 20:489-501.

Hakala, M.; Risteli, L.; Manelius, J.; Nieminen, P. and Risteli, J. (1993):

Increased type I collagen degradation correlates with disease severity in rheumatoid arthritis.
Ann. Rheum. Dis., 52, 866-869.

Halleen, J.M.; Nakasato, Y.R.; Janckila, A.J.; Vaananen, H.K.; Wlton, S.P.; Yam, L.T. (1999):

Clin significance of immunoassays for type 5 tartrate-resistant acid phosphatase.
Clin Chem; 45:2150-7

Halleen, J.M.; Alatalo, S.L.; Hentunen, T.A.; Monkonen, J.M. and Vaaranen, A.K. (2000):

Rapid screening method for osteoclast differentiation in vitro that measure tartrate-resistant acid phosphatase 5b activity secreted into the culture medium.
Clin Chem. 46: 1751-4.

Hampton, L.; Eastell, R. and Colwell, A. (1990):

Urinary collagen crosslinks are highly correlated with radioisotopic measurement of bone resorption.
In: Christiansen C, Overgaard K, editors. Osteoporosis 1990. Copenhagen: Osteopress; p. 469-77.

Hanaoka, H. (1979):

The origin of the osteoclast.
Clin Orthop Rel. Res., 145:252.

Hanson, D.A.; Weis Mae, Bollen A.M.; Maslan, S.L.; Singer, F.R. and Eyre, D.R. (1990):

A specific immunoassay for monitoring human bone resorption: quantitation of type collagen cross linked N-telopeptide in urine.
J. Bone Miner. Res; 7, 1251-1258.

Hart, S.M. and Eastell, R. (1999):

Biochemical markers of bone turnover.
Curr Opin Nephrol Hypertens; 421-7.

Hassager, C. Fabbri-Mabelli G, Christiansen, C. (1993):

The effect of the menopause and hormone replacement therapy on serum carboxyterminal propeptide of type I collagen.
Osteoporosis Int 3:50-2:

Henry, R.J., Cannon D.C. and Winkelman J.W. (1974):

Clinical chemistry, principles and techniques Harper & Row
2nd Ed.

Holtrop, M.E. (1975):

The ultrastructure of bone.
Ann. Clin. Lab. Sci 5:264.

Hreshchyshyn, M.M.; Hopkins, A.; Zylstra, J. and Anbar, M. (1988):

Effect of natural menopause, hysterectomy and oophorectomy on horectomy lumbar spin and femorabl neck.
Bone densities.
Obstet. Gynaecol., 72:631.

Hughes, D.B.; Dallal, G.F.; Krall, E.A. and Harris, S.(1991):

Effect of vitamin D supplementation on winter time and overall bone loss in healthy postmenopausal women.
Ann. Intern. Med., 115-505.

James, I.T, Walne, A.J. Perrett, D.(1996):

The measurement of pyridinium crosslinks: a methodological overview.
Ann Clin Biochem 33:397-420.

Jee, W.S.S. (1983):

The skeletal tissues; In: L. Weiss; L. Lansing (Eds.):
histology: cell and tissue biology,
5th Ed. New York, Elsevier Biomedical, P.33.

Jodar, Gimeno, E. Munoz-Torres M, Escobar,- and Jimenez, F.(1997):

Identification of metabolic bone disease in patients with endogenous hyperthyroidism: role of biological markers of bone turnover.
Calcif Tissue Int; 370-6.

Johnston, C.C. and Epstein, S. (1981):

Clinical, biochemical, radiographic, epidemiological and economic features of osteoporosis.

Ortho. Clin. N. Amer., 12 (3):35-37.

Kelly, P.J. Pocock, N.A.; and Sambrook, P.N. (1989):

Age and menopause-related changes in indices of bone turnover.

J Clin Endocrinol Metab. 69(6):1160-5.

Kerstjens. H.A.; Postma. D.S.; Van Doormaal, J.J.; van zanten a.k.; Brand P.L., Dekhuijzen P.N.; and Koeter, G.H. (1994):

Effect of short-term and long-term treatment with inhaled corticosteroids on bone metabolism in patients with airways obstruction.

Dutch Cnslid study Group. Thorax, 49,652-656.

Kleerekoper, M.; Tolia, K. and parfitt, A.M. (1981):

Nutritional, Endocrine and demographic aspects of osteoporosis.

Ortho. Clin. North. America 12:547.

Kraenzlin, M.E.; Taylor, A.K. and Baylink, D.J. (1989):

Biochemical markers for bone formation and bone resorption: in: Lindh, E. and Thorell, J.L. (Eds.): clinical impact of bone and connective tissue markers.

Academic Press, London. 289-303.

Kusec, V.; Stavljenic-Rukavina, A. and Cubrilo- Turek, M. (1998):

biochemical markers of bone metabolism in postmenopausal women.

Croat. Med. J.Mar.; 39(1):23-7.

Kushida, K.; Tadahaski, M.; Kawana, K. and Inoue, T.(1995):

Comparison of markers for bone formation and resorption in premenopausal and postmenopausal subjects and osteoporotic patients.

J. Clin. Endocrinol. Metab., 80:2447-50.

Kyd, P.A.; Vooght, K.; Kershoff, F.; Thomas, E. and Fairney, A.(1999):

Clinical usefulness of biochemical resorption markers in osteoporosis.

Ann. Clin Biochem 36:483-91

Lane, J.M. and Vigorita, V.M. (1984):

Osteoporosis arth. Cline.
North America, Vol. 15, No.4.

Lane, J.M.; Riley, E.H.; and Wirganowicz, P.Z. (1994):

Osteoporosis: diagnosis and treatment.
Course Lect.; 46:445:58.

La-Orchailurkit, Boonsong, Ongphiphadhana Kul, Noppawanpiaseu, Sunee Saetung, and Radata Rajatanavin (2001):

Biochemical markers of bone turnover and response of bone mineral density to intervention in Early postmenopausal women: an experience in a clinical laboratory.
Clinical chemistry 47:6 1083:1088

Lee, A.J.; Hodges, S. and Eastell, R.(2000):

Measurement of osteocalcin.
Ann Clin Biochem. 37:432-46.

Lindsay, R. (1996):

The menopause and osteoporosis.
Obstetric and gynecol, No. 2, Vol. 8/7, PP. 165-195.

Lundgren, E.; Rastad, J.; Thrufgell, E. and Akerstrom, G. (1997):

Population-based screening for primary hyperparathyroidism with serum calcium and parathyroid hormone values in menopausal women.
Surgery, Mar.; 121(3):298-94.

Manolagas, S.C. and Jilka, R.L. (1995):

Bone marrow, cytokines, and bone remodelling: emerging insights into the pathophysiology of osteoporosis.
N. Engl J. Med 332(5):305-11

Margareta, Hedstrom¹, Jansvensson² and Nilisdalen¹ (2000):

Biochemical Bone markers and bone density in hip fracture patients.
Acta orthop scand; 71(4).409-413.

Marie, P.J. (1982):

Structure, organization and Healing; in: R.L. Cruess (Ed.):
The musculo-skeletal system. Embryology, Biochemistry and physiology.
New York, Churchill Livingstone, P.109

Marshall, P.H.; Grilly, R.G. and Norden, B.E. (1977):

Plasma androstenedione and estrogen levels in normal osteoporotic postmenopausal women.
Br. Med. J. 2:1177.

May, H.; Murphy, S. and Khaw, K.T. (1995):

Bone mineral density and its relationship to skin color in caucasian females.
Eur. J. clin. Invest. Feb, 25(2):85-9

Meanier, P.J.; Delmas, P.D.; Eastell, R. et al. (1999):

Diagnosis and management of osteoporosis in post menopausal women: clinical guidelines. International Committee for osteoporosis clinical. Guidelines. Clin. Ther., 21, 1025-1044.

Midgette, A.S.; and Baron, J.A. (1990):

Cigarette smoking and the risk of natural menopause.
Epidemiology, 1:474.

Morol, Gazzarini, C.; Crivellari, E.; Talamini, R. D.E.; and Bernard, B. (1993):

Biochemical markers for detecting bone metastases in patients with breast cancer.
Clin. Chem., 39:131-134.

Morris, G.L. and Dodd, J.G. (1990) :

Epidermal growth factor receptor mRNA levels in human prostatic tumors and cell lines. J. Urol., 143, 1272-1274.

Nordin, B.E. (1998):

Calcium and osteoporosis.
Nutrition Jul. Aug.; 13(7-8):664-86.

Nordin, G.; Horseman, A.; and Grilly, G. (1980):

Treatment of spinal osteoporosis in postmenopausal woman.
Br. Med. J. 280:451.

Nordin, B.E.; Roberston, A. and Saemark, R.F. (1985):

The relation between calcium absorption, serum dehydroepiandrosterone and vertebral mineral density in postmenopausal women.
J. Clin. Endocrinol. Metab. 60:651:

Owen, M.(1970):

The origin of bone cells.
Int. Rev.cytol. 213.

Paimelo, L., Leirisalo Repo M., Risteli, L., Hakala, M., Helve, T., and Risteli (1994):

Tye I collagen degradation produced in serum of patients with early rheumatoid arthritis, relationship to disease activity and radiological progression in 3 years follow up.
Br.J. Rheumatol 33:1012-1016.

Paul Grech, T.D.; Martin, N.A. and Barrington, P.J. (1985) :

Diagnosis of Metabolic Bone Diseases, 11 th ed., 52-85.

Percherstorfor, M.; Zimmer, Roth, L.; Schilling, T.; Waitage, H.W.; and Schmidt, H.(1995):

The diagnostic value of urinary pyridinum cross-links of collagen, serum total alkaline phosphatase, and urinary calcium excretion in neoplastic bone disease.
J. clin. Endocrinol. Metab.; Jan, 80(1):97-103.

Price, C.P.; (1995):

Multiple forms of human serum alkaline phosphatase: detection and quantitation.
Ann Clin Biochem :355-72.

Price, C.P. Thompson, P.W. (1995):

The role of biochemical tests in the screening and monitoring of osteoporosis.
Am Clin biochem 32-17.

Richard, A.E.; Geoffrey, M.M.S; and Elizabeth, K. (1988):

Bone mass is low in relatives of osteoporotic patients.
Ann. Int. Med. 109:870.

Riggs, B.L.; Melton. III L.J. (1986):

Involutional Osteoporosis.
N. Engl Med 314:1676-1686.

Riggs, B.L.; and Melton, L.J. (1990):

Clinical heterogeneity of involutional osteoporosis: Implications of preventive therapy.
J. Clin. Endocrinol. Metab., 70,1299.

Riggs, B.L.; Wahner, H.W.; Dunn, W.L.; Mazess, R.B.; Offord, K.P.; and Melton, L.J. (1981):

Differential changes in bone mineral density of the appendicular and axial skeleton with aging: relationship to spinal osteoporosis.

J. Clin. Invest 67:328.

Ristelli, L, Ristelli, J.(1993):

Biochemical markers of bone metabolism.

Ann. Med 25:385-93.

Risteli, J.; Elomaa, I.; Niemi, S., Novamo, A. and Risteli, L.(1993):

Radioimmunoassay for other pyridinoline cross-linked carboxy-terminal telopeptide of type I collagen: A new serum marker of bone collagen degradation.

Clin. Chem., 39:635-40

Robins, S.P. (1999):

Biochemical markers of bone metabolism.

CPD bulletin Clin Biochem. 1:116-21

Rodriguez, G.P.; and Claus Walker, J. (1984):

Measurement of hydroxylysine glycosides in urine and its application to osteoporosis.

J. Chromatogr.; 308-65

Rosen, C.J. (1996):

Biochemical markers of bone turnover. In: Rosen C.J, ed. Osteoporosis: diagnostic and therapeutic principles.

Totowa, N: Humana press, 129-41.

Ross, P.D.; Wasnich, R.D. and Vogel, J.M. (1996):

Detection of prefracture spinal osteoporosis using bone mineral absorptiometry.

J. bone. Miner. Res. 3-1

Rude, R.K. and Singer, F.R.; (1991):

Primary hyperparathyroidism; in current diagnosis.

8th. ed., R.B. Conn (Ed.) Philadelphia, W.B. Saunders Co., P. 859.

Satorioa, Conti, A. and Monzanim. (1993a):

New markers of bone and collagen turnover in children and adults with growth hormone deficiency.

Postgrad. Med. J., 69,846-850.

Sartorio, A.; Conti, A.; monzani M, Morabito, F.; and Faglia, G. (1993b):

Growth hormone treatment in adults with G.H deficiency: effect on new biochemical markers of bone and collagen turnover.

J. Edocrinol. Invest., 16,893-898.

Sato, M.; Sasaki, M. and Nagai, Y. (1987):

Increased urinary excretion of collagen metabolites in cadmium metallothionin nephropathy.

Arch. Toxicol.; 61,2:116

Saunders, L.E. (1988):

Specialized connetive tissue, cartilage and bone: general histological principles and primary tissues; in: text atlas of Hostology.

Saunders W.B. Internatioanl Edition, P.168.

Seibel, M.J.; Woitge, H.W. and Pecherstorfew, M. (1996):

Serum immunoreactive bone sialoprotein as a new marker of bone turnover in metaboic and malignant bone disease.

J. Clin Endocrinol Metabl 81:3289-94.

Slovik, D.M.; Adams, J.S.; Neer, R.M. and Hollick, M.F. (1981):

Deficient production of $1.25\text{ (OH)}_2\text{ D}_3$ in elderly osteoporotics.

N. Engl. J. Med., 305:372.

Stevenson, J.C.; Cust, M.P.; and Gangar, K.F. (1990): Effects of transdermal versus oral HRT on bone density in spine and proximal femur in postmenopausal women.

Lancet. 336:1327

Swaminathan, R. (2001) :

Biochemical markers of bone turnover.

Clin. Chem Acta, 313 : 95-105.

Talmage, R.V.; Cooper, C.W.; and Toverud, S.U. (1982): The physiological significance of calcitonin; in: bone and mineral research, Edited by Peck, W.A.

Amestrdam, Excerpta Medica, p. 74-143.

- Sartorio, A.; Conti, A.; monzani M, Morabito, F.; and Faglia, G. (1993b):**
Growth hormone treatment in adults with G.H deficiency: effect on new biochemical markers of bone and collagen turnover.
J. Endocrinol. Invest., 16,893-898.
- Sato, M.; Sasaki, M. and Nagai, Y. (1987):**
Increased urinary excretion of collagen metabolites in cadmium metallothionin nephropathy.
Arch. Toxicol.; 61,2:116
- Saunders, L.E. (1988):**
Specialized connetive tissue, cartilage and bone: general histological principles and primary tissues; in: text atlas of Hostology.
Saunders W.B. Internatioanl Edition, P.168.
- Seibel, M.J.; Woitge, H.W. and Pecherstorfew, M. (1996):**
Serum immunoreactive bone sialoprotein as a new marker of bone turnover in metaboic and malignant bone disease.
J. Clin Endocrinol Metabl 81:3289-94.
- Slovik, D.M.; Adams, J.S.; Neer, R.M. and Hollick, M.F. (1981):**
Deficient production of $1.25\text{ (OH)}_2\text{ D}_3$ in elderly osteoporotics.
N. Engl. J. Med., 305:372.
- Stevenson, J.C.; Cust, M.P.; and Gangar, K.F. (1990):** Effects of transdermal versus oral HRT on bone density in spine and proximal femur in postmenopausal women.
Lancet. 336:1327
- Swaminathan, R. (2001) :**
Biochemical markers of bone turnover.
Clin. Chem Acta, 313 : 95-105.
- Talmage, R.V.; Cooper, C.W.; and Toverud, S.U. (1982):** The physiological significance of calcitonin; in: bone and mineral research, Edited by Peck, W.A.
Amestrdam, Excerpta Medica, p. 74-143.

Taube, T.; Kylmala, T.C.; Lamberq-Allardt, C.; Tammela T.L.J. and Elomaa. I. (1994):

The effect of clodronate on bone in metastatic prostate cancer. Histomorphometric report of a double blind randomised placebo study.

Eur. J. cancer, 30, 751-758

Termine, J.E.; Belcourt, A.B.; and Conn, K.M. (1981):

Mineral and collagen-binding proteins of fetal calf bone.

J. biol. Chem. 256:10403.

Theodorou, S.J.; Theodorou, D.J. and Sartoris, D.J. (2003):

Osteoporosis and Fractures : the size of the problem.

Hosp Med 2003 Feb; 64 (2) : 87-91.

Tietz, N.W.; Textbook of Clinical Chemisty, W.B.; Saunders, Co. and Philadelphia, P.A. (1986) :

PP-52-53 (techniques and procedures to minimize laboratory infections) and pp 478-497 (specimen collection and storage recommendations).

Triffitt, J.T. (1987):

The special proteins of bone tissue.

Clinical science, 72:399-408.

Tromp, A.M.; Ooms, M.E.; Popp-Snijders, C.; Roos, J.C. and Lipo, P.(2000):

Predictors of fractures in elderly women.

Osteoporosis Int; 11:134-40

Uebelhardt, Gineyts, E.; Chapuy M.C. and delmaspd. (1990):

Urinary excretion of pyridinium crosslinks: a new marker of bone resorption in metabolic bone disease.

Bone & Mineral, 8, 87-96.

Valimaki, M.J.; Tahtela, R.; Jones, J.D.; Peterson J.M. and Riggs B.L. (1994):

Bone resorption in healthy and osteoporotic postmenopausal women: comparsion markers for serum carboxy-terminal telopeptide, of type I collagen and urinary pyridinium cross-links.

Eur. J. Endocrinol., 131, 258-262.

Valimaki, M.J.; Tahtela, R.; Jones, J.D.; Peterson J.M. and Riggs B.L. (1998):

Bone resorption in healthy and osteoporotic postmenopausal women: comparison markers for serum carboxy-terminal telopeptide, of type I collagen and urinary pyridinium cross-links.

Eur. J. Endocrinol., 130, 258-262.

Warshwsky, H.(1982):

Embryology and development of skeletal system, in, R.L. Cruess (Ed.): the musculoskeletal system, embryology, biochemistry, physiology.

New York, churchill Livingstone, P. 33

Watts, N.B.(1999):

Clinical utility of biochemical markers of bone remodelling.
Cin. Chem 45:1858-60.

Wichers, M.; Schmidt, E.; Bidlingmaier, F. and Klingmuller, D.(1999):

Diurnal rhythm of cross links in human serum.
Clin chem.

Yagiy, Yoneda, M.; Okada, M.; Omoto, A. and Nakamura, M. (2002):

Apssibility of the prognosis factor of serum cross. Linked carboxy terminal telopeptide region of type I collagen (ICTP) as a marker of bone metastasis.

Kaku Igaku Nov; 39 (4) : 485-91.