

References

VI - REFERENCES

- *A.O.A.C., Sec. 26, 018 and 019 (1975): Association of Official Analytical Chemists (1975). Official methods of Analysis. Chapter 26, Natural poisons, separate revised March 1975. Association of Official Analytical Chemists. Washington, D.C.
- *A.O.A.C. (1980): Official Methods of Analysis of the Association of Official Analytical Chemists. 13th ed. Washington D.C., 20044.
- *A.O.A.C. (1990): Association of official Agricultural chemists. Official Methods of Analysis. 15th ed. A.O.A.C. Washington. U.S.A.
- *Abdel-Fattah, H.M.; Moubasher, A.H. and Swelim, M.A. (1981):

 Studies on air-borne fungi at Qena II-Diurnal fluctuations.

 Zeit. Fur. Allg. Mikro 21:177-179.
- *Abdel-Hafez, S.I.I.; Shoreit, A.A.M.; Abdel-Hafez, A.I.I. and El-Maghraby, O.M.O. (1986): Mycoflora and Mycotoxin-producing fungi of air-dust particles from Egypt. Mycopathologia 93:25-32.
- *AbolAlea, M.F. (1996): Studies on deterioration aspects of some grains during storage and its control. M.Sc. Thesis Faculty of Agriculture. Moshtohor, Zagazig University Egypt.
- *Afify, A.S. (1986): Biochemical studies on soybean enzyme inhibitors during germination. Ph. D. Thesis, Faculty of Agriculture, Cairo University.

- *Ainsworth, G.C. and James, P.W. (1971): Ainsworth and Bishy's Dictionary of the fungi. (6th ed) commonwealth Mycological Institute, kew, surrey, UK, pp. 663.
- *Akinyele, I.O. and Fasaye, O.A. (1988): Nutrient quality of corn and sorghum supplemented with cow peas (U. unguiculata) in the traditional manufacture of Ogi. Journal of Food Science, 53 (6): 1750
- *Amein, A.M. (1970): Survey of fungi associated with sorghum (Sorghum vulgare Pers.) and Fenugreek (Trigonella Foenum-graecumL.) seeds in Assiut province. M.Sc. Thesis, Faculty of Agriculture, Assiut University.
- *Anderson, R.A.; Montgomery, R.R., and Burbridge, L. H. (1969):

 Low fat endosperm fractions from grain sorghum. Cereal

 Science Today.14:366.
- *Anon (1993): Soybean Processing for food Uses. International soybean program (INSOY). Ed. Tanteeratarm, K. Illinols, Champaign.
- *Anon, (1995): Annual book, (central authority of statistics). A.R.E.
- *Anthony, P.P (1977): Cancer of the liver: Pathogenesis and recent etiological factors. Transactions of the Royal Society of Tropical Medicine and Hygiene. 71:466-470.
- *Applewhite, T.T. (1981): Nutritional effects of hydrogenated soy oil.

 Journal of American Oil Chemistry Society, 58: 260-268.

- *Arulnandhy, V.; Senanayake, Y.D.A. (1991): Changes in viability, vigour and chemical composition of soybean seeds stored under the humid tropical conditions. Legume-Research, 1991, 14:3, 135-144; 21 ref.
- *Asao, T. G.; Buchi, N.M.; Abd El-Kader, S.B.; Chang, E.L. Wick and Wogan, G.N. (1965): The structures of aflatoxins B₁ and G₁. Journal of the American Chemistry Society, 87:882-886.
- *Axtell, J.D.; Ejita, A., and Munck, L. (1982): Sorghum nutritional quality-progress and prospects. In proceeding of the International symposium on sorghum in the Eighties, p. 589, ICRISAT, patancheru po, A.P., India.
- *Azzat, K.S. and Pearce, R.S. (1980): Fatty acids of lipid from cultured soybean and rape cells. Phytochemistry. 19:1375.
- *Baker, D.; Neustadt, M.H. and Zeleny, L. (1959): Relationships between fat acidity values and types of damage in grains. Cereal Chemistry, 36: 308.
- *Barakat, M.I.E. (1988): Soybean seed deterioration caused by fungi under storage condition. Ph. D. Thesis, Faculty of Agriculture Cairo, University.
- *Baran, M.; Venglovsky, J.; Valvovcik, J. and Janotikova, I. (1992):

 Maize storage in controlled CO₂ atmosphere.

 Polnohospodarstvo (CSFR), 38 (4): 249.
- *Baruch, H.K. (1961): The air spora of a cowshed. J.Gen. Microbiol. 25:483-491.
- *Bash, G., and Rae, I.D. (1969): The structure and chemistry of aflatoxins. In aflatoxin, ed. L.A. Goldblatt. Academic Press, New York, pp. 55-75

- *Beer, J.V. (1969): Isolation of pathogenic fungi from waterfowl. In "Isolation Methods for Microbiologists" (D.A. Shapton and G.W. Gould, eds), PP. 103-111. Academic press, London & New York.
- *Berk, Z. (1992): Technology of production of edible flours and protein products from soybeans. (FAO, food and Agriculture Organization of the United nations, 1992).
- *Beverdrof, W.D.; Hume, D.J.; Muscat, J.C.; Gostovic, F., and Vrie, Q. Van de. (1987): OAC Scorpio Soybean Canadian Journal of plant science 67(1): 255-256.
- *Bottomley R.A.; Christensen, C.M. and Geddes, W.F. (1950): Grain storage studies. The influence of various temperatures, humidities, and oxygen concentration on mold growth and biochemical changes in stored Yellow corn. Cereal Chemistry, 27: 271-296.
- *Bressani, R. (1981): The role of soybeans in food systems. Journal of American Oil Chemistry Society, 58: 392-400.
- *Brignoli, C.A.; Kinsella, J. E. and Weihrauch, J. L. (1976): V. unhydrogenated fats and oils. Comprehensive evaluation of fatty acids in foods. Journal of American Dietic Association, 68: 244-229.
- *Bullard, W.R.; Garrison, M.V.; Kilburn, S.R. and York, J.O. (1980):

 Laboratory comparisons of polyphenols and their repellent characteristics in bird resistant sorghum grains. Journal of Agricultural and Food Chemistry, 28:1006.

- *Bullerman, L.B. (1974): A screening medium and method to detect several mycotoxins in mold cultures. Milk food Technology, 37 (1): 1-3.
- *Butler, L.G.; Riedl, D.J.; lebryk, D.G. and Blytt, J.H. (1984):

 Interaction of proteins with sorghum tannins mechanism,
 specificity and significant Journal of the American Oil
 Chemistry Society, 61:916.
- *Chakravorty, S.C. (1969): Chemical study of sorghum as a staple food in India. 1. protein content in relation to colour of jowar (sorghum vulagare) grains labdev, part B 7:127-129. (C.F. chem. Abst. 72: 2290E, 1970).
- *Chang, S.S. (1979): Flavor and flavor stability of foods. Journal of American Oil Chemistry Society, 56: 958A.
- *Chang., S.B.; Abd El-Kader, M. M.; Wick E.L. and Wogan, G.N. (1963): Aflatoxin B₂: Chemical identity and biological activity. Science, 142:1191-1192.
- *Cheftel, J.C.; Cuq, J.L., and lorient, D. (1985): Protein alimentaires.

 Tec. and Doc. Lavoisier, Paris. (C.F. Technology of production of edible flours and protein products from soybean. FAO, food and Agriculture Organization of the United nations, 1992).
- *Chibber, B.A.K.; Mertz, E.T. and Axtell, J.D. (1978): Effect of dehulling on tannin content, protein distribution and quality of high and low tannin sorghum. Journal of Agricultural and Food Chemistry, 26:679.

- *Christensen, C.M. (1957): Deterioration of stored grains by fungi.
 Botanical Review, 23:108-134.
- *Christensen C.M., (1967): An note on invasion of durum wheat by storage fungi. Cereal Chem. 44:100-102.
 - *Christensen, C.M. and Dorworth, C.E. (1966): Influence of moisture content, temperature and time of invasion of soybean by storage fungi. Phytopathology. 56: 412.
 - *Christensen, C.M. and Kaufmann, H.H. (1969): Grain storage-the role of fungi in quality loss. Minneapolis: University of Minnesota press, 153 pp. (C.F. Seed pathology 1: 282-297, 1977).
- *Ciegler A.; Peterson, R.E.; Logoda, A.A., and Hall, H.H. (1966):
 Aflatoxin production and degradation by Aspergillus flavus
 in 20 Litre fermentors. Appl. Microbial, 14: 826-833.
- *Clayton, T.A. and Morrison, W.R. (1972): Change in flour lipids during the storage of wheat flour. J. Sci. Fd. Agric, 23:721.
- *Clinton, P.K.S. (1960): Seed-bed pathogen of groundnuts in the Sudan and an attempt control with an artificial testa. Journal of Experimental Agricultural, 28: 211-222.
- *Crawford, D.E. and Gillingham, J.T. (1967): Oil and protein content of soybean grown in South Carolina. S. Carolina Agr. Expt. Sta. Bull. AE 303. (C.F.Soybeans: Chemistry and Technology, Vol. 1-proteins, Rev. nd printing, Ed. Smith, A.K. and Circle, S.J. Westport, Conn. Avipublishing Co., 1978).

- *D'ercole, N. and Nipoti, P. (1979): Microbiological analysis of sorghum seeds. Informatore Fitopatologico 29:7-12 (C.F.R. P.P. 59: 2164, 1980).
- *Davis, N.D. and Diener, U.L. (1970): Environmental factors affecting the production of aflatoxin. In proceeding of the first U.S. Japan Conference on toxic Microorganisms. Ed. M. Herztberg. U.S. Govt. printing office, Washingaton, D.C.
- *Davis, N.D. and Diener, U.L. (1979): Mycotoxins. In food and Beverage Mycology, ed. L.R. Beuchat. AVI Publishing company, West Port, CT, pp. 397-444.
- *Davis, N.D.; Diener, U.L. and Eldridge, D.W. (1966): Production of aflatoxins B, and G, by Asperillus flavus in a semisynthetic medium. Applied Microbiology, 14: 378-380.
- *Deyoe, C.W. and Shellenberger, J.A. (1965): Amino acids and proteins in sorghum. Journal of Agricultural and Food Chemistry, 5:446.
- *Deyoe, C.W.; Shoup, F.K.; Miller, G.D.; Bathurst, J. D.; liang, P.E. Sanford and Murphy, L.S. (1970): Amino acid composition and energy value of immature sorghum grains. Cereal Chemistry, 47: 363-368.
- *Diener, U.L. and Davis, N.D. (1966): Aflatoxin production by isolates of Aspergillus flavus. phytopathology, 56:1390-1393.
- *Diener, U.L. and Davis, N.D. (1968): Effect of environment on afatoxin production in freshly dug peanuts. Tropical Science, 10:22-28.

- *Diener, U.L.; Cole, R.J.; Sanders, T.H.; Royne, G.A.; Lee, L.S., and Klich, M.A., (1987): Epidemiology of aflatoxin formation by Aspergillius flavus. Annual Review of phytopathology. 25, 249-270.
- *Diener, U.L.; Jackson, C.R.; Cooper, W.E.; Stpes, R.J. and Davis, N.D. (1965): Invasion of peanut pods in the soil by Aspergillus flavus. Plant Dis. Reptr., 49:931-935.
- *Dirks, B.M. (1954): The properties of some fungal lipases and their role in grain deterioration. Diss. Abst. 14:458-459 (C.F.R.A.M. 34: 101,1955).
- *Dogget, H. (1988): Sorghum, Utilization of grain sorghum. In: Sorghum ed: longman Scientific & Technical Co. published in U.S. New York, p. 428.
- *Donangelo, C.M.; Trugo, L.C.; Turgo, N.M. and Eggum, B.O. (1995): Effect of germination of legume seeds on chemical composition and on protein and energy utilization in rats. Food Chemistry, 53:23-27.
- *Drinah, B.C.; Nyirend, B. and Vohra, P. (1990): Nutritional improvement of tannin containing sorghum by sodium bicarbonate. Cereal Chemistry, 67 (6): 533.
- *Dube, S.; Shukla, H.S. and Tripathi, S.C. (1988): Changes in sugar and protein contents of wheat due to Aspergilli. Indian phytopathology, 41 (4): 633.
- *Dubois, M.A.; Gilles, J.K.; Hamilton, P.A.; Rebers, P.A. and smith, P.A. (1956): A calorimetric method for determination of sugar and related substances. Analytical Chemistry, 28:350.
- *Dukes, P.D., and Apple, J.L. (1965): Effect of oxygen and carbon dioxide tensions on growth and inoculum potential of

- phytophthora parasitica var. nicotiana. Phytopathology. 55:666-669.
- *Edds, G.T. (1973): Acute aflatoxicosis: A review. Journal of American Veterinary Medical Association. 162:304-308.
- *Eggum, B.O. (1977): Nutritional aspects of cereal proteins. In Genetic Diversity in plant. P. 349-369. Planum press, New York.
- *El-Bazza, Z.E. (1979): Screening and studies on certain toxin-producing fungi in local foods and feeds. M. of Pharma. Sc. Degree. Faculty of Pharmacy, Cairo University, 1-180.
- *El-Bazza, Z.E. (1983): Studies on the effect of gamma irradiation on the growth and toxin production by aflatoxin producing fungi. Ph. D. Thesis. Faculty of Pharmacy, Cairo University, 1-165...
- *El-Gendy, M.M. (1982): Changes which happened in cereals during storage. Food Industries. Part II, Bread Technol., 3rd ed., 480.
- *El-Habbal, M.S.; Nadia, Y. and Salwa, S.G. (1991): Yield and storage ability of soybean seeds as affected by ethephon application and ripening period. Annals of Agricultural-Science, Moshtohor (1991) 29: 1, 241-253; 13 ref.
- *El-Khalafy, H.M. and Meara, M.L. (1970): The composition of Egyptian soybean oils. Grases Aceites (Seville) 21:63-66.
- *Ellis, W.O.; Smith, J.P.; Simpson, B.K. and Oldham, J.H. (1991):

 Aflatoxins in food: occurrence, biosynthesis, effects on organisms, detection, and methods of control. Crit. Rev. Food Sci. Nutr. 30, 403-439.
- *Eppley, R.M. (1968): Screening method for zearalenone, aflatoxins and Ochratoxin A. Journal of AOAC International, 51:74-78.

- *Euclydes, M.P.; Junqueira, R.; Maffia, L. and Kibaaka, C.K. (1983):
 Characterization of sorghum (sorghum bicolor) flours
 processed in mills used for corn and wheat. Protein quality.
 Rev. Cers. 30:199. (C.F. Chem. abst. 100: 21757a).
- *Farag, R.S. (1990): Effects of fungal infection and agrochemical on the chemical composition of some seeds and aflatoxin production. Bulletin of Faculty of Agriculture Cairo, University, 41:1, 43-62.
- *Farag, R. S.; Ali, M.N. and Taha, S.H. (1990): Use of some essential oils as natural preservation for butter. Journal of American Oil Chemistry Society, 68 (3): 188-191.
- *Farag, R.S.; El-Kholony, A.A. and Belal, M. (1981): Influence of some herbicides on the physical and chemical characteristics of soybean oil. Fette-Seifen Anstrichmittel 83: 360.
- *Farag, R.S.; El-leithy, M.A.; Basyony, A.E. and Daw, Z.Y. (1986):

 Effects of varied substrates on aflatoxin production by A.

 parasiticus. Journal of American Oil Chemistry Society, 63:
 1024-1026.
- *Farag, R.S.; Mohsen, S.M.; Khalil, F.A. and Basyony, A.E. (1985):

 Effect of fungal infection on the chemical composition of wheat, soybean and sesame seeds. Bulletin of Faculty of Agriculture Cairo, University, 36 (1): 159.
- *Flood, A.E. and Priestley, C.A. (1973): Two improved method for the determination of soluble carbohydrates. Journal of Science, Food Agriculture, 24: 953.
- *Frazier W.C. (1977): Food microbiology. Tata MaGrawhill Publishing Comp. LTD, New Delhi.

- *Ghali, Y.; Abdel-Samed, A.; Basyony, A.E.; Ibrahim, N. and Mohamed, A.H. (1987): Tannins in sorghum milling fractions in relation to protein digestibility. Egypt Journal of Agronomy, 12:139.
- *Gimeno, A. (1979): Thin layer chromatographic determination of aflatoxins, ochratoxins, sterigmatocystin, Zearalenone Citrinin, T-2 toxin, diacetoxyscirphenol, penicillic acid, patulin, and penitren A. Journal of A.O.A.C., 62:579-585.
- *Glass, R.L.; ponte, J.G.; Christensen, C.M. and Geddes, W. (1959):

 The influence of temperature and moisture level on the behavior of wheat stored in air or nitrogen. Cereal Chem. 36:341-356.
- *Glueck, J.A.; Rooney, L.W.; Rosenow, D.T. and Miller, F.R. (1977):

 Physical and structural properties of field deteriorated (weathered) sorghum grain. Pp. 102-112 in third Annual progress Report, Texas A and M Univ. College Station, Texas, U.S.A (C.F. sorghum Diseases. A world Review, proceedings of the International Workshop. International Grops Research Institute for the semi-Arid Tropics, India, 469pp., 1978).
- *Goldblatt, A.L. (1983): History and natural occurrence of aflatoxins.

 International symposium on mycotoxins Cairo Egypt
 September 6th to 8th p. 33-46.
- *Goodman, J.J. and Christensen, C.M. (1952): Lypolytic activity of fungi isolated from stored corn. Ibid. 29: 299.
- *Guiragossian, V.; Chibber, B.A.K.; Van Scoyoe, S.; Jambunathan, R.; Mertz, E.T. and Axtell, J.D. (1978): Characteristics of protein from normal, high lysine and high tannin sorghums. Journal of Agricultural and Food Chemistry, 26: 219.

- *Gurdev, M.; Ppov, A.; Palaveeva, T.S.; Goranov, K. H. and Goranova, K. (1976): Some studies of oil obtained from soybean Maslo. Sapunena. Prom. St., 12 (1):1. C.F. Chem. Abst. 88, 36055C. (1978).
- *Hadjichristodoulou, A. (1986): Variety and sowing date trials with soybean. Technical Bulletin, Agricultural Research Institute, Cyprus No. 75, 7pp.
- *Hafez, S. A. (1984): Biochemical studies on some soybean varieties.

 M. SC. Thesis. Faculty of Agriculture. Cairo University.
- *Hafez, S.A. (1990): Biochemical studies on soybean. Ph. D. Thesis.

 Faculty of Agriculture Cairo University.
- *Hafez, S.A; Amal. H. Mhamoud and Barakat, M.I.E. (1996): Effect of growth of fungi on the fatty acid composition of stored soybean seeds. Egypt. J. Agric Res., 74 (2), 429-442.
- *Haikerwal, M. and Mathieson, A.R. (1971a): the protein content and amino acid composition of sorghum grain. Cereal Chem. 48:690.
- *Haikerwal, M. and Mathieson, A.R. (1971): the protein content and amino acid composition of sorghum grain. Cereal Chemistry, 48:690-699.
- *Hallalbo, S.A.S. (1977): Chemical studies of naturally occurring unsaponifible matter in some vegetable oil. Ph. D. Thesis Faculty of Agriculture. Cairo University.
- *Hansen, H.N. (1926): A simple method of obtaining single spore culture. Science 64: 384, 1659.

- *Harris, H.B.; Cummins, D.G. and Burns, R.E. (1970): Tannin content and digestibility of sorghum grains as influenced by bagging Journal of Agronomy, 62:633.
- *Harry, E. Sonyder, (1987): Morphology and Chemical composition. Soybean Utilization. Part II. 31.
- *Hartiey, R.D.; Nesbitt, B.F. and O'kelly, J. (1963): Toxic metabolies of Aspergillus favus. Nature (Lond.), 198:1056-1058.
- *Hassan, E.M. (1976): Flour supplementation and its effect on characteristics and nutritive value of bread. MSC Thesis faculty of Agriculture Cairo University.
- *Heathcote, J.G.; Dutton, M.F. and Bbert, J.R.H. (1976): Biosynthesis of aflatoxins. Part II. Chem. Ind., 207-272.
- *Hepperly, P.R. and Sinclair, J.B. (1978): Quality Losses in Phomopsis infected soybean seeds. Phytopathol. 68: 1684.
- *Hesseltine, C.W.; Shotwell, O. L.; Smith, M.L.; Ellis, J.J.; Vandegraft, E. and Shannon, G. (1976): Production of various aflatoxin by strains of the Aspergillus flavus group. Bacteriology Protection Abstract, 68: 2-3.
- *Hesseltine, C.W.; Shotwell, O.L. and Stubblefield, R.D. (1966):

 Aflatoxin formation by Aspergillus flavus. Review of
 Bacteriology, 30: 795-805.
- *Hibbered, C.A.; Wangner, D.G.; Schemm, R.L. Mitchell, E.D.; Hintz, R.L. and Weibel, D. E. (1982): Nutritive characteristics of

- different varieties of sorghum and corn grains. Journal of Animal Science, 55:665.
- *Horowitz-Viassova, Mme L.M. and Livschitz, M.J. (1935): On the question of microbial action on fats. 2bl., Abt. 2: 424-435 (C.F.R.A.M. 15: 42, 1936).
- *Hulse, J.H.; Laing, E.M. and pearson, O.E., (1980a): Sorghum and millets, their compositions and polyphenols in cereal and legumes. International Development Research Center. Ottawa, Canada, 1013 June 1979, Ed. J.H. Hulse, Academic Press London.
- *Hummel, B.C.W.; Cuendet, L.S.; Christensen, C.M. and Geddes, W.F. (1954): Grain storage studies. 13: comparative changes in respiration, viability chemical composition of mold-free and mold-contaminated wheat upon storage. Cereal Chemistry, 31:143-150.
- *Hussein, M.H.A. (1987): production and evaluation of baby food mixtures from some raw and extrded starchy and proteinaceous materials. Ph. D. Thesis-faculty of Agriculture Cairo University.
- *Hyde, H.A. and Williams, D.A. (1949): A census of mould spores in the atmosphere. Nature. Lond. 164:668.
- *Hymowitz, T.; Collins, F.I.; panczner, J. and Walker, W.M. (1972):

 Relation between the content of oil, protein and sugar in soybean seeds. Agronomy Journal, 64: 613-616.
- *Ibrahim, O.S. (1988): Effect of some additives on wheat flour dough and bread characteritisties. M.Sc. Thesis. Faculty of Agriculture Cairo University.

- *Inam, UI.-Haq (1963): Pakistan Journal Science, Research, 15 (4), 123-7.
- *Jadov, S.P. and Rana, B.S., (1983): Tannin content at different stage of grain development in some bird susceptible and resistant sorghum. Current Science, 52:804. (C.F. Chem. Abst. 99: 191820 a).
- *James, H.C. and Phillip, K.H. (1989): Department of Entomology,
 Minnesota Extension Service. University of Minnesota,
 Saint Paul, Minnesota 55108.
- *Jarvis, B. (1971): Factors affecting the production of mycotoxins. J.

 Appl. Bacteriol., 34:199-213.
- *Jood, S. and Kapoor, A.C. (1992): Effect of storage and insect infestation on protein and starch digestibility of cereal grains. Food chemistry 44 (3): 209-212192 (14 ref. En).
- *Jood, S.; Kapor, A.C. and Ram Singh, A. (1993): Available carbohydrates of cereal grains as affected by storage and man. Nutrition 43 (1) 45-54.
- *Kawamura, S. (1967): Quantitative paper chromatography sugars of the cotyledon, hull and hypoetyl of soybeans of selected varieties. Tech. Bull. Fac. of Agric. Kgawa Univ., 15:117-131. (C.F. World Oilseeds: Chemistry, Technology and Utilization. Salunkh; D.K., chavan, J.K.; Adsule, R.N. and kadam, S.S. 1st Ed, publishe by: Van Nastr and Reinhold, New York, U.S.A., 1992).

- *Kennedy, B.W., (1964): Moisture content, mold invasion, and seed viability of stored soybeans. Phytopathology, 54: 771-774.
- *Kent-Jones, D.W. and Amos, A.J. (1967): Modern Cereal chemistry 6th. Ed., the Northen Sublising Co., Liver Pool.
- *Khadzhiiski, T.S.; Stevanov, L.; Savov, I.; Mechenov, G.; Palaveeva, T.S.; Arutyunyan, N. and Kornena, E. (1979): Qualitative characteristics of raw soybean oil. Khrenit-prom., St. 28 (4): 28. C.F. Chem. Abst. 92, 208389 (1980).
- *Klisurov, K.H.; Stoilov, I.; Belcheva, L. and Tankov, D. (1985): Effect of moisture content on storage, processing and biological efficiency in diets for fattening of maize grain preserved in anaerobic conditions. Zhivothov dni-Nauki, 22 (8): 37. (C.F. Maize Abst., 002, 02346, 1986).
- *Krivoruchco, D.; kaba, H.; Sambucetti, M.E. and Sanahuja, J.C. (1979): Maturation time and some seed composition characters affecting nutritive value in soybean varieties. Cereal Chemistry, 56:217.
- *Lacey, J. (1971): The microbiology of moist barley storage in unsealed silos. Ann. Appl. Biol. 69: 187-212.
- *Lacey, J. (1975): Air borne spores in pastures. Trans. Brit. Mycol. Soc. 64 (2): 1-17.
- *Lacey, J. and Lacey, M.E. (1964): Spore concentration in the air of farm buildings. Trans. Brit. Mycol. Soc. 47:547-552.

- *Landers, K.E.; Davis, N.D. and Diener, U.L. (1967): Influence of atmospheric gases on aflatoxin production by Aspergillus flavus_in peanuts. Phytopathology. 57: 1086-1090.
- *Lin, Y.C.; Ayres, J.C. and Kothler, P.E. (1980): Influence of temperature cyclic on production of aflatoxin B₁ and G₁ by Aspergillus parasiticus. Applied and Environmental Microbiology, 40: 333-336.
- *Lortl, H.; Snyder, H. and Gbur, E. (1990): Analysis of single soybean seeds for oil and protein. Journal of American Oil Chemistry Society, 67 (10): 686-688. (C.F. Biological Abstract, 9 (6): 2361).
- *Luddy, F.E.; Barford, R.A.; Herb, S.F. and Magidman, P. (1968): A rapid and quantitative procedure for the preparation of methyl esters of butter oil and other fats. Journal of American Oil Chemistry Society, 45:549-552.
- *Maggon, K.K.; Gopal, S. and Venkitasubramanian, T.A. (1973): Effect of trace metals on aflatoxin production by Aspergillus flavus. Biochemistry Physiology Pflanz., 164: 523-530.
- *Mashaly, R.I., and El-Deeb, S.A. (1983b): Some nutrients as sole energy sources for Aspergillus in aflatoxins production. International symposium on mycotoxins. Cairo Egypt September 6th to 8th p. 485-498.
- *Mashaly, R.I., and El-Deeb, S.A. (1983c): Effect oxygen, Ellumination, and sodium chloride on the aflatoxins produced Aspergillus. International symposium on mycotoxins Cairo Egypt September 6th to 8th p. 477-483.

- *Ministry of Agriculture (1986): Agriculture Economics and Statistics.

 Division, MOA, Cairo, Egypt.
- *Ministry of Agriculture (1992): Agriculture Economics and Statistics.

 Division, MOA, Cairo, Egypt.
- *Mohamed, A.I., and Rangappa, M. (1992): Screeing Soybean (grain and vegetable) genotypes for nutrients and nutritional factors. Plant foods for Human Nutrition 42 (1) 87-96. (C.F. Nutrition Abstract and Review. 63 (7): 4223)
 - *Moubasher, A.H.; Abdel-Fattah, O.M. and Swelim, M.A. (1981a):

 Studies on air-borne fungi at Qena. II-Diurnal fluctuations.

 Zeit. fur Allgem. Mikrob. 21 (3): 177-179.
 - *Moubasher, A.H.; Abdel-Fattah, O.M. and Swelim, M.A. (1981b):

 Studies on air-borne fungi at Qena. I-Seasonal fluctuations.

 Zeit. Fur Allgem. Mikrob. 21 (3): 247-253.
 - *Muindi, P.J. and Thomke, S. (1981): Metabolic studies with laying hens on Tanzanian sorghum grain of different tannin contents. Swed. Journal of Agriculture, 11:17.
 - *Murty, D.S.; Singh, U.; Suryapraksh, S. and Micodemus, K.D. (1985): Soluble sugars in five endosperm types of sorghum. Cereal Chemistry, 62:150.
 - *Nakayama, Y.; Saio, K. and kito, M. (1981): Decomposition of phospholipids in soybean during storage. Cereal Chem. 58 (4): 260.
 - *Nechaev, A. P.; and Sandler, zh, Ya. (1975): Grain lipids (Lipidy Zerna) (book), kolos (Moscow), 159 p.
 - *Nelson, E.P.; Toussoun, A.T., and Marasas, W.F.O. (1983):

 Fusarium species, an illustrated manual for identification.

 The Pennsylvania State University press, USA.
 - *Nelson, N. (1944): A photometric adaptation of the Smogyi method for determination of glucose. Journal of Biological Chemistry, 153:357-380.

- *Neucere, N.J. and Sumrell, G. (1980): Chemical composition of different varieties of grain sorghum. Journal of Agricultural and Food Chemistry, 28:19.
- *Newberne, P.M. and Butler, W.H. (1969): Acute and chronic effect of aflatoxin on the liver of domestic and laboratory animals: A review. Cancer Research. 29:236-250.
- *Newberne, P.M. and Rogers, A.E. (1973): animal model of human disease: primary of hepatocellular carcinoma. Am. J. pathol., 72:137-146.
- *Nget-Hong, T.; Kai-choo, W., and Benito, O.D. (1984): Relationship of tannin levels and trypsin inhibitor activity with the vitro protein digestibility of raw and heat treated Winged bean.

 Journal of Agricultural and Food Chemistry, 32:819-822.
- *Ochiai Yanagi, S.; Galeazi, M.A.M. and Saio, K. (1985): Properties of soybean in model storage studies. Argic. and Biol. Chem. 49 (2): 525-528.
- *Olafson, J.H.; Christensen, C.M. and Geddes, W.F. (1954): Grain storage studies, 15: Influence of moisture content, commercial grade, and maturity on the respiration and chemical deterioration of corn. Cereal Chemistry, 31: 333-340.
- *Oxoid limited (1982): The oxoid manual of culture media, ingredients and other laboratory services. Turnergraphic L td. England 4th Ed.

- *Parish, F.W.; Wiley B.J.; Simmons, E.G. and Long, L. Jr. (1966):

 Production of aflatoxins and kojic acid by species of

 Asperillus and penicillium. Appl. Microbiol., 14:139.
- *Parker, J.C. and Klentworth, G. (1971): Miscellaneous uncommon diseases. In Hand buch der Speziellen pathologischen Anatomie. Vol. 3 Baker, R.D. (ed.). Berlin, Springer-Verlag, pp. 986-987. [C.F.M. Sc. Thesis by N.H. Aziz, Ain Shams Univ., 1982].
- *Paster, N.; Bartov, I.; Menasherov, M.; Padova, R.; Papavizas, C. and Ross, I. (1991): Efficacy of gamma irradiation in preventing mouldiness and preserving the nutritional value of corn grain for btioler chicks. Poultry Science, 70 (4): 823.
- *Paul, Y.; Sharma, B.N. and Bhatia, I.S. (1972): Note on lipids in sorghum. Indian Journal of Agricultural Science, 42:435-436 (C.F. Chem. Abst. 77:85660C).
- *Pitt, J.I. (1979): The genus Penicillium and its teleomorphic states

 Eupenicillium and Talaromyces. Academic Press, London.
- *Pomeranz, Y.; Halton, P. and Peers, F.G. (1956): The effects on flour dough and bread quality of molds grown in wheat and those added to flour in the form of specific cultures. Cereal Chem. 33: 157.
- *Priyadarshini, E. and Tulpule, P.G. (1979): Effect of graded doses of γ- irradiation on aflatoxin production by A.parasiticus in wheat. Fd. Cosmet. Toxicology, 17: 505.
- *Priyadarshini, E. and Tulpule, P.G. (1980): Effect of free fatty acids on aflatoxin production in a synthetic medium. Ibid. 81: 367-369.

- *Richmond, J.W.; Sutcliffe, N.H.; Daniels, N.W.R.; Eggitt, P.W.R. and Coppock, J.B.M. (1962): Factors other than groundnut relating to "turkey 'X' disease". Vet Rec. 74 (18): 544-545.
- *Robertson, J.A; Marrison, W.H. and Burdick D. (1973): Chemcial evaluation of oil from field and storage damaged soybeans.

 J. Amer. Oil chem. Soc. 50:443.
- *Rooney, L.W. (1973): A review of the physical properties, composition and structure of sorghum grain as related to utilization. Indus. Uses Cereals 316-342.
- *Rooney, L.W. (1978): Sorghum and pearl millet lipids. Cereal Chemistry, 55:584.
- *Rooney, L.W.; Blakaly, M.E.; Miller, F.R. and Rosenow, D.T. (1980):

 Polyphenols in cereals and legumes. Hulis, J.H., Ed.,

 International development research center Ottaw Canada p.

 25.
- *Saad, S.A. (1992): Effect of processing and enrichment of some cereal products on their nutritional value. Ph. D. Thesis. Faculty of Agriculture Cairo University.
- *Saio, K.; Kobayakowa, K. and Kito, M. (1982): Protein denaturation during model storage studies of soybean and meals. Cereal Chemistry, 59 (5): 408.
- *Saio, K.; Nikkuni, I.; Ando, Y.; Otsuru, M.; Terauchi, Y. and Kito, M. (1980): Soybean quality changes during model storage studies. Cereal Chem. 57 (2): 77.
- *Saker, Amany, M.A. (1995): Effect of storing African sorghum under different environmental conditions on bread characteristics.

 M.SC. Thesis, of African studies, Natural Resources, Inst. African research and studies, Cairo. University.

- *Sakla, A.B.; Glali, Y.; EL-Farra, A. and Rizk, L.F. (1988b): The effect of environmental conditions on the chemical composition of soybean seeds: Relationship between the protein, oil, carbohydrate and trypsin inhibitor content. Food

 Chemistry, 29: 221-231.
- *Salmon, W.D. and Newberne, P.M. (1963): Occurrence of hepatomas in rats fed diets containing peanut meal as a major source of protein. Cancer Research. 33:571-575.
- *Sargeant, K.; Sheridan, A.; O'kelly, J. and Carnaghan, R.B.A. (1961): Toxicity associated with certain samples of ground nuts. Nature, 192:1096-1097.
- *Sauer, D.B. (1987): Conditions that affect growth of Aspergillus flavus and production of aflatoxin in stored maize. Zuber, M.S.; Lillehoj, E.B. and Renfro, B.L. Aflatoxin in maize, Aproceedings of the workshop. CIMMYT, Mexico, D.F.P. 41-49.
- *Schindler. A.F.; Abadie, A.N. and Simpson, R.E. (1980): Enhanced aflatoxin production by Aspergillus flavus and Aspergillus parasiticus after gamma irradiation of spores inoculum. J. of Food Protec., 43:7-9.
- *Schroeder, H.W. and Boller, R.A. (1973): Aflatoxin production of species and strains of the Aspergillus flavus group isolated from field crops. Appl. Microbiol. 25:885-889.
- *Schroeder, H.W. and H. Hein, Jr. (1967): Aflatoxin production of the toxin in vitro in relation to temperature. Ibid. 15: 441-445.

- *Schroeder, H.W. and Hein, H. J.R. (1968): Effect of diurnal temperature cycles on the production of aflatoxin Ibid. 15: 441-443.
- *Schuller, P.L. and van Egmond, H.P. (1983): Determination of aflatoxin B₁ in feeding stuffs. March., 2-9.
- *Schultz. D.L. and Luedecke, L.O. (1977): Effect of neutral fats and fatty acids on aflatoxin production. Journal of Food protection, 40: 304-306.
- *Seitz, L.M.; Sauer, D.b. and Mohr, H.E. (1982): Storage of high-moisture corn: fungal growth and dry matter loss. Cereal Chemistry, 59 (2):100.
- *Shephered, A.D.; Woodhead, A.H. and kapasi-kakama, J., (1972):

 Cereal processing. In Annual Report, 1972. East African
 Industrial Research Organization, pp 23-57
- *Shih C. N., and Marth E.H. (1975): production of aflatoxin and its partition between the medium and the mycelium of Aspergillus parasiticus during incubation under various conditions. 2 Lebensm. Unters. Forsch., 158: 215:224.
- *Sidky, Hanaa. M. (1986): Formulation and evaluation of baby food mixtures. Ph. D. Thesis. Faculty of Agriculture Cairo University.
- *Sidky, H.M.A. (1995): Nutritional evaluation on soybean extradite as weaning food. Egyptian Journal of Nutrition, 2,125-137.

- *Sikka, K.C.; Gupta, A.K.; Singh, R. and Gupta, P.D. (1978):

 Comparative nutritive value, amino acid content, chemical composition, and digestibility in vitro of vegetable and grain type soybeans. Journal of Agriculture and Food Chemistry, 26: 312-318.
- *Sisodia, B.S.; Apte, B.G. and Solankey, B.C. (1979): Note on the tannin, protein and starch content of some sorghum varieties. Indian Journal of Agriculture Science, 49:135.
- *Sonntage, N.O.V. (1979b):Composition and characteristics of individual fats and oil in "Baley's" industrial oil and fat products Vol. 1.4th ed. (D. Swern. ed.) Willey New-York.
- *Sood, D.R.; Wagle, D.S. and Gupta, V.A. (1977): Varietal difference in the chemical composition of soybean (Glycine max L.).

 Journal of Food Science and Technology, 14:177-179.
- *Sorenson, W.C.; Hesseltine, C.W. and Shotwell, O.L. (1967): Effect of temperature on production of aflatoxin on rice by Aspergillus flavus. Mycopathology Mycology Applied, 33: 49-55.
- *Steel, J.L., (1963): Deterioration of shelled corn during storage as measured by carbon dioxide production. M.Sc. Thesis, lowa State University, U.S.A.
- *Stozky, G. and Goos, R.D. (1965): Effect of high CO₂ and low O₂ tensions on the soil microbiota. Candian. Journal Microbiology, 2:835-868.

- *Stubblefield, R.D.; Shotwell, O.L.; Hesseltine, C.W.; Smith, M.L. and Hall, H.H. (1967): production of aflatoxin on wheat and oats: Measurement with a recording densimeter. Appl. Microbiol., 15:186-190.
- *Sulins, R.D. and Rooney, L.W. (1975): Light and electron microscopic studies of waxy and non-waxy endosperm sorghum varieties. Cereal Chemistry, 52:366.
- *Süss, L. and locatelli, D.P. (1989): Results of practical studies on protection of stored wheat against infestation by insects.

 Tecnica Molitoria 40 (12): 97-986.
- *Suyder, H. E. and kwan, T. W. (1987): Soybean Utilization .1th ed. published by Van Nostrand Reinhold company. Inc. New York.
- *Svoboda, D.; Grady, H.J. and Higginson, J. (1966): Aflatoxin B₁ injury in rat and monkey liver. American Jurnal of Pathology 49:1023-1052.
- *Tadrus (1989): Chemical and biological studies on some baby foods.

 M. Sc. Thesis. Faculty of Agriculture Cairo University.
- *Taha, R. (1974): The biological evaluation of soybean proteins. M. Sc.

 Thesis. Faculty of Agriculture Ain Shams University.
- *Tanteeratarm, K.; Wei, L.S. and Steinberg, M.P. (1989): Effect of soybean maturity of storage stability and process quality.

 Journal of Food Science 54 (3): 593-597.

- *Taria, Ha.; Taira, Hi. and Saito, M. (1971): Influence of location on the chemical composition of soybean seeds. I-protein, oil, carbohydrate, and ash contents. Nippon Sakumatsu Gakkai Kiji 40 (4): 530-544. (C.F. Chem. Abst. 77: 4265C, 1972).
- *Thom, C. and Raper, K.B. (1945): A manual of the Aspergilli.
 Williams and Wilkins Co., Baltimore, Md.
- *Trenk H.L.; Butz, M.E., and Chu, F.S. (1971): Production of aflatoxins in different cereal products by Aspergillus ochraceus. Appl. Microbial., 21:1032-1035.
- *Tsen, C.C.; farrell, E.P.; Hoover, W.J. and Crowley, P.R. (1975):

 Extrucded soy products from whole and dehulled soybeans cooked at various temperatures for bread and cookie fortifications. Cereal foods World 20:9, 413-418.
- *Tsuruta, O. (1987): Quality deterioration of cereal grains caused by fungal infection during storage. Corn and mil imported for feed JARQ-Japan Agricultural Research Quarterly 20 (4): 276. (C.F. Maize Abstract, 004-01767, 1988).
- *Urbanski, E.G.; Wei, S.L. and Nelson, I.A. (1980): Effect of freeze damage on soybean quality and storage stability. J. of Food Sci. 45: 208.
- *Vasi, I. G. and Desai, G.M. (1978): Qualitative and quantitative comparison of amino acids in two varieties of sorghum vulgare. J. Inst. Chem. (India) 50: 141-142 (C.F. Chemical Abstract 91: 18546 W, 1979).
- *Victor, Y. (1978): Protein concentrate from normal and high lysine sorghum, preparation, composition and properties. Journal of Agriculture Food Chemistry, 26: 305.

- *Vogel, A.I. (1975): A text Book of practical organic chemistry. 2nd ed. English language Book Society and Longman Group L td.
- *Waichungo, A. and Holt, P. (1994): High tannin sorghum quality after tannin reduction with aqueous ammonium hydroxide. IFT, Annual Meeting Technical program, Book of Abst. June 25-29, page 109, No. 360-5, Georgia World Congress Center, Atlanta, Georgia U.S.A.
- *Wall, J.S. and Blessin, C.W. (1970): Composition and structure of sorghum grain. Cereal Science Today, 14:264.
- *Wang, H.L.; Swain, E.W. and Kwolek, W.F. (1983): Effect of soybean varieties on the yield quality of tofu. Cereal Chemistry, 60 (3): 245-248.
- *Weber, E.J. (1973): Structure and composition of cereal components as related to their potential industrial utilization. IV. Lipids. In. Indus trial uses of cereals, ed. by y. pomeranz. pp. 161. Proc. Symp. 58. Ann. Meet. Amer. Ass. Cereal Chemistry, St. paul. Minn.
- *Weighrauch, J.L.; kinsella, J.E. and Watt, B.K. (1976):

 Comprehensive evaluation of fatty acids in foods. VI.

 Cereal products. Journal of American Diet. Association,
 68:335.
- *Welsh, R.A. and Buchness, J.A. (1955): Aspergillus endocarditis, inyocarditis and lung abscesses. Am. J. Clin. Pathol. 25: 782-786.
- *Wogan, G.N. and Newbrene, P.M. (1967): Dose response characteristic of aflatoxin B₁ carcinogenesis in the rat. Cancer Res., 27: 2370-2376.

- *Woodhead, S.; padgham, D.E. and Bernays, E.A. (1980): Insect feeding on different sorghum cultivars in relation to cyanide phenolic acid content. Ann. Appl. Binl., 95:151.
- *Youssef, H.H.; Youssef, Y.A.; Abogamra, E.M.; Hammad, H.A. and Refai, M. (1980): Mycological and Serological studies on grain workers in mill factories in Cairo. Egypt. J. Ches. Dis. Tub. 23:33-48.
 - Yousef, S.S. (1981): Soybean seed Quality and stand Establishment, proceedings of Conference for Scientists of Asia J.P. 25-31, Ed. J.B. Sinclair and J. A. Jackobes.
- *Youssef, Y.A. and Abdou, M.H. (1967): Studies on fungus infection of the external ear. I. Mycological and clinical observations. J. Laryng. Otolog. LXXX1, 401-412.
- *Zeleny, L. (1954): Chemical physical and nutritive changes during storage. P. 46-76. In J.A. Anderson and A.W. Alcock (elds.), storage of cereal grains and their products. Am. Assoc. Cereal Chem., St. paul. Minn.
- *Zeleny, L. and Coleman, D.A. (1938): Acidity in cereals and Cereal products, its determination and significance. Cereal Chem. 15: 580-595. J. Amer. Oil chem. Soc. 58:260-268.