

REFERENCES

- Ahmed Y, Van Iddekinge B, Paul C, et al. (1993):** Retrospective analysis of platelet members and volumes in normal pregnancy and in preeclampsia. *Br J. Obstert Gynaecol* 100: 216-219
- Ajne G, Wolff K, Fyhrquist F, et al. (2003):** Endothelin converting enzyme (ECE) activity in normal pregnancy and preeclampsia. *Hypertens Pregnancy* 22: 215-218.
- Al-Ghafra A, Brennecke SP, King RG, et al. (2006a):** Exprssion of adrenomedullin mRNA is altered with gestation and labour in human placenta and fetal membranes. *Clin Sci (lond)*. 110(3): 337-342.
- Al-Ghafra A, Gude NM, Brennecke SP, et al. (2006b):** Increased adrenomedullin protein content and mRNA expression in human fetal membranes but not placental tissues in preeclampsia. *Mol Hum Reprod*. 12(3): 181-186.
- Ando K, Ito Y, Kumada M, et al. (1998):** Oxidative stress increases adrenomedullin mRNA levels in cultured rat vascular smooth muscle cells. *Hypertens Res* 21: 187-191.
- Anko ML and Panula P. (2006):** Regulation of endogenous human NPFF2 receptor by neuropeptide FF in SK-N-MC neuroblastoma cell line. *Journal of Neurochemistry*. **96** 573-584.
- Apodaca CC, Moore KH, Rossignol TM, et al. (2000):** Localization of messenger ribonucleic acid for adrenomedullin and adrenomedullin receptor in the human placenta in normal

pregnancies and pregnancies complicated by oligohydramnios. Am J Obstet Gynecol 183:1213–1219

Aquilin J, Barrnett A, Thompson O, et al. (1999): Second trimester maternal serum inhibin A concentrations as an early marker for preeclampsia. Am J. Obstet Gynecol 181: 131-135.

Ashour AMN, Lieberman ES, Wilkins, et al. (1997): The value of elevated second trimester β -human chorionic gonadotropin in predicting development of preeclampsia. Am J Obstet Gynecol 176: 438-442.

Audibert F, Behchimol Y, Benattar C, et al. (2005): Prediction of preeclampsia or intrauterine growth restriction by second trimester serum screening and uterine Doppler velocimetry. Fetal Diagn there 20: 48-51.

Avrameas S. (1992): Amplification Systems in Immunoenzymatic Techniques. Journal of Immunological Methods, 150: 23-32.

Barron WM, Heckerling P, Hibbard JU, et al. (1999): Reducing unnecessary coagulation testing in hypertensive disorders of pregnancy. Obstet Gynecol 94: 364-368.

Benedetto C, Marozio L, Neri I, et al. (2000): Increased L-citrulline/ L-arginine plasma ratio in sever preeclampsia. Obstet Gynecol 96: 395-398.

Bewley S, Cooper D and Campell S. (1991): Doppler investigation of uteroplacental blood flow resistance in the second trimester: A screening study for preeclampsia and

intrauterine growth retardation. Br J. Obstet Gynaecol 98: 871-875.

Blackburn CA, Keelan JA, Taylor RS et al. (1987): Maternal serum activin A is not elevated before preeclampsia in women who are at high risk. Am J Obstet Gynecol 188: 807-809.

Borghi C, Esposti DD, Immordino V, et al. (2000): Relationship of systemic hemodynamics, left ventricular structure and function, and plasma natriuretic peptide concentrations during pregnancy complicated by preeclampsia. Am J obstet Gynecol 183: 140-143.

Borkakoti N. (1998): Matrix metalloproteases: variations on a theme. Prog Biophys Mol Biol 70:73–94

Branch DW, Porter TF, Rittenhouse L, et al. (2001): Antiphospholipid antibodies in women at risk for preeclampsia am J obstet Gynecol 184: 825-830.

Cameron VA and Fleming AM. (1998): Novel sites of adrenomedullin gene expression in mouse and rat tissues. Endocrinology 139: 2253-2264.

Cappell LC, Seed PT, Briley A, et al. (2002): A longitudinal study of biochemical variables in women at risk of preeclampsia. Am J. Obstet Gynecol 187: 127-133.

Caron C, Goudemaud J, Marey A, et al. (1991): Are haemostatic and fibrinolytic parameters predictors of preeclampsia in pregnancy associated hypertension? Thromb Haemost 66: 410-412.

- Cekmen MB, Balat A, Balat O, et al. (2004):** Decreased adrenomedullin and total nitrite levels in breast milk of preeclamptic women. Clin Biochem.37(2):146-148.
- Chappell L and Bewley S. (1998):** Preeclamptic toxemia: The role of uterine artery Doppler. Br J Obstet Gynaecol 105 : 379-383.
- Charles CJ, Nicholls MG, Rademaker MT, et al. (2001):** Comparative actions of adrenomedullin and nitroprusside: interactions with ANG II and norepinephrine. American Journal of Physiology. 281: R1887-R1894.
- Chavarria ME, Lara-Gonzalez L, Gonzalez- Gleason A, et al. (2003):** Prostacyclin / thromboxane early changes in pregnancies that are complicated by preeclampsia. Am J. Obstet Gynecol 188: 986-992.
- Chini EN, Chini CC, Bolliger C, et al. (1997):** Cytoprotective effects of adrenomedullin in glomerular cell injury: central role of cAMP signaling pathway. Kidney Int 52:917–925.
- Conde-Agudelo A and Belizan JM. (2000):** Risk factors for pre-eclampsia in a large cohort of Latin American and Caribbean women. Br J. Obstet Gynaecol, 107: 75-80.
- Conde-Agudelo A, Villar J and Lindheimer M. (2004):** World health organization systematic review of screening tests for preeclampsia. Obstet Gynecol 104: 1367-1377.
- Conrad KP and Benyo (1997):** Placental cytokines and the pathogenesis of preeclampsia. Am J Reprod Immunol.37: 240-249.

- Cormier-Regard S, Nguyen SV and Claycomb WC. (1998):** Adrenomedullin gene expression is developmentally regulated and induced by hypoxia in rat ventricular cardiac myocytes. *J Biol Chem.* 273: 17787–17792.
- Cuckle H, Sehmi I and Jones R. (1998):** Maternal serum inhibin A can predict preeclampsia. *Br J. Obstet Gynaecol* 105: 1101-1105.
- Cunningham FG, Houth JC, Leveno JK, et al. (2005):** Hypertensive disorders in pregnancy. In: *Williams Obstetrics* 22nd edition. McGraw-Hill Companies, Inc. PP 762-808.
- Dekker GA and Sibai BM. (1998):** Etiology and pathogenesis of preeclampsia: Current concepts. *Am J. Obstet Gynecol* 179: 1359-1363.
- Del Bene R, Lazzeri C, Barletta G, et al. (2000):** Effects of low-dose adrenomedullin on cardiac function and systemic haemodynamics in man. *Clinical Physiology.* 20: 457-465.
- DeWolf F, De Wolf-Peeters C, Brosens I, et al. (1980):** The human placental bed: Electron microscopic study of trophoblastic invasion of spiral arteries. *Am J. Obstet Gynecol* 137: 58-63.
- Di Iorio R, Marinoni E, Scavo D, et al. (1997):** Adrenomedullin in pregnancy. *Lancet.* 349: 328-335.
- Di Iorio R, Marinoni E and Cosmi EV. (1998a):** Adrenomedullin in pre-eclampsia. *Lancet* 351:676–676.
- Di Iorio R, Marinoni E, Letizia C, et al (1998b):** Adrenomedullin, a new vasoactive peptide, is increased in preeclampsia. *Hypertension.* 32: 758-763.

- Di Iorio R, Marinoni E, Letizia C, et al. (1999):** Adrenomedullin production is increased in normal human pregnancy. *Eur J Endocrinol.* 1999; 140: 201–206.
- Di Iorio R, Marinoni E, Letizia C, et al. (2000):** Adrenomedullin is increased in the fetoplacental circulation in intrauterine growth restriction with abnormal umbilical artery waveforms. *Am J Obstet Gynecol* 182:650–654
- Di Iorio R, Marinoni E, Urban G, et al. (2001):** Fetomaternal adrenomedullin levels in diabetic pregnancy. *Horm Metab Res* 33:486–90.
- Di Iorio R, Marinoni E, Letizia C, et al. (2003):** Adrenomedullin in perinatal medicine. *Regul Pept.* Apr15;112(1-3):103-113
- Diedrich F, Renner A, Rath W, et al. (2001):** Lipid hydroperoxides and free radical scavenging enzyme activities in preeclampsia and HELLP (hemolysis, elevated liver enzymes, and low platelet count) syndrome: No evidence for circulating primary products of lipid peroxidation. *Am J. Obstet Gynecol* 185: 166-170.
- Dong YL, Gangula PR, Fang L, et al. (1998):** Uterine relaxation responses to calcitonin gene-related peptide and calcitonin gene-related peptide receptors decreased during labor in rats. *Am J Obstet Gynecol.* 179:497-506
- Ecker J, Karumanchi A, Roberts J, et al. (2003):** First-trimester placental growth factor, SFLT-1, and risk for preeclampsia (abstract). *Am J. Obstet Gynecol* 189: S60-S65.
- Eguchi S, Hirata Y, Iwasaki H, et al. (1994):** Structure-activity relationship of adrenomedullin, a novel vasodilatory

peptide, in cultured rat vascular smooth muscle cells. *Endocrinology* 135: 2454-2458.

Ehrhart BM, Hinson JP, Bornstein SR, et al. (1998): Intraadrenal interactions in the regulation of adrenocortical steroidogenesis [published erratum appears in *Endocr Rev* 1998 Jun;19(3):301]. *Endocr Rev* 19:101–143.

Eto T. (2001): A review of the biological properties and clinical properties and clinical implications of adrenomedullin and proadrenomedullin N-Terminal 20 peptide (PAMP), hypotensive and vasodilating peptides. *Peptides* 22, 1693-1771.

Eto T, Kitamura K and Kato J. (1999): Biological and clinical roles of adrenomedullin in circulation control and cardiovascular diseases. *Clin Experim Pharmacol Physiol* 26: 371-380.

Faas MM, Schuiling GA, Linton EA, et al. (2000): Activation of peripheral leukocytes in rat pregnancy and experimental preeclampsia. *Am J. Obstet Gynecol* 182: 351-360.

Fischer T, Schneider MP, Schobel HP, et al. (2000): Vascular reactivity in patients with preeclampsia and HELLP (Hemolysis, elevated liver enzymes, and low platelet count) syndrome. *Am J. Obstet.* 146: 29-32.

Florio P, Reis FM, Rezzani I, et al. (2003): The addition of activin A and inhibin A measurement to uterine artery Doppler velocimetry to improve the early prediction of pre-eclampsia. *Ultrasound Obstet Gynecol* 21: 165-169

Garayoa M, Martinez A, Lee S, et al. (2000): Hypoxia-inducible factor- (HIF-1) up-regulates adrenomedullin expression in human

tumor cell lines during oxygen deprivation: a possible promotion mechanism of carcinogenesis. *Mol Endocrinol* 12:848–851.

García-Unzueta MT, Berrazueta JR, Pesquera C, et al. (2005): Levels of plasma total adrenomedullin are related with two acute phase inflammatory reactants (fibrinogen and sialic acid) but not with markers of endothelial dysfunction in Type 1 diabetes: Adrenomedullin and vascular risk factors in Type 1 DM . *J Diabetes Complications*.19(3):147-154.

Gervasi MT, Chaiworapongsa T, Pacora P, et al. (2001): Phenotypic and metabolic characteristics of monocytes and granulocytes in preeclampsia. *Am J. Obstet Gynecol* 185: 792-795.

Gratton RJ, Gluszynski M, Mazzuca M, et al. (2003): Adrenomedullin Messenger Ribonucleic Acid Expression in the Placentae of Normal and Preeclamptic Pregnancies .*The Journal of Clinical Endocrinology & Metabolism*. 88 (12): 6048-6055

Grobman WA and Wang EY. (2000): Serum levels of activin A and inhibin A and the subsequent development of preeclampsia. *Obstet Gynecol* 96: 390-395.

Gumusel B, Chang JK, Hao Q, et al. (1996): Adrenotensin: an adrenomedullin gene product contracts pulmonary blood vessels. *Peptides* 17:461–465

Hague S, Zhang L, Oehler MK, et al. (2000): Expression of the hypoxically regulated angiogenic factor adrenomedullin correlates with uterine leiomyoma vascular density. *Clinical Cancer Research*. 3 : 2808-2814.

- Harlow FH, Brown MA, Brighton TA, et al. (2002):** Platelet activation in the hypertensive disorders of pregnancy. *Am J. Obstet Gynecol* 187: 688-691.
- Hata T, Miyazaki K and Matsui K. (1997):** Decreased circulating adrenomedullin in pre-eclampsia. *Lancet* 350:1600-1605.
- Hauth JC and Cunningham FG. (1999):** Pre-eclampsia – eclampsia. in lindheimer M.D.; Roberts J.M. Cunningham F.G (eds): Chesley's Hypertensive Disorders in pregnancy, 2nd ed. Stamford, CT, Appleton & Lange. P169.
- Hayashi M, Hamada Y and Ohkura T. (2004):** Elevation of granulocytemacrophage colony-stimulating factor in the placenta and blood in preeclampsia. *Am J. Obstet Gynecol* 190: 456-459.
- Hayashi Y, Ueyama H, Mashimo T, et al. (2005):** Circulating mature adrenomedullin is related to blood volume in full-term pregnancy. *Anesth Analg.* 101:1816-1820.
- Hayman R, Brockelsby J, Kenny L, et al. (1999):** Preeclampsia ; the endothelium , circulating factor(s) and vascular endothelial growth factor. *J Soc Gynecol Invest.*6: 3-10.
- Hayman R, Warren A, Brockelsby J, et al. (2000):** Plasma from women with pre-eclampsia induced an in vitro alteration in the endotheliumdependent behaviors of myometrial resistant arteies. *Br J. Obstet Gynaecol* 107: 108-111.
- Haynes JM and Cooper ME. (1995) :** Adrenomedullin and calcitonin gene-related peptide in the rat isolated kidney and in the anaesthetised rat: in vitro and in vivo effects. *Eur J Pharmacol* 280: 91-94.

- Hefler LA, Tempfer CB and Gregg AR. (2001):** Polymorphisms within the interleukin-1 β gene cluster and preeclampsia. *Obstet Gynecol* 97: 664-667.
- Hinson JP, Kapas S and Smith DM. (2000):** Adrenomedullin, a Multifunctional Regulatory Peptide. *Endocrine Reviews* 21 (2): 138-167.
- Hippentiel S, Witzenrath M, Schmeck B, et al. (2002):** Adrenomedullin reduces endothelial hyperpermeability. *Circulation Research* 91: 618-625.
- Hjelmqvist H, Keil R, Mathai M, et al. (1997):** Vasodilation and glomerular binding of adrenomedullin in rabbit kidney are not CGRP receptor mediated. *Am J Physiol* 273: R716-R724.
- Hongshi L, Dakour G, Kaufman S, et al. (2003):** Adrenomedullin is decreased in preeclampsia because of failed response to epidermal growth factor and impaired syncytialization. *Hypertension*. 42; 895-899
- Horio T, Nishikimi T, Yoshihara F, et al. (1998):** Production and secretion of adrenomedullin in cultured rat cardiac myocytes and nonmyocytes: stimulation by interleukin-1 β and tumor necrosis factor- α . *Endocrinology* 139: 4576-4580.
- Hoshimoto K, Hayashi M and Ohkura T.(2002):** Mature adrenomedullin concentrations in plasma during pregnancy. *J Matern Fetal Neonatal Med.* Feb;11(2):126-129.

- Ichiki Y, Kitamura K, Kangawa K, et al. (1994):*** Distribution and characterization of immunoreactive adrenomedullin in human tissue and plasma. *FEBS Letters* 338: 6-10.
- Imai T, Hirata Y, Iwashina M, et al. (1995):*** Hormonal regulation of rat adrenomedullin gene in vasculature. *Endocrinology* 136: 1544-1548.
- Ishikawa T, Chen J, Wang J, et al. (2003):*** Adrenomedullin antagonist suppresses *in vivo* growth of human pancreatic cancer cells in SCID mice by suppressing angiogenesis. *Oncogene*. 22: 1238-1242
- Ishimitsu T, Kojima M, Kangawa K, et al. (1994):*** Genomic structure of human adrenomedullin gene. *Biochem Biophys Res Commun* 203:631–639
- Ishimitsu T, Miyata A, Matsuoka H, et al. (1998):*** Transcriptional regulation of human adrenomedullin gene in vascular endothelial cells. *Biochem Biophys Res Commun* 243: 463-470.
- Ishimitsu T, Nishikimi T, Matsuoka H, et al. (1996):*** Behaviour of adrenomedullin during acute and chronic salt loading in normotensive and hypertensive subjects. *Clin Sci Colch* 91:293–298.
- Ishizaka Y, Tanaka M, Kitamura K, et al. (1994):*** Adrenomedullin stimulates cyclic AMP formation in rat vascular smooth muscle cells. *Biochem Biophys Res Commun* 200: 642-646.

- Isumi Y, Shoji H, Sugo S, et al. (1998):** Regulation of adrenomedullin production in rat endothelial cells. *Endocrinology* 139:838–846
- Iwase T, Nagaya N, Fujii T, et al. (2005):** Adrenomedullin enhances angiogenic potency of bone marrow transplantation in a rat model of hindlimb ischemia. *Circulation*. 111: 356-362
- Jacobson SL, Imhof R, Manning N, et al. (1990):** The value of Doppler assessment of the uteroplacental circulation in predicting preeclampsia or intrauterine growth retardation. *Am J. Obstet Gynecol* 162 : 110-115.
- Jerat S, Morrish DW, Davidge ST, et al. (2001):** Effect of adrenomedullin on placental arteries in normal and preeclamptic pregnancies. *Hypertension* 37:227–231
- John JH, Ziebland S, Yudkin P, et al. (2002):** Effects of fruit and vegetable consumption on plasma antioxidant concentrations and blood pressure: A randomized controlled trial. *Lancet* 359:1969-1973.
- Jougasaki M, Rodeheffer RJ, Redfield MM, et al. (1996):** Cardiac secretion of adrenomedullin in human heart failure. *J Clin Invest* 97: 2370-2376.
- Julian M, Cacho M, Mario AG, et al.(2005):** Adrenomedullin: a new target for the design of small molecule modulators with promising pharmacological activities. *Eur J Med Chem* 2005 Aug;40(8):737-750.
- Kanenishi K, Kuwabara H, Ueno M, et al. (2000):** Immunohistochemical adrenomedullin expression is

decreased in the placenta from pregnancies with pre-eclampsia. *Pathol Int* 50:536–540.

Kanenishi K, Kuwabara H, Ueno M, et al. (2001): Change of adrenomedullin concentrations in plasma and amniotic fluid, and human placental adrenomedullin expression with advancing gestation. *Placenta*. 22(2-3):244-250.

Kapas S, Brown DW, Farthing PM, et al. (1997): Adrenomedullin has mitogenic effects on human oral keratinocytes: involvement of cyclic AMP. *FEBS Lett* 418:287–290.

Katano K, Aoki A, Sasa H, et al. (1996): Beta-2- glycoprotein I-dependent anticardiolipin antibodies as a predictor of adverse pregnancy outcomes in healthy pregnant women. *Hum Reprod* 11:509-512.

Kato H, Shichiri M, Marumo F, et al. (1997): Adrenomedullin as an autocrine/paracrine apoptosis survival factor for rat endothelial cells. *Endocrinology* 138:2615–2620

Kato J, Kitamura K and Eto T. (2006): Plasma adrenomedullin level and development of hypertension. *J Hum Hypertens*.20(8):566-570.

Kato J, Kitamura K, Kangawa K, et al. (1996): Receptors for adrenomedullin in human vascular endothelial cells. *Eur J. Pharmacol*. 289: 383- 385.

Kato J, Tsuruda T, Kita T, et al. (2005): Adrenomedullin :A Protective Factor for Blood Vessels Arteriosclerosis, Thrombosis, and Vascular Biology. 25:2480-2485.

- Katsuki A, Sumida Y, Gabazza EC, et al. (2002):** Acute hyperinsulinemia is associated with increased circulating levels of adrenomedullin in patients with type 2 diabetes mellitus. *Eur J Endocrinol*.147(1):71-75.
- Katsuki A, Sumida Y, Urakawa H, et al. (2003):** Increased oxidative stress is associated with elevated plasma levels of adrenomedullin in hypertensive patients with type 2 diabetes. *Diabetes Care*.26(5):1642-1643.
- Keelan JA, Taylor R, Schellenberg JC, et al. (2002):** Serum activin A, inhibin A, and follistatin concentrations in preeclampsia or small for gestational age pregnancies. *Obstet Gynecol* 99: 267-272.
- Kita T, Kitamura K, Kuwasako K, et al. (1998)** Short-term modulation of the renin-angiotensin system does not alter plasma adrenomedullin concentration in humans. *J Hypertension* 16: 2057-2062.
- Kitamura K, Kangawa K, Kawamoto M, et al. (1993a):** Adrenomedullin: a novel hypotensive peptide isolated from human pheochromocytoma. *Biochem Biophys Res Commun* 192: 553-560.
- Kitamura K, Sakata J, Kangawa K, et al. (1993b):** Cloning and characterization of cDNA encoding a precursor for human adrenomedullin. *Biochem Biophys Res Commun*.194: 720-725.
- Kitamura K, Ichiki Y, Tanaka M, et al. (1994a):** Immunoreactive adrenomedullin in human plasma. *FEBS Letters* 341: 288-290.

- Kitamura K, Kangawa K, Ishiyama Y, et al. (1994b):** Identification and hypotensive activity of proadrenomedullin N-terminal 20 peptide (PAMP). *FEBS Letters* 351: 35-37.
- Knapp RG and Miller MC (1992):** clinical epidemiology and dbiostatistics – National series from Willi Egyptian Edition P 209-213.
- Kobayashi K, Kitamura K, Hirayama N, et al. (1996a):** Increased plasma adrenomedullin in acute myocardial infarction. *Am Heart J* 131: 676-680.
- Kobayashi K, Kitamura K, Etoh T, et al. (1996b):** Increased plasma adrenomedullin levels in chronic congestive heart failure. *Am Heart J* 131: 994-998.
- Kobayashi K, Kubota T, Aso T, Hirata Y, et al. (2000):** Immunoreactive adrenomedullin (AM) concentration in maternal plasma during human pregnancy and AM expression in placenta. *European Journal of Endocrinology*, 142 (6): 683-687
- Kohno M, Hanehira T, Kano H, et al. (1996b):** Plasma adrenomedullin concentrations in essential hypertension. *Hypertension* 27: 102-107.
- Kubo A, Iwano M, Minamino N, et al. (1998a):** Measurement of plasma and urinary adrenomedullin in patients with IgA nephropathy. *Nephron* 78:389–394.
- Kubo A, Minamino N, Isumi Y, et al. (1998b):** Adrenomedullin production is correlated with differentiation in human leukemia cell lines and peripheral blood monocytes. *FEBS Lett* 426:233–237

- Kuwasako K, Kitamura K, Onitsuka H, et al. (2002):** Rat RAMP domains involved in adrenomedullin binding specificity. FEBS letters. 519: 113-116
- Lain KY and Roberts JM. (2002):** Contemporary concepts of the pathogenesis and management of preeclampsia. JAMA 287: 383-392.
- Lainchbury JG, Cooper GJS, Coy DH, et al. (1997):** Adrenomedullin: a hypotensive hormone in man. Clinical Science. 92: 467-472.
- Lainchbury JG, Troughton RW, Lewis LK, et al. (2000):** Hemodynamic, hormonal, and renal effects of short-term adrenomedullin infusion in healthy volunteers. Journal of Clinical Endocrinology and Metabolism. 85: 1016-1020.
- Lang MG, Paterno R, Faraci FM, et al. (1997):** Mechanisms of adrenomedullin-induced dilatation of cerebral arterioles. Stroke 28: 181-185.
- Lauria MR, Standley CA, Sorokin Y, et al. (1999):** Adrenomedullin levels in normal and preeclamptic pregnancy at term. J Soc Gynecol Invest 6:318–321
- Lawlor DA, Morton SM, Nitsch D, et al. (2005):** Association between childhood and adulthood socioeconomic position and pregnancy induced hypertension: Results from the Aberdeen children of the 1950s cohort study. J Epidemiol Community Health 59: 49-52.
- Leduc L, Wheeler JM, Kirshon B, et al. (1992):** Coagulation profile in severe preclampsia. Obstet Gynecol 79 : 14-19.

- Levine RJ, Ewell MG, Hauth JC, et al. (2000):** Should the definition of preeclampsia include a rise in diastolic blood pressure of ≥ 15 mm Hg to a level < 90 mm Hg in association with proteinuria ? Am J Obstet Gynecol 183 : 787-791.
- Levine R, Maynard S, Qian C, et al. (2003):** Acute rise in circulating SFLT-1 may herald preeclampsia. Am J Obstet Gynecol 189: S57-S60.
- Levine RJ, Qian C, LeShane ES, et al. (2004):** Two- stage elevation of cell free fetal DNA in maternal sera before onset of preeclampsia Am J. Obstet Gynecol 190: 707-711.
- Lewis LK, Smith MW, Brennan SO, et al. (1997):** Degradation of human adrenomedullin(1–52) by plasma membrane enzymes and identification of metabolites. Peptides 18:733–739.
- Li H, Dakour J, Kaufman S, et al. (2003):**Adrenomedullin Is Decreased in Preeclampsia Because of Failed Response to Epidermal Growth Factor and Impaired Syncytialization. Hypertension. 42:895-899.
- Livingston JC, Park V, Barton JR, et al. (2001):** Lack of association of severe preeclampsia with maternal and fetal mutant alleles for tumor necrosis factor α and lymphotoxin α genes and plasma tumor necrosis α levels. Am J obstet Gynecol 184: 1273-1275.
- Lopez J and Martnez A. (2002):**Cell and molecular biology of the multifunctional peptide, adrenomedullin. Int. Rev. Cytol. 221:1-92.

Lu B, Wang A and Fei Y. (1999): Study of adrenomedullin value in pregnancy induced hypertension patients. *Zhoonghua Fu Chan Ke Za Zhi*.34 (1):17-19.

Luodonpaa M, Leskinen H, Ilves M, et al. (2004): Adrenomedullin modulates hemodynamic and cardiac effects of angiotensin II in conscious rats. *Am J Physiol Regul Integr Comp physiol*. 286(6): R1085-1092.

Luttun A and Carmeliet P. (2003): Soluble VEGF receptor Fltl: The elusive preeclampsia factor discovered? *J Clin Invest* 111: 600-604.

Macara LM, Kingdom JC, Kaufmann P, et al. (1993): Control of fetoplacental circulation. *Fetal Maternal Med Rev*. 5: 167-179.

Macri C, Loup D, McHale M, et al. (1998): Adrenomedullin is widely expressed throughout normal and abnormal reproductive tissues of women - evidence for cyclic regulation. In: Martinez A, Cuttitta F (eds) *Adrenomedullin*. IOS Press, Amsterdam, pp 207-248

Madazli R, Budak E, Calay Z, et al. (2000): Correlation between placental bed piopsy findings, vacular cell adhesion molecule and fibronectin levels in preeclampsia. *Br J Obstet Gynaecol* 107: 514-516.

Makino Y, Shibata K, Makino I, et al. (1999): Expression of adrenomedullin in feto-placental circulation of human normotensive pregnant women and pregnancy-induced hypertensive women. *Endocrinology* 140:5439–5442.

- Makino Y, Shibata K, Makino I, et al. (2001):** Alteration of the adrenomedullin receptor components gene expression associated with the blood pressure in pregnancy-induced hypertension. *J Clin Endocrinol Metab.* 86(10):5079-5082.
- Mallamaci F, Zoccali C, Parlongo S, et al. (1998):** Plasma adrenomedullin during acute changes in intravascular volume in hemodialysis patients. *Kidney Int* 54:1697–1703
- Manten GT, Van der Hoek YY, Marko Sikkema J, et al. (2005):** The role of lipoprotein (a) in pregnancies complicated by pre-eclampsia. *Med Hypotheses* 64: 162-166.
- Marinoni E, Di Iorio R, Letizia C, et al. (1998):** Immunoreactive adrenomedullin in human fetoplacental tissues. *Am J Obstet Gynecol* 179:784-787.
- Marinoni E, Zacharopoulou C, Di Rocco A, et al.(2006):** Effect of Betamethasone In Vivo on Placental Adrenomedullin in Human Pregnancy. *J Soc Gynecol Investig.* 13 (6): 418 – 424.
- Martinez A, Miller MJ, Unsworth EJ, et al. (1995):** Expression of adrenomedullin in normal human lung and in pulmonary tumors. *Endocrinology* 136:4099-4105.
- Martinez A, Weaver C, Lopez J, et al. (1996):** Regulation of insulin secretion and blood glucose metabolism by adrenomedullin. *Endocrinology* 137:2626–2632
- Martinez A, Elsasser TH, Muro CC, et al. (1997):** Expression of adrenomedullin and its receptor in normal and malignant

human skin: a potential pluripotent role in the integument. *Endocrinology* 138:5597–5604.

Martinez A, Cuttitta F and Teitelman G. (1998): Expression pattern for adrenomedullin during pancreatic development in the rat reveals a common precursor with other endocrine cell types. *Cell Tissue Res* 293:95–100.

Martinez A, Vos M, Guedez L, et al. (2002): The effects of adrenomedullin overexpression in breast tumour cells. *Journal of the National Cancer Institute*. 94: 1226-1237.

McLatchie LM, Fraser NJ, Main MJ, et al. (1998): RAMPs regulate the transport and ligand binding specificity of the calcitonin-receptor-like receptor. *Nature*. 393: 333-339.

Meeran K, O'Shea D, Upton PD, et al. (1997): Circulating adrenomedullin does not regulate systemic blood pressure but increases plasma prolactin after intravenous infusion in humans: a pharmacokinetic study. *J Clin Endocrinol Metabol* 82: 95-100.

Minamino N, Shoji H, Sugo S, et al. (1995): Adrenocortical steroids, thyroid hormones and retinoic acid augment the production of adrenomedullin in vascular smooth muscle cells. *Biochem Biophys Res Commun* 211: 686-693.

Minamino N, Kikumoto K, Isumi Y, et al. (2002): Regulation of adrenomedullin expression and release. *Microsc Res Technol* 57:28-39.

Minegishi T, Nakamura M, Abe K, et al. (1999): Adrenomedullin and atrial natriuretic peptide concentrations in normal

pregnancy and pre-eclampsia. *Mol Hum Reprod.* 5:767-770.

Moody TW, Miller MJ, Martinez A, et al. (1997): Adrenomedullin binds with high affinity, elevates cyclic AMP, and stimulates c-fos mRNA in C6 glioma cells. *Peptides* 18:1111–1115.

Mori Y, Takayasu M, Suzuki Y, et al. (1997): Effects of adrenomedullin on rat cerebral arterioles. *Eur J Pharmacol* 330: 195-198.

Moriyama T, Otani T and Maruo T. (2001): Expression of adrenomedullin by human placental cytotrophoblasts and choriocarcinoma JAr cells. *J Clin Endocrinol Metab* 86:3958–3961.

Morris NH, Eaton BM and Dekker G. (1996): Nitric oxide, the endothelium, pregnancy and pre-eclampsia. *Br J. Obstet Gyunaecol* 103: 4-8.

Morrish DW, Linetsky E, Bhardwaj D, et al., (1996): Identification by subtractive hybridization of a spectrum of novel and unexpected genes associated with in vitro differentiation of human cytotrophoblast cells. *Placenta.* 17: 431–441.

Murakami Y, Hattori Y, Taniyama M, et al. (1998): Adrenomedullin regulation by thyroid hormone in the rat. *Life Sci* 63: L337-L342.

Muttukrishna S, Knight PG, Groome, NP, et al. (1997): Activin- A and inhibin-A as possible endocrine markers for preeclampsia. *Lancet* 349: 1285-1289.

- Nagata N, Kato J, Kitamura K, et al. (1998):** Dissociation of adrenomedullin concentrations in plasma and cerebrospinal fluid in pregnant and non-pregnant women. *Eur J Endocrinol* 139:611–614.
- Nagaya N, Nishikimi T, Uematsu M, et al. (1999):** Plasma adrenomedullin as an indicator of prognosis after acute myocardial infarction. *Heart* 81: 483-487.
- Nagaya N, Satoh T, Nishikimi T, et al. (2000):** Hemodynamic, renal and hormonal effects of adreomedullin infusion in patients with congestive heart failure. *Circulation* . 101: 498-503.
- Nagaya N, Goto Y, Satoh T, et al. (2002):** Intravenous adrenomedullin in myocardial function and energy metabolism in patients after myocardial infarction. *Journal of Cardiovascular Pharmacology*. 39: 754-760
- Naicker T, Khedun SM, Moodley J, et al. (2003):** Quantitative analysis of trophoblast invasion in preeclampsia. *Acta. Obste. Gynecal. Scand.* 82 : 722-729.
- Nandha KA, Taylor GM, Smith DM, et al. (1996):** Specific adrenomedullin binding sites and hypotension in the rat systemic vascular bed. *Reg Pept* 62: 145-151.
- National High Blood pressure Education Program (2000):** Working Group Report on High Blood pressure in pregnancy . *Am J. Obstet Gynecol* 183: 51-56.
- Ness RB, Markovic N, Bass D, et al. (2003):** Family history of hypertension, heart disease, and stroke among women who

develop hypertension in pregnancy. *Obstet Gynecol* 102: 1366-1369.

Newman MG, Robichaux AG, Stedman CM, et al. (2003): Perinatal outcomes in preeclampsia that is complicated by massive proteinuria *Am J ObstetGynecol* 188:246-249.

Nikitenko LL, MacKenzie IZ, Rees MC, et al. (2000): Adrenomedullin is an autocrine regulator of endothelial growth in human endometrium. *Mol Hum Reprod.* Sep;6(9):811-819.

Nilsson E, Ros HS, Cnattingius S, et al. (2004): The importance of genetic and environmental effect for pre-eclampsia and gestational hypertension: A family study. *Br J. Obstet Gynaecol* 111: 200-205.

Nishikimi T, Kitamura K, Saito Y, et al. (1994): Clinical studies on the sites of production and clearance of circulating adrenomedullin in human subjects. *Hypertension* 24 : 600-604.

Nishikimi T, Nagata S, Sasaki T, et al. (1997): Plasma concentrations of adrenomedullin correlate with the extent of pulmonary hypertension in patients with mitral stenosis. *Heart* 78: 390-395.

Nishio K, Akai Y, Murao Y, et al. (1997): Increased plasma concentrations of adrenomedullin correlate with relaxation of vascular tone in patients with septic shock. *Crit Care Med* 25: 953-957.

North RA, Taylor RS and Schellenberg JC. (1999): Evaluation of a definition of preeclampsia. *Br J Obstet gynaecol* 106: 767-769.

- Paara MC, Less C, Mann GE, et al. (2001):** Vasoactive mediator release by endothelial cells in intrauterine growth restriction and preeclampsia. *Am J. Obstet Gynecol* 184: 497-499.
- Page NM, Woods RJ, Gardiner SM, et al. (2000):** Excessive placental secretion of neurokinin B during the third-trimester cause preeclampsia. *Nature*. 405: 797-800.
- Parameswaran N, Spielman WS, Pullen M, et al. (1999):** Mechanism of adrenomedullin-stimulated hyaluronic acid release in rat mesangial cells. *Eur J Pharmacol* 370:313–318
- Parkes DG and May CN. (1995):** ACTH-suppressive and vasodilator actions of adrenomedullin in conscious sheep. *J Neuroendocrinol* 7:923–929
- Petrie MC, Hillier C, Morton JJ et al. (2000):** Adrenomedullin selectively inhibits angiotensin II-induced aldosterone secretion in humans. *Journal of Hypertension*. 18: 61-64.
- Pio R, Martinez A, Unsworth EJ, et al. (2001):** Complement factor H is a serum-binding protein for adrenomedullin, and the resulting complex modulates the bioactivities of both partners. *J Biol Chem*. 276: 12292–12300.
- Porstmann T and Kiessig ST. (1992):** Enzyme Immunoassay Techniques, An Overview. *Journal of Immunological Methods*. 150: 5-21.
- Powers RW, Evans RW, Ness RB, et al. (2000):** Homocysteine is increased in preeclampsia but not in gestational hypertension. *J. Soc Gynecol Investig* 7 (1) : 231-235.

- Poyner D, Sexton P, Marshall I, et al.(2002):** International Union of Pharmacology. XXXII. The mammalian calcitonin gene-related peptides, adrenomedullin, amylin, and calcitonin receptors. *Pharmacological Reviews*. 54: 233-246.
- Rademaker MT, Cameron VA, Charles CJ, et al. (2003):** Adrenomedullin and heart failure. *Regulatory Peptides*. 112: 51-60.
- Redman CWG. (1993):** The placenta, preeclampsia and chorionic villitis. In:Redman CWG, Sargent IL , Starkey PM , eds . *The human placenta*. London , England : Blackwell Scientific Publication ; 433-467.
- Redman CWG and Sargent IL. (2003):** Pre-eclampsia, the placenta and the maternal inflammatory response – a review. *Placenta* 17: 21-29.
- Roberts JM. (2000):** Preeclampsia: What we know and what we do not know. *Sermin Perinatol* 24: 24-28.
- Roberts JM and Lain KY. (2002):** Recent insights into the pathogenesis of pre-eclampsia. *Placenta*. 2002; 23: 359–372
- Robinson C and Johnson D. (2003):**A marker for severity of preeclampsia *Am J. Obstet Gynecol* 189: 85-87.
- Ross R. (1999):** Atherosclerosis- an inflammatory disease. *N Engl J. Med* 340: 115-119.
- Ruzicska E, Toth M, Tulassayz et al. (2001):** Adrenomedullin and diabetes mellitus. *Diabetes. Metab Res Rev*. Sep-Oct; 17 (5): 321-329.

- Samson WK. (1999):** Adrenomedullin and the control of fluid and electrolyte homeostasis. *Annu Rev Physiol* 61:363-89: 363-389.
- Samson WK and Murphy TC. (1997):** Adrenomedullin inhibits salt appetite. *Endocrinology* 138: 613-616.
- Samson WK, Murphy T and Schell DA. (1995):** A novel vasoactive peptide, adrenomedullin, inhibits pituitary adrenocorticotropin release. *Endocrinology* 136:2349–2352.
- Sato K, Hirata Y, Imai T, et al. (1995):** Characterization of immunoreactive adrenomedullin in human plasma and urine. *Life Sci* 57:189–194.
- Satoh F, Takahashi K, Murakami O, et al. (1997):** Immunoreactive adrenomedullin in human adrenal glands and adrenal tumors. *Cancer Detect Prevent* 21: 51-54.
- Savvidou MD, Less CC, Parra M, et al. (2002):** Levels of C-reactive protein in pregnant women who subsequently develop pre-eclampsia. *Br J. Obstet Gynaecol* 109: 297-300.
- Schwarz N, Sidhu K, Kapas S, et al. (2004):** Characterization of adrenomedullin receptor (crlr/ramp2) expression in human microvascular endothelial cells *Endocrine Reviews* 8: 38-41.
- Schwarz N, Renshaw D, Kapas S, et al. (2006):** Adrenomedullin increases the expression of calcitonin-like receptor and receptor activity modifying protein 2 mRNA in human microvascular endothelial cells. *J Endocrinol.*190(2):505-514.

- Shindo T, Kurihara H, Maemura K, et al. (2000):** Hypotension and resistance to lipopolysaccharide-induced shock in transgenic mice overexpressing adrenomedullin in their vasculature. *Circulation* 101:2309-2316.
- Shinoki N, Kawasaki T, Minamino N, et al. (1998):** Shear stress down-regulates gene transcription and production of adrenomedullin in human aortic endothelial cells. *J Cell Biochem* 71:109–115.
- Sibai BM. (2003):** Diagnosis and management of gestational hypertension and preeclampsia. *Obstet Gynecol* 102: 181.
- Smith D, Coppock H, Withers D, et al. (2002):** Adrenomedullin: Receptor and Signal Transduction. *Biochemical Society Transactions*. 30: 432-437.
- Staff AC, Ranheim T, Khoury J. et al. (1999):** Increased contents of phospholipids, cholesterol, and lipid peroxides in decidual basalis in women with preeclampsia. *Am J ObstetGynecol* 180: 587-590.
- Stamilio, DM, Sehder, HM, Morgan MA, et al. (2000):** Can antenatal clinical and biochemical markers predict the development of severe preeclampsia? *Am J Obstet Gynecol* 182: 589-592.
- Sugo S, Minamino N, Kangawa K, et al. (1994):** Endothelial cells actively synthesize and secrete adrenomedullin. *Biochem Biophys Res Commun*; 201: 1160 – 1165.
- Sugo S, Minamino N, Shoji H, et al. (1995):** Interleukin-1, tumor necrosis factor and lipopolysaccharide additively stimulate

production of adrenomedullin in vascular smooth muscle cells. *Biochem Biophys Res Commun* 207: 25-32.

Suzuki Y, Yamamoto T, Mabuchi Y, et al. (2003): Ultrastructural changes in omental resistance artery in women with preeclampsia. *Am J Obstet Gynecol* 189: 216-220.

Takacs P, Green KL, Nikaeo A, et al. (2003): Increased vascular endothelial cell production of interleukin-6 in severe preeclampsia. *Am J. Obstet Gynecol* 188: 740-744.

Taniyama M, Kitamura K, Ban Y, et al. (1996): Elevated plasma adrenomedullin level in hyperthyroidism. *European J Clin Invest* 26: 454-456.

Taylor RN and Roberts JM. (1999): Endothelial cell dysfunction. In Lindheimer MD, Roberts JM, Cunningham FG (eds) : Chesley's Hypertensive Disorders in pregnancy 2nd ed. Stamford, CT, Appleton & Lange. P 395.

Thadhani R, Mutter WP, Wolf M. et al. (2004): First trimester placental growth factor and soluble fms-like tyrosine kinase 1 and risk for preeclampsia. *J Clin Endocrinol Metab* 89: 770-773.

Thota C, Gangula RR, Dong Y, Yallampalli C, et al. (2003): Changes in the Expression of Calcitonin Receptor-Like Receptor, Receptor Activity-Modifying Protein (RAMP) 1, RAMP2, and RAMP3 in Rat Uterus During Pregnancy, Labor, and by Steroid Hormone Treatment. *Biol Reprod.* **69**, 1432–1437.

- Treloar SA, Cooper DW, Brennecke SP, et al. (2001):** An Australian twin study of the genetic basis of preeclampsia and eclampsia. *Am J Obstet Gynecol* 184 : 374-377.
- Troughton RW, Frampton CM, Lewis LK, et al. (2001):** Differing thresholds for modulatory effects of adrenomedullin infusion on haemodynamic and hormone responses to angiotensin II and adrenocorticotrophic hormone in healthy volunteers. *Clinical Science*. 101: 103-109.
- Troughton RW, Lewis LK, Yandle TG, et al. (2000):** Hemodynamic, hormone, and urinary effects of adrenomedullin infusion in essential hypertension. *Hypertension*. 36: 588-593.
- Upton PD, Austin C, Taylor GM, et al. (1997):** Expression of adrenomedullin (ADM) and its binding sites in the rat uterus: increased number of binding sites and ADM messenger ribonucleic acid in 20-day pregnant rats compared with nonpregnant rats. *Endocrinology* 138:2508-2514.
- Vesely DL, Blankenship M, Douglass MA, et al. (1996):** Atrial natriuretic peptide increases adrenomedullin in the circulation of healthy humans. *Life Sci*. 59:243–254.
- Von Mandach U, Lauth D and Huch R. (2003):** Maternal and fetal nitric oxide production in normal and abnormal pregnancy. *J Matern Fetal Neonatal Med* 13:22-26.
- Walker JJ. (2000):** Pre-eclampsia. *lancet* 356: 1260-1263.
- Walsh SC. (1994):** Lipid peroxidation in pregnancy. *Hypertens Preg* 13: 1-5.

- Wang X, Yue TL, Barone FC, et al.(1995):** Discovery of adrenomedullin in rat ischemic cortex and evidence for its role in exacerbating focal brain ischemic damage. *Proc Natl Acad Sci* 92: 11480-11484.
- Wang Y, Gu Y, Granger DN, et al. (2002):** Endothelial junctional protein redistribution and increased monolayer permeability in human umbilical vein endothelial cells isolated during preeclampsia. *Am J Obstet Gynecol* 186: 214-217.
- Washimine H, Asada Y, Kitamura K, et al. (1995a):** Immunohistochemical identification of adrenomedullin in human, rat, and porcine tissue. *Histochem Cell Biol* 103: 251-254.
- Washimine H, Yamamoto Y, Kitamura K, et al. (1995b):** Plasma concentration of human adrenomedullin in patients on hemodialysis. *Clin Nephrol* 44: 389-393.
- Weerasekera DS and Peiris H. (2003):** The significance of serum uric acid, creatinine and urinary microprotein levels in predicting pre-eclampsia. *J Obstet Gynaecol* 23: 17-19.
- Wilson C, Nikitenko LL, Sargent IL, et al. (2004):** Adrenomedullin: multiple functions in human pregnancy. *Angiogenesis*. 2004;7(3):203-212.
- Wolf M, Kettle E, Sandler L, et al. (2001):** Obesity and preeclampsia: The Potential role of inflammation. *Obstet Gynecol* 98: 757-762.
- Yallampalli C, Gangula PR, Kondapaka S, et al. (1999):** Regulation of calcitonin gene-related peptide receptors in the rat uterus

during pregnancy, labor and by progesterone. Biol Reprod 61:1023-1030.

Yang S, Zhou M and Chaudry IH. (2002a): Novel approach to prevent the transition from the hyperdynamic phase to the hypodynamic phase of sepsis: Role of adrenomedullin and adrendomedullin binding protein-1. Ann Surg 236 : 625-633.

Yang S, Zhou M, Gowler D E, et al. (2002b): Mechanisms of the beneficial effect of adrenomedullin and adrenomedullin-binding protein-1 in sepsis: down-regulation of proinflammatory cytokines. Crit Care Med 30:2729-2735.

Yeung VT, Ho SK, Nicholls MG, et al. (1996): Adrenomedullin, a novel vasoactive hormone, binds to mouse astrocytes and stimulates cyclic AMP production. J Neurosci Res 46 : 330-335.

Yoneyama Y, Suzuki S, Sawa R, et al.(2002):Relation between adenosine and T-helper1 / T-helper 2 imbalance in women with preeclampsia. Obstet Gynecol 99: 641-645.

Yoshitomi Y, Nishikimi T, Kojima S, et al. (1998):Plasma levels of adrenomedullin in patients with acute myocardial infarction. Cli Sci 94: 135-139.

Zeeman GG, Alexander JM, McIntire DD, et al. (2002): Inhibin-A levels and severity of hypertensive disorders due to pregnancy. Obstet Gynecol 100: 140-145.

Zeeman GG, Hatab M and Twickler DM. (2004): Increased large – vessel cerebral blood flow in severe preeclampsia by magnetic resonance (MR) evaluation. Presented at the

24th Annual Meeting of the society for maternal – fetal Medicine, New Orleans, LA, February 2.

Zhang J, Klebanoff MA and Roberts IM. (2001): Prediction of adverse outcomes by common definitions of hypertension in pregnancy *Obstet Gynecol* 97 : 261-264.

Zhang C, Williams MA, King IB, et al. (2002): Vitamin C and the risk of preeclampsia- results from dietary questionnaire and plasma assay. *Epidemiology* 13: 382-386.

Zhang XQ, Varner M, Dizon-Townson D, et al. (2003): A molecular variant of angiotensinogen is associated with idiopathic intrauterine growth restriction. *Obstet Gynecol* 101: 237-239.

Zhang X, Green KE, Yallampalli C, Dong YL. et al. (2005): Adrenomedullin enhances invasion by trophoblast cell lines. *Biol Reprod.* 73(4):619-626.

Zhao Y, Hague S, Manek S, Zhang L, et al. (1998); PCR display identifies tamoxifen induction of the novel angiogenic factor adrenomedullin by a non estrogenic mechanism in the human endometrium. *Oncogene* 16:409-415

Zhong XY, Laivuori H, Livingston JC, et al. (2001): Elevation of both maternal and fetal extracellular circulating deoxyribonucleic acid concentrations in the plasma of pregnant women with preeclampsia. *Am J. Obstet Gynecol* 184: 414-419.