

REFERENCES

REFERENCES

Adams, R. and Victor, M. (1985):

Principles of neurology (third edition). McGraw-Hill, New York, P. 372.

Adelman, D.C.; Saltiel, E. and Klinenberg, J.R. (1986):

The neuropsychiatric manifestations of systemic lupus erythematosus: An overview.

Semin. Arthritis Rehum., 15: 185-90.

Aida, S. (1993):

Alkaline phosphatase isoenzyme activities in rheumatoid arthritis: hepatobiliary enzyme dissociation and relation to disease activity.

Ann. Rheum. Dis., 55: 520-524.

Aitchenson, C.T.; Pebles, C. and Joslin, H. (1992):

Characteristics of antinuclear antibodies in rheumatoid arthritis.

Arch. Rheumatol., 26: 562.

Appels, A. and Mulder, P. (1988):

Excess fatigue as a precursor of myocardial infarction.

Eur. Heart J., 9: 758-764.

Arnett, F.C. (1992):

Rheumatoid arthritis in: Cecil textbook of Medicine, 19th ed., PP. 1508-1515 W.B. Saunders Company, Philadelphia, London, Toronto, Sydney, Tokyo.

Arnett, F.C.; Edworthy, S. and Block, D.A. (1988):

The 1987 Revised ACR criteria for classification of RA.

Arth. Rheum., 31: 315.

Badley, E.M.; Rasooly, I. and Webster, G.K. (1994):

Relative importance of musculoskeletal disorders a cause of chronic health problems, disability and health care utilization: Findings from 1990 Ontario Health Survey. *J. Rheumatol.*, 21: 505-514.

Baker, G.H.B. (1987):

Psychological factors and immunity.
Journal of Psychosomatic Research, 31: 1-10.

Barendregt, P.J.; Visser, M.R.; Smets, E.M. et al. (1998):

Fatigue in primary sjogren's syndrome.
Ann. Rheum. Dis., 57: 291-295.

Barlow, J.H.; Turner, A.P. and Wright, C.C. (1998):

Long-term outcomes of an arthritis self-management program.
Br. J. Rheumatol., 37: 1315-1319.

Beck, A.T.; Bush, A.J.; Shaw, B.F. and Emery, G. (1979):

Cognitive therapy for depression.
New York: Guilford.

Behan, P. and Behan, W. (1989):

The post viral fatigue syndrome.
Critical Reviews in Neurobiology, PP. 157-178.

Belza, B.L. (1995):

Comparison of self-reported fatigue in rheumatoid arthritis and controls.
J. Rheumatol., 22: 639-43.

Belza, B.L.; Henke, C.J.; Yelin, E.H. et al. (1993):

Correlates of fatigue in older adults with rheumatoid arthritis.
Nurs. Res., 42: 93-99.

Benette, J.C. (1993):

The infectious etiology of rheumatoid arthritis: new considerations.

Arth. Rheum., 21: 53.

Bienestock, H.; Ehrlich, G.E. and Freyberg, R.H. (1963):

Rheumatoid arthritis of the cricoarytenoid joint: a clinicopathological study.

Arthritis Rheum., 6: 48-63.

Bradley, L.A. (1997):

Psychosocial factors. In Klippel J.H., Weyand C.M., Wortmann R.L. (eds.): Primer on the Rheumatic Diseases. Atlanta, Georgia, PP. 413-415.

Bradley, L.A.; Young, L.D.; Anderson, K.O. et al. (1987):

Effects of psychological therapy on pain behaviour of rheumatoid arthritis patients: Treatment outcomes and six-month follow-up.

Arthritis Rheum., 30: 1105.

Brook, R.H.; Ware, J.E.; Davies-Avery, A. et al. (1979):

Overview of adult health status measures fielded in Rand's health insurance study.

Med. Care; 17: 1-55.

Buckley, M.M.; Cohen, B.J.; Clewley, J.P. et al. (1986):

Human parvovirus infection in early rheumatoid and inflammatory arthritis.

Ann. Rheum. Dis., 45: 832.

Butler, S.; Chalder, T. and Wessely, S. (1991):

Cognitive behaviour therapy in chronic fatigue syndrome.

J. of Neurology, Neurosurgery and Psychiatry; 54(2): 153-158.

Byrne, E. and Trounce, I. (1987):

Chronic fatigue and myalgia syndrome: Mitochondrial and glycolytic studies in skeletal muscle.

J. Neurology, Neurosurgery and Psychiatry; 50: 743-746.

Calabrese, L.H.; Davis, M.E. and Wilke, W.S. (1994):

Chronic fatigue syndrome and a disorder resembling Sjogren's syndrome: Preliminary report.

Clin. Infect. Dis., 18 (Suppl. 1): 28-31.

Calin, A.; Edmunds, L. and Kennedy, L.G. (1993):

Fatigue in Ankylosing Spondylitis-why is it ignored?

J. Rheumatol., 20: 991-5.

Carson, D.A. (1982):

Antiglobulin antibodies. In: Panayi, G.S. (ed.) the scientific basis of rheumatology.

Churchill Livingstone, Edinburgh, P. 114.

Cash, J.M. and Klippel, J.H. (1994):

Second-line drug therapy for rheumatoid arthritis.

N. Engl. J. Med., 330: 1368-1375.

Checkley, S.A. and Crammer, J.L. (1977):

Responses to methylamphetamine in depression: a new approach to the noradrenaline depletion hypothesis.

British Journal of Psychiatry; 131: 585-586.

Chen, M. (1986):

The epidemiology of self-perceived fatigue among adults.

Prev. Med., 15: 74-81.

Condemi J.; Branett, E.V. and Atwater, E.C. (1995):

The significance of antinuclear factors in rheumatoid arthritis.

Arth. Rheum., 8: 1080.

- Conn, D.L.; McDuffie, F.C. and Dyck, P.J. (1972):**
Immunopathologic study of sural nerves in
rheumatoid arthritis.
Arthritis Rheum., 15: 135.
- Cox, B.; Blaxter, M. and Buckle, A. (1987):**
The health and lifestyle survey.
Health Promotion research trust, London.
- Crosby, L.J. (1991):**
Factors which contribute to fatigue associated with
rheumatoid arthritis.
Arthritis Care Res., 3: 154-157.
- Dacie, J.V. and Lewis, S.M. (1975):**
Practical haematology, 3rd ed., New York, Grune
Stratton Inc.
- David, A.; Wessely, S. and Pelosi, A. (1988):**
Post-viral fatigue syndrome: Time for a new
approach.
British Med. J., 296: 696-699.
- Decarvalho, A. and Graudal, H. (1980):**
Relationship between radiological and clinical
findings in rheumatoid arthritis.
Acta. Radiological; 21: 797-807.
- Deighton, C.M. and Walker, D.J. (1991):**
The familial nature of rheumatoid arthritis.
Ann. Rheum. Dis., 50: PP 62-65.
- Dunne, J.U.; Carson, D.A. and Spiegelberge, H.L. (1995):**
IgA rheumatoid factor in serum of patients with
rheumatoid arthritis.
Ann. Rheum. Dis., 38: 161.

Edmonds, J.P. (1997):

Rheumatoid arthritis. In: Howe H.S., Feng P.H. (eds.): Textbook of Clinical Rheumatology 1st ed. Singapore, PP. 54-111.

Edward, D. and Harris, J.R. (1993):

Etiology and pathogenesis of rheumatoid arthritis. In: Kelly W.N., Harris E.D., Ruddy S., Sledge C.B. (eds.): Textbook of Rheumatology 4th ed. Philadelphia, London, Toronto, Saunders, PP. 833-873.

Edward, D. and Harris, J.R. (1997):

Clinical features of rheumatoid arthritis. In: Kelly W.N., Harris E.D., Ruddy S., Sledge C.B. (eds.): Textbook of Rheumatology 5th ed. Philadelphia, London, Toronto, Saunders.

El-Shabraway, M.; Abd El-Rahman, H. and Sholah, A. (1985):

Pattern of malnutrition diseases in Kalyoubia villages and its association with socio-economic aspects. Thesis for M.D. Degree in Public Health.

Fairburn, C.G. and Cooper, P.J. (1984):

The clinical features of bulimia nervosa. British Journal of Psychiatry; 144: 238-246.

Farr, M.; Scott, D.L.; Constable, T.J. et al. (1983):

Thrombocytosis of active rheumatoid disease. Ann. Rheum. Dis., 422: 545.

Farrar, D.J.; Locke, S.E. and Kantrowitz, F.G. (1995):

Chronic fatigue syndrome. 1: Etiology and pathogenesis. Behav. Med., 21 (1): 5-16.

Fawzy, F.L.; Cousins, N.; Fawzy, N.W. et al. (1990):

A structured psychiatric intervention for cancer patients. Arch. Gen. Psychiatry; 47: 720-725.

Felson, D.T. (1993):

Epidemiology of rheumatic diseases. In Arthritis and Allied conditions: McCarty D., Koopman W., 12th ed. (1): London, Lea, Philadelphia and Febiger PP. 17-49.

Fifield, J.; Tennen, H.; Reisine, S. and McQuillan, J. (1998):

Depression and the long-term risk of pain, fatigue and disability in patients with rheumatoid arthritis. Arthritis and Rheum., 41 (10): 1851-1857.

Firestein, G.S. (1997):

Etiology and pathogenesis of rheumatoid arthritis. In Kelly W.N., Harris E.D., Ruddy S., Sledge C.B. (eds.): textbook of Rheumatology 5th ed. Philadelphia, London, Toronto, Saunders, PP. 851-897.

Fleming, A.; Crown, J.M. and Corbett, M. (1976):

Early rheumatoid disease I. Onset. Ann. Rheum. Dis., 35: 357.

Fong, S.; Carson, D. and Vanghan, J. (1985):

Rheumatoid factor. In: Gupta S., Talal N. (eds.): Immunology of rheumatic disease, 2nd ed. New York, London, medical book company, PP. 167-197.

Fox, R.I. (1995):

Sjogren's syndrome. In "Treatment of the Rheumatic Diseases". Ed. By Weisman M.H., Weinblatt M.E. PP. 217-240 Philadelphia, Saunders.

Fransen, M. and Edmonds, J. (1997):

Orthopaedic footwear for subjects with rheumatoid arthritis. Arthritis Care and Research, (in press).

Freundlich, B. and Leventhal, L. (1997):

Diffuse pain syndromes. In Klippel J.H., Weyand C.M., Wortmann R.L. (eds.): Primer on the Rheumatic Diseases. Atlanta, Georgia. PP. 123-127.

Fries, J.; Sptiz, P.; Kraines, R. and Holman, H. (1980):

Measurement of patient outcome in arthritis. Arthritis Rheum., 23: 137-145.

Fye, K.H.; Moutsopoulos, H.H. and Talal, N. (1992):

Rheumatic disease. In: Fundenberg H.H., Sities D.P.; Wells J. (eds.): Basic and clinical immunology, 4th ed. Los Angeles, California, PP. 422.

Gardner, D.L. (1986):

Pathology of rheumatoid arthritis. In Scott J.T. (ed.): Textbook of the Rheumatic Diseases. 6th ed. Churchill Livingstone, London, New York, PP. 604-652.

Geirsson, A.J.; Sturfett, G. and Truedsson, L. (1987):

Clinical and serological features of severe vasculitis in RA: Prognostic implications. Ann. Rheum. Dis., 46: 727.

Goldenberg, D.L. (1991):

Fibromyalgia, chronic fatigue syndrome and myofascial pain syndrome. Curr. Opin. Rheumatol., 3(2): 247-258.

Goldman, J. (1988):

Chronic fatigue syndrome. Ann. of Internal Medicine, 109: 166-167.

Gordon, D.A.; Stein, A.L. and Broder, L. (1993):

Articular features of rheumatoid arthritis. Am. J. Med.

- Gravallese, E.M.; Corson, J.M.; Coblyn, J.S. et al. (1989):**
Rheumatoid aortitis: a rarely recognized but clinically significant entity.
Medicine, 68: 95-106.
- Hara, K. (1990):**
Rheumatoid pericarditis: Clinical features and survival.
Medicine, 69: 81-91.
- Hart, F.D. (1990):**
Rheumatoid arthritis: Articular manifestations.
Br. Med. J., 3: 131.
- Harwicke, J. and Squire, J.R. (1992):**
The basis of the erythrocyte sedimentation rate.
Clin. Sci., 2: 333.
- Haylock, P.J. and Hart, L.K. (1979):**
Fatigue in patients receiving localized radiation.
Cancer Nursing; 2: 461-467.
- Hennemann, P.H.; Wang, D.M.K.; Irwin, J.W. and Burrage, W.S. (1955):**
Syndrome following abrupt cessation of prolonged therapy.
Journal of the American Medical Association; 158: 384.
- Hochberg, M.C. (1990):**
Changes in the incidence and prevalence of rheumatoid arthritis in England and Wales, 1970-1982.
Semin. Arthritis Rheum., 19(5): 294-302.
- Holt, P.J. (1993):**
Rheumatoid arthritis. In: Harris N.H. (ed.):
Postgraduate textbook of clinical orthopaedics 3rd ed.
Bristol. John Wright and Son. PP. 443-455.

Hultman, E. and Nilsson, L. (1973):

Liver glycogen as glucose supplying source during exercise. In: Inlkeul J. (ed.): Limiting factors in physical performance.

Stuttgart, George Thieme Verlag. PP. 179-189.

Huyser, B.A.; Parker, J.C.; Phoreson, R. et al. (1998):

Predictors of subjective fatigue among individuals with rheumatoid arthritis.

Arthritis and Rheumatism; 41: 2230-2237.

Jacobsson, L.; Hanson, R.; Knowler, W. et al. (1994):

Decreasing incidence and prevalence of rheumatoid arthritis in pima Indians over a twenty-five year period.

Arthritis and Rheumatism; 37(8): 1158-1165.

Jones, J. and Miller, B. (1987):

The post-viral asthenic syndrome. In: Krustak E, Lipowski Z, Morozov, P. (eds.): Viruses, Immunity and Mental disorders. Plenum, London, PP. 441-451.

Jones, S.D.; Koh, W.H.; Steiner, A. et al. (1996):

Fatigue in ankylosing spondylitis: Its prevalence and relationship to disease activity, sleep and other factors.

J. Rheumatol., 23 (3): 487-490.

Karlsson, J. (1972):

Muscle lactate ATP and CP levels during exercise and after physical training in man.

J. App. Physiol., 33: 199.

Kerstens, P.J.; Boerbooms, A.M.; Jeurissen, M.E. et al. (1992):

Accelerated nodulosis during long term methotrexate therapy for rheumatoid arthritis: an analysis of 10 cases.

J. Rheumatol., 19: 867-871.

Kirkendall, D.T. (1990):

Mobility: Conditioning programs. In Gould J.A. (ed.): Orthopaedic and Sports, 2nd ed. PP. 227-252. Baltimore, Philadelphia, Toronto.

Klippel, J.H.; Weyand, C.M.; Wortmann, R.L. (1997):

Rheumatoid arthritis. In: Primer on the Rheumatic Diseases, 11th ed. P. 155-175.
Published by Arthritis Foundation, Atlanta, Georgia.

Kobashi-Schoot, J.A.M.; Hanewald, G.J.F.P.; Van Dam, F.S.A.M. and Bruning, P.F. (1985):

Assessment of malaise in cancer patients treated with radiotherapy.
Cancer Nursing; 8: 306-314.

Kogi, K.; Saito, Y. and Mitsuhashi, T. (1970):

Validity of three components of subjective fatigue feelings.
J. Sci. Labour., 46: 251-270.

Kremer, J.M.; Alarcon, G.S.; Lightfoot, R.W. et al. (1994):

Methotrexate for rheumatoid arthritis: Suggested guidelines for monitoring liver toxicity.
Arthritis Rheum., 37: 316-328.

Krupp, L.B.; Alvarez, L.A.; Larocca, N.G. and Scheinberg, L.C. (1988):

Fatigue in multiple sclerosis.
Arch. Neurol., 45: 435-437.

Krupp, L.B.; LaRocca, N.C.; Muir-Nash, J. and Steinberg, A.D. (1989):

The fatigue severity scale applied to patients with multiple sclerosis and systemic lupus erythematosus.
Arch. Neurol., 46: 1121-1123.

Lawrence, R.C.; Hochberg, M.C.; Kelsey, J.L. et al. (1989):

Estimates of the prevalence of selected arthritis and musculoskeletal diseases in the United States.
J. Rheumatol., 16: 427-441.

Liang, M.H. (1989):

Compliance and quality of life: Confessions of a difficult patient.
Arthritis Care Res., 2: 571.

Liang, M.H. and Daltroy, L.H. (1985):

The impact of inflammatory arthritis on society and the individual. In: Halder N.M. and Gillings D.B. (eds.): Options for Public Health Programs in Arthritis and Society.
London, Butterworths, PP. 1-16.

Linos, A.; John, W.; Michael, W. and Kurland, L. (1980):

The epidemiology of rheumatoid arthritis in Rochester, Minnesota.
Am. J. of Epi., 3: 87.

Linton, S.J. and Bradley, L.A. (1996):

Strategies for the prevention of chronic pain. In: Gatchel R.J., Turk D.C. (eds). Psychological approaches to pain management: A practitioner's handbook.
New York, Guilford Press, pp 438-457.

Lorig, K.R.; Mazanson, P.D. and Holman, H.R. (1993):

Evidence suggesting that health education in self-management in patients with chronic arthritis has sustained health benefits while reducing health care costs.
Arthritis Rheum., 36: 439-446.

Lynch, S.; Seth, R. and Shah, A. (1989):

Depression and the post-viral fatigue syndrome. Abstracts of the spring quarterly meetings. Royal College of Psychiatrists, Leeds, April 4. (Quoted from Wessely and Thomas, 1990).

Mackay, J.M.; Sim, A.K. and Gardner, D.L. (1993):

Etiology of rheumatoid arthritis.
Ann. Rheum. Dis., 42: 309-311.

- Magni, G.; Marchetti, M.; Moreschi, C. et al. (1993):**
Chronic musculoskeletal pain and depressive symptoms in the National Health and Nutrition Examination.
Epidemiologic follow-up study pain; 53: 163-168.
- Mahowald, M.L. and Dykstra, D. (1997):**
Rehabilitation of patients with rheumatic disease. In Klippel J.H., Weyand C.M., Wortmann R.L. (eds.): Primer on the Rheumatic Diseases. Atlanta, Georgia, PP. 407-412.
- Mahowald, M.W.; Mahowald, M.L.; Burdlie, S.R. and Yetterberg, S.R. (1989):**
Sleep fragmentation in rheumatoid arthritis.
Arthritis Rheum., 32: 974.
- Maisiak, R.; Austin, J.S.; West, S.G. et al. (1996):**
The effect of person-centered counseling on the psychological status on persons with systemic lupus erythematosus or rheumatoid arthritis.
Arthritis Care Res., 9: 60-66.
- Mazzuca, S.A. (1982):**
Does patient education in chronic disease have therapeutic value?
J. Chronic Dis., 35: 521.
- Meenan, R.F.; Gertman, P.M. and Mason, J.H. (1980):**
Measuring health status in arthritis: The Arthritis Impact Measurement Scales.
Arthritis Rheum., 23: 146-152.
- Mengshoel, A.M. and Forre, O. (1993):**
Pain and fatigue in patients with rheumatic disorders.
Clin. Rheumatol., 12 (4): 515-521.

McCarty, D.J. (1993):

Clinical picture of rheumatoid arthritis. In: McCarty D.J., Koopman W.J., eds: Arthritis and Allied Conditions.

Philadelphia: Lea & Febiger, 781: 809.

McDonald, E.M.; Mann, A.H. and Thomas, H.C. (1987):

Interferons as mediators of psychiatric morbidity.

The Lancet; ii: 1175-1178.

Moldofsky, H.; Lue, F.A.; Eisen, J. et al. (1986):

The relationship of interleukin-1 and immune functions to sleep in humans.

Psychosom. Med., 48: 309-318.

Montgomery, G.K. (1983):

Uncommon tiredness among college undergraduates.

J. Con. Clin. Psychol., 51: 517-525.

Mowat, A.G. (1971):

Haematologic abnormalities in RA.

Sem. Arthritis. Rheum., 1: 195.

Mullen, P.D.; Laville, E.A.; Biddle, A.K. and Lorig, K. (1987):

Efficacy of psycho-educational interventions on pain, depression and disability with arthritic adults.

J. Rheumatol., 15: 33.

Nakano, K.K.; Schoene, J.C.; Baker, R.A. and Dawson, D.M. (1978):

The cervical myelopathy associated with rheumatoid arthritis.

Am. Neuro. Assoc., 3: 144.

Nepom, G.T. and Nepom, B.S. (1992):

Prediction of susceptibility to rheumatoid arthritis by human leukocyte antigen genotyping.

Rheum. Dis. Clin. North. Am., 18: 785-792.

- Neuberger, G.B.; Press, A.N.; Lindsley, H.B. et al. (1997):**
Effects of exercise on fatigue, aerobic fitness and disease activity measures in persons with rheumatoid arthritis.
Res. Nurs. Health., 20(3): 195-204.
- Neumark, T.; Domby, M. and Gaspard, G. (1979):**
Ultrastructural studies in rheumatoid polyneuropathy.
Acta. Morphological; 27: 205.
- O'Dell, J.R.; Haire, C.E.; Palmer, W. et al. (1997):**
Treatment of early rheumatoid arthritis with minocycline or placebo: Results of a randomized double-blind, placebo-controlled trial.
Arthritis Rheum., 40: 842-848.
- Orr, W.C. (1983):**
Sleep related breathing disorders.
Chest; 84: 475-480.
- Ostergaard, M. (1996):**
Changes in synovial membrane and joint effusion volumes after intra-articular methylprednisolone. Quantitative assessment of inflammatory and destructive changes in arthritis by MRI.
J. Rheumatol., 23: 1151-1161.
- Otterness, I. (1994):**
The value of C-reactive protein measurement in rheumatoid arthritis.
Sem. Arthritis Rheum., 24: 91-104.
- Overall, J.E. (1980):**
Correlational structure of a new self rating scale for psychiatric patients.
Journal of Clinical Psychology; 36: 616-624.

Pacy, P.J.; Read, M.; Preedy, V. et al. (1988):

Post-absorptive whole body leucine kinetics and quadriceps muscle protein synthetic rate (MPSR) in the post-viral syndrome.

Clinical Science and Molecular Medicine, 15 (Suppl. 19): 36-37.

Paget, S. (1997):

Treatment of rheumatoid arthritis. In: Klippel J.H., Weyand, C.M., Wortmann, R.L. (eds.): Primer of Rheumatic Diseases, 11th ed. Atlanta, Georgia, PP. 168-174.

Pawlikowska, T.; Chalder, T.; Hirsch, S.R. et al. (1994):

Population based study of fatigue and psychological distress.

B.M.J., 308: 763-777.

Pearson, M.G.; Oadiri, M.R. and Finn, R. (1986):

Hypokalemia with chronic hyperventilation syndrome.

British Journal of Clinical Partice; 40: 28-29.

Pearson, P.G. and Byars, G.E. (1956):

The development and validation of a checklist measuring subjective fatigue (report No. 56-115).

School of aviation. USAF, Randolph AFB, Texas.

Pinals, R.S.; Masi, A.T.; Larsen, R.A. and the Subcommittee for Criteria of Remission in Rheumatoid Arthritis of the American Rheumatism Association Diagnostic and Therapeutic Criteria Committee (1981):

Preliminary criteria for clinical remission in rheumatoid arthritis.

Arthritis Rheum., 24: 1308-1315.

Pincus, T. (1993):

Limitations of randomized clinical trials to recognize possible advantages of combinations therapies in rheumatoid arthritis.

Semin. Arthritis Rheum., 23: 11-18.

Pincus, T. and Callahan, I.F. (1992):

Early mortality in RA predicted by poor clinical status.
Bull. Rheum. Dis., 41:4.

Pincus, T.; Callahan, I.F.; Bradley, L.A. et al. (1986):

Elevated MMPI scores for hypochondriasis, depression and hysteria in patients with RA reflect disease rather than psychological status.
Arthritis Rheum., 29: 1456.

Piper, B.F. (1989):

Fatigue: Current basis for practice. In: Funk S., Tornquist E., Wiese R. (eds.).
Springer, New York, PP. 187-198.

Piper, B.F.; Lindsey, A.M.; Dodd, M.J. et al. (1989):

The development of an instrument to measure the subjective dimension of fatigue. In Funk S.G., Tornquist E.M., Wiese R.A. (eds.): Key Aspects of comfort. Management of pain, fatigue and nausea. New York, Springer Publishing Company.

Pither, C. (1989):

Treatment of persistent pain.
Br. Med. J., 299: 1239.

Poteliakhoff, A. (1981):

Adrenocortical activity and some clinical findings in acute and chronic fatigue.
Journal of Psychosomatic Research, 25: 91-95.

Powell, R.; Dolan, R. and Wessely, S. (1990):

Attribution and self-esteem in depression and chronic fatigue syndrome.
J. Psychosomatic Research; 34: 665-673.

Ramasay, A. (1986):

Post-viral fatigue syndrome: The saga of the Royal Free disease.
Gower Medical, London.

Rapaport, R.J.; Kozin, F.; Mackel, S.E. and Jordon, R.E. (1980):

Cutaneous vascular immunofluorescence in RA.
Am. J. Med., 68: 325.

Raskin, R.J.; Schnapf, D.J.; Wolf, C.R.; Killian, P.J. and Lawless, O.J. (1983):

Computerized tomography in evaluation of atlanto-axial subluxation in rheumatoid arthritis.
J. Rheumatol., 10: 32.

Riemsma, R.P.; Rasker, J.J.; Taal, E. et al. (1998):

Fatigue in rheumatoid arthritis: The role of self-efficacy and problematic social support.
Br. Journal of Rheumatol., 37: 1042-1046.

Ritchie, D.M.; Boyle, J.A.; Mcinnes, J.M. et al. (1968):

Clinical studies with an articular index for the assessment of joint tenderness in patients with rheumatoid arthritis.
Journal of Medicine; 37: 393-406.

Robert, J.A. (1985):

Loss of form in young athletes due to viral infection.
British Medical Journal; 290: 357-358.

Rogers, M.P.; Liang, M.H. and Partridge, A.J. (1982):

Psychological care of adults with RA.
Ann. Intern. Med., 96: 344.

Rosenberg, N.L.; Neville, H.E. and Ringel, S.P. (1985):

Tubular aggregates.
Archives of Neurology; 42: 973-976.

Rosner, B. (1990):

Fundamentals of biostatistics, 3rd ed. Boston: PWS, Kent, USA.

Rothfield, N.F. (1993):

Systemic lupus erythematosus: Clinical aspects and treatment. In McCarty D.J., Koopman W.J. (eds.): Arthritis and Allied Conditions: A textbook of Rheumatology, Philadelphia, Lea, Febiger, PP. 1155-1178.

Saag, K.G.; Koehnke, R.; Caldwell, J.R. et al. (1994):

Low-dose, long-term corticosteroid therapy in rheumatoid arthritis and analysis of serious adverse events.

Am. J. Med., 96: 115-123.

Samuels, B.; Lee, J.C.; Engleman, E.P. and Hooper, J.Jr. (1977):

Membranous nephropathy in patients with rheumatoid arthritis: Relationship to gold therapy. Medicine; 57: 319.

Salisbury, M.D. and Nottage, W.M. (1985):

A new evaluation of gross pathologic changes and concepts of rheumatoid articular cartilage degeneration.

Clin. Orthop. Rel. Res., 199: 243.

SAS Institute (1995):

Version 6.11, Cary, North Carolina, USA.

Savage, G. (1875):

Overwork as a cause of insanity.

Lancet; ii: 127 (Quoted from Wessely and Thomas 1990).

Schumacher, H.R. (1988):

Palindromic onset of rheumatoid arthritis.

Arthritis Rheum., 25: P 361.

- Schumacher, H.R.; Klippel, J.H. and Koopman, W.J. (1993):**
Rheumatoid arthritis. In: Primer on Rheumatic disease.
Atlanta, Arthritis Foundation.
- Scott, D.L.; Grindulis, K.A.; Coulton, B.L. et al. (1984):**
Progression of radiological changes in rheumatoid arthritis.
Ann. Rheum. Dis., 43: 8-17.
- Sharpe, M.; Hawton, K.; Seagroatt, V. and Pasvol, G. (1992):**
Follow up of patients presenting with fatigue to an infectious diseases clinic.
British Medical J., 305: 147-152.
- Simpson, R.W.; McGinty, L.; Simon, L. et al. (1984):**
Association of parvoviruses with rheumatoid arthritis of humans.
Science; 223.
- Singsen, B.H. and Rose, C.D. (1995):**
Juvenile rheumatoid arthritis and pediatric spondyloarthropathies. In Weisman M.H., Weinblatt M.E. (eds.): Treatment of the Rheumatic Diseases, Philadelphia, Saunders, PP. 217-240.
- Sjogaard, G. (1986):**
Water and electrolytes shifts during exercise and their relation to muscular fatigue.
Acta. Physiol. Scand., 128 (Suppl. 556): 129-136.
- Smets, E.M.A.; Garssen, B.; Bonke, B. and DeHaes, J.C.J.M. (1995):**
The Multidimensional Fatigue Inventory (MFI) psychometric qualities of an instrument to assess fatigue.
Journal of Psychosomatic Research; 39: 315-325.

Stokes, M.; Cooper, R. and Edwards, R. (1988):

Normal strength and fatigability in patients with effort syndrome.
British Medical J., 297: 1014-1018.

Sukoff, M.H.; Kadin, M.M. and Moran, T. (1994):

Rheumatoid arthritis of the cervical spine.
J. Neurosurgery; 37: 493.

Swezey, R. (1993):

Rehabilitation medicine and arthritis. In McCarty D.J., Koopman W.J. (eds.): Arthritis and allied Conditions, 12th ed. Philadelphia, Lea, Febiger, PP. 887-917.

Tack, B. (1990):

Fatigue in rheumatoid arthritis: Conditions, strategies and consequences.
Arthritis Care Res., 3(2): 65-70.

Tan, E.M.; Cohen, A.S.; Fries, J.F. et al. (1982):

The 1982 revised criteria for the classification of systemic lupus erythematosus.
Arthritis Rheum., 25: 1271-1277.

Thomas, P.K. (1987):

Post viral fatigue syndrome (letter).
The Lancet; i: 218-219.

Tugwell, P.; Pincus, T.; Yocum, D. et al. (1995):

Combination therapy with cyclosporine and methotrexate in severe rheumatoid arthritis: The methotrexate-cyclosporine combination study group.
N. Engl. J. Med., 333: 137-141.

Van der Heijde (1995):

Radiographic progression on radiographs of hands and feet during the 1st 3 years of rheumatoid arthritis measured according to Sharp's method (van der Heijde modification).
J. Rheumatol., 22: 1792.

Van Eijk, L.M.; Kempen, G.I.J.M. and Van Sonderen, F.L.P. (1994):

Een korte schaal voor het meten van sociale steun bij ouderen: de SSL12-I. (A short scale to measure social support among elderly the SSL12-I).

Tijdschr Gerontol. Geriatr., 25: 192-196.

Vertommen, H. and Leyssen, J. (1988):

Vermoeidheid: van onhanteerbaar symptoom tot diagnostisch waardevolle gevoelstoestand.

Tijdschr. Klin. Psych., 18: 35-59.

Watson, R.G. (1965):

A more specific method for detecting and quantitating rheumatoid factors using a simple modification of the rheumatoid arthritis test.

Am. J. Clin. Path., 43: 152.

Watt, I. and Rozman, B. (1996):

The role of MRI in the assessment of the rheumatological disease.

Brit. J. Rheum., 35: Suppl. 3: 1-47.

Weinberger, M.; Tiemey, W.M.; Cowper, P.A. et al. (1993):

Cost-effectiveness of increased telephone contact for patient with osteoarthritis: A randomized, controlled trial.

Arthritis Rheum., 36: 243-246.

Weinblatt, M.E.; Kaplan, H.; Germain, B.F. et al. (1994):

Methotrexate in rheumatoid arthritis: A five-year prospective multicenter study.

Arthritis Rheum., 37: 1492-1498.

Weiss, B. and Laties, V.G. (1962):

Enhancement of human performance by caffeine and amphetamines.

Pharmacology Reviews; 14: 1-36.

Wessely, S. (1990):

The natural history of chronic fatigue and myalgia syndromes. In: Sartorius N. (ed.) Psychological disorders in general medical settings. World Health Organization.

Wessely, S. and Powell, R. (1989):

Fatigue syndromes: A comparison of chronic "post-viral" fatigue with neuromuscular and affective disorders.

J. of Neurology, Neurosurgery and Psychiatry; 52: 940-948.

Wessely, S. and Thomas, P.K. (1990):

The chronic fatigue syndrome-myalgic encephalomyelitis or post-viral fatigue.

Recent Advances in Neurology; 6: 85-131.

Westergren, A. (1926):

Studies of suspension stability of the blood in pulmonary tuberculosis.

Am. Rev. Tuberc., 14: 94.

Weyand, C.M.; Hicok, K.C. and Conn, D.L. (1992):

The influence of HLA DRB1 genes on disease severity in RA.

Ann. Intern. Med., 117: 801.

Wolfe, F.; Cathey, M.A.; Hawley, D.J. et al. (1985):

Clinical trial with r-IFN-gamma in rheumatoid arthritis. In: Biologically Based Immunomodulators in Therapy of Rheumatic Diseases.

Amsterdam, Elsevier; P. 379.

Wolfe, F.; Hawley, D.J. and Wilson, K. (1996):

The prevalence and meaning of fatigue in rheumatic disease.

J. Rheumatol., 23: 1407-1417.

Wolfe, F.; Smythe, H.A.; Yunus, M.B. et al. (1990):

The American College of Rheumatology 1990 criteria for the classification of fibromyalgia: Report of the Multicenter Criteria Committee.
Arthritis Rheum., 33: 160-72.

Yousef, G.E.; Bell, E.J. and Mann, G.P. (1988):

Chronic enterovirus infection in patients with viral fatigue syndrome.
Lancet; I: 146-150.

Yunus, M.B. and Masi, A.T. (1992):

Fibromyalgia, restless leg syndrome, periodic limb movement disorder and psychogenic pain. In: McCarty D.J., Koopman W.J. (eds.): Arthritis and Allied Conditions: A text of Rheumatology, Philadelphia, Lea, Febiger, PP. 1383-1405.

Yunus, M.B.; Masi, A.T. and Aldage, J.C. (1989):

Preliminary criteria for Primary Fibromyalgia Syndrome (PFS): Multivariate analysis of a consecutive of a consecutive series of PFS, other pain patients and normal subjects.
Clin. Exp. Rheumatol., 7: 63-69.

Zigmond, A. and Snaith, R. (1983):

The hospital anxiety and depression scale.
Acta. Psych. Scand., 67: 361-370.

Zvaifler, N.J. (1993):

Etiology and pathogenesis of rheumatoid arthritis. In: McCarty D., Koopman W.: Arthritis and Allied Conditions, 12th ed. London, Lea, Philadelphia, Febiger, PP. 723-761.

المراجع العربية

قائمة حالة - سمة القلق:

تعريب د. أمينة كاظم
دار القلم للنشر والتوزيع

مقياس بك للإكتئاب:

تعريب د. غريب عبدالفتاح
مكتبة النهضة المصرية