

VI. REFERENCES

- Abdel-Tawab, F.M.; M.A. Rashed, R.S. Dhindsa, A. Bahieldin and A. Abo Doma (1998b). Molecular markers for salt tolerance in *Sorghum bicolor*. Inter. Cong. Mol. Genet., 195-204.
- Abdel-Tawab, F.M.; M.A. Rashed, F.M. El-domyati; T.Z. Salam; S.A. Azer and A.F. Khafaga (2001). Marker-assisted selection for salt tolerance in maize (*Zea mays* L.). J. Genet. Cytol., 30: 175-188.
- Abdel-Tawab, F.M.; C.W. Stuber; M.A. Rashed, A. Bahieldin and A.H. Atta (1998c). Genome mapping in maize utilizing recombinant inbreds with RFLP and isozymes markers associated with QTLs affecting yield-related traits under salt stress. Inter. Cong. Mol. Genet., Feb., 21-26. Cairo, Egypt.
- Abou-Khadrah, S.H.; S.A. Abdel-Hafez; F.A. Sorour and A.Z. El-Bably (1999). Influence of irrigation with saline water on wheat yield, yield components and nutrient uptake. Irrigation management and saline conditions. Proceedings of Regional Symposium, JUST, Irbid, Jordan, June 1999. 1999, 87-97; 21 ref.
- Afiah, S.A.N.; E.T. Kishk and A.M. Abdel-Hakim (1999). Genetic analysis of yield and its attributes under two salinity levels in bread wheat (*Triticum aestivum* L.). Annals-of-Agricultural-Science-Cairo. 1999, 44: 1, 309-336; 30 ref.
- Akhtar, J.; J. Gorham; R.H. Qureshi and M. Aslam (1998). Does tolerance of wheat to salinity and hypoxia correlate with root dehydrogenase activities or aerenchyma formation?. Plant-and-Soil. 1998, 201: 2, 275-284; 46 ref.
- Akkaya, M.S.; A.A. Bhagwat and P.B. Cregan (1992). Length polymorphisms of simple sequence repeat DNA in soybean. Genetics, 132:131-139.
- Akkaya, M.S.; R.C. Shoemaker; J.E. Specht; A.A. Bhagwat and P.B. Cregan (1995). Integration of simple sequence repeat DNA markers into a soyabean linkage map. Crop Sci., 35: 1439-1445.
- Alam, S.M.; A.R. Azmi; S.A. Ala; S.S.M. Naqvi and R. Ansari (1998). Effect of aqueous leaf extract of field bindweed (*Convolvulus arvensis* L.) and salinity on growth of wheat. Rachis. 1998, 17: 1-2, 49-51; 16 ref.
- Almansouri, M.; J.M. Kinet and S. Lutts (1999). Compared effects of sudden and progressive impositions of salt stress in three durum

- wheat (*Triticum durum* Desf.) cultivars. Journal-of-Plant-Physiology. 1999, 154: 5-6, 743-752; 38 ref.
- Aouad, A.; M. Mergoun and M. Baaziz (1996). Preliminary study of enzyme polymorphism revealed at the early developmental stages of some cereals. Al Awamia. 1996, publi. 1998, No. 95, 919; 18 ref.
- Ashraf, M. and J.W. O'-Leary (1999). Changes in soluble proteins in spring wheat stressed with sodium chloride. Biologia-Plantarum. 1999, 42: 1, 113-117; 23 ref.
- Bahraei, S. (1996). The use of random amplified polymorphic DNA (RAPD) markers in genetic diversity studies of *Triticum boeoticum* and *T. urartu*. Seed-and-Plant. 1996, 12: 1, 31-43(Pe), 5 (En); 15 ref.
- Baldini, M. and G. P. Vannozzi (1998). Agronomic and Physiological assessment of genotypic variation for drought tolerance in sunflower genotypes obtained from cross. Agricoltura Mediterra, 128,3: 232-240.
- Barakat, M.N. and M.K. El-Haris (1998). *In vitro* selection for salt-tolerant lines in wheat. III. Agronomic evaluation of the progeny under salt stress conditions. Alexandria-Journal-of-Agricultural-Research. 1998, 43: 2, 21-32; 18 ref.
- Barrett, B.A. and K.K. Kidwell (1998). AFLP-based genetic diversity assessment among wheat cultivars from the pacific northwest. Crop-Science. 1998, 38: 5, 1261-1271; 40 ref.
- Barrett, B.A.; K.K. kidwell and P.N. fox (1998). Comparison of AFLP and pedigree-based genetic diversity assessment methods using wheat cultivars from the pacific nor thwest. Crop-Science. 1998, 38:5, 1271-1278; 30 ref.
- Barriga, B.P.; T.J.C. Slebe; S.J. Mansilla (1994). Genetic analysis with RAPD markers in wheat. Agro-Sur. 1994, 22: 2, 133-142; 12 ref.
- Beckmann, J. and M. Sollar (1990). Restriction fragment length polymorphism in genetic improvement. In. B.J. Miflin (ed) Oxford Survey of Plant, Molecular Biology, Vol. 3: 196-250.
- Beckmann, J. and M. Sollar (1993). Restriction fragment length polymorphism in genetic improvement methodology. Mapping and cost Theor. Appl. Genet. 67: 33-43.
- Begum, F. and J.L. Karmoker (1999). Effect of salinity stress on the accumulation and distribution of proline in wheat. Rachis. 1999, 18: 1, 22-25; 19 ref.
- Bohnert, H.J.; D.E. Nelson and R.G. Jensen (1995). Adaption to enviromental stresses. Plant Cell, 7: 1099-1111.

- Bowcock, A.M.; A. Ruiz-linares, J. Tomfohrde; E. Minch; J.R. Kidd and L.L. Cavalli-Sforza (1994). High resolution of human evolutionary trees with polymorphic microsatellites. *Nature*, 368:455-457.
- Bruford, M.W. and R.K. Wayne (1993). Microsatellite and their genetic application to population genetic studies. *Curr. Opin. Genet. Dev.*, 3: 939-943.
- Brunell, M.S.; A.J. Lukaszewski and R. whitkus (1999). Development of arm specific RAPD markers for rye chromosome 2R in wheat. *crop science*. 1999, 39 : 6, 1702- 1706 ; 35 rep .
- Cao, W.; G. Scoles; P. Hucl and R.N. Chibbar (1999). The use of RAPD analysis to classify *Triticum* accessions. *Theoretical-and-Applied-Genetics*. 1999, 98: 3-4, 602-607; 19 ref.
- Cao, wenGuang; G. Scoles; P. Hucl; R.N. chibbar and W.G. Cao. (2000). phylogenetic relationships of five morphological groups of hexaploid wheat (*Triticum aestivum* L. em The 11) based on RAPD analysis. *Genome*. 2000, 43: 4, 724-727;17 ref.
- Castagna, R.; S. Gnocchi; M. Perenzin and M. Heun (1997). Genetic variability of the wild diploid wheat *Triticum urartu* revealed by RFLP and RAPD markers. *Theoretical-and-Applied-Genetics*. 1997, 94: 3-4, 424-430; 42 ref.
- Chabane, K.; J. Valkoun and A.A. Jaradat (1998). Standardization of RAPD marker techniques to determine the diversity of diploid wheat: *Triticum urartu*. *Triticeae III. Proceedings of the Third International Triticeae Symposium, Aleppo, Syria, 4-8 May 1997*. 1998, 155-158; 14 ref.
- Chen, DeMing.; RenPei. Y.u; D.M. Chen; R.P. Y.u (1998). Salt resistance and ionic characteristics of three wheat cultivars under salt stress. *Acta-Pedologica-Sinica*. 1998, 35: 1, 88-94; 7 ref.
- Chen, DeMing.; RenPei. Y.u; D.M. Chen and R.P. Y.u (1998). Salt-resistance and ionic characteristics of three wheat varieties under salt stress. *Acta-Pedologica-Sinica*. 1998, 35: 1, 88-94; 7 ref. cultivars for *in vitro* salt tolerance. *Journal-of-Science-and-Technology-of-Agriculture-and-Natural-Resources*. 1999, 3: 1, 21-34; 27 ref.
- Cao, W.; G. Scoles; P. Hucl and R. Chibbar (1999). The use of RAPD analysis to classify *Triticum* accessions. *Theor. Appl. Genet.*, 4: 602-607.
- Debener, T.; C. Barlets and L. Mattiesch (1996). RAPD analysis of genetic variation between a group of rose cultivars and selected with rose spe. *Mol. Breeding*, 2: 312-326.

- Deragon, J.M. and B.S. Landry (1992). RAPD and other PCR-based analyses of plant genomes using DNA extracted from small leaf disks. *PCR-Methods-and-Applications*. 1992, 1: 3, 175-180; 22 ref.
- Devos, K.M. ; G.J. Bryan ; A.J. collines ; P. stephenson and M.D. Gale (1995). Application of two microsatellite sequences in wheat storage proteins as molecular markers. *Theoretical and Applied Genetics*. 1995, 90: 2, 247-252; 18 ref.
- Devos, K.M. and M.D. Gale (1992). The use of random amplified polymorphic DNA markers in wheat. *Theoretical-and-Applied-Genetics*. 1992, 84: 5-6, 567-572; 14 ref.
- Dhaliwal, H.S.; J.S. Sidhu and J.L. Minocha (1993). Genetic diversity in diploid and hexaploid wheats as revealed by RAPD markers. *Crop-Improvement*. 1993, 20: 1, 17-20; 12 ref.
- Donini, p.; M.I. Elias; S.M. Bougourd and R.M.D. Koebner (1997). AFLP fingerprinting reveals pattern differences between template DNA extrated from Different plant organs. *Genome*. 1997, 40: 4, 521-526; 15 ref.
- Duncan, D. B. (1955), multiple range and multiple F test. *Biometrics*, 11: 1-42.
- Dweikat, I.; S. Mackenzie; M. Levy; H. Ohm (1993) Pedigree assessment using RAPD-DGGE in cereal crop species. *Theoretical-and-Applied-Genetics*. 1993, 85: 5, 497-505; 37 ref.
- El-Sayed, F.F.; A.M.A.S. Ali; M.M. El-Menoufi; R.A. Mitkees; A.A. Hamada; M.A. Mahrous; T.M. Shehab El-Din; M.M. El-Shami; M.A. Mostafa; A.M. Abdel-Ghani; A.H. Abdel-Latif; M.S. Sharshar; M.A. Eid; Eman, M.M. Sadek; A.M. Ali; M.H. Iskandar; A.H. Bassiuni; A.M.K. Eissa; Y.G. Abdel-Gawad; Kadna Hegazi and Y.H. El-Daoudi (1996). An introduction to the new bread wheat cultivar Gemmeiza 3. *J. Agric. Sci. Mansoura Univ.*, 21 (11): 3811-3823.
- Evlagon, D.; I. Ravina and P. Neumann (1990). Interactive effects of salinity and calcium on hydraulic conductivity osmotic adjustment and growth in primary root of maize seedlings. *Israel J. of Botany*, 39: 239-247.
- Fahima, T.; G.L. Sun; A. Beharav; T. Krugman; A. Beiles and E. Nevo (1999). RAPD polymorphism of wild emmer wheat populations, *Triticum dicoccoides*, in Israel. *Theoretical-and-Applied-Genetics*. 1999, 98: 3-4, 434-447; 53 ref.
- Farooq, S.; T.M. Shah; E. Askari; A.A. Zaidi; M. Arif and N. Iqbal (1994). Identification of different wheat genotypes through polymorphism

based on random amplified polymorphic DNA (RAPD). *Pakistan-Journal-of-Botany*. 1994, 26: 2, 373-382; 23 ref.

Fritz, A.K.; T.S. Cox; B.S. Gill and R.G. Sears (1995). Molecular marker-facilitated analysis of introgression in winter wheat X *Triticum tauschii* populations. *Crop-Science*. 1995, 35: 6, 1691-1695; 25 ref.

Fritz, A.K.; T.S. Cox; B.S. Gill and R.G. Sears (1995). Marker-based analysis of quantitative traits in winter wheat X *Triticum tauschii* populations. *Crop-Science*. 1995, 35: 6, 1695-1699; 22 ref.

Gao, WenGuang; P. Hucl; G. Scoles; R.N. Chibbar and W.G. Cao (1998). Genetic diversity within spelta and macha wheats based on RAPD analysis. *Euphytica*. 1998, 104: 3, 181-189; 21 ref.

Ghulam, A.; A. Hussain; A.I. Jaloud; S.A. Al-shammary; S. Karimulla and O.S. Aswald (1997). Effect of saline irrigation on germination and growth parameters of barely (*Hordium vulgare* L.) in pot experiment. *Agric. Water Management*. 34: 2, 125-135.

Gill, K.S.; E.L. Lubbers; B.S. Gill; W.J. Raupp and T.S. Cox (1991). A genetic linkage map of *Triticum tauschii* (DD) and its relationship to the D genome of bread wheat (AABBDD). *Genome*. 1991, 34: 3, 362-374; 60 ref.

Gomaa, A.S.A. (1999). Wheat improvement in Egypt; history and future prospects. Proceed, First PL. Breed. Conf., December 4. Giza. Special Issue of Egypt. J. Plant Breed., Vol. 3: 1-14.

Gupta, P.K. and R.K. Varshney (2000). The development and use of microsatellite markers for genetic analysis and plant breeding with emphasis on bread wheat. *Euphytica*. 2000, 113: 3, 163-185, 7pp. of ref.

Gupta, V. and A.E. Slinkard (1998). Molecular marker assisted wheat improvement in India: a progress report. Proceedings of the ninth international wheat genetics symposium, Saskatoon, Saskatchewan, Canada, 2-7 August 1988, 138-141; 3 ref.

Hakki, E.; C. Savaskan and M. S. Akkaya (2001). Genotyping of Anatolian doubled-haploid durum lines with SSR markers. *Euphytica*, 122:257-262.

Hanna, N.S.; S.R.S. Sabry; M.M. Khodier and M.K. Moshref (1999). Scales for measuring wheat (*Triticum aestivum* L.) salinity tolerance in pot and field experiments. *Annals-of-Agricultural-Science-Cairo*. 1999, 44: 2, 573-587; 15 ref.

He, CongFen.; ZhanJing. Huang; YinZhu. Shen; ZhiYi. Liu; Feng. Bai; C.F. He; Z.J. Huang; Y.Z. Shen; Z.Y. Liu and F. Bai (1996). RAPD

- analysis on different nullisomic lines of wheat cv. "Chinese Spring". *Acta-Agriculturae-Boreali-Sinica*. 1996, 11: 3, 31-34; 5 ref.
- He, S.; H. Ohm and S. Mackenzie (1992). Detection of DNA sequence polymorphisms among wheat varieties. *Theoretical-and-Applied-Genetics*. 1992, 84: 5-6, 573-578; 27 ref.
- Hernandez, P.; A. Martin and G. Dorado (1999). Development of SCARs by direct sequencing of RAPD products: a practical tool for the introgression and marker-assisted selection of wheat. *Molecular-Breeding*. 1999, 5: 3, 245-253; 24 ref.
- Hernandez, P.; M. Hemmat; N.F. Weeden; G. Dorado and A. Martin (1999). Development and characterization of *Hordeum chilense* chromosome-specific STS markers suitable for wheat introgression and marker-assisted selection. *Theoretical-and-Applied-Genetics*. 1999, 98: 5, 721-727; 22 ref.
- Hewstone, O.N. and R.P. Hinrichsen (1994). High molecular weight glutenin subunit composition of Chilean bread wheats (*Triticum aestivum* L.). *Agricultura Tecnica*; 54 (3) 211-218, 24 ref.
- Hohmann, U.; T.R. Endo; K.S. Gill; B.S. Gill (1994). Comparison of genetic and physical maps of group 7 chromosomes from (*Triticum aestivum* L.). *Molecular-and-General-Genetics*. 1994, 245: 5, 644-653; 38 ref.
- Hu, Y.; J.J. Oertli and U. Schmidhalter (1997). Interactive effects of salinity and macronutrient nutrition. 20:9, 1155-1167.
- Iqbal, M.J. and A.L. Rayburn (1994). Stability of RAPD markers for determining cultivar specific DNA profiles in rye (*Secale cereale* L.). *Euphytica*, 75: 215-220.
- Joshi, C.P. and H.T. Nguyen (1993). RAPD (random amplified polymorphic DNA) analysis based intervarietal genetic relationships among hexaploid wheats. *Plant. Sci. Limerick*. 1993, 93: 1-2, 95-103; 33 ref.
- Joshi, C.P. and H.T. Nguyen (1993). Application of the random amplified polymorphic DNA technique for the detection of polymorphism among wild and cultivated tetraploid wheats. *Genome*. 1993, 36: 3, 602-609; 36 ref.
- Katerji, N.; J.W. van. Hoorn; A. Hamdy; M. Mastrorilli and J.W. Van Hoorn (2000). Salt tolerance classification of crops according to soil salinity and to water stress day index. *Agricultural-Water-Management*. 2000, 43: 1, 99-109; 21 ref.
- Keller, M.; C. Karutz; J.E. Schmid; P. Stamp and M. Winzeler (1999). Quantitative trait loci for lodging resistance in a segregating wheat X

- spelt population. *Theoretical and Applied Genetics*. 1999, 98: 6-7, 1171-1182; 51 ref.
- Keller, N.L. and Evers, A.D. (1994). *Technology of cereals*, Fourth Edition. Elsevier Science. Ltd. Oksford, UK.
- Kerepesi, I.; G. Galiba and E. Banyai (1998). Osmotic and salt stresses induced differential alteration in water-soluble carbohydrate content in wheat seedlings. *Journal-of-Agricultural-and-Food-Chemistry*. 1998, 46: 12, 5347-5354; 41 ref.
- Khlestkina, E.K.; D.E.A. Salina; I.N. Leonova; L.I. Laiknova and S.F. Koval (1999). The use of RAPD and STS analyses for marking genes of homologous group 5 chromosomes of common wheat. *Genetika-Moskva*. 1999, 35:10, 1349-1357; 20 ref.
- Khodier, M.M.; S.R.S. Sabry and N.S. Hanna (1999). Screening wheat (*Triticum aestivum* L.) genotypes for salinity tolerance in sandy and clay loam soils. *Annals-of-Agricultural-Science-Cairo*. 1999, 44: 1, 151-157; 11 ref.
- Ko, H.L.; R.J. Henry; R.J. (ed.). Henry and J.A. Ronalds (1994). Rapid cereal genotype analysis. Improvement of cereal quality by genetic engineering. Proceedings of the Royal Australian Chemistry Institute, Cereal Chemistry Division Symposium on Improvement of Cereal Quality by Genetic Engineering, Sydney, Australia, 12-16 September 1993. 1994, 153-157; 7 ref.
- Koebner, R.M.D. and P.K. Martin (1994). RAPDs as molecular markers for the detection of the presence of rye chromosomes in wheat. *Journal-of-Genetics-and-Breeding*. 1994, 48: 1, 85-88; 18 ref.
- Koebner, R.M.D. (1995). Predigestion of DNA template improves the level of polymorphism of random amplified polymorphic DNAs in wheat. *Genetic-Analysis-Techniques-and-Applications*. 1995, 12: 1, 63-67; 15 ref.
- Koller, B.; A. Lehmann; J.M. McDermott and C. Gessler (1993). Identification of apple cultivars using RAPD markers. *Theor Appl. Genet.*, 85: 901-904.
- Konak, C.; R. Yilmaz and O. Arabaci (1999). Salt tolerance of wheat of the Aegean region. *Turkish-Journal-of-Agriculture-and-Forestry*. 1999, 23: Supplement 5, 1223-1229; 14 ref.
- Kondratjev, M.N. and T.A. Rybkina (1998). Response and adaptation of wheat, millet and sorghum to chloride salinity. *Izvestiya-Timiryazevskoi-Sel'skokhozyaistvennoi-Akademii*. 1998, No. 3, 123-139; 28 ref.

- Kong, LingRang.; YuChen. Dong; J.I. Zeng. Jia; L.R. Kong; Y.C. Dong and J.Z. Jia (1998). Random amplified polymorphism of DNA analysis in *Aegilops tauschii*. *Acta-Botanica-Sinica*. 1998, 40: 3, 223-227; 20 ref.
- Kong-Xiuring; Ge- Chumin; Jia- Jizeng; Dong- Yushen; R.R.C. Wang; X.Y. Kong; C.M. Ge; J.Z. Jia and Y.S. Dong (1999). RAPD anlysis of phylogeny for five basic genomes in Agilops. *Acta-Botanica-sinica*. 1999, 41:4, 393-397; 18 ref.
- Labuschagne, M.T.; H. Maartens and A.M. Oberholster (1998). A comparison of HMW-GS, LMW-GS and RAPD fingerprinting for cultivar identification of some South African wheat cultivars. *South-African-Journal-of-Plant-and-Soil*. 1998, 15: 4, 147-150; 14 ref.
- Lagudah, E.S.; D. The; R. Appels; R.F. Eastwood; R.A. McIntosh; S. chandramorhan; D. Hoisington (ed.) and A. McNab (1993). Application of molecular markers and genetic maps: analysis of rust and Heterodera avenae resistance derived from (*Triticum tauschii*). Progress in genome mapping of wheat and related species: proceedings of the 3 rd public workshop of the International Triticea Mapping Initiative, 22-26 September 1992, CIMMYT, Mexico. 1993, 26-28.
- Langridge, P.; K. Chalamers and A.E. Slinkard (1998). Techniques for marker development. proceddings of the ninth international wheat genetics symposium, sakatoon, sakatchewan, canada, 2-7 August 1998. 1988, 107-117; 3 pp. of ref.
- Lee, S.J.; G.A. Penner and K.M. Devos (1995). Characterization of loci containing microsatellite sequences among Canadian wheat cultivars. *Genome*. 1995, 38: 5, 1037-1040; 7 ref.
- Litvinenko, N.A.; H.J. Braun (ed.); F. Altay (ed.); W.K. kronstad (ed.); S.P.S. Beniwal (ed.) and A. McNab (1997). Breeding intensive winter bread wheat varieties for southern Ukraine wheat: prospects for global improvement. Proceedings of the 5th International wheat conference, Ankara, Turkey, 10-14 June 1996. 1997, 29-36; Developments in plant Breeding volume 6; 9 ref.
- Liu, Z.; P.M. Biyashev and M.A. Saghai-Marooof (1996). Development of simple sequence repeat DNA markers and their integration into a barley linkage map. *Theoretical and Applied Genetics*. 1996, 93: 5-6, 869-876; 38 ref.
- Liu, Z.Q.; Y. Pei and Z.J. Pu (1999). Relationship between hybrid performance and genetic diversity based on RAPD markers in wheat, (*Triticum aestivum* L). *Plant-Breeding*. 1999, 118: 2, 119-123; 31 ref.

- Loarce, Y.; G. Hueros and E. Ferrer (1996). A molecular linkage map of rye. *Theoretical and Applied Genetics*. 1996, 93: 7, 1112-1118; 41 ref.
- Ma, Z.Q.; M. Rader and M.E. Sorrells (1996). Frequencies and sequence characteristics of di, tri and tetranucleotide microsatellite in wheat. *Genome*, 29: 123-130.
- Malkawi, H.I.; M.M. Ajlouni; A.A. Jaradat and A.A. Jaradat (1998). Analysis of genetic diversity of wild emmer and durum wheats by RAPDs. *Triticeae III. Proceedings of the Third International Triticeae Symposium, Aleppo, Syria, 4-8 May 1997*. 1998, 165-170; 20 ref.
- Mammadov, J.A. (1999). Evaluation of drought- and salt-resistance during the germination of selected durum wheat lines. *Turkish-Journal-of-Biology*. 1999, 23: 2, 177-186; 18 ref.
- Manabe, M.; T. Ino; M. Kasaya; S. Takumi; N. Mori; I. Ohtsuka and Nakamura (1999). Segregation distortion through female gametophytes in interspecific hybrids of tetraploid wheat as revealed by RAPD analysis. *Hereditas.*, Landskrona, 131(1) : 47-53.
- Sambrook, J.; E.F. Fritsch and T. Maniatis (1989). *Molecular cloning. A Laboratory manual*. Cold Spring Harbor laboratory Publisher, New York.
- McDonald, M.B.; L.J. Elliot and P.M. Sweeney (1994). DNA extraction from dry seeds for RAPD analyses in varietal identification studies. *Seed-Science-and-Technology*. 1994, 22: 1, 171-176; 15 ref.
- Michelmore, R.W.; I. Paran and R.V. Kesseli (1991). Identification of markers linked to disease- resistance genes by bulked segregant analysis. A rapid method to detect markers in specific region by using segregating population. *Proc. Nat. Acad. Sci.*, 88: 9828-9832.
- Mohamed, I.Z.; A.A. Abd El-Raheem; M.H. Iskander and G.F. Detizer (1998). Genetic analysis of some physiological and yield characters under salinity conditions in wheat (*Triticum aestivum* L.). *Egypt. J. Appl. Sci.* 13, 9: 48-63.
- Mohan, M.; S. Nair; A. Bhagwat; T.G. Krishna; M. Yano; C.R. Bhatia and T. Sasaki (1997). Genome mapping, molecular markers and marker-assisted selection in crop plants. *Molecular Breeding*, 3: 87-103.
- Munns, R. (1993). Physiological processes limiting plant growth on saline soils some dogmas and hypotheses. *Plant Cell Environment*, 16: 15-24.
- Munns, R. and H.M. Rawson (1999). Effect of salinity on salt accumulation and reproductive development in the apical meristem of wheat and barley. *Australian-Journal-of-Plant-Physiology*. 1999, 26: 5, 459-464; 21 ref.

- Munns, R. and Termaat, A. (1986). Whole plant responses to salinity. *Aust. J. Plant Physiology*, 13: 143-160.
- Munns, R.; R.A. Hare; R.A. James and G.J. Rebetzke (2000). Genetic variation for improving the salt tolerance of durum wheat. *Australian-Journal-of-Agricultural-Research*. 2000, 51: 1, 69-74; 23 ref.
- Nagaoka, T. and Y. Ogihara (1997). Applicability of inter-simple sequence repeat polymorphisms in wheat for use as DNA markers in comparison to RFLP and RAPD markers. *Theoretical-and-Applied-Genetics*. 1997, 94: 5, 597-602; 22 ref.
- Nelson, J.c.; , A.E. van Deynze; E. Autrique.; M.E. sorrells; Y.H. LU.; M. Merlion; M. Atkinson and P. Leroy (1995). Molecular mapping of wheat. Homoeologous group 2. *Genome* 1995 Vol. 38, no. 3, ppl 516 - 524.
- Nelson, J.c.; , A.E. van Deynze; E. Autrique.; M.E. sorrells; Y.H. LU.; S. Negre.; M. Bernard. and P. Leroy (1995). Molecular mapping of wheat. Homoeologous group 3. *Genome* 1995 Vol. 38, no. 3 pp 525-533.
- Nguyen, H.T.; C.P. Joshi; D. (ed.). Hoisington and A. McNab (1993). RAPD analysis in tetraploid and hexaploid wheats. Progress in genome mapping of wheat and related species: Proceedings of the 3rd Public Workshop of the International Triticeae Mapping Initiative, 22-26 September 1992, CIMMYT, Mexico. 1993, 47-50; 8 ref.
- Nguyen, H.T.; D. (ed.). Hoisington and A. McNab (1993). Wheat genetics research at Texas Tech University. Progress in genome mapping of wheat and related species: Proceedings of the 3rd Public Workshop of the International Triticeae Mapping Initiative, 22-26 September 1992, CIMMYT, Mexico. 1993, 51-52; 12 ref.
- Parker, G.D.; K.J. Chalmers; A.J. Rathjen and P. Langridge (1998). Mapping loci associated with flour colour in wheat (*Triticum aestivum* L.). *Theoretical and Applied Genetics*. 1998, 97: 1-2, 238-245; 42 ref.
- Perenzin, M.; M. Corbellini; M. Acerbi; P. Vaccino; B. Borghi; H.J. Braun (ed.); F. Altay (ed.); W.E. Kronstad (ed.); S.P.S. Beniwal (ed.) and A. McNab (1997). Bread wheat: F₁ hybrid performance and parental diversity estimates using molecular markers. Wheat; prospects for global improvement. Proceeding of the 5th international wheat conference, Ankara, Turkey, 10-14 June 1996. 1997, 339-345; Developments in plant Breeding Volume 6; 26 ref.
- Prasad, M.; R.K. Varshney; A. Kumar; H.S. Balyan; P.C. Sharma; K.J. Edwards; H. Singh; H.S. Dhaliwal; J.K. Roy and P.K. Gupta (1999). A microsatellite marker associated with a QTL for grain

- protein content on chromosome arm 2 DL of bread wheat. Theoretical and applied genetics. 1999, 99: 1-2, 341-345; 29 ref. 1999.
- Pujar, S.; S.A. Tamhankar; V.S. Rao; V.S. Gupta; S. Naik and P.K. Ranjekar (1999). Arbitrarily primed-PCR based diversity assessment reflects hierarchical groupings of Indian tetraploid wheat genotypes. Theoretical and Applied Genetics. 1999, 99: 5, 868-876; 46 ref.
- Qiao, YuHui.; ZhenRong. Y.u; YinSuo. Zhang; JingFeng. Xin; P.M. Driessen; Y.H. Qiao; Z.R. Y.u; Y.S. Zhang and J.F. Xin (1999). Effects of irrigation with light saline water on growth of winter wheat and soil environment in salinized regions. Soils-and-Fertilizers-Beijing. 1999, No. 4, 11-14; 8 ref.
- Rahman, M.; T.A. Malik; M.J. Iqbal; Y. Zafar; K. Alam; Z. Perveen and S. Rehman (1998). DNA marker for salinity resistance. Rice Biotechnology Quarterly. 1998, 35: 12-13; 5 ref.
- Ranjekar, P.K.; H.S. Dhaliwal; V.S. Rao; V.S. Gupta, R. Tiwari; S. Nagarajan (ed), Singh gyanendra (ed) and B.S. Tyagi (1998). Molecular biology approaches to wheat improvement in India Present status and future strategies. Wheat; research needs beyond 2000 Ad. Proceedings of an international conference, 12-14 august 1997, Karnal, India. 1998, 145-159; 5 pp of ref.
- Riede, C.R.; D.J. Fairbanks; W.R. Andersen; R.L. Kehrer and L.R. Robison (1994). Enhancement of RAPD analysis by restriction-endonuclease digestion of template DNA in wheat. Plant Breeding. 1994, 113: 3, 254-257; 7 ref.
- Roy, j. k.; M. Prasad; R.K. Varshney; H.S. Balyan; T.K. Blake; H.S. Dhaliwal; H. singh; K.J. Edwards and P.K. Gupta (1999). Identification of microsatellite on chromosomes 6B and a STS on 7D of bread wheat showing an association with preharvest sprouting tolerance. Theoretical and Applied Genetics. 1993, 99: 12, 336; 39 ref.
- Saghai-Marooof, M.A.; R.B. Biyashev; G.P. Yang; Q. Zhang and R.W. Allard (1994). Extraordinarily polymorphic microsatellite DNA in barley: Species diversity, chromosomal locations, and population dynamics. Proc. Natl. Acad. Sci., USA, 91:5466-5470.
- Salam, A.; P.A. Hollington; J. Gorham; R.G.W. Jones and C. Gliddon (1999). Physiological genetics of salt tolerance in wheat (*Triticum aestivum* L.): performance of wheat varieties, inbred lines and reciprocal F1 hybrids under saline conditions. J. Agronomy and Crop Sci. 1999, 183: 3, 145-156; 39 ref.

- Sanjay, Singh; T.N. Singh and S. Singh (1999).** Root growth of wheat in simulated vertical and lateral splits of layered salt profiles in soil. *Indian J. Plant Physiology*. 1999, 4: 2, 73-78; 17 ref.
- Sayed, Shahram Mirodjagh.; Arzani. Ahmad; S.S. Mirodjagh and a. Arzani (1999).** Assessment of durum wheat (*Triticum turgidum* var. *durum*) cultivars for *in vitro* salt tolerance. *J. Sci. and Technol. Of Agriculture and Natural Resources*. 1999, 3: 1, 21-34; 27 ref.
- Shah, M.M.; Yang Yen; K.S. Gill; P.S. Baenziger and Y. Yen (2000).** Comparisons of RFLP and PCR-based markers to detect polymorphism between wheat cultivars. *Euphytica*. 2000, 114: 2, 135-142; 28 ref.
- Sharon, D.; A. Adato; S. Mhameed; U. Lavi; J. Hillel; M. Gomolka; C. Epplen and J.T. Epplen (1995).** DNA fingerprints in plants using simple- sequence repeat and minisatellite probes. *Hortscience*. 1995, 30: 1, 109-112; 25 ref.
- Sivolap, Yu.M.; L.I. Kutsevich; A.I. Palamarchuk; V.N. Totsky and V.N. Totskii (1997).** Molecular-genetic polymorphism of winter durum wheat determined by polymerase chain reaction with arbitrary primers. *Russian-Agricultural-Sciences*. 1997, publ. 1998, No. 1, 9-13; translated from *Doklady Rossiiskoi Akademii Sel'skokhozyaistvennykh Nauk* (1997) No. 1, 6-8.; 4 ref.
- Slafer, G.A. and E.H. Satorre (1999).** An introduction to the physiology-ecology and physiology of yield determination (Satorre; E.H. and G.A. Slafer Eds.). New York pp. 3-13.
- Slafer, G.A.; Satorre, E.H. and F.H. Andrade (1994).** Increases in grain yield in bread wheat from breeding and associated physiological changes. In genetic improvement of field crops (Slafer, G. A. Ed). Marcel Dekker, inc. New York pp. 1-68.
- Sourdille, P.; M.R. Perretantp; G. Charmet; T. Cadalen; M.H. Tixier; P. Joudrier; M.F. Gautier; G. Branlard; S. Bernard; C. Boeuf; M. Bernard; G.T. Scarascia-Mugnozza (ed.); E. porcedudu (ed.) and M.A. Pagnotta (1999).** Detection of QTL for bread-making quality in wheat using moleuclar markers. Genetics and breeding for crop quality and resistance. Proceedings of the XV EUCARPIA congress, Viterbo, Italy, September 20-25, 1998, 1999, 361-366; 12 ref.
- Steppuhn, H. and K.G. Wall (1999).** Canada's salt tolerance testing laboratory. *Canadian-Agricultural-Engineering*. 1999, 41: 3, 185-189; 10 ref.

- Sun, GenLou; B. Salomon; R.von. Bothmer and R. Von-Bothmer (1997). Analysis of tetraploid *Elymus* species using wheat microsatellite markers and RAPD markers. *Genome*. 1997, 40: 6, 806-814; 54 ref.
- Talbert, L.E.; N.K. Blake; P.W. Chee; T.K. Blake and G.M. Magyar (1994). Evaluation of "sequence-tagged-site" PCR products as molecular markers in wheat. *Theoretical-and-Applied-Genetics*. 1994, 87: 7, 789-794; 19 ref.
- Tyrka, M.; P. Dziadczyk; D. Miazga; C. (ed.). Ceoloni and A.J. Worland (1998). The use of RAPD method in analysis of disomic addition lines 'Grana'-'Dankowskie Zlote'. Proceedings of the 10th EWAC meeting, Viterbo, Italy, 16-19 June 1997. *EWAC-Newsletter*. 1998, 118-121; 14 ref.
- Varshney, P.K.; P.C. Sharma; P.K. Gupta; H.S. Balyan; B. Ramesh; J.K. Roy; A. Kumar and A. Sen (1998). Low level of polymorphism detected by SSR probes in bread wheat. *Plant- breeding*. 1998, 117: 2, 182-184; 15 ref.
- Vierling, R.A. and H.T. Nguyen (1992). Use of RAPD markers to determine the genetic diversity of diploid, wheat genotypes. *Theoretical-and-Applied-Genetics*. 1992, 84: 7-8, 835-838; 10 ref.
- Vijai, Prakash.; EVD. Sastry and V. Prakash (1999). Variation and character associations in wheat grown under moderate salinity. *Agricultural Science Digest Karnal*. 1999, 19: 3, 175-178; 4 ref.
- Walton, M. (1994). Microsatellite DNA: new tool for plant breeder's box. *Seed-World-Des-Plaines*. 1994, 132: 3, 20-21.
- Wang, Z.J.; L. Weber; G. Zhong and S. D. Tanksley (1994). Survey of plant short tandem DNA repeats. *Theor. Appl. Genet.*, 88: 1-6.
- Weber J. L. (1990). Informativeness of human (dC-dA)_n, (dG-dT) polymorphism. *Genomics*, 7: 524-530.
- Welsh, J. and M. McClelland (1990). Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acid Resarch*. 18, 24: 7213-7218.
- Wie, JunZhi.; R.R.C. Wang and J.Z. Wie (1995). Genome- and species-specific markers and genome relationships of diploid perennial species in Triticeae based on RAPD analyses. *Genome*. 1995, 38: 6, 1230-1236; 39 ref.
- William, H.M.; D. Hoisington; R.P. Singh and D. Gonzalez de Leon (1997). Detection of quantitative trait loci associated with leaf rust resistance in bread wheat. *Genome*. 1997, 40:2,253-260,33 ref.
- William, J.C.K.; Anne, R. Kubelik; K. J. Liva; J. A. Rafalski and S. V. Tingey (1990). DNA polymorphisms amplified by arbitrary primers

are useful as genetic markers. *Nucleic Acids Research*, 18, 2: 6531-6535.

- William, M.D.H.M.; L. Gray; A.L. Rayburn and A. Mujeeb-Kazi (1994). Development of DNA markers based on randomly amplified polymorphic sequences in a *Triticum aestivum* L. X *Thinopyrum bessarabicum* amphiploid. *J. Genetics and Breeding*. 1994, 48: 1, 1-5; 12 ref.
- Yogita, Gupta.; E.V.D. Sastry and Y. Gupta (1998). Effect of salinity on combining ability in wheat (*Triticum aestivum*). *Indian J. Agricultural Research*. 1998, 32: 2, 139-143; 5 ref.
- Zhang, J.; H. Nguyen and A. Blum (1999). Genetic analysis of osmotic adjustment in crop plants. *J. Experimental Botany*, 50: 291-302.
- Zhao, X.C., M.R. Sharifiou, G. Godd; P.J. Sharp and A.E. Slinkard (1998). Developing waxy wheat Cultivars: wx null alleles and molecular markers. *Proceedings of the Ninth International wheat Genetics symposium, saskatoon, saskatchewan Canada, 2-7 August 1998*. 1998, 254; 17 ref.
- Zhu, ZhiHua and Z.H.Zhu (1998). Yield responses of wheat cultivars to salt stress at various growth phases. *Crop-Genetic-Resources*. 1998, No. 3, 31-33; 2 ref.
- Zyingila, D.; V. Kleizaite and V. Popenkityte (1998). Analysis of RAPD and isozyme (SOD) polymorphism in cereals. *Biotechnologija augalu selekcijoje. Mokslinė konferencija, Lietuvos žemės ūkio universitetas, 1998 m. spalio 8-9 d.* 1998, 33-37; 10 ref.