

References

- Al-Nawas B, Krammer I and Shah PM (1996).*** Procalcitonin in diagnosis of severe infections. Eur J Med Res; 1:331-333.
- Al-Nawas B, and Shah PM(1997).*** Procalcitonin in acute malaria. Eur J Med Res; 2:206-208.
- Alpern ER, Alessandrini EA, and McGowan KL (2001).*** Serotype prevalence of occult pneumococcal bacteremia. Pediatrics Aug; 108(2): E23 .
- Arch J W,(2002)*** Fever in infants and young children , Ped Adal. Med. 156:512.
- Ardaillou R,. (1970).*** Metabolic clearance rate of radioiodinated human calcitonin in man. J Clin Invest; 49:2345-2352.
- Assicot M, Gendrel D, Carsin H, Raymond J, Guilbaud J , and Bohuon C(1993).*** High serum procalcitonin concentrations in patients with sepsis and infection. Lancet; 341:515-518.
- Avner JR, and Baker MD.(2002).*** Management of fever in infants and children. Emerg Med Clin North Am ; Feb; 20(1): 49-67.
- Bachur RG and, Harper MB(2001).*** Predictive model for serious bacterial infections among infants younger than 3 months of age. Pediatrics ;Aug; 108(2): 311-6.

Baker MD(1999). Evaluation and management of infants with fever.
Pediatr Clin North Am ; Dec; 46(6): 1061-72.

Balow A ., and Sonnen wirth, A.C. (1983). Bacteremia laboratory and clinical aspects . Chales C Thomas , Springfield, III.

Baraff LJ, Oslund S, Prather M. and Fleisher GR(1993). Effect of antibiotic therapy and etiologic microorganisms on the risk of bacterial meningitis in children with occult bacteremia.
Pediatrics; 92:140143.

Baraff LJ, Schriger DL, and Bass JW, et al(1997). Management of the young febrile child. Commentary on practice guidelines.
Pediatrics Jul; 100(1): 134-6.

Baraff LJ(2000). Management of fever without source in infants and children. Ann Emerg Med Dec; 36(6): 602-14.

Bass JW, Steele RW, and Wittler RR, (1993). Antimicrobial treatment of occult bacteremia: a multicenter cooperative study.
Pediatr Infect Dis J Jun; 12(6): 466-73.

Bhrmann , Kliegman and Jenson , and Keith R Powell (2000); Chpter of infectious diseares – Fever without a focus –W.B. Sanders company –Nelson textbook of perdidier PP:742-743.

Becker KL, (1994). Small cell lung carcinoma cell line express mRNA for calcitonin and alpha- and beta-calcitonin gene related peptide. Cancer Lett; 81:19-25.

Becker KL, Nylén E, and Thompson K. (1995). Preferential hypersecretion of procalcitonin and its precursors in

pneumonitis: a cytokine-induced phenomenon. ; (Abstract)
Philadelphia, USA, Endotoxemia and Sepsis Congress

Becker KL, Nylén ES, Cohen R, and Snider RH. Calcitonin(1996).

Procalcitonin, structure, molecular biology, and actions.
Principles of Bone Biology, Academic Press Inc; 1:471-494.

**Benador N., Gendrel D., Siegrist C.A., Greder C., Benador D.,
Bohuon C. and Girardin E. (1997).** Procalcitonin is a marker
of severity of renal lesions in pyelonephritis .15th Annual
meeting of ESPID, 21-23 May, abstr. 34.

**Bertagna XY, Nicholson WE, Pettengill OS, Sorenson GD, Mount
CD and, Orth DN(1978).** Ectopic production of high molecular
weight calcitonin and corticotropin by human small cell
carcinoma cells in tissue culture:evidence for separate
precursors. J Clin Endocrinol Metab; 47:1390-1393.

Black SB, Shinefield HR, and Hansen J (2001). Postlicensure
evaluation of the effectiveness of seven valent pneumococcal
conjugate vaccine. Pediatr Infect Dis J, Dec; 20(12): 1105-7.

Body J.J, (1987) SCC antigen and other tumor markers in lung cancer :
Preliminary results ; Excerpta Medica ; 162-170.

Bracq S, and Machason M (1993). Calcitonin gene expression in
normal human liver. FEBS 1993; 331:14-18.

**Brain SD, Tippins JR, Morris HR, MacIntyre K, and William
TJ(1986).** Potent Vasodilator activity of calcitonin gene-
related peptide in human skin. J Invest Dermatol; 87:533-536.

Browne G,J, Rayan,J,M, and McIntyre, P,(1997). Evaluation of a protocol for selective empirical treatment of fever without localizing signs. Arch dis child; 76:129-33.

Brunkhorst FM, Clark AL, Forycki ZF, and Anker SD(1999).. Pyrexia, procalcitonin, immune activation and survival in cardiogenic shock: the potential importance of bacterial translocation. Int J Card; 72: 3-10.

Brunkhorst FM, Forycki ZF, and WagnerJ(1995). Discrimination of infectious and non-infectious etiologies of the adult respiratory distress syndrome (ARDS) with procalcitonin immunoreactivity. Clin Intens Care; 6:3.

Brunkhorst FM, Heinz U, and Forycki ZF(1998).Kinetics of procalcitonin in iatrogenic sepsis. Intens Care Med; 24:888-892.

Camas P, Leddoux D, Vrinats Y, de Groote D, Franchimont P, and Lamy M. (1992). Cytokine serum level during severe sepsis in human. Il-6 as a marker of severity. Ann Surg; 215:356-362.

Carlet J,(1999). Bacterial sepsis and septic shock: Rapid methods in the detection of sepsis. Infect Dis Clin North Am; Jun 13 (2) 483-494.

Carsin H, Assicot M, Feger F, Roy O, Pennacino I, and Le Bever H, . (1997). Evolution and significance of circulating procalcitonin

levels compared with IL-6, TNFa and endotoxin levels early after thermal injury. *Burns*; 23:218-224.

Castillo MJ, Scheen AJ, and Lefebvre PJ. (1995). Amylin/islet amyloid polypeptide: Biochemistry, physiology, pathophysiology. *Diabet Metab*; 21:3-25.

Chan YL, Tseng CP, Tsay PK, Chang SS, Chiu TF, and Chen JC; Procalcitonin as a marker of bacterial infection in the emergency department an observational study, *Crit Care* Feb;8(1);R 12-20; 2003.

Chiesa C, Panero A, Rossi N, Stegagno M, De Giusti M, and Osborn JF, (1998). Reliability of procalcitonin concentrations for the diagnosis of sepsis in critically ill neonates. *Clin Infect Dis*; 26:664-672.

Chirouze C, Schuhmacher H, Rabaud C, Gil H, Khayat N, Estavoyer JM, May T, and Hoen B; (2002); Low serum procalcitonin level accurately predicts the absence of bacteremia in adult patients with acute fever ;*Clin. Infect. Des* Jul 15;35 (20:156-61

Crain E I. (1990). Urinary tract infections in febrile infants younger than 8 weeks of age. *Pediatrics*;86:363-7.

Dandona P, Nix D, Wilson MF, Aljada A, Love J, and Assicot M, . (1994). Procalcitonin increase after endotoxin injection in normal subjects. *J Clin Endocrinol Metab*; 79:1605-1608.

Davis TME, Assicot M, Bohuon C, St.John A, Li GQ, and Ahn TK. (1994). Serum Procalcitonin concentrations in acute malaria. Trans R Soc Trop Med Hyg; 88:670-671.

Delevaux I, Andre M, Colombier M, Albuissou F, Myltheuc F, Pegue Rj, Piette Jc, and Aumaitre O;(2003); Can procalcitonin measurement help in differentiating between bacterial infection and other kinds of inflammatory processes; Ann Rheum Dis Apr; 62 (4):337-40.

Delevaux I, Andre M, and Colombier M.(2003).Can procalcitonin measurement help in differentiating between bacterial infection and other kinds of inflammatory processes, Ann rheum Dis, Apr. 62 (40):337-40.

Domenech VS, Nylén ES, White JC, Snider RH, Becker KL, and Landmann R, (2001). Calcitonin gene-related peptide expression in sepsis: postulation of microbial infection-specific response elements within the calcitonin I gene promoter. J Invest Med 2001;49:514-521.

Enguix A, Rey C, Concha A, Medina A, Coto D, and Dieguez MA;(2001) Comparison of procalcitonin with C-Reactive protein and serum amyloid for early diagnosis of bacterial sepsis in clinically ill neonates and children. Intensive care Med Jan; 27(1) ;211-5.

Ferriere F (1999) : Procalcitonin , a new marker for bacterial infections. Crit Care Med; Oct 27 (10) : 2172-6.

Fleisher GR, and Bass JW. (1993). Paractice guideline for the management of infants and children 1-36 monthes of age wuth fever without source , Pediatrics; 92:1.

Fleisher GR, Rosenberg N, and Vinci R,. (1994). Intramuscular versus oral antibiotic therapy for the prevention of meningitis and other bacterial sequelae in young, febrile children at risk for occult bacteremia. J Pediatr;124:504-512.

Fleischhack G, Cipic D, Juettner J, Hasan C and, Bode U. (2000). Procalcitonin - a sensitive inflammation marker of febrile episodes in neutropenic children with cancer. Intens Care Med; suppl. 2:202-211.

Fleischhack G, Cipic D, Kambeck I, Ngampolo D, Hasan C, and Bode U. (1997). Procalcitonin - A sensitive marker of severe infections in neutropenic patients. (Abstract) 3rd Int Symp on Febrile Neutropenia Brussels, Dec 10-13.

Galetto-LacourA, and Zamora SA, (2003) . Procalcitonin and C-reactive protein in children with fever without localizing signs of infection seen in referral center. Pediatrics: Nov; 112(5):1054-60.

Gendrel D , Raymond J , Assicot M , Molin F , Bergert M, Badoual J and Bohoun C (1995): Procalcitonin in bacterial and viral meningitis in children . (Abstract).ICAAC 20 Sept.

Gendrel D, Assicot M, Raymond J, Moulin F, Francoual C, Badoual J and Bohoun C (1996) : Procalcitonin as a marker for the early diagnosis of neonatal infection . Journal of pediatrics ; Apr; 128(4): 570-573.

Gendrel D, and Bohoun C.A (2000). Procalcitonin in pediatrics for differentiation of bacterial and viral infections . Intensive care Med. Vol. 26: 178-181.2000.

Gerard Y, Hober D, Assicot M, Alfandari S, Ajana F and, Bourez JM, . (1997). Procalcitonin as a marker of bacterial sepsis in patients infected with HIV-1. J Infect; 35:41-46.

Giebink GS: (2001). The prevention of pneumococcal disease in children.. N Engl J Med ., Oct 18; 345(16): 1177-83.

Grace M, Lee MP, and Marvin B. (1998). risk of bacteremia for febrile young children in post haemophilus influenza type b Era, Arch pediatric Adolesc med.; 152: 624-628.

Gramm HJ, Dollinger P, and Beier W. (1995). Procalcitonin - ein neuer Marker der inflammatorischen Wirtsantwort. Longitudinalstudien bei Patienten mit Sepsis und Peritonitis. Chir Gastroenterol; 11, suppl. 2:51-54.

Guibourdnche J, Bedu A and Petzald L (2002).Biochemical marker in neonatal sepsis, value of procalcitonin in the emergency sitting. Ann Clin Biochem. Mar; 39(pt2): 130-5.

Guven H, Alintop L, and Baydin A , (2002). Diagnostic value of procalcitonin as an early indicator of sepsis, Am J Emerg Med 20: 202-206

Hack CE, De Groote ER, and Felt-Bersma RJF, (1989). Increased plasma levels of interleukin-6 in sepsis. Blood; 74:1704-1710.

Ham J, Ellison ML and, Lymoden J. (1980). Tumor calcitonin. Interaction with specific calcitonin receptors. Biochem J; 190:545-550.

Harper MB, and Fleisher GR (1993): Occult bacteremia in the 3 month old to 3 years old age group. Pediatr Ann Aug; 22(8) : 484, 487-93.

Hathrill M, Tibby SM, Turner C and Murdoch I A , (1999). Diagnostic markers of infection: comparision of procalcitonin with C reactive protein and leucocyte count. Arch dis child ; 81: 471- 421.

Hensel M, Volk T, Döcke WD, Kern F, Tschirna D, and Egerer K,. (1998). Hyperprocalcitoninemia in patients with noninfectious SIRS and pulmonary dysfunction associated with cardiopulmonary bypass. Anesthesiology; 89:93-104 .

Hergert M, Lestin HG, Scherkus M, Brinker K, Klett I, and Stranz G, I. (1998). Procalcitonin in patients with sepsis and polytrauma. Clin Lab; 44:659-670.

Herr SM, Wald ER, Pitetti RD, and Choi SS: (2001). Enhanced urinalysis improves identification of febrile infants ages 60

days and younger at low risk for serious bacterial illness.
Pediatrics Oct; 108(4): 866-71.

Hoberman A, and J Pediatr . (1993). Prevalence of urinary tract infection in febrile infants.;123:17-23.

Housfater P, Grric C, and Benayed S,(2002). Usefullness of procalcitonin as a marker of bacteremia ; Clin. Infect Dis 34:895-901.

Isaacman DJ, Zhang Y, Reynolds EA, and Ehrlich GD: (1998). Accuracy of a polymerase chain reaction-based assay for detection of pneumococcal bacteremia in children. Pediatrics May; 101(5): 813-6.

Jacobs JW, Lund PK, Potts JT, Bell NH, and Habener JF: (1981). Procalcitonin is a glycoprotein. J Biol Chem; 256:2803-2807.

Jaffe DM, and Fleisher GR: (1991). Temperature and total white blood cell count as indicators of bacteremia. Pediatrics;87:670-674.

Jaskiewicz JA and, McCarthy CA: (1993). Evaluation and management of the febrile infant 60 days of age or younger. Pediatr Ann Aug; 22(8): 477-80, 482-3.

Jilma B, Blann A, Pernerstorfer T, and Stohlowetz P,: (1999). Regulation of adhesion molecules during human endotoxamia , Amer J. Resp. Crit. Care Med., 159, 857-863.

Joyce CD. CGRP; (1990). levels are elevated in patients with sepsis. Surgery; (108)1097-1101.

Kaplan SL, Mason EO Jr, and Wald E, : (2002). Six year multicenter surveillance of invasive pneumococcal infections in children. Pediatr Infect Dis J, Feb; 21(2): 141-7.

Klein G.C., (1976). Manual of clinical immunology, Amer . Soc. Microbial . 264.

Kramer MS, and Shapiro ED: (1997). Management of the young febrile child: a commentary on recent practice guidelines. Pediatrics , Jul; 100(1): 128-34.

Kuppersmann N: (1999). Occult bacteremia in young febrile children. Pediatr Clin North Am ,Dec; 46(6): 1073-109.

Kuppersman N, Malley R, Inkelis SH, and Fleisher GR: (1999). Clinical and hematologic features do not reliably identify children with unsuspected meningococcal disease. Pediatrics; 103: E20.

Kuse ER, Langefeld I, Jaeger K, and K  lpmann WR;(2000). Procalcitonin - a new diagnostic tool in complications following liver transplantation. Intens Care Med; suppl. 2:187-192.

Lacour AG, Gervais A, and Zamora SA, ; (2001). Procalcitonin, IL-6, IL-8, IL-1 receptor antagonist and C-reactive protein as identifiers of serious bacterial infections in children with

fever without localising signs. Eur J Pediatr , Feb; 160(2): 95-100.

Lee GM, Fleisher GR, and Harper MB: (2001). Management of febrile children in the age of the conjugate pneumococcal vaccine: a cost-effectiveness analysis. Pediatrics , Oct; 108(4): 835-44.

Lee GM and, Harper MB: (1998). Risk of bacteremia for febrile young children in the post-Haemophilus influenzae type b era. Arch Pediatr Adolesc Med , Jul; 152(7): 624-8.

Lestin F, Lestin HG, Burstein O, Anders O, and Freund M:(1998). Vorläufige Erfahrungen mit Procalcitonin, C-reaktivem Protein, Neopterin, ausgewählten Zytokinen und Hmostaseparametern an Patienten mit Hrsg, O Anders, J Jacob Weller-Verlag, Neckargemünd Malignen hämatologischen Erkrankungen, bei zytostatikainduzierter Neutropenie und Fieber. Hmostase und Entzündung; 1:40-52.

Levi M, and Tencate H, N . Eng. (1999). Dissiminated intravascular coagulation . N . Eng, J. Med; 341: 586-92.

Lewis S,M, Bain B,G and Bates,I, (2001). Practical haematology Department of hematology Hammersmith Hospital, Imperial College School of Medicine, London, UK.

Lietzmann A. Untersuchungen zum Syntheseort und zur Induktion des neuen. (1999). Infektionsparameters Procalcitonin. Dissertation, FAU-Erlangen-Nürnberg.

Marty C, Misset B, Tamion F, Fitting C, Carlet J, and Cavaillon JM. (1994). Circulating interleukin-8 concentrations in patients with multiple organ failure of septic and nonseptic origin. Crit Care Med; 22:673-679.

Meisner M, Lohs T, Hüttemann E, and Reinhart K. (1999). Elimination of procalcitonin and plasma levels during continuous veno-venous hemofiltration in patients with acute renal failure and sepsis. Intens Care Med; 25 Suppl. 15:76.

Meisner M, Lohs T, Hüttemann E, Schmidt J, and Reinhart K. (1999). The plasma elimination rate and urinary secretion of PCT in patients with normal and impaired renal function. Anesthesiology; 91 Suppl. 3A:A236.

Meisner M, Rauschmayer C, and Schmidt J. (1999). Procalcitonin indicates increased risk after cardiovascular surgery. Shock; 12 Suppl.:16-17.

Meisner M Schmidt J, Huettner H, and Tschaikowsky K. (2000). The natural elimination rate of procalcitonin in patients with normal and impaired renal function. Intens Care Med; 26, suppl. 2:212-216.

Meisner M, Tschaikowsky K, Palmaers T, Schmidt J, Mangold G, and Schüttler J. (1997). Comparison of procalcitonin (PCT) and C-reactive protein (CRP) plasma concentrations at different APACHE II scores during the course of sepsis and MODS. Anaesthesiology; 87:243.

Meisner M, Tschaikowsky K, Schnabel S, Schmidt J, Katalinic A, and Schüttler J. (1997). Procalcitonin - Influence of temperature, storage, anticoagulation and arterial or venous asservation of blood samples on procalcitonin concentrations. Eur J Clin Chem Clin Biochem; 35(8):597-601.

Merk H. Beers and Robert Berkow. (1997). Manual of Medical Information – Home Edition, edited by Mark H. Beers and Robert Berkow. Copyright by Merck & Co., Inc., Whitehouse .

Miesner M, Berlin: (1996). Brhams Procalcitonin . A new innovative infection parameter . Biochemical and clinical aspects . Berlin: Brhams Diagnostica IS BN 3-00-000803-9.

Mimoz O, Benoist JF, Edouard AR, Assicot M, Bohuon C, and Samii K. (1998). Procalcitonin and C-reactive protein during the early posttraumatic systemic inflammatory response syndrome. Intens Care Med; 24:185-188.

Monneret G, Labaune JM, Isaac C, Bienvenu F, Putet G, and Bienvenu J. (1997). Procalcitonin and C-reactive protein levels in neonatal infections. Acta Paediatr; 86:209-212.

Morello JA and, Leitch C, Nitz, . (1994). Detection of bacteremia by Difco ESP blood culture system. J Clin Microbiol 1994;32:811-818.

Moulin F, Raymond J, Lorrot M, Mark E, Cost J, Iniguez, Kalifa G,Bohuon C, and Gendrel D (2001); Procalcitonin in children

admitted to hospital with community acquired pneumonia.;
Arch Dis. Child Apr; 84 (4):332-6.

Muller B, White JC, Nylen ES, Snider RH, Becker KL and, Habener JF. (2001) Ubiquitous expression of the calcitonin-I gene in multiple tissues in response to sepsis. J Clin Endocrinol Metab ;86:396-404.

Nadine B ,Claire S,MD;Dominique Gendrel(1998):
procalcitonin is a Marker of severity of Renal lesions in
Pyelonephritis - Pediatrics vol.102 No.6 December , PP: 1422-1424

Nylen E, Jeng J, Jordan MH, Snider R, Thompson K, and Lewis MS, . (1995). Late pulmonary sequela following burns: persistence of hyperprocalcitoninemia using a 1-57 amino acid N-terminal flanking peptide assay. Respir Med; 89:41-46.

Nylen E, Muller B, Snider R, Vath S, Wagner K, White J, Zulewsk H, Vannier E, Habener J, and Becker K. (1999). Pathophysiological significance of calcitonin precursors in sepsis and systemic inflammation. Shock; 12 Suppl.:14.

Nylen E, O'Neill WJ, Jordan MH, Snider R, Moore CF, and Lewis MS, . (1992). Serum Procalcitonin as an index of inhalation injury in burns. Horm Metab Res; 24:439-442.

Nylen E, Snider R, Thompson KA, Rohatgi P, and Becker KL. (1996). Pneumonitis-associated hyperprocalcitoninemia. Am J Med Sci; 312:12-18.

Oberhoffer M, Bogel D, Meier-Hellmann A, Vogelsang H, and Reinhart K. (1996). Procalcitonin is higher in non-survivors during the clinical course of sepsis, severe sepsis and septic shock. *Intens Care Med*; 22:A245.

Oberhoffer M, Bitterlich A, Hentschel T, Meier-Hellmann A, Vogelsang H, and Reinhart K. (1996). Procalcitonin (ProCT) correlates better with the ACCP/SCCM consensus conference definitions than other specific markers of the inflammatory response. *Clin Intens Care*; 7, suppl.1:46.

Oberhoffer M, Karzai W, Meier-Hellmann A, Bogel D, Fassbinder J, and Reinhart K. (1999). Sensitivity and specificity of various markers of inflammation for the prediction of TNF-a and IL-6 in patients with sepsis. *Crit Care Med*; 27(9):1814-1818.

Oberhoffer M, Stonans I, Russwurm S, Stonane E, Vogelsang H, Junker U, Jger L, and Reinhart K. (1999). Procalcitonin expression in human peripheral blood mononuclear cells and its modulation by lipopolysaccharides and sepsis related cytokines in vitro. *J Lab Clin Med*; 134:49-55.

Oberhoffer M, Vogelsang H, Jger L, and Reinhart K. (1999). Katalcalcin and calcitonin immunoreactivity in different types of leukocytes indicates intracellular procalcitonin content. *J Crit Care*; 14:29-33.

Oczenski W , Fitzgerald RD and Scarz S (1998) : Procalcitonin : A new parameter for the diagnosis of bacterial infection in the peri-operative period. *Europ J anesth*; 15: 202-209.

O'Neill WJ, Jordan MH, Lewis MS, Snider R, Moore CF, and Becker KL. (1992). Serum calcitonin may be a marker for inhalation injury in burns. *J Burn Care Rehabil*; 13:605-616.

Pannen B, Robotham J, and Teppo A. (1995). The acute phase response. *New Horiz*; 3:183-197.

Paresh D , David N and , Michael F (1994) :Procalcitonin increase after endotoxin injection in normal subjects – *Journal of clinical endocrinology and metabolism* P: 1605-1608

Patrono C. and Peskar, B.A. (1987). (eds) Radioimmunoassay in Basic and clinical pharmacology. Heidelberg, Springer-Verlag,.

Petitjean S, and Assicot M. (1993). Etude de l'immunoreactivite calcitonine-like au cours des processus infectieux. Diplome d'etudes approfondies de biotechnologie,; Université Paris V:1-29.

Prat C Daninguez J, and Rodrigo C, (2003):Elevated serum procalcitonin values correlate with renal scarring in children with urinary tract infection; *Pediatr. Infect. Dis* May; 22 (5):438-42.

Prat C, Dominguez J, Rodrigo C, Gimenez O, Gali N and Ausina V (2003); Procalcitonin , C-Reactive protein and leukocytic

count in children with lower urinary tract infection ; *Pediatr Infect Dis J* Nov. 22 (11)963-8,.

Pulliam PN, Attia MW, and Cronan KM, : (2001). C-reactive protein in febrile children 1 to 36 months of age with clinically undetectable serious bacterial infection. *Pediatrics* Dec; 108(6):1275-9.

Rau B, Steinbach G, Baumgart K, Gansauge F, Grünert A, and Beger HG. (2000). The clinical value of procalcitonin in the prediction of infected necrosis in acute pancreatitis. *Intens Care Med*; 26, suppl: 2:159-164.

Rau B, Steinbach G, Gansauge F, Mayer M, Grünert A, and Beger HG. (1997). The role of procalcitonin and interleukin-8 in the prediction of infected necrosis in acute pancreatitis. *Gut* 1997; 41:832-840.

Reinhart K, Wiegand-Lohnert C, Grimminger F, Kaul M, Withington S, and Treacher D, et al. (1996). Assessment of the safety and efficacy of the monoclonal anti-tumor necrosis factor antibody-fragment, MAK 195F, in patients with sepsis and septic shock: a multicenter, randomized, placebo-controlled, dose-ranging study. *Crit Care Med*; 24:733-742.

Reith HB , Mittelkotter U , Debus ES , Kussner C and Thiede A (2000): Procalcitonin in early detection of post operative complications . *Ann Biol Clin (Paris)*; Jan – feb 58(1) : 49-59.

Reith HB, Lehmkuhl P, Beier W, and Högl B. (1995). Procalcitonin
- ein prognostischer Infektionsparameter bei der Peritonitis.
Chir Gastroenterol 1995; 11 suppl 2:47-50.

Reith HB, Mittelkötter U, Debus ES, Lang J, and Thiede A. (1997).
Procalcitonin (PCT) immunreactivity in critical ill patients on a
surgical ICU. The Immune Consequences of Trauma, Shock
and Sepsis, Edit. Monduzzi Editore, Bologna; 1:673-677.

**Rosenfeld MG, Mermod JJ, Amara SG, Swanson LW, Sawchenko
PE, and Rivier J, . (1983).** Production of a novel neuropeptide
encoded by the calcitonin gene via tissue-specific RNA
processing. Nature; 304:129-135.

Rothrock SG, Harper MB, and Green SM, I. (1997). Do oral
antibiotics prevent meningitis and serious bacterial infections in
children with Streptococcus pneumoniae occult bacteremia? A
meta-analysis. Pediatrics; 99:438-444.

**Russwurm S, Wiederholt M, Oberhoffer M, Stonans I, Peiker G, and
Reinhart, K. (1999).** Procalcitonin als monozytrter Marker für
die Frühdiagnostik bei septischem Abort. Z Geburtsh Neonatol
1999; 203:1-4.

Staeher M, berfuhr P, Reichart Band, Hammer c (1997).

Differentialdiagnostik der Abstossungsreaktion und Infektion bei
herztransplantierten Patienten: neue Wege mit Zytokinen und
Procalcitonin als Marker. Transplantationsmedizin; 9:44-50 .

Steinhart JM . (1988). Ultrasound screening of healthy infants for urinary tract abnormalities. *Pediatrics*;82:609-614.

Svoldi M, Hirber J, and Lanthaler A,(2001). Procalcitonin in neutropenic and immunocompromised patients , *Brit J Haematol.* 115:53-57.

Teach SJ, and Fleisher GR. (1995). Efficacy of an observation scale in detecting bacteremia in febrile children three to thirty-six months of age, treated as outpatients. *J Pediatr*;126:877-881.

Vadheim CM, Greenberg DP, and Eriksen E, . (1994). Eradication of *Haemophilus influenzae* type b disease in Southern California. *Arch Pediatr Adolesc Med*;148:51-56.

Vincent.J.L (2000): Procalcitonin the marker of sepsis. *Critical care medicin* ; Apr; 28 (4) 1226-8

Wack RP, Demers DM, Bass JW: (1994). Immature neutrophils in the peripheral blood smear of children with viral infections. *Pediatr Infect Dis J* , Mar; 13(3): 228-30[Medline].

Walters M.C.and Abelson H T (1996) ; Interpretations of the complete blood count. *Ped Clin north Am*; Jun; 43 (3) : 599-622.

Wanner GA., Keel M. ,and Steckholzer U. (2000). Relationship between procalcitonin plasma levels and severity of injury , sepsis , organ failure , and mortality in injured patients . *Crit Car Med* . ; 28: 950 – 957.

Whang KT, Steinwald PM, White JC, Nylen ES, Snider RH, and Simon GL, (1998). Serum calcitonin precursors in sepsis and systemic inflammation. J Clin Endocrinol Metab; 83:3296-3301.

Whitcher J, Bienvenu J and Monnert G;(2001) Procalcitonin as an acute phase marker ;Review Article.Ann Clin Biochem:38:483-493.

Wildling E, Pusch F, Aichelburg A, Zimpfer M and., Weinstabl C, (1997). Procalcitonin is elevated in patients after severe injury. Intens Care Med; 23:S62.

Wrenger S, Kahne T, Bohuon C, Weglöhner W, Ansorge S, and Reinhold D. (2000). Aminoterminal truncation of procalcitonin, a marker for systemic bacterial infections, by dipeptidyl peptidase IV CDP IV. FEBS Lett, 466(1): 155-159.

Zwadyk P, Pierson CL, and Young C. (1994). Comparison of Difco ESP and Organon Teknika BacT/Alert continuous monitoring blood culture systems. J Clin Microbiol;32:1273-1279.