




References

- Ackerman, S.; Loegering, D.; Venge, P.; Olsson, I.; Harley, J.; Fauci, A. and Gleich, G. (1983):* Distinctive cationic proteins of the human eosinophil granule: Major basic protein, eosinophilic cationic protein and eosinophil-derived neurotoxin. *J. Immunol.*, 131: 2977.
- Adelroth, E.; Rosenhall, L.; Johansson, S.; Linden, N.; Venge, P. (1990):* I inflammatory cells and eosinophilic activity in asthmatics investigated by broncho alveolar lavage: the effects of antiasthmatic treatment with budenoside or terbutaline. *Am. Rev Respir. Dis.*, 142: 91-9.
- Ahlstedt, S.; Enander, I.; Peterson, O.; Lanner, A. (1993):* The clinical assessment of the inflammatory component in asthma with emphasis on the eosinophils. *Pract. Allergy Immunol*; 8: 149- 54.
- Anderson, H.; Bland, J.; Patek, S. and Peckham, C. (1986):* The natural history of asthma in childhood. *J. Epid. Comm. Hlth.*, 40:121.
- Arima, M. and Kanoh, T. (1999):* Eosinophilic myocarditis is associated with dense deposits of eosinophilic cationic protein (ECP) in endomyocardium of patients with high serum ECP. *Heart*, 81 (6): 669.
- Ayers, J. (1990):* Late onset asthma. *Br. Med. J.*, 300: 1602.

- Azzawi, M.; Bradley, B. and Jeffery, P. (1990):* Identification of activated T lymphocytes and eosinophils in bronchial biopsies in stable atopic asthma. *Am. Rev. Respir. Dis.*, 142: 1407.
- Backman, K.; Jarvinen, P. and Sorva, R. (1993):* Serum eosinophilic cationic protein during treatment of asthma in children. *J. allergy Clin. Immunol.*, 92: 34.
- Badr- El- Din, I.; El- Sawy, O.; El- Azzouni, M.; Badr- El- Din and A. Salem (1999):* eosinophil cationic protein as a serological marker of disease activity in childhood bronchial asthma. *Eastern Mediterranean health journal*, 5, (4), 664- 675.
- Barnes, P. (1986):* Asthma as an axon reflex. *Lancet*, I: 242.
- Beasley, R.; Roche, W.; Roberts, A. and Holgate, S. (1989):* Cellular events in the bronchi in mild asthma and after bronchial provocation. *Am. Rev. Respir. Dis.*, 139: 806.
- Benvensite, J.; Henson, P. and Cochrane, C. (1972):* Leucocyte- dependant histamine release from rabbit platelets: The role of IgE, basophils and platelet-activating factor.
- Boix, E.; Leonidas, D.; Nikolovski, Z.; Nogues, M.; Cuchillo, C. and Acharya, K. (1999):* Crystal structure of eosinophilic cationic protein at 2.4° A resolution. *Biochemistry*, 38: 16794.

References

- Bousquet, J. ; Chanaz, P. ; lacoste J. (1990):* Eosinophilic inflammation in asthma. N engl J. Med ; 323 : 1033-9.
- Bradding, P.; Feather, I. and Howarth, P. (1992):* Interleukin 4 is localized to and released by human mast cells. J. Exp. Med.; 176:1381.
- Bromley, I. and Donaldson, K. (1996):* Binding of *Aspergillus fumigatus* spores to lung epithelial cells and basement membrane proteins: Relevance to the asthmatic lung. Thorax, 51: 1203.
- Brown, R.; Zerhouni, E. and Mitzne, W. (1995):* Airway oedema potentiates airway reactivity. J. Appl. Physiol., 79: 1242.
- Busse, W. (1998):* Current research and future needs in allergic rhinitis and asthma. J. Allergy Clin. Immunol., 101: S424.
- Busse, W. and Parry, D. (1998):* The biology of asthma. In: Fishman's Pulmonary Diseases and Disorders, 3rd edition, Vol. 1. Alfred P. Fishman, Jack A. Elias, Jay A. Fishman, Michael A. Grippi, Larry R. Kaiser and Robert M. Senior, Eds. New York, Section 9, Chapter 47, P. 721.
- Carl, A.; Burtis and Edward R. Ashwood. (2001):* Teitz fundamentals of clinical chemistry, fifth edition volume 1, chapter 10 p. 191.

- Carlson, M.; Hakansson, L.; Peterson, C.; Stalenheim, G. and Venge, P. (1991):* Secretion of granule proteins from eosinophils and neutrophils is increased in asthma. *J. Allergy Clin. Immunol.*, 87: 27.
- Chan- Yeung, M. and Malo, J. (1994):* Aetiological agents in occupational asthma. *Eur. Respir. J.*, 7: 346.
- Cheng, J.; Ott, N.; Peterson, E.; George, T.; Hukee, M.; Gleich, G. and Leiferman, K. (1997):* Dermal eosinophils in atopic dermatitis undergo cytolytic degeneration. *J. Allergy Clin. Immunol.*, 99:683.
- Coyle, A.; Ackerman, S. and Irvin, C. (1993):* Cationic proteins induce airway hyper- responsiveness dependant on charge interactions. *Am. Rev. Respir. Dis.*, 147: 896.
- Coyle, A.; Uchida, D.; Ackerman, S.; Mitzner, W. and Irvin, C. (1994):* Role of cationic proteins in the airway: Hyper-responsiveness due to airway inflammation. *Am. J. Respir. Crit. Care Med.*, 150: S63.
- Cromwell, O.; Hamid, Q. and Corrigan, C. (1992):* Expression and generation of IL-8, IL-6 and granulocyte- macrophage colony- stimulating factor by bronchial epithelial cells and enhancement by- IL-1 beta and tumour necrosis factor- alpha. *Immunology*, 77:330.

- Dempsey, O; Kennedy, G. and Lipworth, B. (2002):*
Comparative efficacy and anti- inflammatory
profile of once- daily therapy with leukotriene
antagonist of low- dose inhaled corticosteroid in
patients with mild persistent asthma. J. Allergy
Clin. Immunol., 109: 68.
- Djukanovic, R.; Roche, W. and Wilson, J. (1990):* Mucosal
inflammation in asthma. Am. Rev. Respir. Dis.,
142: 434.
- Emphy, D.; Lartimen, L.; Jaccob, S.; Gold, W. and Nodel,
J.A. (1976):* Mechanisms of bronchial hyper-
reactivity in normal subjects after upper respiratory
tract infection. Am. Rev. Respir. Dis., 113: 131.
- Ferguesson, D.; Horwood, L. and Shannon, F. (1980):*
parental smoking and respiratory illness in infancy.
Arch. Dis. Child., 55: 358.
- Ferguson, A.; Vaughan, R.; Brown, H. and Curtis, C.
(1995):* Evaluation of serum eosinophilic cationic
protein as a marker of disease activity in chronic
asthma. J. Allergy Clin. Immunol., 95: 23.
- Fredens, K.; Dahl, R. and Venge, P. (1982):* The Gordon
phenomenon induced by eosinophilic cationic
protein and eosinophil protein X. J. Allergy Clin.
Immunol., 70: 361.

- Frigas, E. and Gleich, G. (1986):* The eosinophil and the pathophysiology of asthma. J. Allergy Clin. Immunol., 77: 527.
- Fuller, R. (1992):* Macrophages. Br. Med. Bull., 48: 65.
- Gauchat, J.; Henchoz, S. and Mazzei, G. (1993):* Induction of human IgE synthesis in B cells by mast cells and basophils. Nature, 365: 340.
- Griffin, E.; Hakansson, L., Formgren, H.; Jorgensen, K.; Peterson, C. and Venge, P. (1991):* Blood eosinophil number and activity in relation to lung function in patients with asthma and with eosinophilia. J. Allergy Clin. Immunol., 87: 548.
- Hamann, K.; Ten, R.; Loegering, D.; Jekins, R.; Heise, M.; Schad, C.; Pease, L.; Gleich, G. and Baker, R. (1990):* Structure and chromosome localization of the human eosinophil- derived neurotoxin and eosinophilic cationic protein genes: Evidence for intronless coding sequences in the ribonuclease gene superfamily. Genomics, 7: 535.
- Haslett, C. (2000):* Asthma: Cellular and humoral mechanisms. In: Crofton and Douglas's Respiratory Diseases, 5th edition, Vol. 3. Anthony Seaton, Douglas Seaton and A. Gordon Leitch, Eds. Oxford, Blackwell Science Ltd., Chapter 33, p. 907.

- Hedman, J.; Molianen, E.; Poussa, T. and Nieminen, M. (1999):* Serum ECP and myeloperoxidase but not leukotriene E₄ are associated with bronchial hyper-responsiveness. *Respir. Med.*, 93 (8): 589.
- Herzog, H. (1982):* Clinics and treatment of asthma. In: *Proceedings of the XIth International Congress of Allergology and Clinical Immunology*. J.W. Kerr and M.A. Ganderton, Eds. The Mac Millan Press, Chapter 12, P. 327.
- Holgate, S. (1996):* The immunopharmacology of mild asthma. *J. Allergy Clin. Immunol.*, 98: S7.
- Ishioka, S.; Hozawa, S.; Haruta, Y.; Hiyama, K.; Maeda, A. and Yamakido, M. (1999):* Pranlukast, a cysteinyl leukotriene antagonist, reduces serum eosinophilic cationic protein levels in patients with asthma. *Hiroshima J. Med. Sci.*, 48: 105. *J. Allergy Clin. Immunol.*, 102: 65.
- James, A.; Pare, P. and Hogg, J. (1989):* The mechanisms of airway narrowing in asthma. *Am. Rev. Respir. Dis.*, 139: 242.
- Jang As, Yem CH, Choi IS. (2003) :* Nitric oxide metabolites, eosinophils, and eosinophilic cationic protein in patients with asthma: sputum versus blood. *J korean med sci.*, 18 (4): 489- 93.

- Joseph- Brown, J.; deklerk, N.; Holt, P.; Slay, P. (2004):*
Relationship of asthma, atopy and brochial
responsiveness to serum eosinophil cationic
proteins in early child hood. J Allergy Clin
Immunol. Nov; 114 (5): 1040-5.
- Kaplan, A.; Baeza, M.; Reddigaria, S. and Kuna, P. (1991):*
Histamine- releasing factors. Int. Arch. Allergy
Appl. Immunol., 94: 148.
- Kapp, A.; Czech, W.; Krutmann, J. and Schopf, E. (1991):*
Eosinophilic cationic protein in sera of patients
with atopic dermatitis. J. Am. Acad. Dermatol., 24:
555.
- Kato, Y.; Fujisawa, T.; Terada, A.; Iguchi, K. and Kamiya,
H. (1999):* Mechanisms of eosinophilic cationic
protein release in the serum. Int. Arch. Allergy
Immunol., 120: 160.
- Kay, A. (1991):* Asthma and inflammation. J. Allergy Clin.
Immunol., 87 (5): 893.
- Kaye, M. and Smith, L. (1990):* Effects of inhaled leukotriene
D4 and platelet- activating factor on airway
reactivity in normal subjects. Rev. Respir. Dis.,
141: 993.
- Konig, P. and Godfrey, S. (1978):* Prevalence of exercise-
induced bronchial lability in families of children
with asthma. Arch. Dis. Child., 48: 513.

References

- Kricka, L. (2001):* Principles of immunochemical techniques.
In: Tietz Fundamentals of Clinical Chemistry, 5th
edition, Vol. 1. Carl A.B. and Edward R.A., Eds.,
W.B. Saunders Company, Chapter 10, P. 177.
- Kwon, O.; Collins, P. and Au, B. (1993):* Glucocorticoid
inhibition of TNF- α induced IL-8 gene expression
in human primary cultured epithelial cells. Am.
Rev. Respir. Dis., 147: A752.
- Laitinen, L.; Heino, M.; Laitinen, A.; Kava, T. and
Haahtela, T. (1985):* Damage of the airway
epithelium and bronchial respiratory tract in
patients with asthma. Am. Rev. Respir. Dis., 131:
599.
- Lechner, S. (1983):* Asthma. In: Fundamentals of internal
Medicine, 4th edition, Vol.3. Kage D. and Rose
F.L., Eds. The C.V. Mosby Company, St. Louis,
Chapter 22, p. 1018.
- Lee, T. (1986):* Biological properties of lipoxygenase
products: an overview. In: Asthma: Clinical
Pharmacology and Therapeutic Progress, 3rd
edition. Kay A.B., Ed. Blackwell Scientific
Publications, Oxford, Chapter 15, p. 490.
- Leung, D. and Geha, R. (1987):* Regulation of the human IgE
antibody response. Intern. Rev. Immunol., 2: 75.

- Lonkvist, K.; Hellman, C.; Ludahl, J.; Hallden, G. and Hedlin G. (2001):* Eosinophil markers in blood, serum and urine for monitoring the clinical course in childhood asthma: Impact of budesonide treatment and Withdrawal. *J. Allergy Clin. Immunol.*, 107: 812.
- Lopez, A.; Sanderson, C.; Gamble, J.; Campbell, H.; Young, I. and Vadas, M. (1988):* Recombinant human IL-5 is a selective activator of human eosinophil function. *J. Exp. Med.*, 167: 219.
- Matsumoto, H.; Niimi, A.; Minakuchi, M.; Izumi, T.; (2001):* Serum eosinophil cationic protein levels measured during exacerbation of asthma: characteristics of patients with low titres. *Clin Exp. Allergy*, Apr; 31 (4): 637- 43.
- Mertens, A.; De- Clerck, L.; Moens, M.; Bridts, C. and Stevens, W. (1993):* Study of eosinophil- endothelial adhesion, production of oxygen radicals and release of eosinophilic cationic protein by peripheral blood eosinophils of patients with rheumatoid arthritis. *Clin. Exp. Allergy*, 23 (10): 868.
- Munroford, R. (1983):* Etiology of asthma: A continuing review. *Ann. Allergy*, 50: 43.
- Naclerio, R. and Baroody, F. (1998):* Understanding the inflammatory processes in upper allergic airway disease and asthma. *J. Allergy Clin. Immunol.*, 101: S345.

- Nagy, L.; Lee, T. and Kay, A. (1982):* Neutrophil chemotactic factor activity in antigen induced late asthmatic reaction. N. Engl. J., 306: 497.
- Panettieri, R.; Lazaar, A.; Pure, E. and Albelda, S. (1995):* Activation of c- AMP dependant pathways in human airway smooth muscle cell inhibits TNF- α induced ICAM-1 and VCAM-1 expression and lymphocyte adhesion. J. Immunol. Mar., 154 (5): 2358.
- Patella, V.; Cresenzo, G.; Marino, I.; Genevase, E.; Adt, M.; Gleich, G. and Morone, G. (1996):* Eosinophil granule proteins activate human heart mast cells. J. Immunol., 157 (3): 1219.
- Pepys, J. (1975):* Bronchial provocation tests in etiologic diagnosis and analysis of asthma. Am. Rev. Respir. Dis., 141: 518.
- Peterson, C. and Venge, P. (1987):* Interaction and complex formation between the eosinophilic cationic protein and α_2 - macroglobulin. Biochem. J., 245:781.
- Peterson, C.; Enander, I.; Nystrand, J.; Anderson, A.; Nilsson, L. and Venge, P. (1991):* Radioimmunoassay of human eosinophilic cationic protein (ECP) by an improved method. Establishment.

- Pin, I.; Gibson, P. and Kolendowicz, R. (1992):* Use of induced sputum cell counts to investigate airway inflammation in asthma. *Thorax*, 47:25.
- Pizzichini, E.; Pizzichini, M.; Efthimiadis, A.; Dolovich, J. and Hargreave, F. (1997):* Measuring airway inflammation in asthma: Eosinophils and eosinophilic cationic protein in induced sputum compared with peripheral blood. *J. Allergy Clin. Immunol.*, 99: 539.
- Pronk-Admiral, C.; Haitjema, T.; Horikx, P. and Bartels, P. (2001):* Surplus Value of eosinophil count and ECP to diagnose and monitor asthmatic patients. *Neth. J. Med.*, 58 (1): 9.
- Rankin, J. (1989):* The contribution of alveolar macrophages to hyper-reactive airway disease. *J. Allergy Clin. Immunol.*, 83: 7222.
- Reimert, C.; Mshinda, H.; Hatz, C.; Koombe, Y.; Nkulila, T.; Pouslen, L.; Christensen, N. and Vennervald, B. (2000):* Quantitative assessment of eosinophiluria in *Schistosoma haematobium* infections: A new marker of infection and bladder morbidity. *Am. J. Trop. Med. Hyg.*, 62 (1): 19.
- Roche, W.; Beasley, R.; Williams, J. and Holgate, S. (1989):* Subepithelial fibrosis in the bronchi of asthmatics. *Lancet*, i: 250.

References

- Rogers, D.; Alton, E.; Aursudkij, B.; Boschetto, P.; Dewar, A. and Barnes, P. (1991):* Effect of platelet-activating factor on formation and composition of airway fluid in the guinea pig trachea. *J. Physiol.*, 43: 643.
- Roquet, A; Hallden, G.; Ihre, E.; Hed, J. and Zatterstrom, O. (1996):* Eosinophil activity markers in peripheral blood have high predictive value for bronchial hyper-reactivity in patients with suspected mild asthma. *Allergy*, 52: 482.
- Rose, B.; Hogg, J. and Macklein, P. (1978):* The pathogenesis of bronchial asthma. In: *Immunological Diseases*, 2nd edition, Vol. 2. M. Smater and others, Eds. Little Brown Company, Chapter 47, p. 852.
- Sanderson, C. (1992):* Interleukin-5, eosinophils and disease. *Blood*, 79:1301.
- Sanjar, S.; Smith, D.; Kings, M. and Morley, J. (1990):* Pretreatment with rh- GMCSF, but not rh- IL3, enhances PAF- induced eosinophil accumulation in guinea pig airways. *Br. J. Pharmacol.*, 100: 399.
- Sano, H.; Nakagwa, N. and Nakajima, H. (1995):* Role of vascular cell adhesion molecule-1 and platelet-activating factor in selective eosinophil migration across vascular endothelial cells. *Int. Arch. Allergy. Immunol.*, 107:533.

References

- Scadding, J. (1976):* Definition and clinical categorization. In: The Bronchial Asthma; Mechanisms and Therapeutics, 2nd edition. E.W. Weiss and M.S. Segal, Eds., Little Brown Company, Chapter 2, p.19.
- Schleimer, R.; Sternbinsky, C. and Kaiser, C. (1992):* Interleukin-4 induces adherence of human eosinophils and basophils but not neutrophils to endothelium: Association with expression of VCAM-1. *J. Immunol.*, 148: 1086.
- Seaton, A. and Crompton, G. (2000):* Asthma: Clinical features. In: Crofton and Douglas's Respiratory Diseases, 5th edition, Vol.3. Anthony Seaton, Douglas Seaton and A. Gordon Leitch, Eds. Oxford, Blackwell Science Ltd., Chapter 34, p. 922.
- Sedgwick, J.; Calhoun, W. and Gleich, G. (1991):* Immediate and late airway responses of allergic rhinitis patients to segmental antigen challenge: Characterization of eosinophil and mast cell mediators. *Am. Rev. Respir. Dis.*, 144: 1274. Serum eosinophilic cationic protein during treatment of asthma in children. *J. Allergy Clin. Immunol.*, 92:34.
- Shalit, M.; Sekhsaria, S.; Li, F.; Mauhorter, S.; Mahanti, S. and Malech, H. (1996):* Early commitment to the eosinophil lineage by cultured human peripheral blood CD34⁺ cells: Messenger RNA analysis. *J. Allergy Clin. Immunol.*, 98:344.

- Simon, H.; Weber, M.; Becker, E.; Zilberman, Y.; Blaser, K. and Schaffer, F. (2000):* Eosinophils maintain their capacity to signal and release eosinophilic cationic protein upon repetitive stimulation with the same agonist. *J. Immunol.*, 165: 4069.
- Sinkiewicz, W.; Romanski, B.; Bartuzi, Z.; Zbikowska, M. and Staszynska, M. (1998):* Eosinophilic activation based on measurements of eosinophilic cationic protein in patients with acute myocardial infarction and patients with angina pectoris. *Przegl. Lek.*, 55 (10): 512.
- Stern, M.; Meagher, L.; Savill, J. and Haslett, C. (1992):* Apoptosis in human eosinophils: Programmed cell death in the eosinophil leads to phagocytosis by macrophages and is modulated by IL-5. *J. Immunol.*, 148: 3543.
- Tai, P.; Spry, C.; Paterson, C.; Venge, p. and Olsson, I. (1984):* Monoclonal antibodies distinguish between storage and secreted forms of eosinophilic cationic protein. *Nature*, 309: 182.
- Tang, R. and Chen, S. (2001):* Serum levels of eosinophilic protein and eosinophils in asthmatic children during a course of prednisolone therapy. *Pediatr. Pulmonol.*, 31 (2): 121.

Terr, A. (2001): The atopic diseases. In: Medical Immunology, 10th edition. Tristram G. Parslow, Daniel P. Stites, Abba I. Terr and John B. Imboden, Eds. New York, Lange Medical Books, Chapter 26, P. 349.

The Egyptian Society of Chest Diseases and Tuberculosis (1998): Egyptian guidelines for asthma management. The Egyptian Journal of Chest Diseases and Tuberculosis, 47 (2): S21.

Tosi, M.; Stark, J.M.; Smith, C.; Hamedani, A.; Gruenert, D. and Infeld, M. (1992): Induction of ICAM-1 expression on human airway epithelial cells by inflammatory cytokines: Effects on neutrophil-epithelial cell adhesion. Am. J. Respir. Cell Mol. Biol., 7:214.

Tosmassini, M. et al. (1996): Serum level of eosinophil cationic protein in allergic diseases and natural allergen exposure. Journal of allergy and clinical immunology, 97 (6): 1350- 5.

Vancheri, C.; Mastruzzo, C.; Armato, F.; Tomaselli, V.; Magri, S.; Pistorio, M.; LaMicela, M.; D'Amico, L. and Crimi, N. (2001): Intranasal heparin reduces eosinophil recruitment after nasal allergen challenge in patients with allergic rhinitis. J. Allergy Clin. Immunol., 108: 703.

- Vatrella, A.; Ponticiello, A.; Parrella, R.; Romano, L.; Zofra, S.; DiLeva, A. and Bariffi, F. (1996):* Serum eosinophilic cationic protein (ECP) as a marker of disease activity and treatment efficacy in seasonal asthma. *Allergy*, 51: 547.
- Venge, P. and Bystrom, J. (1998):* Molecules in focus eosinophilic cationic protein. *Int. J. Biochem. Cell Biol.*, 30: 433.
- Venge, P.; Bergstrand, H. and Hakansson, L. (1996):* Neutrophils and eosinophils. In: *Textbook of Rheumatology*, 5th edition. W.N. Kelley, E.D. Harris, S. Ruddy and C.B. Sledge, Eds. Philadelphia, W.B. Saunders Co., Chapter 3, p. 146.
- Villa, J.; Garcia, G.; Ruede, S.; Nogales, A. (1998) :* Serum eosinophilic cationic protein may predict clinical course of wheezing in young children. *Arch Dis Child*; 78: 448 – 452.
- Weller, P. (1997):* Human eosinophils. *J. Allergy Clin. Immunol.*, 100: 283.
- Woolley, K.; Adelroth, E.; Wolley, M.J.; Ellis, R.; Jordana, M. and O'Byrne, P. (1994):* Granulocyte-macrophage colony- stimulating factor, eosinophils and eosinophilic cationic protein in mild asthmatics and non asthmatics. *Eur. Respir. J.*, 7: 1576.

Yamaguchi, Y.; Hayashi, Y. and Sugama, T. (1988):

Highly purified murine IL-5 stimulates eosinophil function and prolongs its in vitro survival: IL-5 is an eosinophil chemotactic factor. *J. Exp. Med.*, 167: 1737.

Young, J.; Peterson, C.; Venge, P. and Cohn, Z. (1986):

Mechanism of membrane damage mediated by human eosinophilic cationic protein. *Nature*, 321 (6070): 613.

Zakrzewski, J.; Barnes, N.; Piper, P. and Costello, J. (1985):

Measurement of leukotrienes in arterial and venous blood from normal and asthmatic subjects by radioimmunoassay. *Br. J. Clin. Pharmacol.*, 19:574.