

References

References

A

Abdalla, S., J. Vila, and M. T. Jimenez de Anta. (1994): Identification of *Salmonella* spp. with Rambach agar in conjunction with the 4-methylumbelliferyl caprylate (MUCAP) fluorescence test. *Br. J. Biomed. Sci.* 51:5-8.

Abd el- Fattah , M.A.; Hassan , E.M.; Shaheen , H.I; Awad, R.A and Girgis, N.I. (1991) : Evaluation of ELISA for detection of salmonella typhi antigens and antibodies . *J. Egypt. Pub. Health. Assoc.* (1,2) : 145 – 157 .

Abraham , G.; Teklu , B.; Gedebu , M.; Selassie , G.H. and Azene, G. (1981) : Diagnostic value of the Widal test . *Trop . Geogr. Med.* 33 (4) : 329-333 .

Adegbola , R.A., Old ,D.C. and Aleksic,S.(1983):Rare MR/K-like haemagglutinins and type 3-like fimbriae of salmonella strains. *FEMS Microbial .Lett.*, 19: 233 - 238.

Adegbola , R.A and Old , D.C.(1987) : Antigenic relationships among type-1 fimbriae of Enterobacteriaceae revealed by immuno electronmicroscopy . *J.Med. Microbial* .,24 : 21:28 .

Aguirre , P.M; Cacho , J.B. and Folgueira , L.R. (1990) : Rapid fluorescence method for screening salmonella spp . From enteric differential agars . *J. Clin . Microbiol* . 28 (1) 148 – 149 .

Anglaret , X., Sylla-Koro , F., Bonard , D., Combe P., Coulibaly , M., Aoussi , E. and Dabis , f. (1997) : Susceptibilites to Co-trimoxazole of pathogens isolated from blood and stool Specimens in Abidjan , Ivory Coast , 1994 to 1996 *J. Clin . Microbiol.*, 35 : 1907 – 1915

d'Aoust, J.-Y.(1991): Psychrotrophy and food-borne salmonella. *Int. J. Food Microbial.*,13 : 207-216.

Appassakij , H.; Bunchuin , N.; Sarasombath , S. and Rungpitarangsi , B. (1987) : Enzyme – linked immunosorbent assay for detection of *Salmonella typhi* protein antigen . *J. Clin Microbiol.* 25 (2) : 273 – 277.

Arcangiolo, M.A., Leroy- Setrin, S., Martel, J.-L and Chaslus - Dancia, E.(2000): Evolution of chloramphenicol resistance with emergence of cross-resistance to florfenicol in bovine *Salmonella typhimurium* strain simplices definitive phage type (DT) 104 .J. Med. Microbiol. , 49: 103- 110.

Arlet ,G. and Philipon , A.(1991): Construction by polymerase chain reaction and intragenic DNA probes for three main types of transferable β -lactamases (TEM,SHV,CARB).FEMS Microbial .Lett ., 82 : 19-26.

Axel Fey,¹ Stefan Eichler,¹ Sébastien Flavier,² Richard Christen,² Manfred G. Höfle,¹ and Carlos A. Guzmán¹ (2004); Establishment of a Real-Time PCR-Based Approach for Accurate Quantification of Bacterial RNA Targets in Water, Using *Salmonella* as a Model Organism Appl Environ Microbiol. 2004 June; 70(6): 3618–3623

B

Baffone, W., Ciaschini, G., pianetti, A., Brandi, G., Casaroli, A. and Bruscolini, F.(2001): Detection of *Escherichia coli* O157:H7 and other intestinal pathogens in patients with diarrheal disease. Eur.J. Epidemiol., 17 :97 - 99.

Baine, W.B., Gangarosa, E . J. and Bennett, J.V.(1973) : Instructional salmonellosis . J. Infect.Dis., 128 : 357-360 . Cited by : Mandal ,B.K. (1996): *Salmonella* Infections. In:Cook,G. (ed.) *Mansons Tropical Diseases*, 12th ed., Chap. 42,PP : 849-863. Saunders Company, Philadelphia.

Baird-Parker, A.C (1991) : food-borne salmonellosis. Lancet Review of food-Borne Illness,55:53-61.

Barker ,R.M (1985) : Utilization of d-tartaric acid by *Salmonella paratyphi* B and *Salmonella Java* : Comparison of anaerobic plate test and turbidity test . J. Hygiene, 95 : 107-114 .

Barret , T.J.; Blake , P.A.; Brown , S.L. and Hoffman , K. (1983) : Enzyme linked immunosorbent assay for detection of human antibodies to *Salmonella typhi* vi- antigen . J . Clin Microbiol . 17 : 526 – 627 .

- Bauernfeind , A., Stempliger , I., Jungwirth , R., Ernst, S. and Casellas, J.M .(1996) :** Sequences of β -lactamase genes encoding CTX-M-1 (MEN-1) and CTX-M-2 and relationship of their amino acid sequences with those of other β -lactamase. *Antimicrob. Agents Chemother.*, 40 : 509 -513.
- Bhutta ZA(1996):.** Impact of age and drug resistance on mortality in typhoid fever *Archives of Disease in Childhood* 1996;75:214-7.
- Bopp, C.A., Brenner, F.w., wells, J. and Strockbine ,N.A. (1999):** *Escherichia, Shigella and Salmonella*.In: Murray,P.R. ,Baron, E.J., Pfaller,M.A., Tenover, F.C.and Yoolken, R.H.(eds).Manual of Clinical Microbiology, 7th th ed.,PP:459-474.ASM Press, Washington DC.
- Boriraj, V., Bangtrakulnonth , A., pornruengwong ,S. and Saitanu, K.(1997):** Demographic data on *Salmonella enteritidis* infection in Thailand , 1990-1995.*Southeast Asian J. Trop. Med. Public Health*, 28: 774-780 .
- Boyd, J.F. and Hart, D.L(1997):**Recent horizontal transmission of plasmids between natural populations of *Escherichia coli* and *Salmonella enterica*.*J.Bacteriol.*,179:1622-1627.
- Bradford, P.A.,Yang,Y., Sahm, D.,Grobe, Gradovsk, D. and stroch, G.(1998):**CTX-M-5 : A novel cefotaxime-hydrolyzing β -lactamase from an outbreak of *Salmonella typhimurium* in Latvia. *Antimicrob. Agents Chemother* .,42:1980-1984.
- Brenner, F.W.(1998):** Modified Kauffmann-white scheme. Centers for Disease Control and prevention. Atlanta.
- Brenner, F.W., Villar ,R.G., Angulo, F.J., Tauxe, R. and Swaminathan, B.(2000):***Salmonella* Nomenclature.*J.Clin.Microbial.*,38:2465-2567.
- Buck , R.L.; Escamilla, J.; Sangalang R.p. and Cross J. H. (1987) :** Diagnostic value of a single pretreatment Widal test in suspected enteric fever cases in the philippines . *Trans . Roy . Soc . Trop. Medi . Hyg.* 81 : 871 – 873 .
- Buissiere, J., Coynault, C.and Le Minor, L.(1977):** Etude des conditions d expression du caracter raffinose chez les *Escherichia coli* et *salmonella*. *Ann. Microbial.*, 128A:167-183.Cited by: Old D.C. and

Threlfall, E.J. (1998): *Salmonella*. In: Blows, A and Duerden, B.I. (eds.) Topley and Wilson's Microbiology and Microbial Infections, Vol. 2, 9th ed., Chap. 41, pp: 969-997. Arnold, London.

Burnens, A.P., Stanley, J., Sechter, I. And Nicolet, J. (1996): Evolutionary origin of a monophasic *Salmonella* serovar 9,12:1,v:- revealed by IS200 profiles and restriction fragment polymorphisms of the *fli B* gene. J. Clin. Microbiol., 34:1641-1645.

C

Cannon, M., Harford, S. and Davies, J.-A. (1990): Comparative study on the inhibitory actions of chloramphenicol, thiamphenicol and some fluorinated derivatives. J. Antimicrob. Chemother., 26:307-317.

Carlsson, H.E.; Lindberg, A.A. and Hammarstrom, S.H. (1972): Titration of antibodies to *Salmonella* O antigens by enzyme linked immunosorbent assay (ELISA) infect. Immun. 6: 703 - 708.

Carlsson, H.E.; Lindberg, A.A.; Hammarstrom, S.H. and Ljunggren, A. (1975): Quantitation of *Salmonella* O- antibodies in human sera by enzyme-linked immunosorbent assay (ELISA). Int. Arch. Allergy Appl. Immunol. 48: 485 - 494.

Cary Endleberg, N.; Barret, T.J. and Fishesr, H. (1983): Identification of a carrier by using Vi Enzyme linked immunosorbent assay serology in an outbreak of typhoid fever on an Indian reservation. J. Clin. Microbiol. 18: 1320-1322.

Casalino, M. and Comanducci, A. (1984): Stability of plasmid content in *Salmonella* wien in late phases of the epidemic history. Antimicrob Agents Chemother., 25:499-501.

Casin, I., Breuil, J., Brisabois, A., Moury, F., Grimont, F. and Collatz, A. (1999): Multidrug-resistant human and animal *Salmonella* typhimurium isolates in France belong predominantly to a DT104 clone with the chromosome and integron-encoded β -lactamase PSE-1. J. Infect. Dis., 179:1173-1182.

- Chaicumpa , W.; Thin Intra.,W. and Khmith, S.(1988):** Detection with monoclonal antibody of Salmonella typhi antigen 9 in specimens from patients. *J. Clin. Microbiol.* 26:1824-1830.
- Chaicumpa , W.; Ruangkunaporn , Y ; Burr , D. and Echeverria , P. (1992) :** Diagnosis of typhoid fever by detection of Salmonella typhi antigen in urine . *J. clin Microbiol .* 30 : 2513-2518 .
- Change, C-T.,You,c.-y., Shen,H.-C., LiA.-M., Chen, C.-Y., Chou, J.-L. .and Huang ,S.-P.(1999):**Recovery of salmonella by using selenite brilliant green sulfa enrichment brth.*J.Clin.Microbiol.,*37:4120-4123.
- Chart, H., Rowe, B., Threlfall, E.J. and Ward, L.R.(1989):**Conversion of Salmonella enteritidis phage type 4 to phage type 7 involves loss of lipopolysaccharide with concomitant loss of virulence .*FEMS Microbiol.Lett.,*60:37-40.
- Cheesbrough, M.(1991) :** Enteric Gram-negative rods and Gram-negative anaerobes. In cheesbrough M.(ed).*Medical laboratory Manual for Tropical countries.* Butter worth Heinemann ltd., Oxford.P:92-185.
- Cheesbrough, M.(2000):**Microbiological tests. In: Cheesbrough, M.(ed.).*District Laboratory Practise in Tropical countries*,Cap.7,PP:1-266. Cambridge university press,VK.
- Christie , A.B. (ed) (1987) :** Typhoid and paratyphoid fevers . In : infectious Diseases : Epidemiology and clinical practice. 4th edn. Churchill Livingstone , New York . Vol . One . Ch. 5, PP. 100 – 164 .
- Clegg, S. and Swenson, D.L.(1994):** Salmonella fimbriae In: Klemm, P., Press, C.R.C. and Boca- Raton, F.L.(eds.) .*Fimriae. Adhesion, Genetics, Biogenesis and vaccines* ,pp:105-113.Arnold,London. .
- Collee, J.G., Miles, R.S. andwatt, B.(1996):** Test for identification of bacteria. In: Collee, J.G., Fraser, A.G, Marmion, B.P. and Simmons, A.(eds.)*Mackie and MacCarthy practical Medical Microbiology*,Vol.114 thed.,PP:131-149.Churchill Living stone, New York.
- Collins, C.H., Lyne, P.M. and Grange, J.M.(1995) :** Salmonella and Shigella. In :Collins and Lynes *Microbiological Methods* ,7th ed., Chap. 28, pp: 314-325.Butterworth Heinemann, Oxford.

Collis, C.M. and Hall, R.M.(1995): Expression of antibiotic resistance genes in the integrated cassette of integrons. *Antimicrob. Agents Chemother.*, 39:155-162.

Cooavdia , Y.M.; Singh , V.; Bhana , R.H. and Modely , N. (1986) : Comparison of passive hemagglutination test with Widal agglutination test for serological diagnosis of typhoid fever in an endemic area . *J. clin Path .* 39 : 680 – 683 .

Change, C-T., You, C.-Y., Shen, H.-C., Li, A.-M., Chen, C.-Y.,J.-L and Huang, S.-P.(1999): Recovery of salmonella by using selenite brilliant green sulfa enrichment broth.*J.Clin.Microbial.*,37:4120-4123.

Cooke, V. M., R. J. Miles, R. G. Price, and A. C. Richardson. (1999): A novel chromogenic ester agar medium for detection of salmonellae. *Appl. Environ. Microbiol.* 65:807-812.

Crichton, P.B. and Yakubu, D.E.(1989): Immunological and genetic relatedness of type-1 and type 2 fimbriae in salmonellas of serotypes Gallinarum, Pullorum and Typhimuri.*J.Appl.Bacteriol.*,67:283-291.

Crump JA, Youssef FG, Luby SP, Wasfy FG, Rangel JM, Oun SA, Mahoney FJ, (2003): Estimating the incidence of typhoid fever and other febrile illnesses in developing countries. *Emerg Infect Dis* 9: 539–544

Crum NF. Current trends in typhoid Fever(2003): *Curr Gastroenterol Rep.*;5(4):279-86.

D

Daniels, E.M. and Schneerson, R.(1989): Characterization of the *Salmonella paratyphi C Vi* polysaccharide. *Infect.Immunol.*,57:3159-3164.

Doern, G.V., Jones, R.V. and Gerlach, E.H.(1995): Multicenter clinical laboratory evolution of a β -lactamase disk assay employing a novel chromogenic cephalosporin. *J.Cline.Microbiol.*,33:1665-1667.

Duguide, J .P .and old ,D.C.(1980): Adhesive properties of Enterobacteriaceae. In: Behachey, E.H., Chapman, R. and Hall, S.(eds.) *Bacterial*

adherence, Receptors and Recognition, Series B, Vol.6, pp:187-217. Arnold, London.

Duguid, J.P.; Marminon, B.P. and Swain, R.H.A. (1984) : Salmonella: 1. Typhoid and paratyphoid fevers. In: Medical Microbiology: Microbial infections. 13th edn. volume one, Ch. 26, PP. 314-319.

Duguid, J.P.(1985): Antigens of type-1 fimbriae. In: Stewart-Tull, D.E.S., Davies, M., Wiley, J. and sons (eds.) Immunology of the Bacterial Cell Envelope, PP:301-318. Chichester.

Duguid, J.P. and Old, D.C.(1994): Introduction: A historical perspective. In: Klemm, P., Press, C.R.C. and Boca-Raton, F.L. (eds.) Fimbriae Adhesion, Genetics, Biogenesis and Vaccines, PP:1-7. Arnold, London.

Duguid, J.P. Anderson, E.S., Alfredsson, G.A., Barker, R. and Old, D.C.(1975): A new biotyping Scheme for Salmonella typhimurium and its phylogenetic significance J. med. Microbiol., 8: 149-166. Cited by Old, D.C. and Threlfall, E.J.(1998): Salmonella. In: Balows, a. and Duerden, B.I.(eds.) Topley and Wilsons Microbiology and microbial Infections, Vol.2, 9th ed., Chap.41, pp:969-997. Arnold, London.

Duguid, J.P. Anderson, E.S., Alfredsson, G.A., Barker, R. and Old, D.C. and Threlfall, E.J.(1998): Salmonella. In: Balows, a. and Duerden, B.I.(eds.) Topley and Wilsons Microbiology and microbial Infections, Vol.2, 9th ed., Chap.41, pp:969-997. Arnold, London.

Dusch, H., and Altwegg, M.(1993): Comparison of Rambach agar, SM-ID medium and Hektoen enteric agar for primary isolation of non-typhi Salmonellae from stool samples. J. Clin. Microbiol., 31:410-412.

Dusch, H. and Altwegg, M.(1995): Evaluation of five new plating media for isolation of salmonella species. J. Clin. Microbiol., 33:802-804.

E

Edelman, R. and Levine, M.M. (1986) : Summary of an international workshop on typhoid fever. Rev. Infect. Dis. 8(3) : 329 – 249.

Eigner, U., R. Reissbrodt, R. Hammann, and A. M. Fahr. (2001): Evaluation of a new chromogenic medium for the isolation and

presumptive identification of *Salmonella* species from stool specimens. *Eur. J. Clin. Microbiol. Infect. Dis.* **20**:558-565 .

Ewing E.H.(1963): An outline of nomenclature for the family Enterobacteriaceae. *Int. Bull. Bacteriol .Nomen.Taxon.*,13:95-110.Cited by :Janda, J. M.and Abbott,S.L.(1998): The Salmonellae .In :Janda ,J. M. and Abbott, S.L(eds.)The Salmonellae In: Janda J.M. and Abbott, S.L.(eds.)The Enterobacteria,1sted.,Chap.6,PP:80-100.Lippincott-Raven, Philadelphia.

Ewing E.H.(1986):Edwards and Ewings identification of Enterobacteriaceae,4 th ed. Elsevier Science publishing Company, New York.

Ewing, E.H. and Edwards, P.R. (1972): Identification of Enterobacteriac.3rded.,BurgesspublishingCompany,Minneapolis.Cited by: Janda, J.M. and Abbott, S.L.(eds.).The Enterobacteria, 1st ed., Chap.6,PP: 80 -100 .Lippincott, Philadelphia.

F

Fadeel, M. A., J. A. Crump, F. J. Mahoney, I. A. Nakhla, A. M. Mansour, B. Reyad, D. E. Melegi, Y. Sultan, E. D. Mintz, and W. F. Bibb.(2004): Rapid diagnosis of typhoid fever by enzyme-linked immunosorbent assay detection of *Salmonella* serotype Typhi antigens in urine. *Am. J. Trop. Med. Hyg.* **70**:323-328.

Falkow, S. and Mekalanos, J.(1990) : Enteric bacilli and Vibrios. In: Davis ,B.D., Bulbecco, R. and Eisen, D.(eds.).*Microbiology*,4 th ed., PP: 561- 587. Lippincott, Philadelphia.

Fallon D, Ackland G, Andrews N, Frodsham D, Howe S, Howells K, . (2003) A comparison of the performance of commercially available chromogenic agars for the isolation and presumptive identification of organisms from urine. *J Clin. Pathol.* Aug;56(8):608-12.

Farmer III, J.J., McWhorter, A.C., Brenner D.J. and Morris, G.K.(1984): The *Salmonella* Arizona group of enterobacteriaceae : Nomenclature ,classification and reporting. *Clin. Microbiol. Newsletter*, 6 : 63-66.

Farmer III, J.J. and Kelly, M.T.(1991): Enterobacteriaceae :Introduction and identification. In: Murray ,P.R., Baron, E.J., Pfaller,

M.A., Tenover, F.C and Yolken, R.H.(eds.).Manual of Clinical Microbiology, 5th ed., Chap.35,PP:361-383.American Society for Microbiology ,Washington, DC.

FarmerIII, J.J.(1995): Enterobacteriaceae : Introduction and identification .In: Murray, P.R., Baron, E.J., Pfaller, M.A., Tenover, F.C. and Yolken, R.H.(eds.).Manual of Clinical Microbiology,6th ed., pp:438-449.American Society for Microbiology Washington, DC.

Feng, P.C.S. and Hartman , PA. (1982) : Fluorogenic assays for immediate confirmation of *Escherichia coli* . Appl . Environ Microbiol. 43 : 1320- 1329.

Finally ,B.B and Falkow, S.(1988): Virulence factors associated with *Salmonella* species. Microbiol .Sci.,5:324:328.

Forbes, B.A, Sahm, D.F. and Weissfeld, A.S. (1998): Enterobacteriaceae. In: Forbes, B. A., Salm, D.F. and Weissfeld, A.S.(eds.).Bailey and scott s Diagnostic Microbiology,10th ed.,Chap.37,PP:510-526.Mosby,Sta.Louis.

Foster, J.W.(1992):Beyond pH homeostasis: The acid tolerance response of *Salmonellae*. ASM News,58:266-270.

Fricker, C.R.(1987): The isolation of *Salmonella* and *Campylobacters*. J. Apple. Bacteriol.,63:99-116.

Frost, J.A., Threlfall, E.J. and Rowe, B(1995): Antibiotic resistance in salmonellas from humans in England and Wales: The situation in 1994.PHLS Microbiol.Dig.,12:131-133.

Frost, J.A., Ward, L.R. and Rwoe, B.(1989):Acquisition of a drug resistance plasmid converts *Salmonella enteritidis* phage type 4 to phage type 24.Epidemiol.Infect.,103:243-248.

G

Gaillot, O., Camillo, P.D., Berche, P., Courcol, R. and Savage, C. (1999): Comparison of CHROMagar *Salmonella* medium and Hektoen enteric agar for isolation of *Salmonellae* from stool samples. J. Clin. Microbiol., 37:762-765.

- Gallardo, F., Ruiz, J., Marco, F., Towner, K.J. and Vila, J.(1999):** Increase in incidence of resistance to ampicillin, chloramphenicol and trimethoprim in clinical isolates of *Salmonellae* serotype Typhimurium with investigation of molecular epidemiology and mechanisms of resistance *J.Med.Microbiol.*,48:367-374.
- Gam , L.H.; Verdugo – Rodriguez , A.; Puthuncheary , S.D. and Pan, T. (1993)** Detection of antibodies against *S. typhi* outer membrane protein (OMP) preparation in typhoid fever patients . *Asian Pac. J. Allergy Immunol.* 11 : 45-52.
- Gazouli, M., Slidorenko, S.V., Tzelepi, E., Kozlova, N.S., Gladin, D.P. and Tzouvelekis, L.S.(1998):** Plasmid-mediated β -lactamase conferring resistance to cefotaxime in a *Salmonella typhimurium* clone found in St. Petersburg, Russia. *J.Antimicrob.Chemother.*,41:119-121.
- Gilchrist, M.J.R (1995):** Enterobacteriaceae: Opportunistic pathogens and other genera. In : Murray, P.R., Baron, E.J. and Tenover, M.A.(eds.). *Manual of Clinical Microbiology*, 6th ed., PP: 440-450. American Society for Microbiology , Washington, DC.
- Gil-Setas, A., Mazon Ramos, A., Martin - salas, Urtiaga -Dominguez, M. and Inza-Elia, M.E.(2002):** Non -typhoid salmonellosis in a basic health area of Navarra, Spain. *Rev.Esp.Salud.Publica.*,76:49-56.
- Giraud, E., Cloeckert, A., Kerboeuf, D. and Chaslus-Dancla, E.(2000):** Evidence for active efflux as the primary mechanism of resistance to ciprofloxacin in *Salmonella enterica* serovar Typhimurium. *Antimicrob. Agents Chemother.*,44:1223-1228.
- Gliman, R.H.; Terminal, M.; Levine, M.M.; Hernandez-Mendoza, P. and Hornick, R.B.(1975):** Relative efficacy of blood, urine, rectal swabs , bone marrow, and rose-spot cultures for recovery of *S. typhi* in typhoid fever. *Lancet* 1 : 1211-1213.
- Glynn, M.K., Bopp, C., Dewitt, W., Dabney, P., Mokhtar, M. and Angulo, F.J.(1998):** Emergence of multidrug-resistant *salmonella enterica* serotype Typhimurium DT104 Infections in the United states. *N. Engl. J.Med.* ,338:1333-1338.
- Godsey, J.H., Matteo, M.R., Shen, D., Tolman, G. and Gohlke, J.R(1981):** Rapid identification of enterobacteriaceae with microbial enzyme activity profiles *J. Clin.Microbiol.* 13:483-490.

Gomez, H.F and Cleary, T.G.(1997): Salmonella species In :Long, S.S., Pickering .L.K and prober, C.G.(eds.) principles and practice of pediatric Infectious Disease , 1st ed.,Chap.155,pp:923-929.Churchill Livingstone, New York .

Gray, L.D.(1995): Escherichia, salmonella, Shigella and Yersinia In: Murray, P.R.,Baron, E.Jand Pfaller, M.A.(eds.)Manual of Clinical Microbiology,6th ed.,PP:450-456.American society for Microbiology, Washington, DC.

Greene, L.C; Appelbaum, p.C. andKellogg, J.A. (1984) : Evaluation of a two hour method for screening pathogens from stool specimens. J. Clin . Microbiol . 20 (2) 285-287.

Greene, S.D.R and Cheesbrought, J.S.(1993): Salmonella bacteremia among young children at a rural hospital in Western Zaire .Ann.Trop.Paediatr.,13:45-53.

Gruenewald, R., R. W. Henderson, and S. Yappow. 1991. Use of Rambach propylene glycol containing agar for identification of Salmonella spp. J. Clin. Microbiol. 29:2354-2356.

Gudmundsdottir S, Hardardottir H, Gunnarsson E. (2003): Subtyping of Salmonella enterica serovar typhimurium outbreak strains isolated from humans and animals in Iceland. J Clin Microbiol. Oct;41(10):4833-5

Gupta , A.K. and Rao, K.M. (1979) : Simultaneous detection of S. typhi antigen and antibody in serum by counter- immunoelectrophoresis for an early and rapid diagnosis of typhoid fever . J. Immunol . Meth. 30 : 349-353 .

Gupta , A.K. and Rao , K.M. (1981) : A simple bactericidal antibody test for serodiagnosis of typhoid fever . J. Immunol . Meth. 42 : 121- 125

H

Hamer, D.h and gorbach, S.L.(1998): Infectious diarrhea and bacterial food poisoning. In : Feldman ,M., Sleisenger and Fordtran,s Gastrointestinal and liver disease, Pathophysiology, Diagnosis and

Management, Vol. 2 , 6th ed., Chap.95, pp; 1594-1632 . Saunders Company, Philadelphia.

Hargrett-Bean, N.t., Pavia, A.T. and tauxe, R.V.(1988): Salmonella isolates from humans in United states,1984-1986.MMWR,37:25-31.

Hashimoto , Y.; Itho , Y.; fujinaga , Y.; Khan , AQ.; and Ezoki, T. (1995): Development of nested PCR based on the ViaB Sequence to detect Salmonella typhi . J. Clin . Microbiol . 33 : 775 – 777

Helmuth, R. and Stephen, R. (1985): Epidemiology of virulence associated plasmids and outer membrane protein patterns with seven common Salmonella serovars Infect .Immunol., 48:175-182.

Hengstler, A., H. Rainer, and A. Marie (1997): Evaluation of BBL CHROM agar Orientation Medium for detection and Presumptive Identification of Urinary Tract pathogens J. Clin. Microbial 35:2773 - 2777.

Hindler, J.a., Howard, B.J., and Keiser, J.F.(1994): Antimicrobial agents and antimicrobial susceptibility testing. In: Howard ,B.J., keiser, J.F., weissfeld, A.S., smith ,T.F and Tilton ,R.C.(eds.)Clinical and pathogenic Microbiology ,2nd ed., Chap.9 pP:145-195.Mosby,St.Louis

Hinder, J.A., Jorgensen, J.H.,(2000)concepts in Antimicrobial susceptibility Testing. In Moham,C.R., Monuselies .G.,(eds.).Diagnostic Microbiology. Chap: 3.,P:59 - 98..

Hirschl , A., Stanek , G.; Rotter , M.; Niemetez , A.H. and Diridl , G. (1983) : Comparison of ELISA (LPS)and Widal reaction (O- antigen) in diagnsosis of Salmonella infections . Zentrable . Barkterial . Mikrobiol . Hyg. I .Abt . Orig . A. 255 : 247 (Abstract)

Hooper, d.c.(1999): Mechanisms of fluorquinolone resistance .Drug Resist. Updates , 2; 38-55.

Hoorfar, J., D. L. Baggesen, and P. H. Porting. 1999. A PCR-based strategy for simple and rapid identification of rough presumptive Salmonella isolates. J. Microbiol. Methods 35:77-8

Hornick , R.B. (ed) (1989) : Typhoid fever . In : Infectious Diseases . 4th edn. J. B. Lippincott company , Philadelphia Ch. 72, PP . 717 – 724 .

Hughes, J.M. and gangarosa, E.J.(1986); Hospital food services ; Role in prevention of nosocomial food - borne disease.In: Bennett , J.V. and Brachman, P.S.(eds.)Hospital Infections , 2nd ed., PP : 257-267. little, Brown , and Company, Boston.

I

Inoue , K.; Mik, K.; Tamura , K. and Sakazaki , R. (1996) : Evaluation of L-pyrrolidonly peptidase paper strip test for differentiation of members of the family Enterobacteriaceae. Particularly Salmonella spp. J. Clin. Microbiol . 34 : 1811-1812.

Ivanoff BN, LevineMM, LambertPH (1994): Vaccination against typhoid fever :Present status. Bulleton of the world Health Organization 1994 ; 72(6) 957-71.

Ivanoff B(1995): Typhoid fever: global situation and WHO recommendations. Southeast Asian J Trop Med Public Health ;26(Suppl 2):1-6.

J

Jamal, W.Y, pal,T., Rotimi, V.O. and chugh, T.D.(1998):Serogroups and antimicrobial susceptibility of clinical isolates of Salmonella Species from a teaching hospital in Kuwait. J. Diarrhoeal Dis.Res.,16 : 180 -186.

Janda, j.m. and Abbott, S.L.(1998) : The Salmonellae. in: Janda, J.M. and Abbott, S.L.(eds.).The Enterobacteria,1st ed.Chap.6,PP :80-100. Lippincott-Raven, Philadelphia.

Joklik, W.K., willett, H.P., Amos, D.B and Wilfert, C.m. (1992); Enterobacteriaceae : salmonella and Shigella : Intestinal Pathogens. In: Joklik, W.K., willett, H.P., Amos, D.B. and Wilfert, C.M.(eds.) Zinsser Microbiology,20th ed., Chap.35, PP:556-565. Appleton and Lange, Norwalk.

Juan Alvarez,¹ Mertxe Sota,² Ana Belén Vivanco,¹ Ildefonso Perales,³ Ramón Cisterna,^{1,2} Aitor Rementeria,¹ and Javier Garaizar^{1;(2004)}
Development of a Multiplex PCR Technique for Detection and Epidemiological Typing of Salmonella in Human Clinical Samples J Clin Microbiol. 2004 April; 42(4): 1734–1738.

June, G.A., Sherrod, P.S., Hammzck, T.S., Amaguana, R.m. and Andrews, W.H.(1995):Relative effectiveness of selenite cystine broth, tetrathionate broth and Rappaport-Vassiliadis medium for the recovery of *Salmonella* from raw flesh and other highly contaminated foods; precollabortive study. *J. AOAC. Int.*,78:375-379.

K

Kariuki, S., Gilks, C., Corkill, J., Kimari, J.,benea, A.,Waiyaki, P. and Hart, C.A.(1996); Multidrug resistant non-typhi *Salmonellae* in Kenya.*J.Antimicrob.Chemother.*,38:425-434.

Kauffmann, F.(1951):*Salmonella* group .In: *Enterobacteriaceae* Munksgaard ,Copenhagen, Denmark. Cited by : Burnens ,A.P., Stanley ,J., Sechter, I.aand Nicolet, J.,(1996);Evolutionary origin of a monophasic *Salmonella* serovar 9,12:1.v:-,revealed by IS200 profiles and restriction fragment polymorphisms of the *fli B* gene. *J. Clin. Microbiol.*, 34 ; 1641-1645.

Kauffmann, F.(1954); *Enterobacteriaceae*. Munksgaard, Copenhagen, Denmark. Cited by Old, D.C.(1992): *Nomenclature of Salmonella*. *J. Med.Microbiol.*,37;361-363.

Kauffmann, F.(1966): The bacteriology of *Enterobacteriaceae*. Munksgaard, Copenhagen, Denmark. Cited by : Brenner, F.W., Villar, R.G., Angulo ,F.J., Tauxe, R. and Swaminathan B. (2000):*Salmonella* Nomeclature. *J. Clin. Microbiol.*, 38: 2465-2567.

Koneman, E.W., Allen, S.D., Janda, M., Schreckenbergr, P.C. and winn, W.C.(1997):The *Enterobacteriaceae*. In; Koneman, E.W., Allen, S.D., Janda, M., Schreckenberger, P.C. and Winn, W.C. (eds.) *Color Atlas and Textbook of Diagnostic Microbiology*, 5th ed., Chap. 4, PP:171-252.

Kurowski, P. B., J. L. Traub-Dargatz, P. S. Morley, and C. R. Gentry-Weeks.(2002): Detection of *Salmonella* spp. in faecal specimens by use of real-time polymerase chain reaction assay. *Am. J. Vet. Res.* 63:1265-1268.

L

Le Minor, L.(1984): Genus III .salmonella lignieres 1900.In;Kreig,N.R. and Holt, J.G.(eds.) Bergeys Manual of Systematic Bacteriology, Vol. 1, PP: 427-458.Williams and Wilkins, Baltimore.

Le Minor, L. and Popoff, M.Y.(1987): Request for an opinion ; Designation of Salmonella enterica spp. as the type and only species of the genus Salmonella. Int.J. Syst.Bacteriol.,37;465-468.

Le Minor, L. and Popoff, M.Y.(1988): Antigenic formulas of the Salmonella serovars, 5th revision , World Health Organization Collaborating Center for Reference and Research on salmonella. Pasteur Institute, Paris, France.

Le Minor, L. and Rohde, r.(1986) : Guidelines for preparation of Salmonella antisera, World Health Organization Collaborating Center for Reference and Research on Salmonella. Pasteur Institute, Paris , France.

Le Minor, L., Veron, M .and popoff, M.Y.(1982): Taxonomie des Salmonella .Ann.Microbiol.,133:223-243.Cited by :Old, D.C.(1992) : Nomenclature of Salmonella.J.Med.Microbiol.,37 ; 361-363.

Leading , A. (1978) :Typhoid and its serology . Brit . Med. J. 1: 389-397.

Lee, L.A., Puh, N.D., Maloney, K., Bean, N.H. and; Tauxe, R.V. (1994) : Increase antimicrobiol- resistant Salmonella infections in the United states , 1989-1990.J.Infect.Dis.,170:128-134.

Leegaard, T.M., Van-Gestel, M.H., Petit, P.L.C and Van de Klundert ,J.A.M. (1996): Antibiotic resistance mechanisms in Salmonella species causing bacteremia in Malawi and Kenya .APMIS,104:302-306.

Lepage, P., Bogaerts, J., Nsengumuremyi, F., Hitimana, D.G., Van-Goethem , C., vandepitte, J. and Butzler , J.P (1984): Sever multiresistant Salmonella typhimurium systemic infections in Central Africa. Clinical features and treatment in a pediatric department .J. Antimicrob. Chemother ., 14(Suppl B); 153-159.

Lesmana, M., Richie, E., Subekti, D., Simanjuatak, C. and Watz ,S, (1997): Comparison of direct-plating and enrichment methods forisolation of Vibrio cholerae from diarrhea patients. J. Clin. Microbiol., 35: 1856-1858.

Levine, M. M. (1979) : shigella and Salmonella diarrheal disease. Trop . Doct. Jan . 15 :4-9 .

Levine, M.M.; Grados, O.; Gilman , R.H.; Woodward, W.W.; Plaza, R.S. and Waldman, W. (1978): Diagnostic value of the Widal test in areas endemic for typhoid fever . Am. J. Trop Med. Hyg. 27 (4) : 759-800.

Lindberg, A.A. (1980); Bacterial Virulence factors with particular reference to salmonella bacteria. Scand .J. Infect.Dis.,24;86-92.

Lindberg, A.A. and Le Minor, L.(1984): Serology of Salmonella In : Bergan , T.(ed.)Methods in Microbiology ,Vol. 15 ,pp; 1-141. Academic press, London.

Liebana E, Gibbs M, Clouting C, Barker L, Clifton-FA, Pleydell E, Abdalhamid B, Hanson ND, Martin Poppe C, Davies RH.(2004): Characterization of beta-lactamases responsive resistance to extended-spectrum cephalosporin Escherichia coli and Salmonella entrica strain from food producing animals in the United Kingdom. Veterinary Laboratories Agency, Department of food Environmental Safety, Defra, Addlestone KT 151 AT.

Little , K.J. and Hartman , P.A. (1983) : Fluorogenic selective and differential medium for isolation of fecal streptococci. J. App. Environ. Microbiol. 45 (2) : 622-627.

Liu, P.Y.F., Gur, D., Hall, L.M.S. and Livermore, D.M.(1992):Survey of prevalence of β -lactamase amongst 1000-gram negative bacilli isolated consecutively at the Royal London hospital .J .Animicrob. Chemother., 30: 429-447.

Lu-Fong M, LudanAC, Martinez MM, Raymundo JG. DotEIA (1991): Aid to the diagnosis of typhoid fever. The Southeast Asian Journal of Tropical Medicine and public Health 1991 ;8:163.

M

Mabilat,C and Courvalin, P.(1990): Development of oligotyping for characterization and molecular epidemiology of TEM β -lactmases in amembersof the family Enterobacteriaceae. Antimicrob.Agens Chemother.,34:2210-2216.

Maddocks, J.L. and Greenan ,M.J.(1975): Technical method.A rapid method for identifying bacterial enzymes .J. Clin. Pathol .28:686-687.

Maddocks, S., T. Olma, and S.Chen.2002.Comparison of CHROM agar Salmonella medium and Xylose-Lysine-desoxycholate and salmonella-shigella agars for the isolation of salmonella strains from stool samples. J. Clin. Microbiol. 40: 2999-3003.

Maiorini, E., Lopez., E.L., Morrow, A.L., Ramirez, F., Procopio, A., Furmanski, S., Woloj, G.M., Miller, G. and Cleary, T.G.(1993): Multiply resistant nontyphoidal Salmonella gastroenteritis in children .pediatr. Infect .Dis.J., 12:139-145.

Makela, P.H.and Stocker , B.A.D.(1969):Genetics of polysaccharide biosynthesis .Annu.Rev .Genet, 3; 291-322.Cited by ;Old , D.C. and Threlfall, E.J.(1998): Salmonella .In : Balows,A. and Duerden ,B.I. (eds.) Topley and Wilson s Microbiology and Microbial Infections ,Vol.2,9th ed., PP; 969-997. Arnold, London.

Malorny, B., J. Hoorfar, C. Bunge, and R. Helmuth. (2003). Multicenter validation of the analytical accuracy of Salmonella PCR: towards an international standard. Appl. Environ. Microbiol. 69:290-296..

Manafi , M. and Sommer , R. (1992) : Comparison of three rapid screening methods for Salmonella spp: MUCAP test , Micro-screen latex and Rambach agar .Lett. App. Microbiol. 14: 163- 168.

Manafi, M. (1996): Fluorogenic and chromogenic enzyme substrates in culture media and identification tests. Int. J. Food Microbiol. 31:45-58

Manafi, M.(2000): New developments in chromogenic and fluorogenic culture media. Int. J. Food Microbiol. 60:205-218

. Manson – Bahr , P.E.C. and Bell, D.R. (eds) (1987) : Typhoid fever (enteric fever) . Manson's Tropical Diseases . 19th edn. Bailliere Tindall, London . Ch. 8 & 22, PP. 194-206 , 448-520.

Martinez-Beltran, J., Negri,C., Morosini, M., Canton, R., Loza, E., Baquero, F., Papaicolaou, G. and Medeiros, A.(1990):acquisition of anew plasmid-mediated β -lactamases in an intrahospital Salmonella arizonae outbreak.In program and abstract of the 30th Interscience

Conference on Antimicrobial agents and Chemotherapy. American Society for Microbiology .Washington, DC.

Matthews, J., M. Chung, and J. Matyas. (2002). Persistent DNA contamination in competitive RT-PCR using cRNA internal standards: identity, quantity, and control. *BioTechniques* 32:1412-1417.

Meideros, A.A., Gillece, E.S, Obrien, T.F.(1981):Distribution of plasmid type β -lactamases in ampicillin-resistance *Salmonella* from humans and animals in the United States .In; Levy, S.B., Close, R.D. and Loening, E.L.(eds.).*Molecular Biology, Pathogenicity and Ecology of Bacterial Plasmids*,PP:634-640. plenum press, NewYork.

Metzler , J. and Nachamkin , I. (1988) : Evaluation of a latex agglutination test for the detection of *Salmonella* and *Shigella* spp. by using broth enrichment. *J. Clin Microbiol* . 26 (12) : 2501-2504.

Mikhail , I.A.; Sanborn , W.R. and Sippel J.E. (1983) : Rapid , economical diagnosis of enteric fever by a blood clot culture coagglutination procedure. *J. Clin. Microbial*. 17 (3) : 564-565.

Miller, S.I. and pegues, D.A.(2000): *Salmonella* species including *Salmonella typhi*. In: Mandell, G.L., Bennett, J.E. and Dolin, R.(eds.) *Mandell, Douglas and Bennett s Principles and practice of Infectious Diseases* ,Vol. 2,5th ed., PP: 2344-2363. Churchill Livingstone, London.

Mobilat, C. and Courvalin, P.(1990) : Development of oligotyping for characterization and molecular epidemiology of TEM β -lactamases in members of the family Enterobacteriaceae .*Antimicrob. Agents Chemother.* ,34; 2210-2216.

Molbak, K., Baggesen,D.-L Aerstrup, F.M., Ebbesen, J.M. and Wegener, H.C.(1999):An outbreak of multi -drug resistant and quinolone resistant *Salmonella enterica* serotype Typhimurium DT104. *N. Engle .J. Med.*, 341;1420-1425.

Molbak, K., Gerner-Smith, P. and Wegener, H.C. (2002) : Increasing quinolone resistance in *Salmonella enterica* serotype Enteritidis. *Emerg. Infect. Dis.*,8:514-515.

use of CHROM agar Orientation medium in conjunction with simple biochemical tests .J. Clin. Microbiol .,38; 4586-4592.

Old , D.C. (1990) : Solmonella . In : Collee J. G.Duguid JP, Fraster AG, Marmion BP (eds). Mackie & McCartney's Practical Medical Microbiology ,14th edn. Churchill Livingstone, Edinburgh Ch. 29, PP. 456-481.

Old ,D.C(1992): Nomenclature of Salmonella. J. Med. Microbiol., 37;361-363.

Old, D.C(1996): Salmonella In: Collee, J.G., Fraser, A.G., Marmion, Microbiology, 14th ed., Chap.21, PP:385-404.Chrchill Livingstone, New York.

Old, D. Cand Adegbola, R.A.(1985): Antigenic relationships among type-3 finbriae of Enterobacteriaceae revealed by immunoelectronmicroscopy . J. Med. Microbiol.,20;113-123.

Old, D.C and Threlfall , E.J.(1998):Salmonella .In: Balows,a . and Duerden,B.L.(eds.).Topley and Wilson s Microbiology and .microbial Infections,Vol.2,9th ed.,PP;969-997. Arnold, London.

Old, D.C., Rankin, S.C. and Crichton, P.B (1999): Assessment of strain relatedness among Salmonella serotypes Salinatis , Duisburg and Sandiego by biotyping,ribotyping,IS200 fingerprinting and pulsed-field gel electrophoresis.J.Clin.Microbiol.,37:1687-1692.

P

Pan , T.-M., Wang , T.-K. and Lee , C.L. (1997) : Food – borne disease out breaks due to bacteria in Taiwan , 1986 to 1995 J. clin. Microbiol ., 35 : 1260 - 1262

Pang, T. and Puthucheary , S.D. (1983) : Significance and value of the Widal test in the diagnosis of typhoid fever in an endemic area. J. Clin. Pathol 36: 471-475.

Parker, M.T. (1990) : Enteric Infection : typhoid and paratyphoid fever In : Parker M.T. and Duerden , B.I. (eds). Topley & Wilson's Principles of Bacteriology , Virology and Immunity . 8th edn. Vol .3, Arnold , London . Ch. 22, pp 424-442.

Pavia, A.T., Shipman, L.D., Wells, J.G., pühr, N.D., Smith, J.D., Mckinley T.W. and Tauxe, R.V. (1990); , Epidemiologic evidence that antimicrobial exposure decreases resistance to infection by antimicrobial sensitive Salmonella. *J.Infect.Dis.*,161:255-260.

Payne, D.J. and Cramp, R.(1992);Determination of Id 50 values for 35 plasmid-mediated β -lactmases. Abstract of the first International Congress on the Management of Infection,10:65.

Pearson , R. and Guerrant , R. (1990) : Enteric fever and other causes of abdominal symptoms with fever . In : “ Principles and practice of infectious Diseases” 3rd edn. Churchill Livingstone, New York. Part 2. ch. 85. PP. 880-893.

Perry, J. D., M. Ford, J. Taylor, A. L. Jones, R. Freeman, and F. K. Gould. (1999): ABC medium, a new chromogenic agar for selective isolation of Salmonella spp. *J. Clin. Microbiol.* 37:766-768 .

Philippon, A., Fournier, G., Cornel, E., Paut, G., Le Minor, L. and Nevot, P.(1984): β -lactmases of ampicillin-resistant Salmonella. *Ann. Microbiol.*, 135:229-238.

Poole, K.(2000):Efflux-mediated resistance to fluroquinolone in gram-negative bacteria, *Antimicrob. Agents chemother.*,44;2233-2241.

Popoff, M. Y., and Le Minor, L.(1985):Expression of antigenic factor O:54 is associated with the presence of plasmid in salmonella *Ann Inst . Pasteur Microbiol.*,136:169-179.

Popoff, M. Y., and Le Minor, L.(1992): Antigenic formulas of the salmonella serovars, 6th Revision , world Health Organization Collaborating Center for references and Research on salmonella Pasteur Institute ,Paris, France.

Popoff, M. Y., and Le Minor, L.(1997): Antigenic formulas of the salmonella serovars 7th Revision ,world Health Organization Collaborating Center for references and Research on salmonella Pasteur Institute, Paris, France.

Popoff, M. Y., J. Bockemühl, and F. W. Brenner. (2000): Supplement 1998 (no. 42) to the Kauffmann-White scheme. *Res. Microbiol.* 151:63-65.

Punjabi NH. Cost;(1998) evaluation of typhoid fever in Indonesia
.Medical Journal of Indonesia 1998;7(S):90-3TR.

Q

Qrskoy, I., Cerskoy, F., Jann, K.(1977): Serology, chemistry and genetics of O and K antigens of *Escherichia coli*. Bacteriological Reviews 41:667-710.Cited by Ols, D.C.(1996):*Salmonella* In : Collee, J.G., Fraser, A.G., Marmion, B.P. and Simmons, A.(eds.) Mackie and McCartney practical Medical Microbiology, 14th Ed., Chap. 21, pp: 385-404. Churchill Livingstone, New York.

R

Rao, G.G., Ozerek, A.E. and Jeanes, A.(2001):Rational protocols for testing faeces in the investigation of sporadic hospital-acquired diarrhea. J. Hosp. Infect.,47:79-83.

Rambach, A. (1990). New plate medium for facilitated differentiation of *Salmonella* spp. from *Proteus* spp. and other enteric bacteria. Appl. Environ. Microbiol. 56:301-303

Rashtchian, A. and Curiale, M.S. (1989) : DNA probe assays for detection of *Campylobacter* and *Salmonella*. In Swaminathan, B. and Prakash, G. (eds).Nucleic acid and Monoclonal Antibody probes . Churchill Livingstone, New York. PP.221-239.

Reeves,M.W.,Evins,G.M.,Heiba,A.A.,Plikaytis,B.D.andFarmerIII,J.J.

(1989); Clonal nature of *salmonella typhi* and its genetic relatedness to other *Salmonella* as shown by multilocus enzyme electrophoresis and proposal of *salmonella bongori*.J.Clin.Microbiol.,27:313-320.

Robert, C., and P. Cuschieri 2003 Comparison of *salmonella* chromagar medium with DCLS Agar for Isolation of *Salmonella* species from stool specimens. J. Clin. Microb. 41(7): 3229-32.

Robertson, P.P. and Wahab, M.F.A. (1970) : Influence of chloramphenicol and ampicillin on antibody response in typhoid-paratyphoid fever . Ann , Intern . Med. 72: 219-224

- Rockhill, R.C.; Rumans , L.W.; Lesmana, M. and Dennis, D. T. (1980) :** Detection of *S. typhi* D, vi and d antigens by slide coagglutination in urine from patients with typhoid fever . *J. Clin. Microb .*11 (3) : 213-216.
- Rodrigue , D.C., Taxue, R.V. and Rowe , B. (1990) :** international increase in *Salmonella enteritidis* : A new pandemic *Epidemiol. Infect.*, 105:21-27.
- Rodriguez – Noriaga, E., Andrade- villaneurva , J. and Amaya-Tapla, G. (1989) :** Quinolones in the treatment of *Salmonella* carriers . *Rev. Infect, Dis.*, 11 (Supl. 1) : 1179-1187.
- Rohde, R. and Aleksic, S. (1975) :** Profuse fimbriae conferring O-inagglutinability to several strains of *S. typhimurim* and *S. enteritidis* isolated from pasta products :Cultural , Morphological and serological experiments . *Zentrable. Bakteriolog. Parasitenkd. Infektionskr. Hyg. Abt. I. Orig.*, 230 : 38-50 . Cited by : Old , D.C. and Threlfall , E.J. (1998) : *Salmonella*. In : Balows, A. and Duerden, b.I. (eds.). *Topley and Wilson's Microbiology and Microbial Infections* ,Vol.2, 9th ed., PP: 969-997 . Arnold, London .
- Rowe B, world LR, Threlfall EJ (1997):** Multidrug-resistance *Salmonella typhi*: a worldwide epidemic. *Clinical Infectious Diseases*1997 ; 24(Suppl 1): S106-9.
- Rubin , F.A. (1990) :** Nucleic acid probes for the identification of *Salmonella* .In : Macario , A.J.L (ed).*Gene Probes for Bacteria* . Churchill Livingstone , New York. PP. 323-352.
- Rubin , F.A. and Kopecko, D.J. (1989) :** Nucleic acid probes for detection of clinically significant bacteria In : Swaminathan , B. and Paraksh , G. (eds).. *Nucleic Acid and Monoclonal Antibody Probes*. Churchill Livingstone, New York. PP. 185 – 219.
- Rubin, F.A.; Kopecko, D.J. ; Sack , R.B. and Sudarmone , P. (1988) :** Evaluation of a DNA Probe for identifying *Salmonella typhi* in Peruvian and Indonesian bacterial Isolates . *J. Infect. Dis.* 157: 1053.
- Rubin, F.A; Kopecko, D.J., Noon , K.F. and Baron L.S. (1985):** Development of a DNA probe to detect *S. typhi* . *J. Clin. Microbiol .* 22: 600-605.
- Rubin, F.A.; McWhirter, P.D. and Burr D. (1990) :** Rapid diagnosis of typhoid fever through identification of *S. typhi* within 18hours of

specimen acquisition by culture of the mononuclear cell-platelet fraction of blood . J. Clin . Microbiol . 28 (4) 825- 827.

Ruiz, J.; Varela, M.C.; Sempere, M.A. and Oliva, J.(1991) : presumptive identification of *Salmonella enterica* using two rapid tests. Eur. J. Clin . Microbiol. Infect. Dis. 10: 649-354

Ruiz , J., Núñez, M.L., Climent ,C., Sympere , M.A. and Gómez, J. (1995) : Utilización del agar Rambach para la detección *Salmonella* en heces a partir de un caldo de enriquecimiento. Comparación de metodología. Enferm. Infecc. Microbiol. Clin.,13: 213-217 Cited by Ruiz, J., Núñez, M.L., Díaz, J., Lorente, I. M. Pérez, J. and Gómez, J. (1996a): comparison of five plating media for isolation of *Salmonella* species from human stools . J. Clin .Microbiol., 34: 686-688.

Ruiz, J., Núñez, M.L., Díaz, J., Lorente, I. M. Pérez, J. and Gómez, J. (1996a): comparison of five plating media for isolation of *Salmonella* species from human stools . J. Clin .Microbiol., 34: 686-688.

Ruiz , J.R., Núñez, M.L., Díaz, J. M Sempere , M.A., Gomez, J. and Usera , M. A. (1996b) : Note : comparison of media for the isolation of lactose – positive *Salmonella* . J. Appl Bacteriol ., 81: 571- 574. .

Ruiz, J., M. L. Nunez, I. Lorente, J. Perez, E. Simarro, J. Gomez. 1996. Performance of six culture media for isolation of *Salmonella* species from stool samples. Eur. J. Clin. Microbiol. Infect. Dis. 15:922-26.

Ruiz M, Rodriguez JC, Sirvent E, Escribano I, Cebrian L, Royo G. (2003) Usefulness of different techniques in the study of the epidemiology of salmonellosis. APMIS Sep;111(9):848-56.

S

Sadallah , F.; Brighthouse , G. and Drager- Dayal , R.(1990): Production of specific monoclonal antibodies to *Salmonella typhi* flagellin and possible application to immunodiagnosis of typhoid fever . J. Infect . Dis.161 : 59-64.

Saha SK, Baqui AH, Hanif M, Darmstadt GL, Ruhulamin, Nagatake T, Santosham M, Black R(2001): Typhoid fever in Bangladesh: implications for vaccination policy. The pediatric Infectious Disease Journal;20:521-4.

- Sallustio , S., Rubin , R.H. and Golderg, M.B. (1998) :** Salmonella . In: Gorabch , S.L., Bartlett, J.G. and Blacklow, N.R. (eds.) .In: Infectious Diseases , 2nd ed., Chap. 205 , PP: 1796-1803. Saunders Company, Philadelphia.
- Sanborn , W.R.; Lesmana, and Edwards . E.A. (1980) :**Enrichment culture coagglutination test for rapid , low- cost diagnosis of salmonellosis .J. Clin Microb.12(2) : 151 –155.
- dos Santos, S.M. and Kupek, E.(2000):** Serial outbreaks of food-borne diseasein Blumenau, Brazil caused by Salmonella enteritidis. Braz. J. Infect.Dis., 4:275-278 .
- Sarasombath, S.; Lertmemongkolchai, G, and Banchuin, N.(1988):** Characterization of monoclonal antibodies to protein antigen of samonella typhi . J. Clin. Microbiol. 26 (3): 508- 512.
- Selander, R.K. and Musser , J.M. (1990) :** Population genetics of bacterial pathogenesis . In : Iglewski ,B.11. and Clark, V.L. (eds.). Molecular Basis of Bacterial Pathogenesis , vol. 11, PP: 11-36. Academic press, London.
- Selander, R.K., Beltran , P., Smith , N.H., Helmuth , R., Rubin. F.A., Kopecko, D.J., Ferris , K., Tall ,B.D., Cravioto, A. and Musser , J. M. (1990) :** Evolutionary genetic relationships of clones of Salmonella serovars that cause human typhoid and other enteric fevers. Infect . Immunol., 58: 2262-2275.
- Sevenungsson , B.; Jobeck, H. and Lindberg, A.A. (1979) :** Diagnosis of Salmonella infections specificity of indirect immunofluorescence for rapid identification of S. enteritidis and usefulness of ELISA. J. Infect . Dis 140: 927-936.
- Shannon , K., Williamas, H., King ,A. and Phillips , 1 .(1990) :** Hyperproduction of TEM-1 β - lactamase in clinical isolates of Salmonellae. FEMS microbiology Letters , 55: 319- 323.
- Shannon , K., and French, G.(1998):**Multiple antibiotic-resistant salmonella. Lancet ,352 : 490 – 491.
- Shaokat , S., Ouelette , M., Sirot , D., Joly , B. and Cluzel , R. (1987) :** Spread of SHV-1 β - lactamase in Salmonellae isolated from fecal samples in Africa. Antimicrob. Agents Chemother., 31: 943-945.

Shaprio, R.L., Kumar, L., Phillips-Howard, P., Wells, J.g., Adcock, P., Brooks, J., Ackers, M.L. Ochieng, J.B., Mintz, E., Wahlquist, S., Waiyaki, P and Slutsker, L. (2001): Antimicrobial-resistant bacterial diarrhea in rural western Kenya. *J. Infect. Dis.*, 183: 1701 – 1704.

Shaw, W.V. (1984): Bacterial resistance to chloramphenicol. *Br. Med. Bull.*, 40: 36-41.

Shehabi, A.A. (1981): The value of a single Widal test in the diagnosis of acute typhoid fever. *Trop. Geogr. Med.* 33(2): 113-116

Shehabi, A.A. (1995): Extra-intestinal infections with multiply drug resistant *Salmonella typhimurim* in hospitalized patients in Jordan. *Eur. J. Clin. Microbiol. Infect. Dis.*, 14: 448 – 451.

Silverman, M. and Simon, M. (1980): Phase variation: Genetic analysis of switching mutants. *Cell*, 19: 845-854.

Simarro, E., Navarro, F., Ruiz, J., Miro, E., Gómez, J. and Mirelis, B. (2000): *Salmonella enterica* serovar Virchow with CTX-M-like β -lactamase in Spain. *J. Clin. Microbiol.*, 38: 4676-4678.

Sinha A, Sazawal S, Kumer R, Sood S, Reddaiah VP, Singh B, Rao M, Naficy (1999): A Clemens J, Bhan MK. Typhoid fever in children aged less than 5 years. *Lancet* 1999; 354: 734 – 737.

Sippel, J.e.; Mamy, H.K, H.K.; Weiss, e.; Joseph, S.W. and Beasley W.J. (1978): Outer membrane protein antigens in an ELISA for *Salmonella enteric* fever and meningococcal *Salmonella*. Enteric fever and meningococcal *Salmonella*. *J. Clin. Microbiol.* 7 (4): 3782-378.

Sippel, J.E.; Bukhari, N.; Awan, M.B., Krieg, R. and Duncan J. F. (1989): Indirect IgG and IgM ELISA and IgM capture ELISA for detection of antibodies to LPS in adult typhoid fever patients in Pakistan. *J. Clin. Microbiol.* 27(6): 1298-1302.

Sivadasan, K.; Kurien, B. and John, T.J. (1984): Rapid diagnosis of typhoid fever by antigen detection. *Lancet* I: 134-135.

Slack, R.C.B. (1997): Diagnostic procedures in Greenwood, D., Slack, R.C.B. and Peutherer, J.F: Medical microbiology, fifteenth edition. Chapter: 64, P: 616-23 Churchill Livingstone.

Song, J.H.; Cho, H.; Park, M.Y. and Pai, C.H. (1993): Detection of *Salmonella typhi* in the blood of patients with typhoid fever by polymerase chain reaction. *J. Clin. Microbiol.* 31: 1439-1443.

Sonja J. Olsen,¹ Jim Pruckler,¹ William Bibb,¹ Nguyen Thi My Thanh,² Tran My Trinh,² Nguyen Thi Minh,³ Sumathi Sivapalasingam,¹ Amita Gupta,¹ Phan Thu Phuong,⁴ Nguyen Tran Chinh,⁵ Nguyen Vinh Chau,⁵ Phung Dac Cam,⁴ and Eric D. Mintz¹ (2004):Evaluation of Rapid Diagnostic Tests for Typhoid Fever *J Clin Microbiol.* ; 42(5): 1885–1889.

Spencer, H.L., Donnelly, M.T. and Teahon, K.(2001): Culturing of stool samples from hospital inpatients. *Lancet*, 358:151-152.

Stam F, Romkens TE, Hekker TA, Smulders YM. (2003): Turtle-associated human salmonellosis. *Clin Infect Dis.* 37:167-9.

Stites, D.P.; Stobo, J.D.; Fudenberg, H.H. and Wells, J.V. (eds) (1982): Basic and Clinical Immunology . 4th edn. Lange Medical Publications, California. Ch.22. PP.326.

Sue, L.H., Chiu, C.H., Kuo, A.I., Chia, L.H., Sun, C.F., Leu, H.S. and WU, T.L.(2001): Secular trends in incidence and antimicrobial resistance among clinical isolates of salmonella at a university hospital in Taiwan, 1983-1999. *Epidemiol. Infect.*, 127:207-213.

Swaminathan, B.(2000):salmonella Nomenclature *J. Clin. Microbiol.*, 38;2465-2567.

T

Taylor , W.L. and Schelhart , D. (1965) : Isolation of Shigella . I. Xylose lysine agar : ew media for isolation of enteric pathogens . *Am . J. Clin Pathol.*, 44 : 471 – 475 : cited by : chang , C-T., You , C.- Y., Shen , H.- C., Li, A.,- M., Chen , C.-Y Chou , J.- L. and Huang, S.-P. (1999) : Recovery of Salmonella by using selenite brilliant green sulfa enrichment broth . *J. Clin. Microbiol.*, 37: 4120-4123.

Thomson , C.J. and Amyes , S.G.B. (1992) : TRC-1 Emergence of a clavulanic acid- resistant TEM β - lactamase in clinical strain . *FEMS microbial . Lett.*, 91 : 113 – 118.

Threlfall , E.J. (2002) : Antimicrobial drug resistance in Salmonella : Problems and perspectives in Food – and water – borne infections . *FEMS Microbiol . Rev.*, 26 : 141-148.

Legakis , N.J. (1994) molecular epidemiology of ampicillin – resistant clinical isolates of salmonella enteritidis . J. Clin Microbiol ., 32: 13322 – 1325.

Villasante , P.A; Merino, F.J.; Perez , T.; Guevara, C.L. and Velasco, A.C. (1987) : Rapid automated method for screening of enteric pathogens from stool specimens. J. Clin. Microbiol. 25(3) : 584-585.

Voller , A.; Bidwell , D.E. and Bortlet , A.(1979) : The enzyme immunosorbent assay (ELISA) . J. Infect . Dis . 140 : 615 – 622.

W

Walker , R.A., Saunders , N., Lawson , A.J., Lindsay , E.A., Dassama, m., Ward , L.R. Woodward , j.J., Davies , R.H.mLiebana, E.and Threlfall, E.J. (2001) : Use of a light Cyler gyrA mutation assay for rapid identification of mutations conferring decreased susceptibility of ciprofloxacin in multiresistant Salmonella enterica serotype typhimurim DT104 isolates. J. Clin Microbiol., 39: 1443-1448.

Wall , P.G. (1995) : Transmission of multiresistant strains of Salmonella typhimurim from cattle to man. Vet Rec., 136 : 591-592.

WainJ, DiepTS, HoVA, WalshAM, HoaTTN, ParryCM, White NJ. (1998): Quantitation of bacteria in blood of typhoid fever patients and relationship between counts and clinical features, transmissibility, and antibiotic resistance. Journal of Clinical Microbiology 1998 ; 36:1683-7.

Wasfy,M.O.,Oyofa,B.A.,David,J.C.,Ismail,T.F.,ElGendy,A.M.,Mohran,Z.S.,Sultan,Y.and Peruski, L.F.(2000): Isolation and antibiotic susceptibility of salmonella, Shigella and Campylobacter from acute enteric infections in Egypt .J. Health Poul. Nutr., 18: 33-38.

World Health Organization, 2003. The World Health Report 2003:The diagnosis, treatment, and prevention of typhoid fever. Geneva: World Health Organization.

W.H.O (1987) : Program for control of diarrheal diseases .In: Manual for Laboratory Investigations of Acute Enteric Infections.P.9.

Wuiff, C., Masen, M., Baggesen, D.-L.and Aarestrup, F.M.(2000): Quinolone resistance among salmonella enterica from cattle, broilers and swine in Denmark. Microb. Drug Resist.,6:11-7.